

 Open access • Journal Article • DOI:10.1038/NREND0.2016.193

An updated view of hypothalamic–vascular–pituitary unit function and plasticity

— [Source link](#) 

Paul Le Tissier, Pauline Campos, Pauline Campos, Pauline Campos ...+9 more authors

Institutions: University of Edinburgh, French Institute of Health and Medical Research, University of Montpellier, Centre national de la recherche scientifique ...+2 more institutions

Published on: 01 May 2017 - Nature Reviews Endocrinology (Nature Publishing Group)

Topics: Neuroendocrinology, Pituitary gland, Anterior pituitary and Hypothalamus

Related papers:

- [Cellular in vivo imaging reveals coordinated regulation of pituitary microcirculation and GH cell network function.](#)
- [Anterior pituitary cell networks.](#)
- [Existence of long-lasting experience-dependent plasticity in endocrine cell networks](#)
- [Ion Channels and Signaling in the Pituitary Gland](#)
- [Related pituitary cell lineages develop into interdigitated 3D cell networks](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/an-updated-view-of-hypothalamic-vascular-pituitary-unit-2d24m6dun7>

An updated view of hypothalamic–vascular–pituitary unit function and plasticity

Le Tissier, Paul; Campos, Pauline; Lafont, Chrystel; Romano, Nicola; Hodson, David;
Mollard, Patrice

DOI:

[10.1038/nrendo.2016.193](https://doi.org/10.1038/nrendo.2016.193)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Le Tissier, P, Campos, P, Lafont, C, Romano, N, Hodson, D & Mollard, P 2017, 'An updated view of hypothalamic–vascular–pituitary unit function and plasticity', *Nature Reviews Endocrinology*, vol. 13, pp. 257-267. <https://doi.org/10.1038/nrendo.2016.193>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the accepted manuscript before final editorial and formatting changes were carried out. The final version of record is available at: <http://dx.doi.org/10.1038/nrendo.2016.193>

Checked for eligibility: 05/09/2016.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A 21st century perspective on hypothalamic–vascular–pituitary unit function

[Au: our journal style is to write 21st century in full as ‘twenty first’, which would take the title over the length limits for this type of article. We also use ‘perspective’ for a specific opinion-based article. To reduce the length of the title and avoid the use of perspective, I suggest the following title modification:

“An updated view of hypothalamic–vascular–pituitary unit function”. OK?]

[Au: you use a mixture of hypothalamo and hypothalamic in the main text. I suggested consistently using on hypothalamic, changes throughout main text also OK?]

Paul Le Tissier¹, Pauline Campos^{2,3,4}, Chrystel Lafont^{2,3,4}, Nicola Romanò¹, David J. Hodson^{5,6} and Patrice Mollard^{2,3,4}

¹Centre for Integrative Physiology, University of Edinburgh, Edinburgh, EH8 9XD, UK. ²CNRS, UMR-5203, Institut de Génomique Fonctionnelle, F-34000 Montpellier, France; ³INSERM, U661, F-34000 Montpellier, France; ⁴Université de Montpellier, UMR-5203, F-34000 Montpellier, France; ⁵Institute of Metabolism and Systems Research and Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham, Edgbaston, B15 2TT, UK. ⁶Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, B15 2TH, UK. [Au: please include the street name for all addresses]

Correspondence to

Paul.LeTissier@ed.ac.uk

patrice.mollard@igf.cnrs.fr

Abstract [Au: edits OK?]

The discovery of novel functional adaptations of the hypothalamus and anterior pituitary gland for physiological regulation has transformed our understanding their interaction. The activity of a small number of hypothalamic neurons can control complex hormonal signalling, which is disconnected from a simple stimulus and subsequent hormone secretion relationship but is dependent on physiological status. The interrelationship of hypothalamic neurons and pituitary cells with the vasculature has an important role in determining the pattern of neurohormone exposure. Cell in the pituitary gland form networks with distinct organizational motifs that are related to the duration and pattern of output. These networks are modified in different physiological states and can persist after cessation of demand, which results in enhanced function [Au: edit OK?]. Consequently, the hypothalamus and pituitary can no longer be considered as having a simple stratified relationship: they form a tripartite system with the vasculature, which must function in concert for appropriate hypothalamic regulation of physiological processes, in particular reproduction [Au: edit OK? I think it's important

to mention reproduction in the abstract here]. An improved understanding of the mechanisms underlying these regulatory features has implications for current and future therapies that correct defects in the hypothalamic–pituitary axis. In addition, recapitulating proper network organization will be an important challenge for regenerative stem cell treatment.

[Au:For your information, H1 and H3 refer to the level of heading and will be removed before proofs are made. H1 subheadings can have max 38 characters inc spaces. H3 subheads can be of any length. Subheads have been edited to fit these limits, where indicated]

[Au: I think the Review would benefit from a figures introducing the basics of the concepts you discuss. I think this will be important to increase the impact of the paper, and for our less informed readers who are endocrinologists, but not particularly familiar with the neuroendocrine aspects. I have included the details of the figures you might wish to consider adding at the end of this document. I have also indicated in the main text where I think new figures might be useful additions. When you submit your revisions, please submit any new figures and I will work with our art editor to generate the first versions. We can revise these before making the final proof. If you wish to discuss these in greater detail before submitting, please do not hesitate to contact me]

[H1] Introduction [Au: I have moved the second section (originally called ‘Hypothalamic–pituitary axes’ of your original draft to here to serve as an introduction (the opening section is always titled ‘Introduction’ in our style guide). I think this text is more appropriate here and sets up the rest of the article nicely. Please also include the appropriate references in this section]

To maximize reproductive success, via the appropriate timing of ovulation, lactation or body growth, the output of several hypothalamic–pituitary axes are dramatically altered **[Au: edit OK?]**. These adaptive changes occur over differing time scales, with varying frequencies and levels of predictability **[Au: is there a general review you can cite here?]**. For example, the increase in growth hormone (GH) output at puberty is largely predictable **[Au: meaning the levels are predictable or simply that it occurs?]**. On a relatively short time scale **[Au: meaning measured in days?]**, the surge in luteinizing hormone (LH) secretion required for oestrus is an acute change that occurs regularly once every reproductive cycle and, in humans, continues for years in the absence of pregnancy. On a longer time scale **[Au: meaning months/years?]**, the increase in prolactin required for lactation is maintained for a variable time (which depends on when offspring are weaned) and recurs at each pregnancy, but is unpredictable before gestation **[Au: edit OK to avoid repetition of ‘pregnancy’?]**. These large changes in output require modification of both hypothalamic and pituitary function, but are reversed after pregnancy, which likely reflects the fact that these processes need to be

repeated in subsequent pregnancies [Au: edit OK is this what you?]. A mechanistic understanding of these alterations in hypothalamic–pituitary function is fundamental to interpret and treat defects that lead to endocrine diseases [Au: please be more specific here as to the diseases you mean in this context to set the scene for your review]. In this Review, we will focus on three pituitary axes that have roles in driving changes in physiology; the gonadotropin, prolactin and growth hormone axes [Au: addition OK?]. The level of our understanding varies for each of these axes and the features that might serve as general principles will be highlighted in the text [Au: edit OK?].

[H1] Beyond stimulus-secretion coupling [Au: I think you could include a simple figure in this section on stimulus-secretion as an addition to BOX 1. And based on the summary figure you have already submitted. I think these concepts will work well in graphical form for our less informed readers]

The speed of communication between the brain and peripheral tissues is highlighted by muscle contraction, which requires the transfer of electrical signals from axons via the neuromuscular junction [Au: edit OK? Is this what you mean here?]. This sequence of events, known as excitation-contraction coupling¹, takes <1 s in mammals and is highly plastic [Au: please elaborate on what you mean by plastic in this instance]. Similarly, in the hypothalamus just a few thousand neurons can also send signals to the periphery, in this case toward the median eminence (ME) via a specialized neurohaemal junction [Au: edits OK?]. Here, nerve terminal depolarization, either originating from the perikarya² or the terminal itself³, allows the sufficiently rapid entry of calcium ions to trigger exocytosis of neurohormones towards the first loop of the portal fenestrated capillaries [Au: please reference this statement here]. This rapid (< 1 s) sequence of events was termed ‘stimulus-secretion coupling’ due to the clear similarities with excitation-contraction coupling^{4,5}. Soon after release, neurohormones pervade the second loop of fenestrated capillaries within the downstream pituitary gland, before binding to cognate receptors on endocrine cells to induce pituitary hormone exocytosis through a second ‘stimulus-secretion coupling’ event⁶⁻⁹. [Au: I think here might be a nice place to include a figure detailing these events for our less informed readers. A simple outline/schematic would be sufficient and that will set the scene for understanding the different axes later in the text]

In the second half of the twentieth century (and building on Harris’ work on the hypothalamus–pituitary axis [Au: please reference this statement with a reference to

Harris' work for our less informed readers) the analogy between excitation-contraction and stimulus-secretion coupling was developed further **[Au: edit OK?]**¹⁰. However, important and fundamental differences exist between the two processes. Specifically, in the hypothalamus, endogenous **[Au: circadian]** rhythms exist and the time scale for pituitary hormone release is much longer (measured in minutes to several hours) **[Au: edit OK? Please reference this statement]**. Nevertheless, the analogy with neural control of locomotor activity led to a generally accepted model of hypothalamic regulation of pituitary function. Specifically, the excitation of specific hypothalamic neuron populations, determined by higher brain centres and peripheral feedback, is relayed as an unmodified series of signals to drive balanced pituitary hormone output **[Au: please reference this statement]**. The release of neurohormones and subsequent transportation and the effects on target cells were previously considered to be passive events in the regulation of pituitary hormone secretion, with only variation in the number of endocrine cells seeming to affect response levels^{11,12}. Similarly, the alterations in gene expression and cell proliferation, which support maintenance of hormone output, were simply considered a correlated response to hypothalamic regulation of secretion **[Au: please reference this statement]**.

In the early twentyfirst century, a series of paradigm shifts in our understanding of the hypothalamic–pituitary system was established as a consequence of newly developed tools and techniques **[Au: such as?]** for use in genetically modified mice **[Au: please reference this statement with some examples]**. The use of these methods have shown that both the pituitary gland and portal system can no longer be considered as static structures simply responding to neurohormonal regulation (BOX 1) **[Au: edits OK to shorten sentence? I think the text box should be cited later in the text as more of a summary section]**. In addition, hypothalamic neuron function has been found to be more dynamic than initially thought, which might, rather than changes in excitation, contribute to modifications in its regulation of the pituitary under different physiological states **[Au: edit OK? Please reference this statement]**.

[H1] Gonadotroph axis

The reproductive system is critically dependent upon pulsatile secretion of gonadotrophin-releasing hormone (GnRH) and LH; however, the understanding of pulse generation has been hampered by the complexity of the regulatory mechanisms, many of which are lost in *in vitro* preparations **[Au: please include the references here also as examples]**. Investigators

working in the late 1980s using pituitary portal bleeding and microdialysis documented the pulsatile nature of GnRH release into the portal vasculature of the sheep, monkey and rat²⁵⁻³¹, and showed a strong correlation between GnRH and LH pulses^{28,32,33}. However, the scattered distribution and relative paucity of GnRH cell bodies limited the investigation of the cellular events that lead to pulsatile secretion of LH *in vivo*. In the past few years, the development of optogenetic techniques in rats and mice and an ultra-sensitive ELISA capable of measuring LH levels in small whole blood samples³⁴ has enabled investigators to dissect the GnRH neuron excitation parameters that generate LH pulses³⁵. In these studies, the stimulation of just 60 GnRH neurons can trigger short-lived increases in LH secretion that resemble endogenous pulses **[Au: Edits OK? please reference this statement]**. Given the critical importance of GnRH neurons to the survival of all mammalian species, a degree of functional redundancy within this cell population is expected. Indeed, activating ~5% of the GnRH hypophysiotropic neurons seems to be sufficient to generate an LH pulse **[Au: please reference this statement]**. This finding is consistent with studies in which just 10% of the GnRH neuron population is sufficient to maintain pulsatile LH secretion^{36,37}. When the timing and frequency of stimulation is varied a brief (2 min) optogenetic stimulation at high frequency (10 Hz) evokes an LH pulse, whereas shorter periods and lower frequencies cannot elicit LH output that resembles endogenous pulses³⁵. **[Au: edit OK?]** This finding was also the case for a bursting pattern of stimulation **[Au: meaning of LH secretion? How is a bursting pattern defined in this context?]**, which had been assumed to be effective for pulse generation and the focus of many previous studies³⁸ **[Au: meaning that this mechanism was the only one thought to be necessary of LH output?]**. Whether such a stimulatory signal exists *in situ* **[Au: meaning within the organism?]** and where its origin might be is unknown, although a ‘GnRH pulse driver’ might be located in the mediobasal–hypothalamus, specifically at the level of neurons co-expressing kisspeptin, neurokinin B, as well as dynorphin A (so-called ‘KNDy’ neurons)^{39,40}. **[Au: Edits OK? perhaps you can expand on that specific evidence here for clarify?]**

Pulsatile secretion of GnRH requires synchronization **[Au: of the secretion and/or pulsatility itself? What do you mean in this context?]** within the GnRH neuron population. While the cell bodies of GnRH neurons are scattered throughout the basal forebrain, their projections have dendrodendritic bundling, that is they share synapses **[Au: edit OK, is this what you mean?]**⁴¹, and become highly concentrated around the ME **[Au: please reference this statement]**. Fascinatingly, these projections simultaneously receive and integrate synaptic

inputs — they possess both axonal and dendritic characteristics, leading to their description as ‘dendrons’, before finally acquiring an axonal morphology within the ME and ramifying into numerous terminals that appose blood vessels⁴². **[Au: I think this could be represented in the Figure also. It will then also give context to the vasculature aspect of the article]** Dendrons might be an ideal location for putative afferent axons to modulate the excitability of multiple GnRH neuron dendrites, and for multiple GnRH neurons to align their firing pattern, which thereby provides a potential mechanism for their synchronized activity directly in the mediobasal hypothalamus **[Au: is this your opinion?]**. An additional source of pulse synchronization is in the ME, where hypophysiotropic GnRH neurons terminate within the external zone close to endothelial cells of the portal vasculature⁴³. Endothelial cells in the ME might modulate GnRH release through nitric oxide secretion (which has been reviewed elsewhere⁴⁴) **[Au: edit OK? I would include some of the original reference here also to reduce the number of other reviews cited]**. At the ME, nitric oxide is spontaneously released from an endothelial source and follows a pulsatile and cyclic pattern of secretion⁴⁵, and inhibition of nitric oxide synthesis **[Au: by local do you mean specifically in the ME or a wider area?]** can disrupt reproductive cyclicity⁴⁶. Conversely, in the GnRH neuron perikarya, basal nitric oxide synthase activity might provide the tonic inhibition of the GnRH neural system required to maintain nadir levels of LH **[Au: edit OK?]**⁴⁷. **[Au: this mechanism to be represented in the figure also?]**

Once released into the ME, the transport of GnRH to the pituitary, and the pattern of gonadotroph exposure to the neurohormone, have been largely assumed to represent a simple linear process **[Au: please reference this statement]**. However, the use of fluorescent tracking using 4 kDa dextran, which mimics the size of most hypothalamic neurohormones, has shown that the diffusion processes, both at the level of the ME and the pituitary capillaries, are complex and non-linear⁷. **[Au: edit OK?]** Consequently, the portal vessel network might function as a ‘physical integrator’, enabling neurohormones to be transferred from the ME to the gonadotroph within a few seconds **[Au: can you cite a reference here?]**. Once in the blood stream, the moderately rapid clearance rate **[Au: can you define this rate here?]** of LH underlies **[Au: meaning it generates it?]** the specific asymmetric pulse shape of this hormone, which is characterized by a fast increase immediately followed by a slower decrease³⁴. Importantly, a faithful delivery of the pulsatile pattern of GnRH secretion to the pituitary is crucial for gonadotroph function⁴⁸ **[Au: please include some of the original key references for this finding]**. For example, high GnRH pulse frequencies (>1 pulse per h)

activate LH production, whereas low frequencies (<1 pulse per 2–3 h) preferentially induce follicle-stimulating hormone (FSH) synthesis and release⁴⁹. Overall, the intricate relationships between pulsatile GnRH release, secretory competency of the pituitary gonadotrophs and regulatory mechanisms within the vasculature, generate the rhythmic fluctuations in LH secretion.

[H3] GnRH and LH surge generation

The GnRH/LH surge mechanism is sex specific and normally occurs only in women⁵⁰⁻⁵² **[Au: Please include the original references and only one Review here; it might be interesting to add a note about when it might occur in men]**. During the oestrous cycle, increasing concentrations of plasma oestrogen alter feedback to the GnRH neuronal afferent networks and gonadotrophs from negative to positive to induce the gonadotrophin surge **[Au: please reference this statement]**. That the oestrogen-responsive kisspeptin neurons in the rostral periventricular area of the third ventricle have a critical role in enabling ovulation in rodents by activating GnRH neurons is now well accepted⁵³. Importantly, the relative contribution of the hypothalamic and pituitary levels to the oestrogen-induced gonadotropin surge seems to be species-dependent, with the latter the predominant mechanism in human and non-human primates⁵⁴. In the female sheep, the GnRH surge is composed of high-frequency pulsatile events superimposed on a constantly elevated level of GnRH release, although whether the surge is driven by a fundamentally altered pattern of GnRH secretion⁵⁵, or by a simple increase in the frequency of pulsatile secretion is unclear **[Au: edit OK?]**⁵⁶. This huge increase in GnRH secretion continues for a period of 24 h, considerably longer than the duration of the LH surge it induces, before returning to a strictly episodic pattern of release^{31,55,57}. The firing pattern of GnRH neurons needed to generate the GnRH/LH surge is unknown. However, that the prolonged firing of an increased number of GnRH neurons is required for the secretion of surge levels, compared with that required for a pulse, is a reasonable assumption **[Au: edit OK? Have I retained your meaning here?]**. Indeed, mice with 10% of the normal GnRH neuronal content failed to ovulate **[Au: meaning 90% were ablated? How so?]**, but cyclicity was restored when approximately 30% of the GnRH population was present^{36,37}. **[Au: please include a little more detail of how these studies were conducted]**

In addition to the putative change in GnRH population electrical activity, anatomical changes are found within the external zone of the ME where GnRH nerve terminals are ensheathed by tanycytes^{58,59}. The cellular conformation changes with fluctuating oestrogen profiles throughout the oestrous cycle. For example, in rats, semaphorin-7a-dependent structural

remodelling of tanycytes occurs during the preovulatory surge, resulting in release of the engulfed axons and direct access of GnRH nerve terminals to the portal vasculature [Au: please reference this statement also]. By contrast, fenestrated endothelial cells of the hypothalamic–hypophyseal portal vessels release semaphorin-3A, which is thought to induce GnRH neuron axonal growth and sprouting within the ME as a function of the oestrous cycle⁶⁰. [Au: also data from rats?] These mechanisms are likely to enable the generation of high concentrations of GnRH, which evoke the GnRH/LH surge, to be released into the pituitary portal circulation⁶¹⁻⁶³.

Within the pituitary, the distinct network organization of gonadotrophs¹⁴ and their large scale reorganization during puberty¹⁵ suggests that homotypic cell organisation [Au: meaning the cells reorganize according to their type? Please expand on what you mean here. Can this be included in the figure?] might have a functional role [Au: in puberty itself?]. However, this reorganisation [Au: edit OK? Is this what you mean here?] has not been studied in detail in the other pituitary axes to date. Although the dynamic gonadotroph responses at the time of the proestrous surge have not yet been described *in vivo*, static snapshots [Au: meaning ‘individual readings’?] in rats and sheep suggest that changes occur in gonadotrophin subunit expression, granule distribution and GnRH receptor abundance^{64,65}. [Au: edit OK?] However, mRNA sequencing of the anterior pituitary glands from women [Au: of what age?] reveal that genes regulating [Au: gonadotroph?] secretion, blood pressure and cell adhesion were all enriched during proestrus⁶⁶. Likewise, immortalized cell lines and cells in pituitary slices both extend cellular processes and increase their cellular movement at [Au: stages that replicate puberty?] puberty¹⁵ and following GnRH stimulation⁶⁷. These findings suggest that changes in the relationship of the gonadotroph network with the vasculature might modify the secretory response of gonadotrophs^{68,69}. [Au: edit OK?]

[H3] Clinical relevance

The mechanisms that underlie both pulsatile secretion and surge generation of LH have important implications for the treatment of infertility in women [Au: edit OK?]. For example, polycystic ovarian syndrome (PCOS), the most common anovulatory cause of infertility⁷⁰ and affects >100 million women worldwide, is associated with a dysregulation of the normal pattern of LH secretion [Au: please reference the latter part of this statement also]. Whether the origin of this multi-factorial disorder is at the level of the hypothalamic–pituitary axis is unknown⁷¹, but PCOS is characterized by increases in GnRH pulse frequency [Au: of

how much? Can this be quantified? and sensitivity of the pituitary gland to the neurohormone^{72,73}. Consequently, potential interventions that modify the dynamics of GnRH output, its transport to the ME or its actions in the pituitary might have implications for the treatment of PCOS **[Au: is this opinion?]**. This is also the case for congenital hypogonadotropic hypogonadism **[Au: we avoid using two letter abbreviations where possible. HH removed OK?]**, which results from a pituitary or a hypothalamic defect with or without anosmia **[Au: please reference this statement for clarity also]**. Several novel gene mutations that are associated with this disorder have been identified, including those encoding neuropeptides **[Au: such as?]**, transcription factors **[Au: such as?]** and G-protein coupled receptors**[Au: such as?]** **[Au: please give examples here]**⁷⁴. To induce female fertility, hypogonadotropic hypogonadism of pituitary origin can be reversed by subcutaneous injections of FSH followed by human chorionic gonadotrophin or LH to trigger ovulation **[Au: please reference this statement]**. Conversely, hypogonadotropic hypogonadism of hypothalamic origin can be treated using GnRH pumps to restore pituitary hormone secretion **[Au: please reference this statement]**. Pulsatile GnRH has the advantage of decreasing the risk of multiple pregnancy and ovarian hyperstimulation syndrome⁷⁵. In both situations, the pulsatility of GnRH or the rhythmic secretion of FSH and/or **[Au: OK?]** LH is required to obtain sufficient follicular maturation and proper ovulation⁷⁶. Advances in understanding of GnRH secretion and its interactions with LH are essential for designing novel, and indeed modifying existing, therapies for hypogonadotropic hypogonadism. For example, some patients **[Au: roughly how many? Is there a % you can include here?]** with this disease, who are yet to undergo treatment, have transient phases of normal fertility⁷⁷ **[Au: edit OK?]**. The underlying mechanisms and relevant therapeutic interventions to maintain this phenomenon might be elucidated by further investigation of pulsatility and rhythmicity **[Au: edit OK?]**.

[H1] The prolactin axis

The prolactin axis is unique among the pituitary hormonal systems, as in men and women who have not previously had sexual intercourse, it can be considered a system primed for activation by tonically inhibited by hypothalamic dopamine⁷⁸ **[Au: edit OK? Is this what you mean?]**. **[Au: 'In this situation?']** Low concentrations **[Au: can this be quantified?]** of circulating prolactin are maintained by short-loop feedback, with prolactin receptor-mediated stimulation of dopamine neuron firing rate leading to an increase in catecholamine

production⁷⁹ and output². The timescale of the secretory response for prolactin (~10–20 min) requires the coordination of multiple neurons **[Au: please reference this statement]**. Specifically, tuberoinfundibular dopamine neurons undergo long-term coordinated changes in firing rate, which correlate with episodic secretion **[Au: of prolactin?]** in mice². **[Au: please expand on this finding. How do you mean long-term and what are the changes in the firing rate? I think you need to include a little more explanation here]** Gap junctions and local dendritic dopamine release have been proposed to mediate this activity^{80,81}, and integration of single cell firing rates seems to be involved in the generation of longer DA **[Au: DA meaning dopamine?]** release output events (N. Romano and P. Mollard, unpublished data).

Variations in prolactin output occur in virgin female rats as a surge at proestrus, which coincides with that of LH⁸². Prolactin also increases following vaginal stimulation of rodents **[Au: meaning it is true in mice and rats? Or has this only been shown in rats?]** as twice daily surges⁸³. These surges are coordinated by signals from the suprachiasmatic nucleus⁸³, most likely through the actions of vasoactive-intestinal peptide⁸⁴. At the level of the pituitary, lactotrophs form a network of honeycomb motifs that allow the congregation of cells along the fine pituitary capillary network⁸⁵. **[Au: can this be represented in a figure summarising the Prolactin axis?]** This organization supports low levels of cell-cell coordination¹⁶, with a small proportion **[Au: how many? Can this be defined?]** of cells acting as coordinating nodes by functionally connecting distant ensembles **[Au: please reference this latter part of the statement]**. In addition to synchronizing Ca²⁺ activity, cellular organization also mediates the coordination of gene transcription^{86,87}, with gap junction signalling enabling local correlation of bursts of transcriptional activity that are otherwise randomly timed **[Au: please reference this latter part of the statement]**. This mechanism resembles quorum-sensing where apparently random systems display complex activity as long as the components (in this case the cells) can interact, and might contribute to hormone gene expression and cell proliferation^{87,88,89}. **[Au: edit OK?]** How gap junctions might orchestrate this mechanism remains unknown; however, regenerative Ca²⁺ waves and Ca²⁺-regulation (*e.g.* through NFAT) could be one possibility⁹⁰. **[Au: I think you need to expand on this highlighted (in grey) statement — e.g. why specifically could it be a possibility — or remove it altogether if this is too speculative]**

[H3] Increased prolactin output during lactation

The long-term requirement for large increases of circulating prolactin in lactation is associated with a decrease in dopamine output, which begins in late pregnancy⁹¹ and is coincident with a surge of pituitary prolactin secretion [Au: please reference this latter statement – 91 also? Edit OK to simplify sentence and avoid repetition?]. The dopamine tone needs to be strongly decreased throughout lactation to enable the necessary increase in circulating prolactin, and is mediated by a decrease in phosphorylation of tyrosine hydroxylase, the rate-limiting enzyme for dopamine synthesis⁹¹. This mechanism is not the result of a reduced feedback of prolactin on dopamine neurons, which remain electrically responsive at the level of the cell body, but rather, neuronal firing becomes uncoupled from dopamine secretion². [Au: edit OK? have I retained your meaning here?s] Remarkably, the reduction in dopamine tone is accompanied by the production of opioids^{92,93}, which might enable these neurons to stimulate prolactin secretion [Au: is this latter statement an opinion?].

In concert with changes in the hypothalamic inhibition of prolactin secretion, substantial alterations occur in the pituitary to support the 10–50-fold increase in prolactin secretion that is required for milk production in mammals⁹³ [Au: edit OK?]. In humans and rats, this [Au: milk production?] is generally accompanied by proliferations of lactotrophs, with some evidence of hypertrophy [Au: of the lactotrophs?], although these results [Au: meaning the hypertrophy results?] are based on 2D histological studies [Au: edit OK?]⁹⁴. By contrast, in lineage tracing and FACS studies in mice, lactotrophs become hypertrophied during lactation and increase their volume threefold rather than simply increasing in number [Au: edit OK?]⁹⁵. Other investigators have confirmed these findings, and also showed that the lactotroph network *in situ* becomes highly-connected during lactation, which is associated with the strength of the suckling stimulus¹⁶. [Au: edit OK?] This increase in structural connectivity leads to a substantial increase [Au: of how much?] in the proportion of the subpopulation of lactotrophs that function as coordinating nodes and orchestrate increased output of prolactin [Au: please reference this statement].

[H3] *Memory of prolactin demand after weaning* [Au: edit OK?]

At weaning [Au: edit OK? I think the term weaning is well-enough understood to omit an explanation], a rapid decrease in prolactin secretion occurs as a result of a return of dopamine inhibition⁹⁶. In rodents^{97,98} and humans⁹⁹ basal prolactin secretion is reduced below that of virgin animals, which might reflect an enhanced pituitary response to dopamine inhibition¹⁰⁰. Strikingly, and despite this reduction in basal prolactin secretion, lactotrophs

remain enlarged and well-connected **[Au: with each other? Or generally within the network? Please clarify here]** at both the structural and functional levels¹⁶, with an increased proportion of cells acting as nodes **[Au: what is the proportion? What does it change from and to? What is the specific function of the nodes at this stage? Possibly to be included in a new figure on the prolactin axis]**, which persists for many months after lactation has ceased **[Au: edit OK, is this what you mean? please reference this statement]**. Such hard-wiring or ‘memory’ of previous stimuli, which was previously thought to only exist for neurons and immune cells, leads to a lactotroph network behaviour **[Au: what do you mean specifically by ‘lactotroph network behaviours’ in this instance?]** during subsequent lactations **[Au: meaning after subsequent births?]**, which drives even higher concentrations of prolactin¹⁶. This mechanism is independent of reproductive experience *per se*, since it can be prevented by reducing the suckling demand **[Au: please reference this statement]**.

[H3] Clinical relevance

The dysregulation of the prolactin axis, owing to either pituitary adenomas¹⁰¹ or as an adverse effect of treatment with antipsychotic drugs¹⁰², leads to impaired fertility. Hyperprolactinaemia is the second most common cause of infertility in women after PCOS **[Au: please reference this statement; how many individuals does hyperprolactinaemia affect?]**. A clear understanding of the interactions that lead to altered dopamine output and the response of the pituitary might help to identify novel treatment strategies for this disease. For example, further understanding the regulation of prolactin release by dopamine might enable the reduction in the doses of commonly-used dopamine receptor agonists, as their safety in the treatment of prolactinomas has been questioned¹⁰³. **[Au: why so? Please include a little extra detail]** In rodent studies, prolactin seems to affect multiple neuroendocrine axes **[Au: such as? Please include the axes here for clarity]**⁷⁷, and these warrant further study to determine the potential effects of its over-secretion in humans. For example, hyperprolactinaemia might lead to changes in GnRH neuron activity via interactions with the GPR54/kisspeptin pathway in mice¹⁰⁴ and GnRH pulsatility has been reinstated in mice with physiological hyperprolactinaemia by administration of kisspeptin¹⁰⁵. **[Au: edit OK?]** By contrast, in studies using sheep with a lower prolactin dose **[Au: lower dose than what? And how much lower?]**, no effects on hypothalamic kisspeptin expression have been seen **[Au: in response to this treatment/administration please clarify here? edit OK?]**¹⁰⁶. An improved understanding of these pathways could aid the development of treatments for women with

hyperprolactinaemia that is resistant to dopamine agonists **[Au: please provide a reference for this dopamine agonist-resistant hyperprolactinaemia].**

[H1] The GH axis [Au: suggest a new figure for this section including the organisation and plasticity of the GH axis]

In humans and animals in which it can be measured **[Au: such as?]**, pulsatile GH output is present from birth^{107,108,109}. However, the output is markedly increased at puberty when sexually-dimorphic body growth occurs¹¹⁰. **[Au: edit OK?]** Since the discoveries of GH releasing hormone (GHRH) and somatostatin that control GH secretion from pituitary somatotrophs¹¹¹⁻¹¹³, a remarkable advancement our understanding of GH pulse generation during critical physiological windows has taken place **[Au: edit OK?]**.

[H3] Pulsatile GHRH output

Using genetically-modified mouse models with GHRH neurons marked with green fluorescent protein¹¹⁴, several investigators have defined the mechanisms that underlie pulse generation using *ex vivo* slices of brain **[Au: edit OK?]**. Before puberty, GHRH neurons are excitable neuroendocrine neurons with complex synaptic inputs¹¹⁵. These early stages of hypothalamic development ensure appropriate regulation of the somatotroph axis, as in the Ames dwarf mice in which loss of GH leads to a compensatory increase in GHRH cell number¹¹⁶, **[Au: edit OK?]** and programming effects of steroids on GHRH cell number and gene expression¹¹⁷ **[Au: please expand on this latter statement. What specifically do you mean by ‘programming’ and how does the cell number and gene expression change?]**. Modification of these synaptic inputs and electrical properties over the first 6 postnatal weeks correlates with and likely drives, at least in part, increased pituitary GH output and sexual dimorphism¹¹⁸. In all these studies, the intrinsic hourly rhythms of GHRH neuronal activity predicted by simulation studies of *in vivo* GH pulsatility have not been indentified^{115,119}. However, somatostatin can generate GHRH neuron pulsatile output by delaying oscillations of action potential firing via a recurring inhibition of inhibitory GABAergic interneurons (that is, inhibition of inhibition)¹¹⁹. **[Au: edit OK? Have I retained your meaning here?]** Consequently, somatostatin can both acutely inhibit the excitability of GHRH neurons and also promote their patterned output together with more sustained GHRH neuron stimulation in response to other stimuli in the brain (for example, acetylcholine¹¹⁸) and peripheral tissues (such as, ghrelin)¹¹³. **[Au: edit OK?]** **[Au: can these processes be represented in the a figure on the GH axis? E.g. to highlight the plasticity of these neurons during puberty?]**

[H3] Modification of pituitary somatotroph output

No full description of the *in vivo* dynamics of GHRH and somatostatin neurons and their regulation of pituitary somatotrophs exist [Au: edit OK?]. This event can be viewed as a three-step process: delivery of neurohormone to target cells; cellular secretory responses to regulation; and entry of pituitary hormone into the peripheral circulation [Au: please reference this statement. Or is this based on opinion?]. *In vivo* imaging of the mouse portal system and somatotroph network have provided insights into the first step in this process and the role of the vasculature in shaping the pattern of exposure of the pituitary to hypothalamic neuropeptides⁷. Delivery of neuropeptides such as GHRH to the somatotroph network, which extends throughout the pituitary gland, follows specific vascular/capillary routes and results in specific temporal patterned regulation rather than a homogenous ‘immersion’ [Au: by ‘immersion’ meaning the whole pituitary exposure to the secretagogue?] of the pituitary parenchyma in secretagogue [Au: please reference this statement]. In addition, the initial stimulation by GHRH evokes a coordinated enhancement of oxygen to the stimulated somatotroph network via increased capillary blood flow, which provides fuel for energy-depleting secretory responses [Au: please reference this statement]. Indeed, the GH [Au: neuronal?] network is spatially organized versus the vasculature [Au: Do you mean that the GH network is organized in the same spatial manner as the vasculature? Do you mean all of the vasculature in this instance? please reference this statement], which suggests an important role of local oxygen regulation on GH release and that this may also be the case for other pituitary axes [Au: is there any evidence to support this highlighted statement? If not I suggest you remove it]. The second step in pituitary regulation has been characterized using *ex vivo* data from acute pituitary slices where the somatotroph network organization is preserved, in which the homotypic network-mediated coordination of stimulation, triggering long-lasting GH secretion²⁰. These studies have also shown that network organization is likely to have a major role in the increased GH output at puberty. In particular, the GH network undergoes large changes in the volume to surface ratio [Au: of what specifically?] that coincides with the onset of puberty, before gradually returning to normal pre-pubertal levels by [Au: postnatal?] day 100 in mice [Au: please reference this statement]. Such changes correlate with sexually dimorphic patterns of output¹³. [Au: please expand on this statement, what do you mean precisely here?] Sex steroid manipulation also leads to rapid and dramatic increases in cell motility that remodels the network organization¹²⁰ [Au: please can you expand on the details of this finding here]. [Sentences rearranged to improve

flow OK? Have I retained you meaning here? These findings highlight the importance of somatotroph network organization and its plasticity in the generation of pituitary somatotroph output. The vasculature also has an important role in somatotroph output, where *in vivo* imaging shows that capture of secreted GH is a controlled event where the perivascular space acts as a gate-keeper for hormone entry into the capillary lumen¹²⁰. The relationship between neuronal network organization and the vasculature in the pituitary is, therefore, central to the delivery of incoming hypothalamic signals, and the build-up of GH pulses within capillaries.

[H3] Clinical relevance

GH deficiency resulting from congenital defects or acquired following traumatic brain injury, pituitary tumours or cranial irradiation^{121,122}, is commonly treated with a daily subcutaneous dose of recombinant GH in childhood to increase growth rate **[Au: please reference this latter statement]**. However, considerable uncertainty exists regarding the optimal dosage or regime, and current treatments **[Au: specifically GH injections?]** do not fully recapitulate the physiological pattern of GH secretion **[Au: please reference this statement]**. One potential therapy is repopulation of the pituitary with stem cells, which has been investigated in the mouse **[Au: addition OK? was this approach successful?]**¹²³. However, such approaches would require the recapitulation of the normal cellular organization to achieve normal pulsatility and homeostatic regulation **[Au: edit OK?]**.

Patients with acromegaly, which results from a GH secreting pituitary adenoma, frequently have glycaemic disorders: a lack of GH pulsatility modifies lipolysis, whereas overall GH hypersecretion can induce insulin resistance¹²⁴. Consequently, an improved understanding of the mechanisms that determine the pattern of GH output might help to define new therapeutic options for dyslipidaemia or diabetes mellitus. In addition, GH pulsatility also has an important role in a subgroup of patients who have clinical acromegaly with increased insulin-like growth factor 1 **[Au: abbreviation used once removed OK?]**, but unaltered mean 24-h GH concentration compared with healthy controls^{125,126}. Altered GH pulsatility might explain the clinical presentation of this sub-group of patients, and our new understanding of the mechanisms underlying patterning of pituitary output might explain the abnormal GH axis function in these individuals and warrants further investigation **[Au: edit OK?]**.

[H1] Conclusions

The examples in this Review provide new insights into regulation of three hypothalamic–pituitary axes and demonstrate that these mechanisms are not a simple relay of stimulus–

secretion coupled events [Au: edit OK?]. The disconnection or modulation of hypothalamic excitation and neurohormone release, and an active role of the vasculature and pituitary in the network-mediated modification of responses, demonstrates that this view of the hypothalamic–pituitary system is over-simplistic (FIG. 1). [Au: we avoid first citing figures in the conclusions, but I think the addition of new figures as suggested above we negate the need for this Figure here and these points can be introduced earlier. You can then refer back to the earlier figures here in the conclusions] [Au: sentence removed here to avoid repetition] Given that the hypothalamic arcuate nucleus contains no more than a few thousand parvocellular neurons, the rapid development of techniques for interrogating neuronal function should enable the characterization of this structure’s regulation and output. Such studies will be invaluable for the deeper understanding of mammalian physiology, as this region controls a much larger panel of known body functions than any other brain region. In the vasculature and pituitary gland, many of the mediators of functional modulation remain unknown. Although loss of Prop1, which is essential for pituitary hormonal cell development, leads to loss of organ vascularization¹²⁷, studies of the processes leading to normal vascularization and its modification have not been reported to date. Similarly, hypothalamic and steroid factors have been shown to regulate cell network organization^{19,20,120} but the underlying mechanisms and mediating factors are yet to be identified. The alteration of the gonadotroph network in mice with a developmental block of corticotroph terminal differentiation¹⁴, suggests that the mechanisms will be complex and involve interaction between multiple pituitary cell types. [Au: I suggest moving this information (highlighted in grey) into the preceding sections where appropriate. We try to avoid including new information like this in the conclusions, which should be for summing up the key points. These findings might have more impact in the earlier sections]

The regulation of pulsatile pituitary secretion must now be considered as an integration of hypothalamic, vasculature and pituitary regulation, which has further implications for the understanding of disease [Au: edit OK?]. For example, the identification of kisspeptin has provided an exciting new target for the treatment of infertility¹²⁸. The uncoupling of neuronal excitation and hormone output also has deeper implications such as in the case of ageing [Au: addition OK? Is this what you mean here?]. For example, the reduction in GH output with age might be due to a failure of neurohormone secretion from GHRH neurons that are excited under normal conditions [Au: edit OK?]¹²⁹. The human pituitary gland can be accessed by transphenoidal surgery, which makes this structure ideal target organ for regenerative therapy

[Au: please reference this statement]. The pituitary neuronal networks and their relationship with the vasculature must be considered for such therapy and the microenvironment clearly has an important role in the regulation of the pituitary gland, which might also affect the development of tumours [Au: edit OK?] ¹³⁰. Finally, as pituitary networks are sensitive to peripheral regulation and their modification can persist for extended periods, they are a potential target for endocrine disrupting chemicals (EDCs). Indeed, the identification of EDC [Au: bisphenol A?]-mediated changes in expression of *ICAM5* [Au: which specifically?] in the pituitary¹³¹ has led us to speculate that some EDC effects might be mediated by changes in pituitary organization¹³². These possibilities require further investigation for the understanding of both the aetiology and treatment of diseases associated with pituitary hormones [Au: edits OK?].

[Au: Please re-reference to reflect editing changes (your citation manager should do this automatically on your machine) and put into our format– there should be an option to display bibliography in the style of ‘Nature Reviews’ in your reference manager program. Specifically, please remove the doi designations from the end of the references, which are not needed, unless an article has yet to be assigned a full page number and volume reference]

Example reference styles

1. Author, A. B. & Author, B. C. Title of the article. *Nat. Cell Biol.* 6, 123–131 (2001).
2. Author, A. B. Title of the article. *Nat. Struct. Mol. Biol.* 7, 101–109 (2003).
3. Author, A. B., Author, Z. X. & Author, B. C. Title of the article. *EMBO J.* 25, 3454–3461 (2006).
4. Author, A. B. in *Title of Book* (ed. Surname, I. N.) 75–98 (Publisher, 2000).
5. Author, A. B. Title of the article. *Website title* complete URL (year).

If a reference has six or more authors, only the first author should be listed followed by *et al.*

- 1 Sandow, A. Excitation-contraction coupling in skeletal muscle. *Pharmacol Rev* 17, 265-320 (1965).
- 2 Romano, N. *et al.* Plasticity of hypothalamic dopamine neurons during lactation results in dissociation of electrical activity and release. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 33, 4424-4433, doi:33/10/4424 [pii] 10.1523/JNEUROSCI.4415-12.2013 (2013).

- 3 Smith, J. T. *et al.* Kisspeptin is essential for the full preovulatory LH surge and stimulates GnRH release from the isolated ovine median eminence. *Endocrinology* **152**, 1001-1012, doi:10.1210/en.2010-1225 (2011).
- 4 Douglas, W. W. Stimulus-secretion coupling: the concept and clues from chromaffin and other cells. *Br J Pharmacol* **34**, 451-474 (1968).
- 5 Drouva, S. V., Epelbaum, J., Laplante, E. & Kordon, C. Calmodulin involvement on the Ca⁺⁺-dependent release of LHRH and SRIF in vitro. *Neuroendocrinology* **38**, 189-192 (1984).
- 6 Tse, A., Tse, F. W., Almers, W. & Hille, B. Rhythmic exocytosis stimulated by GnRH-induced calcium oscillations in rat gonadotropes. *Science* **260**, 82-84 (1993).
- 7 Lafont, C. *et al.* Cellular in vivo imaging reveals coordinated regulation of pituitary microcirculation and GH cell network function. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 4465-4470, doi:10.1073/pnas.0902599107 (2010).
- 8 Thomas, P., Wong, J. G., Lee, A. K. & Almers, W. A low affinity Ca²⁺ receptor controls the final steps in peptide secretion from pituitary melanotrophs. *Neuron* **11**, 93-104 (1993).
- 9 Tse, A. & Lee, A. K. Voltage-gated Ca²⁺ channels and intracellular Ca²⁺ release regulate exocytosis in identified rat corticotrophs. *J Physiol* **528 Pt 1**, 79-90 (2000).
- 10 Stojilkovic, S. S., Tabak, J. & Bertram, R. Ion channels and signaling in the pituitary gland. *Endocrine reviews* **31**, 845-915, doi:10.1210/er.2010-0005 (2010).
- 11 Raisman, G. An urge to explain the incomprehensible: Geoffrey Harris and the discovery of the neural control of the pituitary gland. *Annual review of neuroscience* **20**, 533-566, doi:10.1146/annurev.neuro.20.1.533 (1997).
- 12 Moenter, S. M. Leap of Faith: Does Serum Luteinizing Hormone Always Accurately Reflect Central Reproductive Neuroendocrine Activity? *Neuroendocrinology* **102**, 256-266, doi:10.1159/000438790 (2015).
- 13 Bonnefont, X. *et al.* Revealing the large-scale network organization of growth hormone-secreting cells. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 16880-16885, doi:10.1073/pnas.0508202102 (2005).
- 14 Budry, L. *et al.* Related pituitary cell lineages develop into interdigitated 3D cell networks. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 12515-12520, doi:10.1073/pnas.1105929108 (2011).
- 15 Alim, Z. *et al.* Gonadotrope plasticity at cellular and population levels. *Endocrinology* **153**, 4729-4739, doi:10.1210/en.2012-1360 (2012).
- 16 Hodson, D. J. *et al.* Existence of long-lasting experience-dependent plasticity in endocrine cell networks. *Nat Commun* **3**, 605, doi:10.1038/ncomms1612 (2012).
- 17 Le Tissier, P. R. *et al.* Anterior pituitary cell networks. *Front Neuroendocrinol* **33**, 252-266, doi:10.1016/j.yfrne.2012.08.002 (2012).
- 18 Mollard, P., Hodson, D. J., Lafont, C., Rizzoti, K. & Drouin, J. A tridimensional view of pituitary development and function. *Trends in endocrinology and metabolism: TEM* **23**, 261-269, doi:10.1016/j.tem.2012.02.004 (2012).
- 19 Waite, E. *et al.* Different degrees of somatotroph ablation compromise pituitary growth hormone cell network structure and other pituitary endocrine cell types. *Endocrinology* **151**, 234-243, doi:10.1210/en.2009-0539 (2010).
- 20 Sanchez-Cardenas, C. *et al.* Pituitary growth hormone network responses are sexually dimorphic and regulated by gonadal steroids in adulthood. *Proceedings*

- of the National Academy of Sciences of the United States of America **107**, 21878-21883, doi:10.1073/pnas.1010849107 (2010).
- 21 Featherstone, K. *et al.* Spatially coordinated dynamic gene transcription in living pituitary tissue. *Elife* **5**, e08494, doi:10.7554/eLife.08494 (2016).
- 22 Deneff, C. Paracrinicity: the story of 30 years of cellular pituitary crosstalk. *Journal of neuroendocrinology* **20**, 1-70, doi:10.1111/j.1365-2826.2007.01616.x (2008).
- 23 Schaeffer, M., Hodson, D. J., Lafont, C. & Mollard, P. Functional importance of blood flow dynamics and partial oxygen pressure in the anterior pituitary. *Eur J Neurosci* **32**, 2087-2095, doi:10.1111/j.1460-9568.2010.07525.x (2010).
- 24 Schaeffer, M., Hodson, D. J., Lafont, C. & Mollard, P. Endocrine cells and blood vessels work in tandem to generate hormone pulses. *J Mol Endocrinol* **47**, R59-66, doi:10.1530/JME-11-0035 (2011).
- 25 Dierschke, D. J., Bhattacharya, A. N., Atkinson, L. E. & Knobil, E. Circoral oscillations of plasma LH levels in the ovariectomized rhesus monkey. *Endocrinology* **87**, 850-853, doi:10.1210/endo-87-5-850 (1970).
- 26 Carmel, P. W., Araki, S. & Ferin, M. Pituitary stalk portal blood collection in rhesus monkeys: evidence for pulsatile release of gonadotropin-releasing hormone (GnRH). *Endocrinology* **99**, 243-248, doi:10.1210/endo-99-1-243 (1976).
- 27 Knobil, E. The neuroendocrine control of the menstrual cycle. *Recent Prog Horm Res* **36**, 53-88 (1980).
- 28 Clarke, I. J. & Cummins, J. T. The temporal relationship between gonadotropin releasing hormone (GnRH) and luteinizing hormone (LH) secretion in ovariectomized ewes. *Endocrinology* **111**, 1737-1739, doi:10.1210/endo-111-5-1737 (1982).
- 29 Levine, J. E., Pau, K. Y., Ramirez, V. D. & Jackson, G. L. Simultaneous measurement of luteinizing hormone-releasing hormone and luteinizing hormone release in unanesthetized, ovariectomized sheep. *Endocrinology* **111**, 1449-1455, doi:10.1210/endo-111-5-1449 (1982).
- 30 Caraty, A., Locatelli, A. & Martin, G. B. Biphasic response in the secretion of gonadotrophin-releasing hormone in ovariectomized ewes injected with oestradiol. *The Journal of endocrinology* **123**, 375-382 (1989).
- 31 Moenter, S. M., Brand, R. C. & Karsch, F. J. Dynamics of gonadotropin-releasing hormone (GnRH) secretion during the GnRH surge: insights into the mechanism of GnRH surge induction. *Endocrinology* **130**, 2978-2984, doi:10.1210/endo.130.5.1572305 (1992).
- 32 Levine, J. E. New concepts of the neuroendocrine regulation of gonadotropin surges in rats. *Biol Reprod* **56**, 293-302 (1997).
- 33 Karsch, F. J., Bowen, J. M., Caraty, A., Evans, N. P. & Moenter, S. M. Gonadotropin-releasing hormone requirements for ovulation. *Biol Reprod* **56**, 303-309 (1997).
- 34 Steyn, F. J. *et al.* Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice. *Endocrinology* **154**, 4939-4945, doi:10.1210/en.2013-1502 (2013).
- 35 Campos, P. & Herbison, A. E. Optogenetic activation of GnRH neurons reveals minimal requirements for pulsatile luteinizing hormone secretion. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 18387-18392, doi:10.1073/pnas.1415226112 (2014).
- 36 Kokoris, G. J., Lam, N. Y., Ferin, M., Silverman, A. J. & Gibson, M. J. Transplanted gonadotropin-releasing hormone neurons promote pulsatile luteinizing hormone

- secretion in congenitally hypogonadal (hpg) male mice. *Neuroendocrinology* **48**, 45-52 (1988).
- 37 Herbison, A. E. Estrogen positive feedback to gonadotropin-releasing hormone (GnRH) neurons in the rodent: the case for the rostral periventricular area of the third ventricle (RP3V). *Brain research reviews* **57**, 277-287, doi:10.1016/j.brainresrev.2007.05.006 (2008).
- 38 Jasoni, C. L., Romano, N., Constantin, S., Lee, K. & Herbison, A. E. Calcium dynamics in gonadotropin-releasing hormone neurons. *Front Neuroendocrinol* **31**, 259-269, doi:10.1016/j.yfrne.2010.05.005 (2010).
- 39 Lehman, M. N., Coolen, L. M. & Goodman, R. L. Minireview: kisspeptin/neurokinin B/dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology* **151**, 3479-3489, doi:10.1210/en.2010-0022 (2010).
- 40 Navarro, V. M. *et al.* Role of neurokinin B in the control of female puberty and its modulation by metabolic status. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 2388-2397, doi:10.1523/JNEUROSCI.4288-11.2012 (2012).
- 41 Campbell, R. E., Gaidamaka, G., Han, S. K. & Herbison, A. E. Dendro-dendritic bundling and shared synapses between gonadotropin-releasing hormone neurons. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 10835-10840, doi:10.1073/pnas.0903463106 (2009).
- 42 Herde, M. K., Iremonger, K. J., Constantin, S. & Herbison, A. E. GnRH neurons elaborate a long-range projection with shared axonal and dendritic functions. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 12689-12697, doi:10.1523/JNEUROSCI.0579-13.2013 (2013).
- 43 Prevot, V. *et al.* Evidence that members of the TGFbeta superfamily play a role in regulation of the GnRH neuroendocrine axis: expression of a type I serine-threonine kinase receptor for TGRbeta and activin in GnRH neurones and hypothalamic areas of the female rat. *Journal of neuroendocrinology* **12**, 665-670 (2000).
- 44 Bellefontaine, N. *et al.* Nitric oxide as key mediator of neuron-to-neuron and endothelia-to-glia communication involved in the neuroendocrine control of reproduction. *Neuroendocrinology* **93**, 74-89, doi:10.1159/000324147 (2011).
- 45 Knauf, C. *et al.* Evidence for a spontaneous nitric oxide release from the rat median eminence: influence on gonadotropin-releasing hormone release. *Endocrinology* **142**, 2343-2350, doi:10.1210/endo.142.6.8073 (2001).
- 46 De Seranno, S. *et al.* Vascular endothelial cells promote acute plasticity in ependymogial cells of the neuroendocrine brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **24**, 10353-10363, doi:10.1523/JNEUROSCI.3228-04.2004 (2004).
- 47 Hanchate, N. K. *et al.* Kisspeptin-GPR54 signaling in mouse NO-synthesizing neurons participates in the hypothalamic control of ovulation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 932-945, doi:10.1523/JNEUROSCI.4765-11.2012 (2012).
- 48 McArdle, C. A. & Roberson, M. S. in *Knobil and Neill's Physiology of Reproduction (Fourth Edition)* 335-397 (Academic Press, 2015).
- 49 Tsutsumi, R. & Webster, N. J. GnRH pulsatility, the pituitary response and reproductive dysfunction. *Endocrine journal* **56**, 729-737 (2009).

- 50 Cooke, B., Hegstrom, C. D., Villeneuve, L. S. & Breedlove, S. M. Sexual
differentiation of the vertebrate brain: principles and mechanisms. *Front*
Neuroendocrinol **19**, 323-362, doi:10.1006/frne.1998.0171 (1998).
- 51 Simerly, R. B. Organization and regulation of sexually dimorphic neuroendocrine
pathways. *Behavioural brain research* **92**, 195-203 (1998).
- 52 Simerly, R. B. Wired for reproduction: organization and development of sexually
dimorphic circuits in the mammalian forebrain. *Annual review of neuroscience* **25**,
507-536, doi:10.1146/annurev.neuro.25.112701.142745 (2002).
- 53 Herbison, A. E. Control of puberty onset and fertility by gonadotropin-releasing
hormone neurons. *Nat Rev Endocrinol*, doi:10.1038/nrendo.2016.70 (2016).
- 54 Plant, T. M. A comparison of the neuroendocrine mechanisms underlying the
initiation of the preovulatory LH surge in the human, Old World monkey and
rodent. *Front Neuroendocrinol* **33**, 160-168, doi:10.1016/j.yfrne.2012.02.002
(2012).
- 55 Moenter, S. M., Caraty, A., Locatelli, A. & Karsch, F. J. Pattern of gonadotropin-
releasing hormone (GnRH) secretion leading up to ovulation in the ewe:
existence of a preovulatory GnRH surge. *Endocrinology* **129**, 1175-1182 (1991).
- 56 Clarke, I. J. Variable patterns of gonadotropin-releasing hormone secretion during
the estrogen-induced luteinizing hormone surge in ovariectomized ewes.
Endocrinology **133**, 1624-1632, doi:10.1210/endo.133.4.8404603 (1993).
- 57 Caraty, A. *et al.* Nature and bioactivity of gonadotropin-releasing hormone
(GnRH) secreted during the GnRH surge. *Endocrinology* **136**, 3452-3460 (1995).
- 58 Kozlowski, G. P. & Coates, P. W. Ependymoneuronal specializations between
LHRH fibers and cells of the cerebroventricular system. *Cell and tissue research*
242, 301-311 (1985).
- 59 King, J. C. & Rubin, B. S. Dynamic alterations in luteinizing hormone-releasing
hormone (LHRH) neuronal cell bodies and terminals of adult rats. *Cellular and*
molecular neurobiology **15**, 89-106 (1995).
- 60 Giacobini, P. *et al.* Brain endothelial cells control fertility through ovarian-steroid-
dependent release of semaphorin 3A. *PLoS Biol* **12**, e1001808,
doi:10.1371/journal.pbio.1001808 (2014).
- 61 King, J. C. & Letourneau, R. J. Luteinizing hormone-releasing hormone terminals
in the median eminence of rats undergo dramatic changes after gonadectomy, as
revealed by electron microscopic image analysis. *Endocrinology* **134**, 1340-1351
(1994).
- 62 Prevot, V. *et al.* Definitive evidence for the existence of morphological plasticity in
the external zone of the median eminence during the rat estrous cycle:
implication of neuro-glio-endothelial interactions in gonadotropin-releasing
hormone release. *Neuroscience* **94**, 809-819 (1999).
- 63 Parkash, J. *et al.* Semaphorin7A regulates neuroglial plasticity in the adult
hypothalamic median eminence. *Nat Commun* **6**, 6385,
doi:10.1038/ncomms7385 (2015).
- 64 Clarke, I. J. Evidence that the switch from negative to positive feedback at the
level of the pituitary gland is an important timing event for the onset of the
preovulatory surge in LH in the ewe. *The Journal of endocrinology* **145**, 271-282
(1995).
- 65 Barkan, A. L., Regiani, S. R., Duncan, J. A. & Marshall, J. C. Pituitary gonadotropin-
releasing hormone receptors during gonadotropin surges in ovariectomized-

- estradiol-treated rats. *Endocrinology* **112**, 1042-1048, doi:10.1210/endo-112-3-1042 (1983).
- 66 Qiao, S. *et al.* Molecular Plasticity of Male and Female Murine Gonadotropes Revealed by mRNA Sequencing. *Endocrinology* **157**, 1082-1093, doi:10.1210/en.2015-1836 (2016).
- 67 Bliss, S. P. *et al.* Signaling complexes associated with the type I gonadotropin-releasing hormone (GnRH) receptor: colocalization of extracellularly regulated kinase 2 and GnRH receptor within membrane rafts. *Molecular endocrinology* **21**, 538-549, doi:10.1210/me.2006-0289 (2007).
- 68 Tibolt, R. E. & Childs, G. V. Cytochemical and cytophysiological studies of gonadotropin-releasing hormone (GnRH) target cells in the male rat pituitary: differential effects of androgens and corticosterone on GnRH binding and gonadotropin release. *Endocrinology* **117**, 396-404, doi:10.1210/endo-117-1-396 (1985).
- 69 Hodson, D. J. *et al.* Coordination of calcium signals by pituitary endocrine cells in situ. *Cell calcium* **51**, 222-230, doi:10.1016/j.ceca.2011.11.007 (2012).
- 70 Padmanabhan, V. Polycystic ovary syndrome--"A riddle wrapped in a mystery inside an enigma". *J Clin Endocrinol Metab* **94**, 1883-1885, doi:10.1210/jc.2009-0492 (2009).
- 71 Francou, M. *et al.* Characterization of pituitary cell populations in rats with induced polycystic ovaries. *Cells Tissues Organs* **188**, 310-319, doi:10.1159/000123202 (2008).
- 72 Roland, A. V. & Moenter, S. M. Reproductive neuroendocrine dysfunction in polycystic ovary syndrome: insight from animal models. *Front Neuroendocrinol* **35**, 494-511, doi:10.1016/j.yfrne.2014.04.002 (2014).
- 73 Cardoso, R. C., Puttabyatappa, M. & Padmanabhan, V. Steroidogenic versus Metabolic Programming of Reproductive Neuroendocrine, Ovarian and Metabolic Dysfunctions. *Neuroendocrinology* **102**, 226-237, doi:10.1159/000381830 (2015).
- 74 Pitteloud, N., Durrani, S., Raivio, T. & Sykiotis, G. P. Complex genetics in idiopathic hypogonadotropic hypogonadism. *Front Horm Res* **39**, 142-153, doi:10.1159/000312700 (2010).
- 75 Christin-Maitre, S., de Crecy, M. & Groupe Francais des pompes a Gn, R. H. [Pregnancy outcomes following pulsatile GnRH treatment: results of a large multicenter retrospective study]. *J Gynecol Obstet Biol Reprod (Paris)* **36**, 8-12, doi:10.1016/j.jgyn.2006.12.001 (2007).
- 76 Boehm, U. *et al.* Expert consensus document: European Consensus Statement on congenital hypogonadotropic hypogonadism--pathogenesis, diagnosis and treatment. *Nat Rev Endocrinol* **11**, 547-564, doi:10.1038/nrendo.2015.112 (2015).
- 77 Sidhoum, V. F. *et al.* Reversal and relapse of hypogonadotropic hypogonadism: resilience and fragility of the reproductive neuroendocrine system. *J Clin Endocrinol Metab* **99**, 861-870, doi:10.1210/jc.2013-2809 (2014).
- 78 Grattan, D. R. & Kokay, I. C. Prolactin: a pleiotropic neuroendocrine hormone. *Journal of neuroendocrinology* **20**, 752-763, doi:10.1111/j.1365-2826.2008.01736.x (2008).
- 79 Arbogast, L. A. & Voogt, J. L. Hyperprolactinemia increases and hypoprolactinemia decreases tyrosine hydroxylase messenger ribonucleic acid

- levels in the arcuate nuclei, but not the substantia nigra or zona incerta. *Endocrinology* **128**, 997-1005, doi:10.1210/endo-128-2-997 (1991).
- 80 Stagkourakis, S., Kim, H., Lyons, D. J. & Broberger, C. Dopamine Autoreceptor Regulation of a Hypothalamic Dopaminergic Network. *Cell Rep*, doi:10.1016/j.celrep.2016.03.062 (2016).
- 81 Lyons, D. J., Horjales-Araujo, E. & Broberger, C. Synchronized network oscillations in rat tuberoinfundibular dopamine neurons: switch to tonic discharge by thyrotropin-releasing hormone. *Neuron* **65**, 217-229, doi:10.1016/j.neuron.2009.12.024 (2010).
- 82 Freeman, M. E., Reichert, L. E., Jr. & Neill, J. D. Regulation of the proestrus surge of prolactin secretion by gonadotropin and estrogens in the rat. *Endocrinology* **90**, 232-238, doi:10.1210/endo-90-1-232 (1972).
- 83 Larsen, C. M. & Grattan, D. R. Prolactin-induced mitogenesis in the subventricular zone of the maternal brain during early pregnancy is essential for normal postpartum behavioral responses in the mother. *Endocrinology* **151**, 3805-3814, doi:10.1210/en.2009-1385 (2010).
- 84 Mai, L. M., Shieh, K. R. & Pan, J. T. Circadian changes of serum prolactin levels and tuberoinfundibular dopaminergic neuron activities in ovariectomized rats treated with or without estrogen: the role of the suprachiasmatic nuclei. *Neuroendocrinology* **60**, 520-526 (1994).
- 85 Egli, M., Bertram, R., Sellix, M. T. & Freeman, M. E. Rhythmic secretion of prolactin in rats: action of oxytocin coordinated by vasoactive intestinal polypeptide of suprachiasmatic nucleus origin. *Endocrinology* **145**, 3386-3394, doi:10.1210/en.2003-1710 (2004).
- 86 Featherstone, K. *et al.* Pulsatile patterns of pituitary hormone gene expression change during development. *Journal of Cell Science* **124**, 3484-3491, doi:jcs.088500 [pii] 10.1242/jcs.088500 (2011).
- 87 Harper, C. V. *et al.* Dynamic organisation of prolactin gene expression in living pituitary tissue. *Journal of Cell Science* **123**, 424-430, doi:123/3/424 [pii] 10.1242/jcs.060434 (2010).
- 88 Long, T. *et al.* Quantifying the integration of quorum-sensing signals with single-cell resolution. *PLoS Biol* **7**, e68, doi:10.1371/journal.pbio.1000068 (2009).
- 89 Weber, W. *et al.* Streptomyces-derived quorum-sensing systems engineered for adjustable transgene expression in mammalian cells and mice. *Nucleic Acids Res* **31**, e71 (2003).
- 90 Mangel, M. & Bonsall, M. B. Phenotypic evolutionary models in stem cell biology: replacement, quiescence, and variability. *PloS one* **3**, e1591, doi:10.1371/journal.pone.0001591 (2008).
- 91 Andrews, Z. B., Kokay, I. C. & Grattan, D. R. Dissociation of prolactin secretion from tuberoinfundibular dopamine activity in late pregnant rats. *Endocrinology* **142**, 2719-2724, doi:10.1210/endo.142.6.8196 (2001).
- 92 Ciofi, P. *et al.* Plasticity in expression of immunoreactivity for neuropeptide Y, enkephalins and neurotensin in the hypothalamic tubero-infundibular dopaminergic system during lactation in mice. *Journal of neuroendocrinology* **5**, 599-602 (1993).
- 93 Merchenthaler, I. Induction of enkephalin in tuberoinfundibular dopaminergic neurons during lactation. *Endocrinology* **133**, 2645-2651, doi:10.1210/endo.133.6.7694844 (1993).

- 94 Castrique, E., Fernandez-Fuente, M., Le Tissier, P., Herman, A. & Levy, A. Use of a prolactin-Cre/ROSA-YFP transgenic mouse provides no evidence for lactotroph transdifferentiation after weaning, or increase in lactotroph/somatotroph proportion in lactation. *Journal of Endocrinology* **205**, 49-60, doi:JOE-09-0414 [pii] 10.1677/JOE-09-0414 (2010).
- 95 Le Tissier, P. R., Hodson, D. J., Martin, A. O., Romano, N. & Mollard, P. Plasticity of the prolactin (PRL) axis: mechanisms underlying regulation of output in female mice. *Advances in experimental medicine and biology* **846**, 139-162, doi:10.1007/978-3-319-12114-7_6 (2015).
- 96 Guillou, A. *et al.* Assessment of Lactotroph Axis Functionality in Mice: Longitudinal Monitoring of PRL Secretion by Ultrasensitive-ELISA. *Endocrinology* **156**, 1924-1930, doi:10.1210/en.2014-1571 (2015).
- 97 Anderson, S. T. *et al.* Mechanisms underlying the diminished sensitivity to prolactin negative feedback during lactation: reduced STAT5 signaling and up-regulation of cytokine-inducible SH2 domain-containing protein (CIS) expression in tuberoinfundibular dopaminergic neurons. *Endocrinology* **147**, 1195-1202, doi:10.1210/en.2005-0905 (2006).
- 98 Byrnes, E. M. & Bridges, R. S. Lactation reduces prolactin levels in reproductively experienced female rats. *Horm Behav* **48**, 278-282, doi:10.1016/j.yhbeh.2005.03.006 (2005).
- 99 Musey, V. C., Collins, D. C., Musey, P. I., Martino-Saltzman, D. & Preedy, J. R. Long-term effect of a first pregnancy on the secretion of prolactin. *The New England journal of medicine* **316**, 229-234, doi:10.1056/NEJM198701293160501 (1987).
- 100 Byrnes, E. M. & Bridges, R. S. Reproductive experience and expression of dopamine D(2) receptor mRNA: a possible mechanism for reduced prolactin secretion in primiparous rats. *Journal of neuroendocrinology* **19**, 773-778, doi:10.1111/j.1365-2826.2007.01586.x (2007).
- 101 Wong, A., Eloy, J. A., Couldwell, W. T. & Liu, J. K. Update on prolactinomas. Part 1: Clinical manifestations and diagnostic challenges. *J Clin Neurosci* **22**, 1562-1567, doi:10.1016/j.jocn.2015.03.058 (2015).
- 102 Holt, R. I. & Peveler, R. C. Antipsychotics and hyperprolactinaemia: mechanisms, consequences and management. *Clin Endocrinol (Oxf)* **74**, 141-147, doi:10.1111/j.1365-2265.2010.03814.x (2011).
- 103 Wong, A., Eloy, J. A., Couldwell, W. T. & Liu, J. K. Update on prolactinomas. Part 2: Treatment and management strategies. *J Clin Neurosci* **22**, 1568-1574, doi:10.1016/j.jocn.2015.03.059 (2015).
- 104 Brown, R. S., Herbison, A. E. & Grattan, D. R. Prolactin regulation of kisspeptin neurones in the mouse brain and its role in the lactation-induced suppression of kisspeptin expression. *Journal of neuroendocrinology* **26**, 898-908, doi:10.1111/jne.12223 (2014).
- 105 Sonigo, C. *et al.* Hyperprolactinemia-induced ovarian acyclicity is reversed by kisspeptin administration. *J Clin Invest* **122**, 3791-3795, doi:10.1172/JCI63937 (2012).
- 106 Li, Q., Rao, A., Pereira, A., Clarke, I. J. & Smith, J. T. Kisspeptin cells in the ovine arcuate nucleus express prolactin receptor but not melatonin receptor. *Journal of neuroendocrinology* **23**, 871-882, doi:10.1111/j.1365-2826.2011.02195.x (2011).

- 107 Hodson, D. J., Townsend, J. & Tortonese, D. J. Characterization of the effects of prolactin in gonadotroph target cells. *Biology of Reproduction* **83**, 1046-1055, doi:biolreprod.110.084947 [pii] 10.1095/biolreprod.110.084947 (2010).
- 108 Adcock, C. J. *et al.* The use of an automated microsampling system for the characterization of growth hormone pulsatility in newborn babies. *Pediatr Res* **42**, 66-71, doi:10.1203/00006450-199707000-00011 (1997).
- 109 Coxam, V., Davicco, M. J., Robelin, J. & Barlet, J. P. Growth hormone secretory pattern and somatomedin C plasma concentrations in newborn calves. *J Dev Physiol* **9**, 113-121 (1987).
- 110 Davicco, M. J. *et al.* Growth hormone (GH) secretory pattern and GH response to GH-releasing factor (GRF) or thyrotropin-releasing hormone (TRH) in newborn foals. *J Dev Physiol* **19**, 143-147 (1993).
- 111 Robinson, I. C. A. F. & Hindmarsh, P. C. in *Comprehensive Physiology* (John Wiley & Sons, Inc., 2010).
- 112 Tannenbaum, G. S. Genesis of episodic growth hormone secretion. *J Pediatr Endocrinol* **6**, 273-282 (1993).
- 113 Steyn, F. J., Tolle, V., Chen, C. & Epelbaum, J. Neuroendocrine Regulation of Growth Hormone Secretion. *Compr Physiol* **6**, 687-735, doi:10.1002/cphy.c150002 (2016).
- 114 Balthasar, N. *et al.* Growth hormone-releasing hormone (GHRH) neurons in GHRH-enhanced green fluorescent protein transgenic mice: a ventral hypothalamic network. *Endocrinology* **144**, 2728-2740, doi:10.1210/en.2003-0006 (2003).
- 115 Gouty-Colomer, L. A. *et al.* Specific involvement of gonadal hormones in the functional maturation of growth hormone releasing hormone (GHRH) neurons. *Endocrinology* **151**, 5762-5774, doi:10.1210/en.2010-0298 (2010).
- 116 Romero, M. I. & Phelps, C. J. Identification of growth hormone-releasing hormone and somatostatin neurons projecting to the median eminence in normal and growth hormone-deficient Ames dwarf mice. *Neuroendocrinology* **65**, 107-116 (1997).
- 117 Chowen, J. A., Frago, L. M. & Argente, J. The regulation of GH secretion by sex steroids. *Eur J Endocrinol* **151 Suppl 3**, U95-100 (2004).
- 118 Baccam, N. *et al.* Dual-level afferent control of growth hormone-releasing hormone (GHRH) neurons in GHRH-green fluorescent protein transgenic mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **27**, 1631-1641, doi:10.1523/JNEUROSCI.2693-06.2007 (2007).
- 119 Osterstock, G. *et al.* Somatostatin triggers rhythmic electrical firing in hypothalamic GHRH neurons. *Sci Rep* **6**, 24394, doi:10.1038/srep24394 (2016).
- 120 Schaeffer, M. *et al.* Influence of estrogens on GH-cell network dynamics in females: a live in situ imaging approach. *Endocrinology* **152**, 4789-4799, doi:10.1210/en.2011-1430 (2011).
- 121 Alatzoglou, K. S., Webb, E. A., Le Tissier, P. & Dattani, M. T. Isolated growth hormone deficiency (GHD) in childhood and adolescence: recent advances. *Endocrine reviews* **35**, 376-432, doi:10.1210/er.2013-1067 (2014).
- 122 Pekic, S. & Popovic, V. Alternative causes of hypopituitarism: traumatic brain injury, cranial irradiation, and infections. *Handbook of clinical neurology* **124**, 271-290, doi:10.1016/B978-0-444-59602-4.00018-6 (2014).

- 123 Andoniadou, C. L. *et al.* Sox2(+) stem/progenitor cells in the adult mouse pituitary support organ homeostasis and have tumor-inducing potential. *Cell stem cell* **13**, 433-445, doi:10.1016/j.stem.2013.07.004 (2013).
- 124 Olarescu, N. C. & Bollerslev, J. The Impact of Adipose Tissue on Insulin Resistance in Acromegaly. *Trends in endocrinology and metabolism: TEM* **27**, 226-237, doi:10.1016/j.tem.2016.02.005 (2016).
- 125 Dimaraki, E. V., Jaffe, C. A., DeMott-Friberg, R., Chandler, W. F. & Barkan, A. L. Acromegaly with apparently normal GH secretion: implications for diagnosis and follow-up. *J Clin Endocrinol Metab* **87**, 3537-3542, doi:10.1210/jcem.87.8.8658 (2002).
- 126 Holly, J. M. *et al.* Inter-relations between growth hormone, insulin, insulin-like growth factor-I (IGF-I), IGF-binding protein-1 (IGFBP-1) and sex hormone-binding globulin in acromegaly. *Clin Endocrinol (Oxf)* **34**, 275-280 (1991).
- 127 Ward, R. D., Stone, B. M., Raetzman, L. T. & Camper, S. A. Cell proliferation and vascularization in mouse models of pituitary hormone deficiency. *Molecular endocrinology* **20**, 1378-1390, doi:10.1210/me.2005-0409 (2006).
- 128 Narayanaswamy, S. *et al.* Subcutaneous infusion of kisspeptin-54 stimulates gonadotrophin release in women and the response correlates with basal oestradiol levels. *Clin Endocrinol (Oxf)*, doi:10.1111/cen.12977 (2015).
- 129 Alonso, G. *et al.* Selective alteration at the growth-hormone-releasing-hormone nerve terminals during aging in GHRH-green fluorescent protein mice. *Aging Cell* **6**, 197-207, doi:10.1111/j.1474-9726.2007.00276.x (2007).
- 130 Andoniadou, C. L. *et al.* Identification of novel pathways involved in the pathogenesis of human adamantinomatous craniopharyngioma. *Acta neuropathologica* **124**, 259-271, doi:10.1007/s00401-012-0957-9 (2012).
- 131 Eckstrum, K. S., Weis, K. E., Baur, N. G., Yoshihara, Y. & Raetzman, L. T. Icam5 Expression Exhibits Sex Differences in the Neonatal Pituitary and Is Regulated by Estradiol and Bisphenol A. *Endocrinology* **157**, 1408-1420, doi:10.1210/en.2015-1521 (2016).
- 132 Le Tissier, P. R. & Mollard, P. Bisphenol A Effects on Gonadotroph Function: Disruption of Pituitary Cell-Cell Communication? *Endocrinology* **157**, 1324-1325, doi:10.1210/en.2016-1117 (2016).

Competing interests statement

The authors declare no competing interests

Acknowledgements [Au: edits OK?]

We thank F. Castinetti (Marseille [Au: which institute? Aix-Marseille Universite?]) for helpful comments and suggestions. P.L.T. was supported by a grant from the Biotechnology and Biological Sciences Research Council, UK (BB/N007026/1). N.R. was supported by a Medical Research Council, UK project grant (MR/J008893/1). D.J.H. was supported by an R.D. Lawrence Fellowship, Diabetes UK (12/0004431), European Foundatin for the Study of Diabetes /Novo Nordisk Rising Star and Birmingham Fellowships, a Medical Research Council, UK project grant (MR/N00275X/1), Imperial Confidence in Concept and Wellcome Trust Institutional Support Awards, both UK. P.M. was supported by funding from the

Agence Nationale de la Recherche (ANR 12 BSV1 0032-01, ANR-15-CE14-0012-01), Institut National de la Santé et de la Recherche Médicale, Centre National de la Recherche Scientifique, Université de Montpellier, Fondation pour la Recherche Médicale (DEQ20150331732), and IPAM-Biocampus of Montpellier and France-Bioimaging, all in France.

Author contributions [Au: OK?]

The authors contributed equally to all aspects of the preparation of this article

Key points [Au: edits OK?]

- The activity of hypothalamic neurons is modified by inputs leading to heterogeneous activity; a small proportion of the total population can drive pituitary hormone pulsatility
- Neurohormone output can vary following neuron excitation according to physiological status, which might also lead to declining neuroendocrine output with age
- The release of hypothalamic factors into the blood is modified by alterations in the juxtaposition of nerve terminals with the vasculature and tanycytes in the median eminence
- Cells in the pituitary gland form homotypic networks, the organization of these relationships with the vasculature are distinct for each endocrine axis, which modifies responses to regulatory factors and patterns of output in response to demand [Au: edit OK?]
- Reorganisation of the pituitary network can store long-term memories of increased output and learn to increase function on repeated challenge
- Understanding the importance of coordinated hypothalamic–vasculature–pituitary function provides new understanding of a range of endocrine axes defects and targets for novel therapies

[Au: a list of suggested Figures and content:

Figure 1 – General introduction of the relationship between the 3rd ventricle, median eminence, portal vessels, vasculature and pituitary. We have a range of in-house options that can be adapted for this purpose. – I can send you an example or two if you wish

Figure 2 – The gonadotroph axis – to include the organisation of the GnRH/LH network in the hypothalamic/pituitary/vasculature and with factors that might influence (e.g. NO). Might also include the homotypic organisation at puberty. E.g. a ‘before’ and ‘after’ panel

Figure 3 – The prolactin axis – could include a representation of the lactotrophs and their development into networks – e.g. an expansion of what is represented in the bottom left of current Figure 1.

Figure 4 – The GH axis – to include a graphical view of this axis and could include how this axis is modified to develop the pulsatility or the modification of pituitary somatotroph output

If simpler, perhaps proposed Figures 2, 3 and 4 could be combined into a single, multi-panel figure if you wish?

Figure 5 – summary of our current understanding – based on your submitted figure 1, although some of these elements can be included in the above figure

Figure 1 | Regulation of the hypothalamic-pituitary axes. In the past decade, a more complex relationship between the hypothalamus and pituitary than previously appreciated has emerged. **a)** At the level of the parvocellular neurons, complex inputs modify the excitation of neurons, which can vary coupling with neurohormone output at terminals of the median eminence (ME) through modification of intracellular pathways. **b)** Alterations in tanycyte ensheathment and the anatomical location of neuron terminals modify their interactions with the vasculature, changing the dynamics of neurohormone release. **c)** In the pituitary gland, changes in blood flow in the portal circulation alter the pattern of exposure of pituitary cells to neurohormone and nutrient supply to facilitate secretion. **d)** Cells of the pituitary gland are organised into intermingled networks with distinct morphological features, which can be altered to meet physiological demand, and relationships with the vasculature.

[Au: I have made a number of edits to the text in this box to adhere to journal style and the overall flow of the article. However, I think some of these concepts could be represented in the suggested figures]

Box 1 | Organization of pituitary cells into homotypic networks

During the past decade, we and others have shown that the pituitary somatotroph¹³, corticotroph¹⁴, gonadotroph^{14,15} and lactotroph¹⁶ lineages form 3D organized cell networks (reviewed in^{17,18}) with the following features:

- *The lineages have distinct developmental programmes*
 - Placement of each endocrine cell network occurs at distinct stages of pituitary organogenesis, before expansion in early post-natal life **[Au: please include the references here]**.
- *The networks have distinct motifs that might relate to function.*
 - For example, the organization of somatotrophs as clusters linked with strands¹³, while lactotrophs form a honeycomb structure¹⁶. **[Au: how do these relate to function specifically? Or is this just an association?]**
- *Hypothalamic factors generate network motifs*
 - For example, the loss of growth hormone-releasing hormone (GHRH) itself leads to isolated somatotrophs, whereas somatotroph ablation with intact GHRH stimulation results in clusters of cells that are isolated from each other¹⁹.
- *The pituitary networks have functional relevance.*
 - Critical roles have been described for pituitary networks in the integration, amplification and propagation of the neurohaemal hypothalamic signals that arrive via the median eminence. For example, the male somatotroph network responds to GHRH input with large, coordinated and oscillatory Ca²⁺ rises that outlast stimulus to drive large excursions in hormone secretion²⁰.
- *Endocrine and non-endocrine cells interact to form homotypic networks* **[Au: edit OK? Is this what you mean here?]**.
 - For example, gap-junctions mediate coupling **[Au: of what cells specifically?]** with folliculostellate cells^{16,21}, paracrine and autocrine interactions^{17,22} as well as the relationship between networks and the vasculature^{7,23,24}. These interactions might help to alter **[Au: somatotroph or simply hormone?]** output. The distinct network motifs of lactotrophs are aligned with the fine pituitary capillary network, while somatotrophs form strands and clusters along the same vessels¹⁷. Similarly, gonadotrophs are in close approximation to one or more blood vessels via their protrusions¹⁵, whereas the corticotroph network has a looser arrangement¹⁴. **[Au: meaning these differences will shape how the neurohormones are released e.g. in terms of timing?]**

Author biographies [Au: edits OK?]

Paul Le Tissier is a Senior lecturer in the Centre for Integrative Physiology, University of Edinburgh, UK. His primary research interests are the understanding of hypothalamic regulation of the anterior pituitary gland, how this mechanism maintains and alters the function of the different cell populations to ensure appropriate output throughout life and modelling how dysregulation leads to pathology.

Pauline Campos has recently become a postdoctoral fellow in the laboratory of Patrice Mollard, at the Institute of Functional Genomics of Montpellier, France. She completed her PhD in the Center for Neuroendocrinology at the University of Otago, New Zealand, and has been focused on developing cutting-edge tools to study the electrical activity of hypothalamic neurons underlying pituitary hormone pulsatility. Her current work aims at understanding the central controls of thyroid function in conscious animals

Chrystel Lafont is an INSERM engineer in the Institute of Functional Genomics of Montpellier, France. She is the technical leader of a small animal *in vivo* imaging facility that develops cellular *in vivo* imaging techniques. Her research interests look at the role of the vasculature and pituitary cell networks in the build-up of hormone pulses.

Nicola Romanò is a postdoctoral fellow at the University of Edinburgh, UK. His main interest is the study of the cellular and systemic mechanisms of pulsatile hormone secretion in neuroendocrine axes, and how they correlate and influence physiological function. His research interests span from analysis of activity patterns of hypothalamic neurons to structural and functional analysis of pituitary cell networks, such as, the spatio-temporal analysis of cell activity.

David Hodson is a Senior Birmingham Fellow, Diabetes UK–RD Lawrence Fellow and European Foundation for the Study of Diabetes/Novo Nordisk Rising Star Fellow. He studied Veterinary Medicine at the University of Bristol, UK, before becoming a group leader at Imperial College London and latterly at the University of Birmingham, UK. His laboratory is developing innovative optical methodologies that enable the mapping and manipulation of endocrine cells directly within their tissue context.

Patrice Mollard is the head of the Department of Physiology and CNRS Research Director at the Institute of Functional Genomics of Montpellier, France. Dr Mollard's laboratory pioneered the study of large-scale functional organization of endocrine cell types within the pituitary gland in the early 2000s. His research is now exploring how pituitary cell networks function together with hypothalamic inputs and vasculature in health and disease.