

## Analysis of *bla*<sub>SHV-12</sub>-carrying *Escherichia coli* clones and plasmids from human, animal and food sources

Carla Andrea Alonso<sup>1</sup>, Geovana Brenner Michael<sup>2,3</sup>, Jun Li<sup>2,4</sup>, Sergio Somalo<sup>1</sup>, Carmen Simón<sup>5</sup>, Yang Wang<sup>4</sup>, Heike Kaspar<sup>6</sup>, Kristina Kadlec<sup>2</sup>, Carmen Torres<sup>1\*</sup> and Stefan Schwarz<sup>2,3</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology, University of La Rioja, Logroño, Spain; <sup>2</sup>Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Neustadt-Mariensee, Germany; <sup>3</sup>Department of Veterinary Medicine, Institute of Microbiology and Epizootics, Centre of Infection Medicine, Freie Universität Berlin, Berlin, Germany; <sup>4</sup>College of Veterinary Medicine, China Agricultural University, Beijing, China; <sup>5</sup>Faculty of Veterinary Science, University of Zaragoza, Zaragoza, Spain; <sup>6</sup>Federal Office of Consumer Protection and Food Safety (BVL), Berlin, Germany

\*Corresponding author. Tel: +34-941299750; Fax: +34-941299721; E-mail: carmen.torres@unirioja.es

Received 17 November 2016; returned 16 December 2016; revised 3 January 2017; accepted 15 January 2017

**Objectives:** This study aimed at characterizing 23 *Escherichia coli* isolates from various sources and their respective *bla*<sub>SHV-12</sub>-carrying plasmids and sequencing one of these plasmids completely.

**Methods:** Isolates were typed by XbaI-PFGE, MLST and PCR-based phylotyping. Transformed *bla*<sub>SHV-12</sub>-carrying plasmids were examined by replicon typing, S1-nuclease, conjugation, EcoRI-HindIII-BamHI digests and plasmid MLST. Co-located resistance genes and integrons as well as the *bla*<sub>SHV-12</sub> genetic environment were analysed by PCR and sequencing. One IncI1 plasmid was sequenced completely using HiSeq 2500 and gap closure by PCRs and Sanger sequencing.

**Results:** Among the 23 SHV-12-positive *E. coli*, some isolates from different sources showed the same characteristics: ST23/phylogroup A (human, dog, livestock), ST57/D (wild bird, chicken meat) and ST117/D (chicken meat, chicken). All *bla*<sub>SHV-12</sub> genes were horizontally transferable via 30–120 kb plasmids of incompatibility groups IncI1 (*n* = 17), IncK (*n* = 3), IncF (*n* = 1), IncX3 (*n* = 1) and a non-typeable plasmid. IncK plasmids, indistinguishable in size and restriction patterns, were found in isolates from different sources (ST57/D, meat; ST131/B2, meat; ST57/B1, dog). The IncI1-*bla*<sub>SHV-12</sub>-carrying plasmids were mostly assigned to plasmid ST (pST) 26 and pST3. Three plasmids showed novel pSTs (pST214, pST215). The majority of the IncI1 transformants exhibited resistance to β-lactams, chloramphenicol and streptomycin (in relation with a class 1 integron containing an *estX-psp-aadA2-cmlA1-aadA1-qacI* gene cassette array), and to tetracycline. A novel *bla*<sub>SHV-12</sub> environment was detected and whole plasmid sequencing revealed a Tn21-derived-*bla*<sub>SHV12</sub>-ΔTn1721 resistance complex.

**Conclusions:** Results from this study suggest that the dissemination of *bla*<sub>SHV-12</sub> genes occurs by vertical (clonal) and horizontal transfer, the latter mainly mediated through IncI1 multidrug-resistance plasmids.

### Introduction

The WHO has defined third- and fourth-generation cephalosporins as critically important antimicrobial agents in human medicine.<sup>1</sup> In Gram-negative bacteria resistance to these antimicrobials has become a major health problem associated with the production of ESBLs such as those of the TEM, SHV and CTX-M families. The presence of ESBL-producing *Escherichia coli* has been widely reported in not only humans but also food,<sup>2</sup> pets,<sup>3</sup> livestock<sup>4,5</sup> and even wildlife.<sup>6</sup>

The CTX-M family is currently the most prevalent worldwide, but other ESBLs, such as SHV-12, remain important among pathogens causing nosocomial and community-acquired infections in

many Southern European and Asian countries and have also been reported in *E. coli* isolated from livestock and wild birds.<sup>4,6–10</sup> Furthermore, SHV-12 was reported as the most prevalent enzyme detected in ESBL-producing Enterobacteriaceae from retail chicken meat and poultry in both Germany and Spain.<sup>11,12</sup>

Although several studies have examined mobile elements carrying *bla*<sub>CTX-M</sub> genes,<sup>13,14</sup> fewer data are available for the *bla*<sub>SHV-12</sub> gene. Thus, the aim of this study was to characterize a collection of *bla*<sub>SHV-12</sub>-positive *E. coli* from different sources and geographical origins and their corresponding *bla*<sub>SHV-12</sub>-carrying plasmids to gain insight into the presence and dissemination of this ESBL gene. Furthermore, we determined the complete sequence of a plasmid

harbouring the *bla*<sub>SHV-12</sub> gene in addition to several other resistance genes.

## Materials and methods

### Bacterial collection, susceptibility testing and clonal characterization of *bla*<sub>SHV-12</sub>-positive isolates

Twenty-three *bla*<sub>SHV-12</sub>-positive *E. coli* from different sources and origins were analysed. Sources of the Spanish isolates are as follows: (i) wild birds ( $n = 4$ ; starling, cuckoo, two storks): cloacal samples collected in the Aragón Reference Centre of Wildlife Recovering (La Alfranca, 2014);<sup>6</sup> (ii) dogs ( $n = 3$ ): faeces of healthy dogs from different kennels (Logroño, 2009); (iii) chicken meat samples ( $n = 4$ ) collected from different supermarkets (Logroño, 2011); (iv) chickens ( $n = 5$ ): faeces of chickens from different slaughterhouses ( $n = 4$ ) and a liver sample from a diseased animal ( $n = 1$ ) (Spain, 2003);<sup>10</sup> and (v) humans ( $n = 3$ ): faecal samples of patients admitted to a Spanish hospital (June–July 2008). Sources of the German isolates: tissue samples of diseased livestock birds ( $n = 4$ ; duck, turkey, two chickens) raised on different farms, collected by the German national resistance monitoring programme (GERM-Vet) (2010–11).

All isolates were tested for susceptibility to ampicillin, amoxicillin/clavulanate, ceftazidime, ceftriaxone, ceftiofur, imipenem, nalidixic acid, ciprofloxacin, gentamicin, amikacin, tobramycin, streptomycin, chloramphenicol, compound sulphonamides, trimethoprim/sulfamethoxazole and tetracycline by disc diffusion according to the CLSI criteria.<sup>15</sup> ESBL production was verified by a double-disc synergy test. *E. coli* ATCC 25922 served as a control strain.

Carriage of the *bla*<sub>SHV-12</sub> gene was confirmed by PCR and sequencing.<sup>16</sup> Genetic diversity of the *bla*<sub>SHV-12</sub>-positive isolates was analysed using PCR-based phylotyping, MLST and XbaI-macrorestriction followed by PFGE.<sup>17,18</sup> A dendrogram, for the analysis of the XbaI-PFGE patterns, was generated using GelJ version 1.3 (UPGMA algorithm; Dice coefficient; 1% tolerance).<sup>19</sup>

### Transfer and characterization of *bla*<sub>SHV-12</sub>-carrying plasmids

Plasmids were transferred by conjugation and electrotransformation using the sodium azide-resistant *E. coli* J53 strain and electro-competent *E. coli* TOP10 as recipient cells, respectively.<sup>20</sup> Transconjugants and transformants were selected on LB agar supplemented with ceftazidime (1 mg/L) and sodium azide (200 mg/L) or with ceftazidime (1 mg/L), respectively.

Plasmids were characterized by PCR-based replicon typing, S1 nuclease digestion followed by PFGE and restriction fragment length polymorphism using the EcoRI, HindIII or BamHI endonucleases.<sup>13,21</sup> IncI1 plasmids were subtyped by plasmid MLST (pMLST).<sup>22</sup>

### Antimicrobial resistance genes, integrons and *bla*<sub>SHV-12</sub> genetic environment

Genes associated with resistance to  $\beta$ -lactams (*bla*<sub>OXA</sub>, *bla*<sub>CTX-M</sub>, *bla*<sub>SHV</sub>, *bla*<sub>TEM</sub>), aminoglycosides [*aac*(3)-I, *aac*(3)-II, *aac*(3)-III, *aac*(3)-IV, *strA*, *strB*], phenicols (*cmlA*, *floR*, *catB3*), quinolones [*qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, *aac*(6')-Ib-cr, *qepA*, *oqxA*-*oqxB*], sulphonamides (*sul1*, *sul2*, *sul3*) and tetracycline [*tet*(A-E)] were tested by PCR in all original isolates and *bla*<sub>SHV-12</sub>-positive transformants.<sup>23,24</sup>

The presence of *intI1* and *intI2* genes, the variable region of the integrons and the genetic structure of their 3'-conserved segments (CSs) were determined by PCR and sequencing.<sup>25,26</sup> The variable region of the class 1 integron carried by *E. coli* C526 was annotated and submitted to the GenBank database (KU317749).

To elucidate the *bla*<sub>SHV-12</sub> genetic environment, a PCR strategy was carried out using previously reported primers.<sup>27,28</sup> To characterize the uncommon downstream region of the *bla*<sub>SHV-12</sub> gene in *E. coli* isolate 101689, a newly

designed primer was used (DEOR\_ge1: 5'-AGGGTACCGCTTCCCAATC-3'). Its design was based on the draft sequence of the *bla*<sub>SHV-12</sub>-carrying plasmid (pCAZ460, *E. coli* 101689) (data not shown).

### Sequencing of *bla*<sub>SHV-12</sub>-carrying plasmids

Plasmid sequencing of two *bla*<sub>SHV-12</sub>-carrying plasmids pCAZ590 (*E. coli* 111918, from a chicken) and pCAZ460 (*E. coli* 101689, from a broiler) was performed using a HiSeq 2500, which produced 150 bp paired-end reads (Berry Genomics Company, Beijing, China). A draft assembly of the sequences was conducted using the CLC Genomics Workbench 5 (CLC Bio, Aarhus, Denmark); the assembly algorithm works by using de Bruijn graphs. The gap closure was performed by PCR and Sanger sequencing for pCAZ590. The draft sequence of pCAZ460 was used for the characterization of the *bla*<sub>SHV-12</sub> genetic environment and the incompatibility group.

A functional annotation of pCAZ590 was done using the RAST Prokaryotic Genome Annotation Server, which was manually curated using the following bioinformatics tools: Artemis software, IS finder ([www-is.bio.toul.fr](http://www-is.bio.toul.fr)) and Swiss-Prot database (<http://www.uniprot.org>). The EMBOSS Needle alignment tool ([http://www.ebi.ac.uk/Tools/psa/emboss\\_needle/](http://www.ebi.ac.uk/Tools/psa/emboss_needle/)) was used for sequence comparison. A circular map of the plasmid was made using DNAPlotter.<sup>29–31</sup>

The *bla*<sub>SHV-12</sub> genetic environment of *E. coli* 101689 and the full-length sequence of plasmid pCAZ590 (*E. coli* 111918) were deposited in the EMBL database under accession numbers LT621755 and LT669764, respectively.

## Results

### Molecular typing of SHV-12-positive *E. coli* isolates

Table 1 shows the molecular diversity of the *bla*<sub>SHV-12</sub>-positive *E. coli* collection. The 23 *bla*<sub>SHV-12</sub>-carrying isolates displayed 13 clones (ST/phylogroup), with ST23/A ( $n = 5$ ), ST57/D ( $n = 3$ ), ST453/B1 ( $n = 3$ ), ST117/D ( $n = 2$ ) and ST405/D ( $n = 2$ ) as the most common ones. Three isolates from different wild bird species belonged to ST453 and were indistinguishable by XbaI-PFGE (pattern A).<sup>6</sup> The same was true for two ST23/A isolates from different dogs (pattern B) and two poultry isolates (pattern C). ST405 isolates from human origin were closely related (patterns E–E1) (Figure S1, available as Supplementary data at JAC Online).

Some *bla*<sub>SHV-12</sub>-positive *E. coli* isolates obtained from different sources shared the same characteristics: ST23/A (human, dog, duck, turkey), ST57/D (wild bird, chicken meat) and ST117/D (chicken meat, chicken). Additionally, isolates belonging to ST57 (wild bird, chicken meat, dog) showed closely related XbaI-PFGE patterns (D, D1, D2, D3) (Figure S1).

### Transformation, conjugal transfer of ESBL-encoding genes and plasmid characterization

At least two replicon types were detected in each of the 23 *bla*<sub>SHV-12</sub>-positive *E. coli*, with IncI1, IncFIB and IncF being the most common ones. All *bla*<sub>SHV-12</sub> genes were located on 30–120 kb plasmids of the incompatibility groups IncI1 ( $n = 17$ ), IncK ( $n = 3$ ), IncF ( $n = 1$ ) and non-typeable plasmids ( $n = 2$ ) and were transferable by transformation. Using the draft sequence one of these non-typeable plasmids, pCAZ460 (*E. coli* 101689), was assessed as an IncX3 plasmid by the PlasmidFinder server.<sup>32</sup> The ST131/B2 *E. coli* isolate harboured two ESBL genes located on different plasmids: *bla*<sub>SHV-12</sub> was detected on a 75 kb IncK plasmid (Table 2) and *bla*<sub>CTX-M-1</sub> on a 100 kb IncI1 plasmid (data not shown).

**Table 1.** Molecular typing, resistance phenotype, integrons and resistance genes of *bla*<sub>SHV-12</sub>-positive *E. coli* isolates

<i>E. coli</i> isolate (source <sup>a</sup> , origin <sup>b</sup> )	Year of isolation	ST (CC)	Phylogroup	PFGE pattern	ESBL	Resistance phenotype to non-β-lactams <sup>c</sup>	Class 1 integron		Class 2 integron		Resistance genes (outside the integron)
							<i>intI1</i> / <i>3'-CS</i> <sup>d</sup>	variable region	<i>intI2</i>	variable region	
C7377 (W, Sp)	2014	57 (350)	D	D3	SHV-12	CHL-CIP-NAL-STR-SUL-SXT-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	+	dfrA1-sat2-aadA1	tet(B)
C7385 (W, Sp)	2014	453 (86)	B1	A	SHV-12	CHL-CIP-GEN-NAL-STR-SUL-TET-TOB	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), oac(3)-II
C7394 (W, Sp)	2014	453 (86)	B1	A	SHV-12	CHL-CIP-GEN-NAL-STR-SUL-TET-TOB	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), oac(3)-II
C7401 (W, Sp)	2014	453 (86)	B1	A	SHV-12	CHL-CIP-GEN-NAL-STR-SUL-TET-TOB	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), oac(3)-II
C4746 (M, Sp)	2011	57 (350)	D	D2	SHV-12	CIP-NAL-STR-SUL-SXT-TET	-/-	-	-	dfrA1-sat2-aadA1	tet(B), sul2
C4748 (M, Sp)	2011	57 (350)	D	D1	SHV-12	CIP-NAL-STR-SUL-SXT-TET	-/-	-	+	IS10-dfrA1-sat2-aadA1	tet(B), aadA, sul2
Ph461 (M, Sp)	2011	117	D	H	SHV-12	CIP-NAL-SUL-SXT-TET	-/-	-	-	-	tet(A), sul2, bla <sub>TEM-1</sub>
C4745 (M, Sp)	2011	131	B2	G	SHV-12, CTX-M-1	NAL-SUL-TET	-/-	-	-	-	tet(A), sul2
C2585 (D, Sp)	2009	23 (23)	A	B	SHV-12	CHL-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A)
C2578 (D, Sp)	2009	23 (23)	A	B	SHV-12	CHL-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A)
C2575 (D, Sp)	2009	57 (350)	B1	D	SHV-12	CIP-NAL-STR-SUL-SXT-TET	+/+	aadA1	+	dfrA1-sat2-aadA1	tet(A), tet(B), sul2
C1536 (H, Sp)	2008	23 (23)	A	F	SHV-12	CHL-CIP-NAL-STR-SUL-SXT-TET	(i) +/+ (ii) +/-	(i) dfrA1-aadA1 (ii) estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), sul2
C1537 (H, Sp)	2008	405 (450)	D	E	SHV-12	CHL-CIP-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), tet(B)
C1538 (H, Sp)	2008	405 (450)	D	E1	SHV-12	CHL-CIP-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), tet(B)
C353 (L, Sp)	2003	155 (155)	B1	H	SHV-12	CHL-STR-SUL	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	-
C515 (L, Sp)	2003	1564	A	I	SHV-12	CHL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), sul2, bla <sub>TEM-1</sub>
C508 (L, Sp)	2003	2001	D	J	SHV-12	CHL-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(B), bla <sub>TEM-1</sub>
C526 (L, Sp)	2003	362	D	K	SHV-12	SUL-TET	+/+	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), tet(B), sul2
C537 (L, Sp)	2003	616 (155)	B1	L	SHV-12	SUL-TET	-/-	qacG-aadA6-qacG	-	-	tet(A), tet(B), sul3
101689 (L, Ge)	2010	117	D	M	SHV-12	NAL	-/-	-	-	-	qmrS1
101908 (L, Ge)	2010	23 (23)	A	C	SHV-12	CHL-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A)
101985 (L, Ge)	2010	23 (23)	A	C	SHV-12	CHL-NAL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A)
111918 (L, Ge)	2011	371 (350)	D	N	SHV-12	CHL-STR-SUL-TET	+/-	estX-psp-aadA2-cmiA1-aadA1-qacI	-	-	tet(A), sul1

<sup>a</sup>W, wild bird; M, chicken meat; D, dog; H, human; L, livestock bird (poultry).<sup>b</sup>Sp, Spain; Ge, Germany.<sup>c</sup>CHL, chloramphenicol; CIP, ciprofloxacin; GEN, gentamicin; NAL, nalidixic acid; TOB, tobramycin; STR, streptomycin; SUL, compound sulphonamides; SXT, trimethoprim/sulfamethoxazole; TET, tetracycline.<sup>d</sup>Class 1 integrons displaying atypical 3'-CS were identified with the negative result for the investigation of the usual structure of 3'-CS and they were associated with IS440-sul3. The positive results for the 3'-CS indicate the class 1 integrons displaying a usual 3'-CS, which were, as expected, associated with qacEA1-sul1.

**Table 2.** Characteristics of *bla*<sub>SHV-12</sub>-carrying plasmids in the studied *E. coli* collection

E. coli isolate (source <sup>a</sup> , origin <sup>b</sup> )	Year of isolation	ST/ phylogroup	Replicon type	replicon type	IncI1 pMLST (ST/CC)	size (kb)	conjugation frequency	genetic environment of <i>bla</i> <sub>SHV-12</sub>	class 1 integron		other co-located resistance genes
									variable region	int1/3'-CS	
C7377 (W, Sp)	2014	57/D	FIB, F, I1	I1	214	100	2.2 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	-
C7385 (W, Sp)	2014	453/B1	FIB, F, I1	I1 <sup>c</sup>	29 (26)	115	3.5 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C7394 (W, Sp)	2014	453/B1	FIB, F, I1	I1 <sup>c</sup>	29 (26)	115	5.2 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C7401 (W, Sp)	2014	453/B1	FIB, F, I1	I1 <sup>c</sup>	29 (26)	115	3.7 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C4746 (M, Sp)	2011	57/D	B/O, FIB, F, I1	I1	3 (3)	90	5.6 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	-
C4748 (M, Sp)	2011	57/D	FIB, F, K	K <sup>d</sup>	-	75	1.9 × 10 <sup>-4</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	-
Ph461 (M, Sp)	2011	117/D	FIB, F, I1	I1	3 (3)	90	3.5 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	-
C4745 (M, Sp)	2011	131/B2	FIB, F, I1, K	K <sup>d</sup>	-	75	9.6 × 10 <sup>-5</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	-
C2585 (D, Sp)	2009	23/A	FIB, F, I1	I1	26 (26)	110	3.1 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C2578 (D, Sp)	2009	23/A	FIB, F, I1	I1	26 (26)	110	2.0 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C2575 (D, Sp)	2009	57/B1	FIB, F, I1, K, P	K <sup>d</sup>	-	75	7.9 × 10 <sup>-5</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	-
C1536 (H, Sp)	2008	23/A	FIB, F, I1	F	-	90	1.7 × 10 <sup>-2</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C1537 (H, Sp)	2008	405/D	FIA, F, I1	I1	215	110	7.4 × 10 <sup>-4</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C1538 (H, Sp)	2008	405/D	FIA, F, I1	I1	215	110	6.9 × 10 <sup>-4</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
C353 (L, Sp)	2003	155/B1	F, I1	I1	178	110	8.7 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	-
C515 (L, Sp)	2003	1564/A	A/C, FIB, F, I1	I1	3 (3)	100	8.3 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	-
C508 (L, Sp)	2003	2001/D	FIB, F, I1	I1	3 (3)	105	2.0 × 10 <sup>-3</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	-
C526 (L, Sp)	2003	362/D	FIB, F, I1, Y	I1	26 (26)	110	1.5 × 10 <sup>-2</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	qacG- <i>aadA6</i> - <i>qacG</i>	tet(A)
C537 (L, Sp)	2003	616/B1	FIA, FIB, F, I1, Y	I1	26 (26)	105	2.0 × 10 <sup>-2</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	-/-	-	tet(A)
101689 (L, Ge)	2010	117/D	FIB, F, I1	X3	-	45	4.4 × 10 <sup>-6</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>ΔdeoR</i>	-/-	-	qnrS1
101908 (L, Ge)	2010	23/A	FIB, F, I1	NT <sup>e</sup>	-	30	NC <sup>f</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	-
101985 (L, Ge)	2010	23/A	FIB, F, I1	I1	26 (26)	110	1.2 × 10 <sup>-7</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)
111918 (L, Ge)	2011	371/D	FIB, F, I1	I1	95	120	1.0 × 10 <sup>-4</sup>	IS26- <i>bla</i> <sub>SHV-12</sub> - <i>deoR</i>	+/-	estX- <i>psp</i> - <i>aadA2</i> - <i>cmiA1</i> - <i>aadA1</i> - <i>qacI</i>	tet(A)

<sup>a</sup>W, wild bird; M, chicken meat; D, dog; H, human; L, livestock bird (poultry).

<sup>b</sup>Sp, Spain; Ge, Germany.

<sup>c-d</sup>These plasmids show indistinguishable patterns after EcoRI, HindIII or BamHI digestion.

<sup>e</sup>NT, non-typeable.

<sup>f</sup>NC, non-conjugative (under the tested conditions, please see the Materials and methods section).

Among the 17 IncI1 *bla*<sub>SHV-12</sub>-carrying plasmids, 3 of them showed novel plasmid STs (pSTs): 1 plasmid carried a transversion in *ardA* (pST214) and 2 other plasmids a novel allele combination (pST215) (Table 2). The three *E. coli* isolates from wild birds belonging to ST453/B1 carried closely related IncI1 plasmids (indistinguishable restriction patterns) (Figure S2). The IncI1/pST26 plasmids harboured by ST23/A isolates from dogs ( $n = 2$ ) and the IncI1/pST215 plasmids carried by ST405/D isolates from humans ( $n = 2$ ) had the same size and resistance genotype, and showed related EcoRI, HindIII and BamHI restriction patterns.

IncK plasmids were found in isolates from different sources (ST57/D, chicken meat; ST131/B2, chicken meat; ST57/B1, dog), but showed equal sizes and EcoRI, HindIII or BamHI restriction patterns (Figure S2). These plasmids carried no additional resistance genes or integrons.

Conjugal transfer of the ESBL phenotype was demonstrated in all isolates except one (101908). IncK plasmids exhibited lower conjugation frequencies than IncI1 plasmids ( $10^{-5}$ – $10^{-4}$  versus  $10^{-4}$ – $10^{-2}$ ) (Table 2).

### Co-located resistance genes and integrons

All original isolates of the studied collection showed multidrug-resistance phenotypes (resistance to antimicrobials of  $\geq 3$  different classes), except one (solely resistant to  $\beta$ -lactams and nalidixic acid) (Table 1). The German isolate 101689 (chicken, 2010) was the only one carrying a plasmid-mediated quinolone-resistance gene, specifically *qnrS1*. This gene was co-located with *bla*<sub>SHV-12</sub> on a 45 kb IncX3 plasmid. Although different genes encoding resistance to tetracycline [*tet(A)*, *tet(B)*] and sulphonamides (*sul1*, *sul2*, *sul3*) were identified among the original isolates, we only found *tet(A)* and *sul3* genes (*sul1* in one isolate harbouring a plasmid-borne class 1 integron) among *bla*<sub>SHV-12</sub>-carrying transformants (Table 2). Overall, four resistance profiles were identified among *bla*<sub>SHV-12</sub>-carrying transformants:  $\beta$ -lactams/tetracycline/chloramphenicol/streptomycin (10 of 23),  $\beta$ -lactams/chloramphenicol/streptomycin (5 of 23),  $\beta$ -lactams/tetracycline (2 of 23) and exclusively  $\beta$ -lactams (6 of 23).

Regarding integrons, 19 of 23 isolates carried class 1 and/or 2 integrons. Class 2 integrons were present in four ST57 isolates. Two different gene cassette (GC) arrays were detected: *dfrA1-sat2-aadA1* ( $n = 3$ ) and a recently described structure (IS10-*dfrA1-sat2-aadA1*).<sup>33</sup> Among *intI1*-positive isolates, three carried classic class 1 integrons containing different GC arrangements: *aadA1* ( $n = 1$ ), *dfrA1-aadA1* ( $n = 1$ ) and *qacG-aadA6-qacG* ( $n = 1$ ). The latter integron, reported as In812 in the INTEGRALL database, was co-located on the same plasmid as the *bla*<sub>SHV-12</sub> gene and was first reported in Enterobacteriaceae in this study. The 59-base element (*attC*) of the GC *aadA6* was truncated by the insertion of the second *qacG* gene. The coding region of both *qacG* cassettes was identical. Thirteen isolates and their transformants harboured a class 1 integron lacking the 3'-CS and containing a large array of GCs (*estX-psp-aadA2-cmlA1-aadA1-qacI*) (Table 1).

### *bla*<sub>SHV-12</sub> genetic environment

Regarding the *bla*<sub>SHV-12</sub> flanking regions, IS26 was located 73 bp upstream and the putative *deoR* transcriptional regulator gene 20 bp downstream of the *bla*<sub>SHV-12</sub> gene.

In isolate 101689, the *deoR* gene was truncated at position 698 (reverse direction) by the insertion of a 445 bp DNA segment preceding an IS26 element. This fragment contained two ORFs, encoding a hypothetical protein and a putative ArsR family transcriptional regulator. The 17 nucleotides located at the 3'-end of this putative *arsR* gene overlapped with the IS26 left inverted repeat (IRL) found downstream of the 445 bp segment (Figure S3).

### Characteristics of the sequenced IncI1 plasmid pCAZ590


The completely sequenced plasmid pCAZ590 comprised 117 387 bp and displayed an average G + C content of 51.7% (Figure 1a). Replication, transfer and leading regions were highly similar to other IncI1 plasmids, with some insertions/deletions suggesting recombination between related plasmids. The entire region involved in conjugal transfer (*tra/trb* genes) was closely related (99.0% identity) to that of the archetypal IncI1 plasmid R64 (accession no. AP005147). Larger portions of the backbone shared high identity (99.0%) with plasmids PDM04 (NZ\_CP013224.1), pSH1148\_107 (NC\_019123.1) and pSD107 (NC\_019137.1) from different *Salmonella enterica* strains.

Plasmid pCAZ590 presented a large accessory module (26 728 bp) associated with antimicrobial resistance, located between the replication and the ColIb colicin immunity regions. This resistance module comprised a Tn21-derived transposon in which an atypical class 1 integron, the *bla*<sub>SHV-12</sub> gene and flanking elements (IS26-*deoR*) and a  $\Delta$ Tn1721 transposon were inserted. It is located in pCAZ590 in the antisense orientation, but it is shown in (Figure 1a) and described in the text in the sense orientation to facilitate comparisons.

The Tn21-derived region carried the left and right Tn21 terminal IRs, the genes involving its own transposition (*tnpA*, *tnpR*, *tnpM*), the terminal imperfect IRi of class 1 integron In2 and the class 1 integrase *intI1* gene. However, almost the whole structure of integron In2 was missing, solely a fragment of the *tniA* gene (615 bp) was identified. Instead of In2, an atypical integron was found, whose arrangement included the standard 5'-CS (*intI1* gene), the GC array *estX-psp-aadA2-cmlA1-aadA1-qacI* and the genetic platform IS440-*sul3-yqkA-yusZ- $\Delta$ mef(B)-IS26*. The *mef(B)* gene, which encodes a macrolide-efflux protein, was found disrupted by the IS26. This atypical class 1 integron was detected in most of the isolates of our collection. The segment IS26-*bla*<sub>SHV-12</sub>-*deoR* was followed by a  $\Delta$ Tn1721, encoding resistance to tetracycline. The  $\Delta$ Tn1721 contained the two characteristic regions of Tn1721, the first corresponding to the genes involved in the production of a putative chemotaxis protein (*orf1*) and transposition (*tnpR*, *tnpA*) was complete. However, the second region contained the tetracycline transcriptional regulator and resistance genes [*tetR*, *tet(A)*] and a *pecM*-like gene, but the truncated transposase ( *$\Delta$ tnpA*) and the terminal inverted repeat IRRII were missing. From the mercury resistance module (*merRTPCAD*), typically located in Tn21, only a fragment of *merR* (120 bp) was found downstream of the  $\Delta$ Tn1721- *$\Delta$ tniA* (Figure 1b).

### Discussion

This study focused on the investigation of the *bla*<sub>SHV-12</sub> gene, which codes for one of the most prevalent ESBLs. The *bla*<sub>SHV-12</sub>-positive isolates were collected over different periods of time and from



**Figure 1.** (a) Circular map of plasmid pCAZ590 (accession number LT669764) and (b) linear illustration of the complex multidrug-resistance region of pCAZ590 and a comparative analysis of this region, Tn21 (accession number AF071413) and Tn1721 (accession number X61367). Some relevant genes are labelled. (a) Second inner ring shows the fragments of truncated genes and the forward and reverse coding sequences are shown in the third and fourth inner rings, respectively. They are shown as arrows (the direction of transcription is indicated by the arrowheads) and are coloured according to their function as shown in the legend. Names of functional regions are shown in the outer ring. First inner ring shows a plot of the GC skew (yellow, above average; purple, below average). To facilitate comparisons, in (b) the sequence is shown according to the orientation described for Tn21 and Tn1721, although in pCAZ590 it is found in the opposite orientation. Coding reading frames are shown as arrows (the direction of transcription is indicated by the arrowheads) and are coloured as described in (a). ISs are presented as boxes and the arrows within the boxes indicate the transposition genes. Vertical lines represent the IRs of ISs, transposons or integron In2. This figure appears in colour in the online version of JAC and in black and white in the print version of JAC.

different sources and geographical areas. The repeated occurrence of distinct clones in different sources/years of isolation suggests the presence of potentially 'epidemic' clones with relatively high stability over time, including the closely related resistance plasmids carried by them.

Some of the most common clones (ST23/A, ST57/D) detected in this study have been associated with ESBL phenotypes and are widely spread among different environments, including clinical settings.<sup>34</sup> A member of the epidemic ST131/B2 clone, associated with CTX-M production, was detected in one chicken meat sample harbouring both *bla*<sub>SHV-12</sub> and *bla*<sub>CTX-M-1</sub> genes. According to the high prevalence of the SHV-12 enzyme among non-ST131 *E. coli* in Spain, the occurrence of a horizontal *bla*<sub>SHV-12</sub> transfer event from local non-ST131 isolates to the epidemic ST131 clone has been suggested.<sup>34</sup> This hypothesis is supported by our findings, which confirm a high similarity between IncK plasmids detected in isolates belonging to ST131 and other STs, reinforcing the possibility of a horizontal transfer of this *bla*<sub>SHV-12</sub>-carrying plasmid to the ST131 lineage.

The *bla*<sub>SHV-12</sub> gene was mainly associated with IncI1 plasmids. These plasmids seemed to be easily transferred by conjugation with high efficiency, which may explain their predominance in different ecosystems.<sup>22,35</sup> Although a considerable diversity was found, IncI1/pST3 and IncI1/pST26 appeared to be the dominant subtypes. In agreement with previous reports,<sup>22,35</sup> all the IncI1/pST3 plasmids harbouring *bla*<sub>SHV-12</sub> were detected in poultry or poultry-derived meat suggesting a potential association between this variant and poultry. Conversely, *bla*<sub>SHV-12</sub>-carrying IncI1/pST26 plasmids and related subtypes belonging to clonal complex (CC) 26 (like pST29), were associated with a wide host range contributing to the spread of SHV-12-encoding genes among different environments and geographical areas. In fact, pST29 detected in wild bird isolates (2014) and the novel pST215, identified in two commensal *E. coli* isolates from humans (2008), may reflect a possible diversifying evolutionary process.

It is also remarkable that, in contrast to IncK-positive transformants, all susceptible to non-β-lactam antimicrobials, 14 *bla*<sub>SHV-12</sub>-carrying IncI1 plasmids and 1 non-typeable plasmid showed a multidrug-resistance phenotype associated (i) with the presence of the same atypical class 1 integron containing *aadA2-cmlA1-aadA1* GCs, and (ii) frequently, with the co-location of the *tet(A)* gene. In particular, the structure of this atypical integron (*intI1-estX-psp-aadA2-cmlA1-aadA1-qacI-IS440-sul3*) appears to be identical to that found by other authors on *bla*<sub>SHV-12</sub>-carrying IncI1 plasmids. It is usually embedded in Tn21-derived transposons and is globally distributed among Enterobacteriaceae from different environments.<sup>36</sup>

As some authors have suggested and as confirmed by the pCAZ590 plasmid sequence, the high prevalence of this atypical integron seems to be associated with its downstream linkage to IS26, which constitutes the highly conserved upstream flanking region of the *bla*<sub>SHV-12</sub> gene (Figure 1b).<sup>36,37</sup> The presence of these genetic platforms plays an important role in the persistence of SHV-12-producing isolates due to their capability to promote the selection of these ESBL genes under the selective pressure imposed even by antimicrobial agents other than β-lactams.

Regarding genetic environments, it is noteworthy that 22 of 23 isolates showed an identical genetic structure flanking the *bla*<sub>SHV-12</sub> gene. However, the novel described downstream

environment revealed the truncation of the putative *deoR* transcriptional regulator gene by a genetic structure containing two ORFs preceding an IS26. Such a genetic structure has been shown to be truncating other genes in different regions of many plasmids [e.g. pYD626E (KJ933392) and pSRC119-A/C (KM670336)], revealing its high mobility potential. This may be due to the ability of IS26 to mobilize neighbouring genes by misidentifying short sequences as its alternative left-hand IR. The insertion of this genetic structure may have important implications due to the putative composite transposon formed, which could facilitate the exchange *en-bloc* of the ESBL gene (Figure S3).

As a final remark, the first described Portuguese *E. coli* isolate carrying an IS10 within a class 2 integron<sup>33</sup> and the one identified in this study belonged in both cases to ST57, suggesting that this specific class 2 integron may be clonally disseminated.

Although the present study provides important insights into the understanding of the dynamics and the molecular background of *bla*<sub>SHV-12</sub>-carrying *E. coli* isolates, future studies, using larger numbers of isolates, are needed to identify other potential epidemic clones/plasmids. Moreover, the long sampling period (2003–14) may represent a drawback due to the rapid evolution of ESBL genes. However, based on our findings, it seems unlikely that the molecular background of *bla*<sub>SHV-12</sub>-carrying clones/plasmids has changed dramatically over the sampling period.

Overall, this study revealed that some SHV-12-producing *E. coli* isolates from different sources showed identical ST/PFGE profiles or carried highly similar plasmids. These observations suggest that both clonal and plasmid transfer facilitate the spreading of *bla*<sub>SHV-12</sub> ESBL genes. Horizontal dissemination was mainly driven by IncI1 plasmids showing rather conserved co-located resistance genes.

## Acknowledgements

We thank Roswitha Becker, Ute Beermann and Regina Ronge for excellent technical assistance.

## Funding

The work performed in the University of La Rioja was partially supported by project SAF2016-76571-R from the Ministerio de Economía y Competitividad (MINECO) of Spain and Fondo Europeo de Desarrollo Regional (FEDER) and also by project 2016/00042/001 of the University of La Rioja, Spain. The work conducted in the Friedrich-Loeffler-Institut was financially supported by the German Federal Ministry of Education and Research (BMBF) (grant no. 01KI1313D-RESET II). The work conducted by J. L. and Y. W. was supported by the National Basic Research Program of China (2013CB127200). C. A. A. has a pre-doctoral fellowship from the MINECO of Spain (BES-2013-063105).

## Transparency declarations

None to declare.

## Supplementary data

Figures S1 to S3 are available as Supplementary data at JAC Online.

## References

- 1 WHO. *Critically Important Antimicrobials for Human Medicine: 3rd Revision*. 2011. [http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf).
- 2 Ben Said L, Jouini A, Klibi N et al. Detection of extended-spectrum  $\beta$ -lactamase (ESBL)-producing Enterobacteriaceae in vegetables, soil and water of the farm environment in Tunisia. *Int J Food Microbiol* 2015; **203**: 86–92.
- 3 Wieler LH, Ewers C, Guenther S et al. Methicillin-resistant staphylococci (MRS) and extended spectrum- $\beta$ -lactamases (ESBL)-producing Enterobacteriaceae in companion animals: nosocomial infections as one reason for the rising prevalence of these potential zoonotic pathogens in clinical samples. *Int J Med Microbiol* 2011; **301**: 635–41.
- 4 Pouget JG, Couthino FJ, Reid-Smith RJ et al. Characterization of *bla*<sub>SHV</sub> genes on plasmids from *Escherichia coli* and *Salmonella enterica* isolates from Canadian Food Animals (2006–2007). *Appl Environ Microbiol* 2013; **79**: 2864–6.
- 5 Michael GB, Freitag C, Feßler AT et al. Emerging issues in antimicrobial resistance of bacteria from food-producing animals. *Future Microbiol* 2015; **10**: 427–43.
- 6 Alcalá L, Alonso CA, Simón C et al. Wild birds, frequent carriers of extended-spectrum  $\beta$ -lactamase (ESBL) producing *Escherichia coli* of CTX-M and SHV-12 types. *Microb Ecol* 2016; **72**: 861–9.
- 7 Perilli M, Dell'Amico E, Segatore B et al. Molecular characterization of extended-spectrum  $\beta$ -lactamases produced by nosocomial isolates of Enterobacteriaceae from an Italian nationwide survey. *J Clin Microbiol* 2002; **40**: 611–4.
- 8 Yagi T, Kurokawa H, Shibata N et al. A preliminary survey of extended-spectrum  $\beta$ -lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* and *Escherichia coli* in Japan. *FEMS Microbiol Lett* 2000; **184**: 53–6.
- 9 Díaz MA, Hernández-Bello JR, Rodríguez-Baño J et al. Diversity of *Escherichia coli* strains producing extended-spectrum  $\beta$ -lactamases in Spain: second nationwide study. *J Clin Microbiol* 2010; **48**: 2840–5.
- 10 Briñas L, Moreno MA, Teshager T et al. Monitoring and characterization of extended-spectrum  $\beta$ -lactamases in *Escherichia coli* strains from healthy and sick animals in Spain in 2003. *Antimicrob Agents Chemother* 2003; **49**: 1262–4.
- 11 Kola A, Kohler C, Pfeifer Y et al. High prevalence of extended-spectrum  $\beta$ -lactamase-producing Enterobacteriaceae in organic and conventional retail chicken meat, Germany. *J Antimicrob Chemother* 2012; **67**: 2631–4.
- 12 Egea P, López-Cerero L, Gómez-Sánchez MC et al. Increased raw poultry meat colonization by extended spectrum  $\beta$ -lactamase-producing *Escherichia coli* in the south of Spain. *Int J Food Microbiol* 2012; **159**: 69–73.
- 13 Schink AK, Kadlec K, Schwarz S. Analysis of *bla*<sub>CTX-M</sub>-carrying plasmids from *Escherichia coli* isolates collected in the BfT-Germ Vet study. *Appl Environ Microbiol* 2011; **77**: 7142–6.
- 14 Dolejska M, Villa L, Hasman H et al. Characterization of IncN plasmids carrying *bla*<sub>CTX-M-1</sub> and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. *J Antimicrob Chemother* 2013; **68**: 333–9.
- 15 Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing: Twenty-fifth Informational Supplement M100-S25*. CLSI, Wayne, PA, USA, 2015.
- 16 Jouini A, Vinué L, Ben Slama K et al. Characterization of CTX-M and SHV extended-spectrum  $\beta$ -lactamases and associated resistance genes in *Escherichia coli* strains of food samples in Tunisia. *J Antimicrob Chemother* 2007; **60**: 1137–41.
- 17 Clermont O, Bonacorsi S, Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 2000; **66**: 4555–8.
- 18 Arbeit RD, Arthur M, Dum RD et al. Resolution of recent evolutionary divergence among *Escherichia coli* from related lineages: the application of pulsed field gel electrophoresis to molecular epidemiology. *J Infect Dis* 1990; **161**: 230–5.
- 19 Heras J, Domínguez C, Mata E et al. GeJ—a tool for analyzing DNA fingerprint gel images. *BMC Bioinformatics* 2015; **16**: 270.
- 20 Kadlec K, Schwarz S. Analysis and distribution of class 1 and class 2 integrons and associated gene cassettes among *Escherichia coli* isolates from swine, horses, cats and dogs collected in the BfT-GermVet monitoring study. *J Antimicrob Chemother* 2008; **62**: 469–73.
- 21 Carattoli A, Bertini A, Villa L et al. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005; **63**: 219–28.
- 22 García-Fernández A, Chiaretto G, Bertini A et al. Multilocus sequence typing of Inc11 plasmids carrying extended-spectrum  $\beta$ -lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. *J Antimicrob Chemother* 2008; **61**: 1229–33.
- 23 Vinué L, Sáenz Y, Martínez S et al. Prevalence and diversity of extended-spectrum  $\beta$ -lactamases in faecal *Escherichia coli* isolates from healthy humans in Spain. *Clin Microbiol Infect* 2008; **15**: 954–7.
- 24 Ruiz E, Sáenz Y, Zarazaga M et al. *qnr*, *aac(6′)-Ib-cr* and *qepA* genes in *Escherichia coli* and *Klebsiella* spp.: genetic environments and plasmid and chromosomal location. *J Antimicrob Chemother* 2012; **67**: 886–97.
- 25 Kadlec K, Kehrenberg C, Schwarz S. Molecular basis of resistance to trimethoprim, chloramphenicol and sulphonamides in *Bordetella bronchiseptica*. *J Antimicrob Chemother* 2005; **56**: 485–90.
- 26 Sáenz Y, Vinué L, Ruiz E et al. Class 1 integrons lacking *qacEA1* and *sul1* genes in *Escherichia coli* isolates of food, animal and human origins. *Vet Microbiol* 2010; **144**: 493–7.
- 27 Pitout JD, Thomson KS, Hanson ND et al. Plasmid-mediated resistance to expanded-spectrum cephalosporins among *Enterobacter aerogenes* strains. *Antimicrob Agents Chemother* 1998; **42**: 596–600.
- 28 Diestra K, Juan C, Curiao T et al. Characterization of plasmids encoding *bla*<sub>ESBL</sub> and surrounding genes in Spanish clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae*. *J Antimicrob Chemother* 2009; **63**: 60–6.
- 29 Aziz RK, Bartels D, Best AA et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 2008; **9**: 75.
- 30 Rutherford K, Parkhill J, Crook J et al. Artemis: sequence visualization and annotation. *Bioinformatics* 2000; **16**: 944–5.
- 31 Carver T, Thomson N, Bleasby A et al. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* 2009; **25**: 119–20.
- 32 Carattoli A, Zankari E, García-Fernández A et al. PlasmidFinder and pMLST: *in silico* detection and typing of plasmids. *Antimicrob Agents Chemother* 2014; **58**: 3895–903.
- 33 Jones-Dias D, Manageiro V, Martins AP et al. New class 2 integron In2-4 among Inc11-positive *Escherichia coli* isolates carrying ESBL and PMA $\beta$  genes from food animals in Portugal. *Foodborne Pathog Dis* 2016; **13**: 36–9.
- 34 Oteo J, Cercenado E, Fernández-Romero S et al. Extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* as a cause of pediatric infections: report of a neonatal intensive care unit outbreak due to a CTX-M-14-producing strain. *Antimicrob Agents Chemother* 2012; **56**: 54–8.
- 35 Accogli M, Fortini D, Giufrè M et al. Inc11 plasmids associated with the spread of CMY-2, CTX-M-1 and SHV-12 in *Escherichia coli* of animal and human origin. *Clin Microbiol Infect* 2013; **19**: E238–40.
- 36 Curiao T, Cantón R, Garcillán-Barcia MP et al. Association of composite IS26-*sul3* elements with highly transmissible Inc11 plasmids in extended-spectrum  $\beta$ -lactamase producing *Escherichia coli* clones from humans. *Antimicrob Agents Chemother* 2011; **55**: 2451–7.
- 37 Kiiru J, Butaye P, Goddeeris BM et al. Analysis for prevalence and physical linkages amongst integrons, *ISEcp1*, *ISCR1*, Tn21 and Tn7 encountered in *Escherichia coli* strains from hospitalized and non-hospitalized patients in Kenya during a 19-year period (1992–2011). *BMC Microbiol* 2013; **13**: 109.