

231465

UCRL JC 127249
UCRL-JC-
PREPRINT


Analysis of Thomsen Parameters for Finely Layered VTI Media

James G. Berryman
Patricia A. Berge
Lawrence Livermore National Laboratory

Vladimir Grechka
Colorado School of Mines

This paper was prepared for submittal to the
Society of Exploration Geophysicists
Dallas, Texas
November 2-7, 1997

April 1997



This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Analysis of Thomsen parameters for finely layered VTI media

James G. Berryman, Lawrence Livermore Lab; Vladimir Grechka, Colorado School of Mines; and Patricia A. Berge, Lawrence Livermore Lab

SUMMARY

The range of Thomsen's anisotropy parameters ϵ and δ for vertical transversely isotropic (VTI) media when the anisotropy is due to fine layering of isotropic elastic materials is considered. We show that ϵ lies in the range $-3/8 \leq \epsilon \leq \frac{1}{2} [\langle v_p^2 \rangle \langle v_p^{-2} \rangle - 1]$, for finely layered media having constant density; smaller positive and all negative values of ϵ occur for media with large fluctuations in the Lamé parameter λ . We show that $\text{sign}(\delta) = \text{sign}(\langle v_p^{-2} \rangle - \langle v_s^{-2} \rangle \langle v_s^2/v_p^2 \rangle)$ for constant density media, so δ can be either positive or negative. Among all theoretically possible random media, positive and negative δ are equally likely in finely layered media limited to two types of constituent layers. Layered media having large fluctuations in Lamé λ are the ones most likely to have positive δ . Since Gassmann's results for fluid-saturated porous media show that the effects of fluids influence only the λ Lamé constant, not the shear modulus μ , these results suggest that positive δ occurring together with positive but small ϵ may be indicative of changing fluid content in layered earth.

INTRODUCTION

Two primary goals of seismic reflection processing are: (1) to image geologic structure and (2) to provide information about lithology for interpretation. The process used to achieve the second goal is made complex by the fact that the same seismic velocity may result from several different combinations/mixtures of materials in the earth, *i.e.*, the possible causes of the observed behavior are often nonunique. It is therefore necessary to explore the possible range of seismic velocities that can occur within the set of circumstances deemed most likely to occur in the earth at the site of interest.

Fine horizontal layering (*i.e.*, layers with thickness small compared to the wavelength of the seismic wave) is known to result in vertical transverse isotropy (VTI) - wherein wave speeds vary with angle in such media, but are uniquely determined by the angle from the vertical. There has continued to be some doubt about the range of anisotropy parameters possible in such media. Here we will correct some common errors found in the literature. We show that Thomsen's parameter ϵ can be negative and actually has a greater negative range than indicated in previous published work. We also show that Thomsen's parameter δ can be positive in finely layered media (contrary to some erroneous claims that have appeared in the literature), and furthermore that regions having both positive δ and posi-

tive but small ϵ are likely to be regions of rapid spatial variation in fluid content. We use Monte Carlo simulations to establish the existence of both positive δ and negative ϵ , and analysis of Backus averaging formulas to clarify when such behavior occurs.

ANISOTROPIC ELASTIC MEDIA

For an isotropic elastic medium, the stiffness tensor has the form

$$c_{ijkl} = \lambda \delta_{ij} \delta_{kl} + \mu (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}), \quad (1)$$

depending on only two parameters (the Lamé constants, λ and μ), this tensor can have up to 21 independent constants for general anisotropic elastic media. The stiffness tensor has pairwise symmetry in its indices such that $c_{ijkl} = c_{jikl}$ and $c_{ijkl} = c_{ijlk}$, which will be used later to simplify the resulting equations.

A commonly used simplification of the notation for elastic analysis is given by introducing the strain tensor, where $e_{ij} = \frac{1}{2}(u_{i,j} + u_{j,i})$. Then, using one version of the Voigt convention, in which the pairwise symmetries of the stiffness tensor indices are used to reduce the number of indices from 4 to 2 using the rules $11 \rightarrow 1$, $22 \rightarrow 2$, $33 \rightarrow 3$, 23 or $32 \rightarrow 4$, 13 or $31 \rightarrow 5$, and 12 or $21 \rightarrow 6$, we replace $\sigma_{ij} = c_{ijkl} e_{kl}$ by the usual 6×6 system. For VTI materials, $c_{11} = c_{22} \equiv a$, $c_{12} \equiv b$, $c_{13} = c_{23} \equiv f$, $c_{33} \equiv c$, $c_{44} = c_{55} \equiv l$, and $c_{66} \equiv m$. There is also one further constraint on the constants that $a = b + 2m$, following from rotational symmetry in the $x_1 x_2$ -plane.

BACKUS AVERAGING

Backus [1962] presents an elegant method of producing the effective constants for a thinly layered medium composed of either isotropic or anisotropic elastic layers. This method applies either to spatially periodic layering or to random layering, by which we mean either that the material constants change in a nonperiodic (unpredictable) manner from layer to layer or that the layer thicknesses may also be random. For simplicity, we will assume that the layers are isotropic.

The results are

$$a = \left\langle \frac{\lambda}{\lambda + 2\mu} \right\rangle^2 \left\langle \frac{1}{\lambda + 2\mu} \right\rangle^{-1} + 4 \left\langle \frac{\mu(\lambda + \mu)}{\lambda + 2\mu} \right\rangle, \quad (2)$$

$$c = \left\langle \frac{1}{\lambda + 2\mu} \right\rangle^{-1} \quad (3)$$

$$f = \left\langle \frac{\lambda}{\lambda + 2\mu} \right\rangle \left\langle \frac{1}{\lambda + 2\mu} \right\rangle^{-1}, \quad (4)$$

Thomsen parameters for layered VTI media

$$l = \left\langle \frac{1}{\mu} \right\rangle^{-1} \quad (5)$$

and

$$m = \langle \mu \rangle. \quad (6)$$

One very important fact that is known about these equations is that they reduce to isotropic results with $a = c$, $b = f$, and $l = m$, if the shear modulus is a constant, regardless of the behavior of λ .

THOMSEN PARAMETER γ

The two shear moduli must satisfy

$$1 \leq \langle \mu \rangle \left\langle \frac{1}{\mu} \right\rangle = \frac{m}{l}, \quad (7)$$

because the well-known Cauchy-Schwartz inequality $\langle \alpha\beta \rangle^2 \leq \langle \alpha^2 \rangle \langle \beta^2 \rangle$ gives this result when $\alpha = \mu^{1/2}$ and $\beta = 1/\mu^{1/2}$. Equality applies in the Cauchy-Schwartz inequalities only when $\alpha = \text{const} \times \beta$, which implies in the present circumstances that μ must be constant for $l = m$. But this is precisely the condition mentioned earlier for the layer equations to be isotropic, so we generally exclude this case from consideration. The implication of (7) for Thomsen's γ is that

$$\gamma = \frac{m-l}{2l} \geq 0. \quad (8)$$

THOMSEN PARAMETER ϵ

An important anisotropy parameter for qP-waves is Thomsen's parameter ϵ , defined by

$$\epsilon = \frac{a-c}{2c}. \quad (9)$$

Formula (2) for a may be rewritten to show that

$$2\epsilon = \left[\langle \lambda + 2\mu \rangle \left\langle \frac{1}{\lambda + 2\mu} \right\rangle - 1 \right] - \left[\left\langle \frac{\lambda^2}{\lambda + 2\mu} \right\rangle \left\langle \frac{1}{\lambda + 2\mu} \right\rangle - \left\langle \frac{\lambda}{\lambda + 2\mu} \right\rangle^2 \right], \quad (10)$$

We find that the first bracket on the right hand side is again in Cauchy-Schwartz form showing that it always makes a positive contribution unless $\lambda + 2\mu = \text{constant}$, in which case it vanishes. Similarly, the second term always makes a negative contribution unless $\lambda = \text{constant}$, in which case it vanishes.

Fluctuations in λ in the earth have important implications for oil and gas exploration. Gassmann's well-known results [Gassmann, 1951] show that, when isotropic porous elastic media are saturated with any fluid, the fluid has no mechanical effect on the shear modulus μ , but can have a significant effect on the bulk modulus $K = \lambda + \frac{2}{3}\mu$, and therefore on λ . Thus, observed variations in μ have no direct information

about fluid content, while observed variations in λ , especially if they are large variations, may contain important clues about variations in fluid content. So the observed structure of ϵ in (10) strongly suggests that small positive and all negative values of ϵ may be important indicators of significant fluctuations in fluid content.

If the finely layered medium is composed of only two distinct types of isotropic elastic materials and they appear in the layering sequence with equal spatial frequency, then we find that

$$2\epsilon = (\mu_2 - \mu_1) \frac{(\lambda_2 - \lambda_1) + (\mu_2 - \mu_1)}{(\lambda_1 + 2\mu_1)(\lambda_2 + 2\mu_2)}. \quad (11)$$

This result agrees with Postma [1955] except for an obvious typographical error in the denominator of his published formula. This formula shows clearly that if $\mu_1 = \mu_2$ then Thomsen parameter ϵ is identically equal to zero as expected. Also, if $\mu_1 \neq \mu_2$ but $\lambda_1 = \lambda_2$, then (11) implies $\epsilon \geq 0$, as we inferred from (10).

Now, we can use this formula to deduce the smallest possible value of the right hand side of (11). The shear moduli must not be equal, so without loss of generality we suppose that $\mu_2 > \mu_1$. Then, the numerator is seen to become negative by taking λ_2 towards negative values and $\lambda_1 \rightarrow +\infty$. The smallest value λ_2 can take is determined by the bulk modulus bound $\lambda_2 + \frac{2}{3}\mu_2 \geq 0$. So we may set $\lambda_2 = -\frac{2}{3}\mu_2$ in both the numerator and denominator. This choice also makes the factor $\lambda_2 + 2\mu_2 = \frac{4}{3}\mu_2$ as small as possible in the denominator, thus helping to magnify the effect of the negative numerator as much as possible. The result so far is that

$$2\epsilon = \frac{3}{4} \left(\frac{\mu_2 - \mu_1}{\mu_2} \right) \left(\frac{-\lambda_1 + \mu_2/3 - \mu_1}{\lambda_1 + 2\mu_1} \right) \quad (12)$$

The parameter λ_1 may vary from $-\frac{2}{3}\mu_1$ to plus infinity. At $\lambda_1 = -\frac{2}{3}\mu_1$, the second expression in parentheses is positive; but, this expression is also a monotonically decreasing function of λ_1 and approaches -1 as $\lambda_1 \rightarrow +\infty$. Thus, the smallest value of Thomsen's parameter ϵ is given by

$$\epsilon = -\frac{3}{8} \frac{\mu_2 - \mu_1}{\mu_2} \geq -\frac{3}{8}. \quad (13)$$

This result differs by a factor of 2 from the corresponding result of Postma [1955], which was obtained improperly by allowing three of the four elastic constants to vanish and also using a physically motivated but unnecessary restriction that both λ_1 and λ_2 must be nonnegative.

THOMSEN PARAMETER δ

Thomsen's parameter δ is defined by

$$\delta = \frac{(f+l)^2 - (c-l)^2}{2c(c-l)}. \quad (14)$$

Thomsen parameters for layered VTI media

This parameter is considerably more difficult to analyze than either γ or ϵ for various reasons, some of which we will enumerate shortly.

Because of the controversy surrounding the sign of δ for finely layered media, we have performed a series of Monte Carlo simulations with the purpose of establishing the existence or nonexistence of layered models having positive δ . The simulations to be presented here were performed on models limited to two types of layers. Another set of simulations was performed having three types of layers, since it is known [Backus, 1962] that this is the most general case that needs to be considered, but we will not present that work here. The set of constant density models were chosen randomly by specifying the possible values of v_p to lie in some range $v_p^{min} \leq v_p \leq v_p^{max}$. The values chosen for the limiting wave speeds in the example shown in Figure 1 were $v_p^{min} = 2.5$ km/sec and $v_p^{max} = 5.5$ km/sec. The range of the shear wave velocity was similarly specified by constraining the ratio v_s/v_p to lie within a range $r_{min} \leq v_s/v_p \leq r_{max}$, where for the case being presented $r_{min} = 0.35$ and $r_{max} = 0.65$. The density was chosen to be $\rho = 2670$ kg/m³, but any choice of constant density would have resulted in the same dimensionless Thomsen parameters. For comparison, note that the density of quartz is about 2650 kg/m³ and the density of calcite is about 2720 kg/m³ [e.g., Wilkens *et al.*, 1984].

Figure 1 shows the Monte Carlo results for δ plotted versus η from 500 random models, where

$$\eta = \frac{\epsilon - \delta}{1 + 2\delta}. \quad (15)$$

The parameter η has been shown to be useful in seismic processing by Tsvankin and Thomsen [1994] and Alkhalifah and Tsvankin [1995]. It is particularly useful in the present context because it is known that for layered media this parameter will always be positive because $\epsilon - \delta$ will always be positive [Postma, 1955; Backus, 1962; Berryman, 1979]. We see that most models are clustered around the origin $(\eta, \delta) = (0, 0)$, but there is no question that a large fraction of the models have positive δ . For this particular method of generating the models, we find that about 25% of the two-layer models have positive δ .

Encouraged by the Monte Carlo simulation results, we have analyzed δ in light of the Backus formulas in order to determine whether it was possible to understand those circumstances in which $\delta > 0$ are most likely to occur. Using the formulas (2)-(6) derived by Backus [1962], it is not hard to show that one natural form of the expression for δ is

$$\delta = 2 \left\langle \frac{\lambda + \mu}{\lambda + 2\mu} \right\rangle \left\langle \frac{\lambda + \mu}{\mu(\lambda + 2\mu)} \right\rangle^{-1} \times$$




Figure 1: Crossplot of δ versus η for 500 random examples of layered media according to the algorithm in the text.

$$\left[\left\langle \frac{1}{\mu} \right\rangle \left\langle \frac{\lambda + \mu}{\lambda + 2\mu} \right\rangle - \left\langle \frac{\lambda + \mu}{\mu(\lambda + 2\mu)} \right\rangle \right]. \quad (16)$$

It is easy to show that the prefactors are all always positive. So the sign of δ is determined by the expression in brackets. Using simple algebraic manipulations, this expression can be rewritten in a number of different but useful forms, including

$$\begin{aligned} \left\langle \frac{1}{\mu} \right\rangle \left\langle \frac{\lambda + \mu}{\lambda + 2\mu} \right\rangle - \left\langle \frac{\lambda + \mu}{\mu(\lambda + 2\mu)} \right\rangle = \\ \left\langle \frac{1}{\lambda + 2\mu} \right\rangle - \left\langle \frac{1}{\mu} \right\rangle \left\langle \frac{\mu}{\lambda + 2\mu} \right\rangle. \end{aligned} \quad (17)$$

The manipulations required to arrive at such results will be described in more detail elsewhere. The important point about this sequence of inequalities is that each successive one can be used to prove something general about the sign of δ for layered media. In each case, we note that if some multiplicative factor is constant in all layers of the finely layered medium, then that factor may be removed from the averages. The second line shows that if $\lambda + 2\mu = \text{constant}$, then again δ is nonpositive. The very first expression may also be used in a slightly different way to arrive at a general result, $\delta = 0$ if Poisson's ratio is constant, for in that case the ratio λ/μ is also constant and so $(\lambda + \mu)/(\lambda + 2\mu)$ is constant.

It is straightforward to show for layered media having only two constituents that the last expression in (17) implies that δ will be positive if

$$\frac{\lambda_1}{\lambda_2} > \frac{\mu_1}{\mu_2} > 1, \quad (18)$$

Thomsen parameters for layered VTI media

or if $\frac{\lambda_1}{\lambda_2} < \frac{\mu_1}{\mu_2} < 1$, showing that the fluctuations in the Lamé constant λ must be greater than those in the shear modulus μ to observe positive δ .

As discussed earlier, v_s/v_p ratios are decreased by the presence of pores, flat cracks, or the addition of clay, dolomite, feldspar, or calcite to silicic rocks. Any of these factors can cause fluctuations in λ without greatly changing μ .

For layered media with constant density, the final expression in (17) shows that

$$\text{sign}(\delta) = \text{sign}(\langle v_p^{-2} \rangle - \langle v_s^{-2} \rangle \langle v_s^2/v_p^2 \rangle). \quad (19)$$

THOMSEN PARAMETER BEHAVIOR

The results concerning the signs of the Thomsen parameters obtained in the preceding sections are summarized in Table 1.

TABLE 1. Behavior of anisotropy parameters as the layer material elastic parameters vary.

Thomsen ϵ, δ	$\lambda = \text{const}$ $\mu \neq \text{const}$	$\lambda + 2\mu = \text{const}$ $\mu \neq \text{const}$	$\nu = \text{const}$ $\lambda, \mu \neq \text{const}$
ϵ	≥ 0	≤ 0	≥ 0
δ	≤ 0	≤ 0	0

The nonnegativity of γ and $\epsilon - \delta$ for layered models are well-known. The fact that ϵ can be either positive or negative, and the circumstances leading to negative values has been little appreciated before. A quick glance at the Table seems to indicate that δ is either zero or negative and this is perhaps why there has been so much confusion about the possibility of $\delta > 0$ for layered models. We have shown that the correct inference about positive δ follows rather simply from the the result expressed in the last column of the Table. Since $\delta = 0$ when Poisson's ratio is constant or when $\mu = \text{constant}$, it is clear that we must have (for example) $\mu_1/\mu_2 > 1$ and $\lambda_1/\mu_1 \neq \lambda_2/\mu_2$ if δ is to be positive. The second expression can be rearranged to $\lambda_1/\lambda_2 \neq \mu_1/\mu_2 > 1$. Now we see that there are two possible cases, either $\lambda_1/\lambda_2 > \mu_1/\mu_2 > 1$ or $\lambda_1/\lambda_2 < \mu_1/\mu_2$. The first case results in positive δ , while the second case does not when $\mu_1/\mu_2 > 1$. This also shows that in perfectly random layered media one expects $\delta > 0$ to account for 50% of the models. In our simulation we found $\delta > 0$ accounted for only about 25% of the models produced, but this apparent discrepancy has been traced to a bias in the particular algorithm we used to generate the models. In any case, the earth does not have to obey perfectly random statistics and there is no reason to suppose that real layered earth will conform to these statistical considerations. Our main point is merely that $\delta > 0$ is entirely possible and quite understandable for layered

earth models, and so it is not at all surprising that $\delta > 0$ is often observed in real data examples.

DISCUSSION

The results obtained here show that Thomsen's parameter ϵ is smallest when the variation in the layer λ Lamé parameter is large, independent of the variation in the shear modulus μ . This result is therefore important for applications to porous layers containing pore fluids for which Gassmann's [1951] equation shows that the effects of fluids influence only the λ Lamé constants, not μ . Similarly, we find that δ is positive in finely layered media having large variations in λ . Thus, the regions of positive but small ϵ when occurring together with positive δ may be useful indicators of rapid spatial changes in fluid content in the layers. This result cannot, however, be considered in isolation, as there may be other effects in nonlayered media that produce similar values of the Thomsen parameters, and these should be explored and enumerated before any definitive statement can be made.

ACKNOWLEDGMENTS

We thank Leon Thomsen and Ilya Tsvankin for suggesting the problem. Work of JGB and PAB performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore National Laboratory under contract No. W-7405-ENG-48 as part of the Advanced Computational Technology Initiative (ACTI), funded through the DOE Office of Energy Research within the Office of Basic Energy Sciences, Division of Engineering and Geosciences.

REFERENCES

- Alkhalifah, T., and Tsvankin, I., 1995, Velocity analysis for transversely isotropic media: *Geophysics*, **60**, 1550-1566.
- Backus, G. E., 1962, Long-wave elastic anisotropy produced by horizontal layering: *J. Geophys. Res.*, **67**, 4427-4440.
- Berryman, J. G., 1979, Long-wave elastic anisotropy in transversely isotropic media: *Geophysics*, **44**, 896-917.
- Gassmann, F., 1951, Über die Elastizität poroser Medien: *Veierteljahrsschrift der Naturforschenden Gesellschaft in Zürich*, **96**, 1-23.
- Postma, G. W., 1955, Wave propagation in a stratified medium: *Geophysics*, **20**, 780-806.
- Thomsen, L., 1986, Weak elastic anisotropy: *Geophysics*, **51**, 1954-1966.
- Tsvankin, I., and Thomsen, L., 1994, Nonhyperbolic reflection moveout in anisotropic media: *Geophysics*, **59**, 1290-1304.
- Wilkens, R., Simmons, G., and Caruso, L., 1984, The ratio V_p/V_s as a discriminant of composition for siliceous limestones: *Geophysics*, **49**, 1850-1860.

Technical Information Department • Lawrence Livermore National Laboratory
University of California • Livermore, California 94551

