

ANALYSIS ON *Alu-I* GROWTH HORMONE (GH^{Alu-I}) GENE IN BALI CATTLE

Jakaria and R. R. Noor

Faculty of Animal Science, Bogor Agricultural University,
Darmaga Campus, Bogor 16680, West Java Indonesia
Corresponding E-mail: jakaria@ipb.ac.id

Received May 01, 2011; Accepted May 28, 2011

ABSTRACT

The research was conducted to identify *Alu-I* locus of growth hormone (GH) gene in Bali cattle by using 232 blood samples collected from Bali and Lombok islands. PCR-RFLP and sequencing methods were used to detect the polymorphism and nucleotide sequence at *Alu-I* locus of GH gene. The result showed that Bali cattle from Bali island has one genotype (LL genotype), whereas Bali cattle originating from Lombok island has two genotypes, namely LL and VV genotypes, respectively. The L and V allele frequencies from Bali and Lombok islands were 1.00 and 0.00; 0.99 and 0.01, respectively. The sequencing result of Bali cattle LL genotype showed an AGCT sequence of enzyme *Alu-I* restriction site. Based on polymorphic informative content (PIC) value, it can be concluded that *Alu-I* locus of Bali cattle from Bali and Lombok islands were monomorphic and polymorphic, respectively.

Keywords: Bali cattle, growth hormone gene, monomorphic, polymorphic

INTRODUCTION

The Bali cattle breed is one of the existing indigenous cattle breeds (Aceh, Pesisir, Madura, Java and Bali) in Indonesia. Although no official historical records exists, it is generally accepted that the Bali cattle is the domesticated direct descendant of the wild Banteng (*Bos sondaicus*, *Bos javanicus*, *Bos bantinger*) still surviving as an endangered species in three National Wild Reservation Parks (Ujung Kulon, Baluran and Blambangan) in Java (Martoyo 2003). As animal genetic resources, some molecular genetic studies have been reported in Indonesian cattle breeds including Bali cattle using microsatellite DNA (nuclear genome) and mitochondrial genome markers (Handiwirawan *et al.*, 2003; Nijman *et al.*, 2003; Abdullah, 2008; Ugglu, 2008; Mohammad *et al.*, 2009). However, molecular genetic marker based on coding sequence or candidate gene approach is limited and still need more in depth study of the existence of Indonesian cattle breeds especially Bali cattle.

The candidate gene approach is purposeful when a gene is known to function in such a way that it may explain genetic variation in traits of interest. The growth hormone (GH) gene is a candidate gene for body weight and weight gain in cattle since it plays a fundamental role in growth regulation (Silveira *et al.*, 2008). The GH

gene is considered an attractive candidate gene for use as a beef and milk production marker due to its role in galactopoietic metabolism and the growth process. Bovine growth hormone (bGH) gene is localized in chromosome 19 (Hediger *et al.*, 1990), and consists of five exons separated by interval introns (Gordon *et al.*, 1983).

The effects of some GH gene polymorphisms have been widely studied in beef cattle and the proximity between some of these polymorphisms, which can be characterized using different restriction enzymes, suggests a strong linkage between them. The presence of the *Alu-I* restriction site corresponds to the presence of the amino acid leucine (L) at position 127 in the polypeptide chain of cattle GH, whereas the absence of this site indicates the presence of valine (V) at the same position (Lucy *et al.*, 1991; Switonski, 2002). The study of GH gene *Alu-I* locus have been reported in Bavarian Simmental cattle (Schlee *et al.*, 1994), Bali, Madura and Benggala cattle (Sutarno *et al.*, 2002), Brahman cattle (Beauchemin *et al.*, 2006), Angus and Shorthorn cattle (Barendse *et al.*, 2006), Iranian cattle (Zakezadeh *et al.*, 2006), and West Sumatra Pesisir cattle (Jakaria *et al.*, 2007).

The aim of this research was to identify the polymorphism of growth hormone (GH) gene *Alu-I* locus in Bali cattle from Bali and Lombok islands.

MATERIAL AND METHODS

Blood Sample and DNA Extraction

The 232 blood samples of Bali cattle were obtained from Bali Cattle Breeding Center (180 samples) and Center for Regional Artificial Insemination (20 samples) at Bali island, whereas 32 samples from Lombok island were collected from small holder farmers. Blood sampling were performed by veterinarians taken via jugular vein contain 5 ml blood samples and then preserved in ethanol absolute. Blood samples were extracted according Sambrook *et al.* (1989) and dissolved in TE solution. The quality and quantity of the total genome were analyzed using spectrophotometer and 1% agarose gel electrophoresis.

Amplification of GH Gene *Alu-I* Locus and Genotyping

The GH gene *Alu-I* locus was analyzed by using the Polymerase Chain Reaction-Restriction Fragment Length Polymorphism (PCR-RFLP) method. PCR amplification conditions were as follows: 94°C for 2 min, followed by 30 cycles of denaturation at 94°C for 30 s, 60°C annealing for 45 s and extension at 72°C for 1 min, and final extension at 72°C for 5 min. A 211 bp fragment of partial exon 5th in GH gene was amplified by using primer forward 5'-GCT GCT CCT GAG GGC CTT C-3' and reverse 5'-CAT GAC CCT CAG GTA CGT CTC CG-3' (Reis *et al.*, 2001). The PCR product was digested at 37°C for overnight by *Alu-I* enzyme (AGCT restriction site sequence) and then, the digestion product was separated by Mupid horizontal electrophoresis.

Sequence of GH *Alu-I* Fragment

The GH *Alu-I* fragment was done by sequencer machine of ABI Prims 3100-Avant Genetic Analyzer (PT Charoen Pokphan Indonesia). The sequence material used was PCR product of GH *Alu-I* fragment, primer *forward*, QIA-Quick PCR Purification Kit-Qiagen, 125 mM EDTA, ethanol absolute, 70% ethanol and Hi-Di *Formamide*.

Data Analysis

PCR-RFLP data was analyzed by allele frequency (Nei 1987). The allele frequency was calculated by formula as :

$$p = \frac{2(AA) + (Aa)}{2N}, q = \frac{2(aa) + (Aa)}{2N}$$

Where, p is the L allele frequency, q is the V allele frequency and N is the total number of individual tested.

Polymorphic Informative Content (PIC) value was estimated by calculated (Botstein *et al.*, 1980) as :

$$PIC = 1 - \sum_{i=1}^n p_i^2 - \sum_{i=1}^{n-1} \sum_{j=i+1}^n 2p_i^2 p_j^2$$

Where p_i is the population frequency of the i^{th} allele and n is the number of alleles per marker. Sequences result were analyzed by Molecular Evolutionary Genetic Analysis (MEGA4) software program with alignment explorer/clustal method (Kumar and Tamura, 2006).

RESULTS AND DISCUSSION

Allele Frequencies of GH Gene *Alu-I* Locus

The following DNA restriction fragments result was obtained only one LL genotype of Bali cattle from Bali island, while Bali cattle from Lombok island was obtained two alleles, those were LL and VV, respectively. Based on genotyping data of GH *Alu-I* locus indicated 160 bp and 51 bp for the LL genotype, 211, 160 and 51 bp for the LV genotype and 211 bp (no digestion) for the VV genotype (Figure 1 and 2).

The genotype and allele frequencies of GH *Alu-I* locus for Bali cattle from Bali and Lombok islands are presented in Table 1. LL genotype and L allele frequencies of Bali cattle from Bali and Lombok islands were 1.00 and 0.97, respectively. The high L allele frequency was observed in both Bali cattle from Bali and Lombok islands. Based on the results obtained, the GH gene *Alu-I* locus is monomorphic in Bali cattle from Bali island, whereas polymorphic in Bali cattle from Lombok island. Nei (1987) stated that monomorphic is allele frequency level equal or less than 0.99 (99%), whereas polymorphic is allele frequency level equal or less than 0.01 (1%). This means that the Bali cattle was found on the Bali island were uniform, unlike the Bali cattle on the Lombok island were various in GH *Alu-I* locus. GH *Alu-I* locus diversity (polymorphic) will give an importance subject in animal breeding program that can be used as a marker assisted selection (MAS) when association with the economic traits.

There was a tendency that L allele frequency of GH-*Alu-I* locus in hump cattle is (*Bos indicus*) higher than humpless cattle (*Bos taurus*) (Table 2). Base on result obtained that the L allele




Figure 1. PCR Product of *GH Alu-I* Locus Detected by Agarose Gel Electrophoresis. M: marker (100 bp); line 1-10 : sample number




Figure 2. Genotyping Results of *GH Alu-I* Locus Detected by Agarose Gel Electrophoresis. (+): positive control (PCR product); LL: genotype

Table 1. Genotype Number and Allele Frequency of Bali Cattle from Bali and Lombok Islands

Originated	n	Genotype			Allele	
		LL	LV	VV	L	V
Bali island	200	200	0	0	1.00	0.00
Lombok island	32	31	0	1	0.97	0.03

n = number of animal

frequency is higher than the V allele in Bali cattle from Bali and Lombok islands. The highest L allele in Bali cattle population from Bali island is caused by natural selection, no crossing program and conservation area. While Bali cattle from Lombok island showed a crossbreeding program intensively. Crossbreeding programs introduced V allele, because the Limousine and Simmental cattle that used as artificial insemination (AI) semen carried V alleles. Jakaria *et al.* (2009) states that V allele frequency in Limousine and Simmental cattle were 0.182 and 0.306, respectively.

Polymorphic Informative Content (PIC) Value

PIC value result presented in Table 3. PIC value showed that the *GH-Alu-I* locus is no informative (monomorphic) in Bali cattle especially Bali cattle from Bali island, while Bali cattle from Lombok island is moderate informative (polymorphic). Based on the estimation PIC value, it could be concluded that *GH Alu-I* marker was not effective for genetic diversity information in Bali cattle from Bali island. Jakaria *et al.* (2007) also reported that PIC value was low in West Sumatra Coastal cattle using PCR-RFLP marker, in contrast on the PIC

Table 2. Allele Frequency Distribution of GH-*Alu-I* Locus in Humpless and Hump Cattle Breeds

Cattle breed	n	Allele Frequencies		Authors
		L	V	
Portuguese Beef cattle	195	0.76	0.24	Reis <i>et al.</i> (2001)
Mazandrani	97	0.91	0.09	Zakezadeh <i>et al.</i> (2006)
Angus	527	0.77	0.23	Barendse <i>et al.</i> (2006)
Shorthorn	500	0.76	0.24	Barendse <i>et al.</i> (2006)
Brahman	324	1.00	0.00	Beauchemin <i>et al.</i> (2006)
Nellore	79	1.00	0.00	Curi <i>et al.</i> (2006)
West Sumatra Pesisir	133	0.99	0.01	Jakaria <i>et al.</i> (2007)
Bali cattle from Bali island	200	1.00	0.00	Present study
Bali cattle from Lombok island	32	0.97	0.03	Present study

n = number of animal

Table 3. Estimation of Polymorphic Informative Content (PIC) Value in Bali Cattle from Bali and Lombok Islands

Originated	n	PIC Value
Bali island	200	0.000
Lombok island	32	0.333

n = number of animal

10	20	30	40	50	60
TTGTTTCCTC	CTTGGCAGGA	<u>GCTGGAAGAT</u>	GGCACCCCCC	GGGCTGGGCA	GATCCTCAAG
70	80	90	100	110	120
CAGACCTATG	AACAATTTGA	CACAAACATG	CGCAGTGACG	ACGCGCTGCT	CAAGAACTAC
130	140	150	160	170	180
GGTCTGCTCT	CCTGCTTCCG	GAAGGACCTG	CATAAGACGG	AGACGTACCT	GAGGGTCATG

Figure 3. Sequences Results of GH *Alu-I* Locus Detected by Sequencer Machine. AGCT= *Alu-I* enzyme restriction site

value estimation using microsatellite DNA marker where PIC value has high informative loci in Spanish sheep (Arranz *et al.*, 2001). Thus, PCR-RFLP maker was less informative than microsatellite DNA marker.

Estimation of PIC value was a parameter that used as an informative marker. Botstein *et al.* (1980) reported that the criteria of PIC value divided in three group namely low ≤ 0.25 , moderate $0.25 < PIC < 0.5$, and high ≥ 0.5 . Furthermore, explained that the PIC is not only can be used to determine an informative marker, but also can be used to determine the presence or absence of the polymorphic allele.

Sequencing of GH *Alu-I* locus

Sequencing results of GH *Alu-I* locus obtained *Alu-I* enzyme restriction site (AGCT) (Figure 3) that has a polymorphism in the GH gene. Zhang *et al.* (1993) and Lucy *et al.* (1993) reported that GH gene *Alu-I* locus occurred mutation between cytosine (AGCT) to guanine (AGGT) respectively. Jakaria *et al.* (2009) also reported a point mutation at GH gene of 5th exon from cytosine (C) to guanine (G) in Indonesian Limousine, Simmental and Bali cattle. Mutation of this gene have been described in dairy cattle (Lagziel and Soller, 1999) and beef cattle

(Barendse *et al.*, 2006) to affect important production and reproduction traits.

CONCLUSION

This study can be concluded that high frequency of L allele on Bali cattle from Bali and Lombok islands. GH *Alu-I* locus was found monomorphic in Bali cattle from Bali island, while polymorphic in Bali cattle from Lombok island. Whereas AGCT sequence *Alu-I* enzyme restriction site was found in GH gene exon 5th of Bali cattle from Bali island.

ACKNOWLEDGMENTS

Authors thank to Higher Education General Directorate that supported a funding in this research (Fundamental research by contract number 85/13.24.4./SPK/BG-PD/2009). Authors also thank to Masykur, M.Si. for collecting blood samples of Bali cattle in Lombok island.

REFERENCES

- Abdullah, M.A.N. 2008. Karakterisasi genetik sapi Aceh menggunakan analisis keragaman fenotipik, daerah D-Loop DNA mitokondria dan DNA mikrosatelit. Dissertation. Program Pascasarjana, Institut Pertanian Bogor. Indonesia.
- Arranz, J.J., Y. Bayon and F.S. Primitivo. 2001. Genetic variation at microsatellite loci in Spanish sheep. *Small Ruminant Research* 39:3-10
- Barendse, W., R.J. Bunch, B.E. Harrison and M.B. Thomas. 2006. The growth hormone 1 GH1:c.457C>G mutation is associated with intramuscular and rump fat distribution in large sample of Australian feedlot cattle. *Anim. Genet.* 37:211-214
- Beauchemin, V.R., M.G. Thomas, D.E. Franke and G.A. Silver. 2006. Evaluation of DNA polymorphisms involving growth hormone relative to growth and carcass characteristics in Brahman steers. *Genet. Mol. Res.* 5(3):438-447
- Botstein, D., R.L. White, M. Skolnick and R.W. Davis. 1980. Construction of a genetic linkage map in human using restriction fragment length polymorphisms. *Am. J. Hum. Genet.* 32:314-331
- Curi, R.A., D.A. Palmieri, L. Suguisawa, H.N. de Oliveira, A.C. Silveira and C.R. Lopes. 2006. Growth and carcass traits associated with *GH1/Alu-I* and *POU1F1/HinfI* gene polymorphisms in Zebu and crossbred beef cattle. *Genet. Mol. Biol.* 29:56-61
- Gordon, D.F., D.P. Quick and R.C. Erwin. 1983. Nucleotide sequence of the bovine growth hormone chromosomal gene. *Mol. Cell Endocrinol.* 33:81095
- Handiwirawan, E., R.R. Noor and Muladno. 2003. The use of HEL9 and INRA035 microsatellites as specific markers for Bali cattle. *Arch. Tierz., Dummerstorf* 46 (6) : 503-512
- Hediger, R., S.E. Johnson, W. Barendse, R.D. Drinkwater, S.S. Moore and J. Hetzel. 1990. Assignment of the growth hormone gene locus to 19q26 qter in cattle and to 11q25 qter in sheep by in-situ hybridization. *Genome* 8:171-174
- Jakaria, D. Duryadi, R.R. Noor, B. Tappa and H. Martojo. 2007. Evaluasi keragaman genetik gen hormone pertumbuhan sapi Pesisir Sumatera Barat menggunakan penciri PCR-RFLP. *Media Peternakan* 30:1-10
- Jakaria, R. R. Noor, H. Martojo, D. Duryadi and B. Tappa. 2009. Distribution of growth hormone (GH) gene *Msp-1* and *Alu-I* Loci Polymorphism in Indonesian Cattle Breeds. Proceeding The First International Seminar in Animal Industry and Food Safety, Bogor-Indonesia. November 22-24, 2009. P 42-47
- Kumar, S. and K. Tamura. 2006. MEGA4: Molecular Evolutionary Genetics Analysis Software. Arizona State University. Arizona USA.
- Lagziel, A. and M. Soller. 1999. DNA sequence of SSCP haplotypes at the bovine growth hormone [bGH] gene. *Animal Genetics* 30, 362-365
- Lucy, M.C., S.D. Hauser, P.J. Eppard, G.G. Krivi, J.H. Clark, D.E. Bauman and R.J. Collier. 1993. Variants of somatotropin allele in cattle: Gene frequencies in major dairy breeds and associated milk production. *Dom. Anim. Endocrinol.* 10: 325-333
- Martojo, H. 2003. Indigenous Bali Cattle: The Best Suited Cattle Breed for Sustainable Small Farm in Indonesia. The Chinese Society of Animal Science. 112 Farm Road. Hsinhua. Tainan. Taiwan.
- Mohamad, K., M. Olsson, H.T.A. van Tol, S. Mikko, B.H. Vlamings, G. Andersson, H.R.quez-Marti, B. Purwantara, R.W. Paling, B. Colenbrander and J.A. Lenstra. 2009. On the

- origin of Indonesian cattle. *PloS One*. 4(5):e9450
- Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press. New York.
- Nijman I.J., M. Otsen, E.L. Verkaar, C. de Ruijter and E. Hanekamp. 2003. Hybridization of banteng (*Bos javanicus*) and zebu (*Bos indicus*) revealed by mitochondrial DNA, satellite DNA, AFLP and microsatellites. *Heredity*. 90:10–16
- Reis, C., D. Navas, N. Pereira and A. Cravador. 2001. Growth hormone Alu-I polymorphism analysis in eight Portuguese *bovine* breeds. *Arch. Zootec.* 50:41-48
- Sambrook, J., E.F. Fritsch and T. Maniatis. 1989. *Molecular Cloning; a Laboratory Manual*. CSH Laboratory Press. USA.
- Schlee, P., R. Graml, O. Rottmann and F. Pirchner. 1994. Influence of growth hormone genotypes on breeding values of Simmental bulls. *J. Anim. Breed. Genet.* 111:253-256
- Silveira, L.G.G., L.R. Furlan, R.A. Curi, A.L.J. Ferraz, M.M. de Alencar, L.C.A. Regitano, C.L. Martins, M.B. Arrigoni, L. Suguisawa, AC. Silveira and H.N. Oliveira. 2008. Growth hormone 1 gene (GH1) polymorphisms as possible markers of the production potential of beef cattle using the Brazilian Canchim breed as a model. *Genet. Mol. Bio.* 31(4):874-879
- Sutarno, A. Junaidi, A. Purwoko and N.I. Lelana. 2002. Identification and characterization of the polymorphic growth hormone gene of the Bali cattle, the Madura cattle and Ongole cattle. *Biodivesitas* 3 (1):169-173
- Switonski, M. 2002. Molecular genetics in beef cattle breeding – A review. *Anim. Sci. Pap. Rep.* 20:7-18
- Uggla, C.M. 2008. Investigating genetic variability within specific indigenous Indonesian cattle breeds. Dissertation SLU, Sweden.
- Zakezadeh, S., G. Rahimi, S.R. Mirae-Ashtiani, A. Nejati-Javaremi, M. Moradi-Shahrbabak, P. Reineeke, M. Reissmann, A.A. Masoudi, C. Amirinia, and S.A. Mirhadi. 2006. Analysis of bovine growth hormone gene polymorphism in three Iranian native breeds and Holstein cattle by RFLP-PCR. *Biotech.* 5:385-390
- Zhang, H.M., D.R. Brown, S.K. Denise and R.L. Ax. 1993. Polymerase chain reaction restriction fragment length polymorphism analysis of the *bovine* somatotropin gene. *J. Anim. Sci. (Abstract)* (71):2276