



Research Article

© 2021 Deguma et al..  
This is an open access article licensed under the Creative Commons  
Attribution-NonCommercial 4.0 International License  
(<https://creativecommons.org/licenses/by-nc/4.0/>)

Received: 30 March 2021 / Accepted: 22 May 2021 / Published: 8 July 2021

## Analyzing Heart Disease Mortality of Filipino: From Statistical Modeling to Health and Lifestyle Education Implications

Jabin J. Deguma

Emerson D. Peteros

Reylan G. Capuno

Ricardo Q. Ybañez

Danilo F. Cebe

Helen O. Revalde

Regina E. Sitoy

Melona C. Deguma

College of Education,  
Cebu Technological University,  
Cebu City, Philippines

DOI: <https://doi.org/10.36941/ajis-2021-0102>

### Abstract

*This paper contributes an interdisciplinary cross-over in studying heart disease mortality of Filipino. First, it validates previous studies on heart diseases via ascertaining statistical models of analyses that estimate heart disease mortality predictability in the Philippines. It then proceeds to understand the implications of the issue through promoting health and lifestyle education. To do this, first, the report from the Epidemiology Bureau of the Department of Health (EBDOH) on mortality cases of diseases of the heart in the Philippines. Based on the statistical analyses, time series analysis suggested that the growth of heart disease mortality in the Philippines followed a quadratic trend. Moreover, symbolic regression (SR) analysis revealed that heredity has more significant influence over lifestyle between the identified factors. Based on the proposed models, this paper implies furthering community-oriented health and wellness programs as practical means to avoid untimely deaths brought by the said diseases.*

**Keywords:** heart diseases mortality, time series, symbolic regression, health and lifestyle guidance

### 1. Introduction

The World Health Organization (WHO) (2017a) declared that heart diseases had taken millions of lives every year while causing thirty-one percent of deaths worldwide. Hajar (2016) claimed that heart disease mortality is expected to grow to more than 23.6 million by 2030. Heart diseases include coronary heart diseases, cerebrovascular disease, peripheral arterial disease, rheumatic heart disease,

congenital heart disease, deep vein thrombosis, and pulmonary embolism [2]. In the data of WHO (2017b), the Philippines has reached 122, 950 or 19.86% of total death caused by coronary heart disease. Consequently, the country ranked 28th in the survey of countries' death caused by the said disease. Simultaneously, the Department of Health (DOH) (2018) reported that heart diseases, among other illnesses, caused a fatality in the country. As written by Palaniappan et al. (2010), Filipino immigrants in the United States of America have a second highest heart disease threat after Indians. Consequently, heart diseases have become a recurrent health problem that every Filipino must be aware of regardless of age and gender.

The struggle to address the rise of deaths due to heart disease begins with addressing the very factors that may have caused it. As purported by Kannel et al. (2004), a risk factor is any condition or personal behavior that increases the chance to develop a particular disease, which may advance to fatality. The severity of a risk factor will be more likely to create one (Enas, 2001). As reported by CADI Research Foundation (2012a), hereditary and lifestyle choices actively promote heart diseases.

Being genetically predisposed or susceptible may cause individuals to be at high risk of acquiring heart diseases despite not being born with it. Reilly et al. (2011) purported that current scientific undertakings show heredity disposes of a patient with heart diseases to a probable heart attack. The Harvard Medical School (2017) cited that heart diseases can be passed down through families. It added that genetic risk scores were more accurate for predicting heart attacks than any other risk factors. Agarwal (2001) added that predisposition to heart diseases at an early age increases by heredity as standard behavioral modes are observed in members of the same family. Various contemporary studies found that heredity is an independent risk factor being considered in the family history of heart diseases (Higgins, 2000; Kraus, 2000; Pereira et al., 2000; Leander et al., 2001; Wood, 2001; de Giorgis, 2009). Hence, an individual's possibility to acquire heart diseases increases due to a family history of such conditions.

Moreover, such cannot discount the significance of lifestyle to affect a person to acquire heart diseases. The studies of Karvonen (1989), Rowe & Kahn (1998), Smedley & Syme (2000), and Ruston & Clayton (2002) found out that lifestyle also contributes to the growing number of heart disease mortality. Heart diseases is also a lifestyle-related non-communicable disease. An unhealthy lifestyle like smoking, eating unhealthy foods, and lack of exercise leads to acquiring heart diseases (DOH, 2018). As purported by Idler & Benyamini (1997), Singer & Ryff (2001), and Niiranen & Vasan (2016), excess rates of chronic diseases, particularly the conditions of the heart, are affected by social and behavioral factors including lifestyle, diet, levels of physical activity, or gender and cultural stress responses. Filipino immigrants in the United States tend to cope with unhealthy eating and smoking to be relieved from work and family stress, thereby making them prone to acquiring heart diseases (U.S. Department of Health and Human Services, 2010).

## 2. Objectives of the Study

Relative to the literature, we draw an assumption that heredity and lifestyle are two salient factors that could have contributed to the growing number of deaths due to the risk of getting heart diseases. In this context, this paper primarily intends to explore the trend of heart disease mortality in the Philippines. It validates previous studies on heart diseases via ascertaining statistical models of analyses that estimate heart disease mortality predictability in the Philippines. It also identifies which among the two identified risk factors, heredity and lifestyle, greatly influenced heart disease mortality in the Philippines by providing models to predict through symbolic regression analysis. It then proceeds to understand the implications of the issue through promoting health and lifestyle education.

## 3. Materials and Methods

This study retrieved essential data through data mining. Data mining is the process of taking out datasets to ascertain patterns and institute associations through data analysis (Rouse, 2008;

Witten, 2011; Deguma et al, 2018). Witten et al. (2011) furthered that data mining's benefit allows uncovering hidden patterns and relationships, which aids in making predictions. However, data mining must consider data privacy as an ethical consideration (Iles, 2013). According to Davis and Patterson (2012), data in itself is ethically neutral. In the Philippines, the responsible use of data for research purposes is a valid ground (National Privacy Commission, n.d.).

This study utilized the Epidemiology Bureau of the Department of Health (EBDOH) (2014) on mortality cases of diseases of the heart in the Philippines featured the demographics, natality, morbidity and notifiable diseases, and mortality statistics from 1964 to 2014. The report also included the mortality of diseases of the heart in the Philippines. Due to the difference in population volume for the past fifty years, this study used data from 1995 to 2014 to determine the trend of heart disease mortality, as shown in Table 1.

This study employed descriptive methods via time-series analyses using Minitab ® and Symbolic Regression using Eureqa Pro®. The data on the number of mortality cases of heart diseases in the Philippines is evaluated first through a trend analysis, which provides three models, namely linear, quadratic and exponential growth. Time series analysis is done to determine which model provides the least Mean Absolute Percentage Error (MAPE) and forecasts the trend of heart disease mortality in the Philippines for the past twenty years using Minitab ®. Moreover, the study also identified which among the two factors, hereditary or lifestyle, contributed significantly to heart disease mortality. Symbolic regression (SR) analysis using Eureqa Pro ® is done to look for formulas that provide the least Mean Absolute Error (MAE) for hereditary and lifestyle factors. SR is a method that predicts relationships and connects variables while providing a model that is not assumed beforehand. According to Barmpalexis et al. (2011), SR operates through genetic programming (GP). GP generates populations of equations patterned after the fittest principle's survival (Koza, 1994; Koza et al., 2003; Barmpalexis, 2011) GP searches for the fittest equation from a series of runs in every cycle called generation by setting a target expression. Through GP, SR structured patterns from a dataset of variables to find the "fittest" combination model prediction (Claveria, 2016). The current SR experiment used the same data of the Epidemiology Bureau of the Department of Health on heart disease mortality from 1995 to 2014. This experiment will find the fittest formula with the least Mean Absolute Error (MAE), which could provide a minimal error in arriving at model estimates (Loce & Dougherty, 1995; De Myttenaere et al., 2016).

**Table 1:** Reported Number of Mortality Cases of Diseases of the Heart in the Philippines from 1995 to 2014 by EBDOH (2014)

Year	Number of Mortality Cases of Diseases of the Heart	Year	Number of Mortality Cases of Diseases of the Heart
1995	50,252	2005	77,060
1996	53,865	2006	83,081
1997	54,787	2007	88,314
1998	55,830	2008	92,133
1999	58,574	2009	100,908
2000	60,417	2010	102,936
2001	62,950	2011	107,294
2002	70,861	2012	112,581
2003	67,696	2013	118,740
2004	70,861	2014	132,825

## 4. Results and Discussion

### 3.1 Structuring the Models for Analyses

The data from EBDOH (2014) on heart disease mortality from 1995 to 2014 provided a twenty-year

time series analysis using linear, quadratic, and exponential growth as shown in Figures 1, 2, and 3. As indicated in the graphs, the number of heart disease mortality has significantly increased over the past twenty years. As shown in the figures, a remarkable difference among the provided by the three models. Results suggested that the Quadratic Trend Model has the least error to map the trend of Filipinos' heart disease mortality. The Quadratic Trend Model could provide the fittest trend analysis of the graph's behavior over the Linear Trend and Exponential Growth Models. With a quadratic trend, the time series values are constantly dependent on variables that could affect the trend's movement, thus changing over time (Wu & Zhao, 2007; Jebb & Tay, 2017). Hence, acknowledging that the mortality cases of heart diseases in the Philippines follow a quadratic trend furthered the importance of identifying the risk factors (e.g., lifestyle and heredity) as variables that could affect the rising and falling of the model trend.




Figure 1: Linear Trend Model for Heart Diseases Mortality in the Philippines from 1995-2014 with a MAPE of 5




Figure 2: Exponential Trend Model for Heart Diseases Mortality in the Philippines from 1995-2014 with a MAPE of 3




Figure 3: Quadratic Trend Model for Heart Diseases Mortality in the Philippines from 1995-2014 with a MAPE of 2

The study advanced at generating the fittest equations for estimating the predictability of heart disease mortality through symbolic regression (SR) using another software, Eureqa Pro<sup>®</sup>. The first target expression used to predict lifestyle is as follows:  $y=f(t)$ , where the number of cases of heart disease mortality ( $y$ ) is affected by the function of time, which is represented by lifestyle ( $t$ ). The lifestyle of individuals is generally associated with the function of time wherein one unit of time ( $t$ ) is equivalent to a one-unit change in lifestyle of the people (Monk et al., 2002; Sakano et al., 2009; Feldman et al., 2017; Brinkman et al., 2018). The target expression presumed that lifestyle is a factor that contributes to heart disease mortality. The result of symbolic regression analysis for the target expression provides a fit formula that follows:

$$y = 5.13e4 + 416t + 177t^2 + 203t^3 \sin(t) - 1.36e3 t \sin(t) - 2.42e3 \cos(158t) - 6.27t^3 \sin(t)$$

The generated equation result showed a quadratic trend manifested by the presence of  $t^2$ . The *sine* and *cosine* functions indicated correction factors and seasonal variations, which contributed to the formula's accuracy to predict the number of heart disease mortality in the Philippines. The model generates a Mean Absolute Error (MAE) of 819.0044.

Moreover, the next target expression used to predict heredity as a cause of heart diseases mortality is as follows:  $y=f(\text{delay}(y,1), \text{delay}(y,2), \text{delay}(y,3))$ , where the number of cases of heart diseases mortality ( $y$ ) is affected by the function of the previous number of those who die due to heart diseases ( $y$ ). The number of heart diseases death ( $y$ ) is affected up to the third-degree consanguinity, thereby looking at heredity as a factor that causes heart disease mortality (Bailleul et al., 1995; Singh et al., 1999; Wienke et al., 2001; Glowinska et al., 2002; Zdravkovic et al., 2002; Daniels, 2011). The result of symbolic regression analysis for the target expression provides a formula that follows:

$$y = 970 + 0.919 \text{delay}(y, 1) + 0.136 \text{delay}(y, 3) + 2.66e3 \sin(2 \text{delay}(y, 2)) + \frac{-332}{\sin(1.09 + \text{delay}(y, 2))} + \frac{-846}{\sin(\sin(5.17 + \text{delay}(y, 3)))}$$

The presence of *sine* function indicated fluctuations within the cycle, which contributed to the accuracy of predicting the number of heart disease mortality in the Philippines, resulting in a Mean Absolute Error (MAE) of 544.07409.

The study progressed at evaluating the MAE of both heredity and lifestyle factors by getting the sum of the total MAE, which is 1,363,07849. The MAE of the heredity affected lifestyle and vice versa. Hence, by dividing the MAE of either of the factors over the total MAE, the result shows the computed value of the factors contributing to heart disease mortality. The study revealed that the influence of lifestyle as a factor has a computed value of 39.92%, whereas heredity has 60.08%. Thus, SR revealed that heredity has more significant influence over lifestyle on heart disease mortality in the Philippines for the past twenty years.

#### 4. Furthering the Implication Based on the Proposed Models

The Sustainable Development Goals (SDG) of the United Nations Development Programme (2015) aimed to increase life expectancy and to reduce mortality from chronic diseases and mental illnesses by promoting good health and well-being for all people. Good health promotes a sound economy while being beneficial to individual lives (National Academy on an Aging Society 1999a, 1999b). As validated in this paper, a quadratic trend model provided the fittest equation with the least MAPE for the mortality cases of diseases of the heart in the Philippines from 1995 to 2014. It supports the claim of Tuomilehto et al. (1984) that starting from 1963 to 1976, there an increase in the cardiovascular disease death rate among Filipinos. This increasing trend is prevalent in many low-income and middle-income countries like the Philippines (Epping-Jordan et al., 2005). Epping-Jordan et al. (2005) added that the reason behind such a phenomenon is the rapid social and environmental changes that cause the increase of risk factors of heart disease mortality. According to the study of WHO (2002), such phenomenon can be prevented through a healthy lifestyle, which becomes uncommon to people living in third world countries. Therefore, it is recommended that supportive policies for a healthy

lifestyle could be enacted to reduce the increasing number (WHO, 2005).

Although this study provided that lifestyle is a lesser risk factor that leads to heart disease mortality, adopting a positive lifestyle change can effectively prevent the 39.92% risk factor of heart disease mortality and improve one's life quality. According to CADI Research Foundation (2012b), a critical choice of lifestyle is needed to sharpen the chances of developing heart diseases. Through DOH, the Philippines government must adopt health promotion programs like lifestyle modification, including proper diet, avoidance of vices such as smoking, heavy drinking, etc., and regular exercises that reduce the chance by 39.92% of getting heart diseases. As suggested by Barbara et al. (2003), lifestyle guidance is encouraged through training to help the individual make and maintain the right choices of a healthier lifestyle. It is thereby reducing the risk through the individual's ability to manage the self through correct lifestyle choices. However, lifestyle intervention must be culturally tailored to be effectively implemented (Leake et al., 2011). Wallace et al. (2008) maintained that a culturally tailored cardiovascular disease prevention curriculum was effective in reducing the mortality rate. Also, Hinohara et al. (1982) recommended strengthening community-oriented health and wellness programs by government and private sectors. These programs must be focusing on reducing lifestyle factors through education, which is tailored specifically to address the most prevalent needs of a particular community. Hence, education must also focus on health promotion and protection and disease prevention (Fitch & Blue, 1982).

## 5. Conclusion and Recommendation

In conclusion, heart disease mortality among Filipinos is increasing for the past twenty years, from 1995 to 2014, if nothing is positively and effectively done to change the trend. The rise of such health phenomenon is greatly affected by two identified factors like heredity and lifestyle. The former is seen to be of more significant influence than the latter. However, modern technology advances at determining an individual's genetic predisposition to heart diseases, the rising trend of heart disease mortality is affected by Filipinos' unhealthy lifestyle. Hence, the change of lifestyle is significantly recommended to lessen the increasing trend of heart disease mortality. Overall, it is imperative to invest in health and wellness, which will eventually lead to better monetary, social, and environmental returns, good health and well-being for all people, and the Sustainable Development Goals' attainment.

## References

- Agarwal, D.P. (2001). *Genetic Predisposition to Cardiovascular Diseases*. International Journal of Human Genetics, 1:4, 233-241
- Bailleul, S., Couderc, R., Rossignol, C., et al. (1995). *Lipoprotein (a) in childhood: relation with other atherosclerosis risk factors and family history of atherosclerosis*. Clin Chem. 41(2):241-245.
- Barbara, W., Yee, K., Nguyen, H.T., & Ha, M. (2003). *Chronic Disease Health Beliefs and Lifestyle Practices among Vietnamese Adults*. Women & Therapy, 26:1-2, 111-125.
- Barmapalexis, P., Kachrimanis, K., Tsakonas, A., & Georgarakis, E. (2011). *Symbolic regression via genetic programming in the optimization of a controlled release pharmaceutical formulation*. Chemometrics and Intelligent Laboratory Systems, 107, 75-82. Doi: 10.1016/j.chemolab.2011.01.012
- Brinkman, S., Voortman, T., Kiefte-de Jong, J. C., van Rooij, F. J., Ikram, M. A., Rivadeneira, F., ... & Schoufour, J. D. (2018). The association between lifestyle and overall health, using the frailty index. *Archives of gerontology and geriatrics*, 76, 85-91.
- CADI Research Foundation. (2012a). *Heart Disease Overview*. Retrieved from <http://www.cadiresearch.org/topic/understanding-heart-disease/overview>
- CADI Research Foundation. (2012b). *Risk Factor Overview*. Retrieved from <http://www.cadiresearch.org/topic/risk-factors/risk-factors-overview>
- Claveria, O., Monte, E., & Torra, S. (2016). Quantification of survey expectations by means of symbolic regression via genetic programming to estimate economic growth in Central and Eastern European economies. *Eastern European Economics*, 54(2), 171-189.
- Daniels, SR. (2011). *Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents: Report from National Heart, Lung and Blood Institute*. Retrieved from [http://www.nhlbi.nih.gov/guidelines/cvd\\_ped/index.html](http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.html)

- Davis, K., & Patterson, D. (2012). *Ethics of Big Data*. Retrieved from Safari Books.
- de Giorgis, T., Giannini, C., Scarinci, A., et al. (2009). *The family history of premature cardiovascular disease as a sole and independent risk factor for increased carotid intima-media thickness*. *J Hypertens*, 27(4):822-828.
- De Myttenaere, A., Golden, B., Le Grand, B., & Rossi, F. (2016). Mean absolute percentage error for regression models. *Neurocomputing*, 192, 38-48.
- Deguma, J., Peteros, E., Case, M., & Igot, V. (2018). Violence against Women and Gender Equality in the Philippines: Are they Related?. *Journal of Educational and Human Resource Development*, 6, 68-81.
- Department of Health. (2018). *Cardiovascular Disease*. Retrieved from <https://www.doh.gov.ph/cardiovascular-disease>
- Enas, E.A. (2001). *Lipoprotein (a) is an important genetic risk for coronary artery disease in Asian Indians*. *The American Journal of Cardiology*. DOI: 10.1016/S0002-9149(01)01659-9
- Epidemiology Bureau Department of Health. (2014). *The 2014 Philippine Health Statistics*. Retrieved from [https://www.doh.gov.ph/sites/default/files/publications/2014PHS\\_PDF.pdf](https://www.doh.gov.ph/sites/default/files/publications/2014PHS_PDF.pdf)
- Epping-Jordan, J., Galea, G., Tukulunga, C., & Beaglehole, R. (2005). *Preventing chronic diseases: Taking stepwise action*. Retrieved from <https://pdfs.semanticscholar.org/e49c/7ee982ef681650360a004f905bf709c6b18e.pdf>. DOI:10.1016/S0140-6736(05)67454-
- Feldman, A. L., Long, G. H., Johansson, I., Weinehall, L., Fhärm, E., Wennberg, P., ... & Rolandsson, O. (2017). Change in lifestyle behaviors and diabetes risk: evidence from a population-based cohort study with 10 year follow-up. *International Journal of behavioral nutrition and physical activity*, 14(1), 39.
- Fitch, J.A., & Blue, B.E. (1982). *From an Educational Perspective: New Approaches to Cardiovascular and other Chronic Diseases Suggest a Bright Future for School and Community Health Education*. *Health Education*, 13:1, 35-38.
- Glowinska, B., Urban, M., & Koput, A. (2002). *Cardiovascular risk factors in children with obesity, hypertension, and diabetes: lipoprotein (a) levels and body mass index correlate with a family history of cardiovascular disease*. *Eur J Pediatr*. 161(10):511-518.
- Hajar, R. (2016). *Framingham contribution to cardiovascular diseases*. *Heart Views*, 17:78-81.
- Harvard Medical School. (2017). *The genetics of heart disease: An update*. *Harvard Health Publishing*. Retrieved from <https://www.health.harvard.edu/heart-health/the-genetics-of-heart-disease-an-update>
- Higgins, M. (2000). *Epidemiology and prevention of coronary heart disease in families*. *Am J Med*, 108: 387-395.
- Hinohara, S., Hinohara, M., & Takagi, H. (1982). *A Study on the Evaluation of Health and Lifestyle Habits in a Rural Area of Japan*. *Med. Inform. Vol. 7, No.3, 229-234*.
- Idler, E.L., & Benyamini, Y. (1997). *Self-rated health and mortality: A review of twenty-seven community studies*. *Journal of Health and Social Behavior*, 38, 21-37.
- Iles, H.R. (2013). *Ethics of Data Mining: a New Zealand Survey*. Retrieved from <https://www.researchgate.net/publication/264673019>.
- Jebb, A. T., & Tay, L. (2017). Introduction to time series analysis for organizational research: Methods for longitudinal analyses. *Organizational Research Methods*, 20(1), 61-94.
- Kannel, W.B., D'Agostino, R.B., Sullivan, L., & Wilson, P.W. (2004). *Concept and usefulness of cardiovascular risk profiles*. *American Heart Journal*. <https://doi.org/10.1016/j.ahj.2003.10.022>
- Karvonen, M.J. (1989). *Determinants of Cardiovascular Diseases in the Elderly*. *Annals of Medicine* 21: 3-12.
- Koza, J.R. (1994). *Genetic programming as a means for programming computers by natural selection*. *Stat. Comput.* 4, 87-112.
- Koza, J.R., Keane, M.A., Streeter, M.J., Mydlowec, W., Yu, J., & Lanza, G. (2003). *Genetic Programming IV: Routine Human-competitive Machine Intelligence*. Kluwer Academic Publisher, USA, pp. 1-3.
- Kraus, W.E. (2000). *Genetic approaches to the investigation of genes associated with coronary heart disease*. *Am Heart J*, 140: S27-35.
- Leake, A., Bermudo, V., Jacob, J., Jacob, M., & Inouye, J. (2011). *Health is Wealth: Methods to Improve Attendance in a Lifestyle Intervention for a Largely Immigrant Filipino-American Sample*. *Immigrant Minority Health*. 14:475-480
- Leander, K., Hallqvist, J., Reuterwall, C., Ahlbom, A., & de Faire, U. (2001). *The family history of coronary heart disease, a strong risk factor for myocardial infarction interacting with other cardiovascular risk factors: results from the Stockholm Heart Epidemiology Program (SHEEP)*. *Epidemiology*. 12(2):215-221.
- Loce, R. P., & Dougherty, E. R. (1995). Mean-absolute-error representation and optimization of computational-morphological filters. *Graphical Models and Image Processing*, 57(1), 27-37.
- Monk, T. H., Frank, E., Potts, J. M., & Kupfer, D. J. (2002). A simple way to measure daily lifestyle regularity. *Journal of sleep research*, 11(3), 183-190.

- National Academy on an Aging Society. (1999a, February). *Is demography destiny?* The Public Policy and Aging Report, 9 (4), 1-14.
- National Academy on an Aging Society. (1999b, March). *Facts on low literacy skills*. FS-99-3.
- National Privacy Commission. (n.d.). *Republic Act 10173-Data Privacy Act of 2012*. Retrieved from <https://www.privacy.gov.ph/data-privacy-act/>.
- Niiranen, T.J. & Vasan, R.S. (2016). *Epidemiology of cardiovascular disease: recent novel outlooks on risk factors and clinical approaches*. Expert Review of Cardiovascular Therapy, 14:7, 855-869.
- Palaniappan, L.P., Araneta, M.R., Assimes, T.L., et al. (2010). *Call to action: cardiovascular disease in Asian Americans: a science advisory from the American Heart Association*. HHS Public Access. doi:10.1161/CIR.0b013e318f22af4.
- Pereira, M.A., Schreiner, P.J., Pankow, J.S., Williams, R.R., Higgins, M., Province, M.A., & Rao, D.C. (2000). *The Family Risk Score for coronary heart disease: associations with lipids, lipoproteins, and body habitus in a middle-aged bi-racial cohort: The ARIC study*. Ann Epidemiol. 10: 239-245.
- Reilly, M.P., Li, M., He, J., et al. (2011). *Identification of ADAMTS7 as a novel locus for coronary atherosclerosis and association of ABO with myocardial infarction in the presence of coronary atherosclerosis: two genome-wide association studies*. Lancet. 377(9763):383-392.
- Rouse, M. (2008). *Data Mining*. Retrieved from <https://searchsqlserver.techtarget.com/definition/data-mining>.
- Rowe, J.W., & Kahn, R.L. (1998). *Successful Aging*. New York, NY: Pantheon Books.
- Ruston, A. & Clayton, J. (2002). *Coronary heart disease: Women's assessment of the risk-a qualitative study*. Health, Risk & Society, 4:2, 125-137.
- Sakano, N., Wang, D. H., Takahashi, N., Wang, B., Sauriasari, R., Kanbara, S., ... & Ogino, K. (2009). Oxidative stress biomarkers and lifestyles in Japanese healthy people. *Journal of Clinical Biochemistry and Nutrition*, 44(2), 185-195.
- Singer, B.H., & Ryff, C.D. (2001). *New horizons in health: An integrative approach*. Washington, DC: National Academy Press.
- Singh, J. P., Larson, M. G., O'Donnell, C. J., Tsuji, H., Evans, J. C., & Levy, D. (1999). Heritability of heart rate variability: the Framingham Heart Study. *Circulation*, 99(17), 2251-2254.
- Smedley, B.D., & Syme, S.L. (2000). *Promoting health: Intervention strategies from social and behavioral research*. Institute of Medicine, Washington, DC: National Academy Press.
- Tuomilehto, J., Morelos, S., Yason, J., Guzman, S., & Geizerova, H. (1984). *Trends in cardiovascular Diseases Mortality in the Philippines*. International Journal of Epidemiology. Vol. 13, No.2
- U.S. Department of Health and Human Services. (2010). *Cardiovascular Risk in the Filipino Community: Formative Research from Daly City and San Francisco, California*. Retrieved from <https://www.nhlbi.nih.gov/files/docs/resources/heart/filipino.pdf>
- United Nations Development Programme. (2015). *Sustainable Development Goals*. Retrieved from <http://www.undp.org/content/undp/en/home/sustainable-development-goals.html>
- Wallace MF., Fulwood, R., & Alvarado, M. (2008). *NHLBI step-by-step approach to adapting cardiovascular training and education curricula for diverse audiences*. *Prev Chronic Dis*. 2008; 5(2). Retrieved from [http://www.cdc.gov/pcd/issues/2008/apr/07\\_0201.htm](http://www.cdc.gov/pcd/issues/2008/apr/07_0201.htm).
- Wienke, A., Holm, N. V., Skytthe, A., & Yashin, A. I. (2001). The heritability of mortality due to heart diseases: a correlated frailty model applied to Danish twins. *Twin Research and Human Genetics*, 4(4), 266-274.
- Witten, I.H., Eibe, F., & Holmes, G. (2011). *Data Mining: Practical Machine Learning Tools and Techniques*, (3rd ed.). Burlington, MA: Morgan Kaufmann.
- Wood, D. (2001). *Established and emerging cardiovascular risk factors*. Am Heart J, 141 (Suppl): S49-57.
- World Health Organization. (2002). *Diet, nutrition, and the prevention of chronic diseases: WHO Technical Report Series 916*. Geneva: World Health Organization.
- World Health Organization. (2005). *Preventing chronic diseases: a vital investment*. Geneva: World Health Organization, 2005. Retrieved from [http://www.who.int/chp/chronic\\_disease\\_report/en/index.html](http://www.who.int/chp/chronic_disease_report/en/index.html)
- World Health Organization. (2017a). *Cardiovascular Disease*. Retrieved from [http://www.who.int/cardiovascular\\_diseases/world-heart-day-2017/en/](http://www.who.int/cardiovascular_diseases/world-heart-day-2017/en/).
- World Health Organization. (2017b). *Genes and Humans Disease*. Retrieved from <http://www.who.int/genomics/public/geneticdiseases/en/index3.html>.
- Wu, W. B., & Zhao, Z. (2007). Inference of trends in time series. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(3), 391-410.
- Zdravkovic, S., Wienke, A., Pedersen, N. L., Marenberg, M. E., Yashin, A. I., & De Faire, U. (2002). Heritability of death from coronary heart disease: a 36-year follow-up of 20 966 Swedish twins. *Journal of internal medicine*, 252(3),247-254.