



Anesthetic efficacy of primary and supplemental buccal/lingual infiltration in patients with irreversible pulpitis in human mandibular molars: a systematic review and meta-analysis

Alpa Gupta¹, Aarushi Sahai¹, Vivek Aggarwal², Namrata Mehta¹, Dax Abraham¹, Sucheta Jala¹, Arundeep Singh¹

¹Department of Conservative Dentistry and Endodontics, Manav Racha Dental College, Faridabad, India

²Department of Conservative Dentistry and Endodontics, Faculty of Dentistry, Jamia Millia Islamia, New Delhi, India

Achieving profound anesthesia in mandibular molars with irreversible pulpitis is a tedious task. This review aimed at evaluating the success of buccal/lingual infiltrations administered with a primary inferior alveolar nerve block (IANB) injection or as a supplemental injection after the failure of the primary injection in symptomatic and asymptomatic patients with irreversible pulpitis in human mandibular molars. The review question was “What will be the success of primary and supplemental infiltration injection in the endodontic treatment of patients with irreversible pulpitis in human mandibular molars?” We searched electronic databases, including Pubmed, Scopus, and Ebsco host and we did a comprehensive manual search. The review protocol was framed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) checklist. We included clinical studies that evaluated and compared the anesthetic outcomes of primary IANB with primary and/or supplementary infiltration injections. Standard evaluation of the included studies was performed and suitable data and inferences were assessed. Twenty-six studies were included, of which 13 were selected for the meta-analysis. In the forest plot representation of the studies evaluating infiltrations, the combined risk ratio (RR) was 1.88 (95% CI: 1.49, 2.37), in favor of the secondary infiltrations with a statistical heterogeneity of 77%. The forest plot analysis for studies comparing primary IANB + infiltration versus primary IANB alone showed a low heterogeneity (0%). The included studies had similar RRs and the combined RR was 1.84 (95% CI: 1.44, 2.34). These findings suggest that supplemental infiltrations given along with a primary IANB provide a better success rate. L’Abbe plots were generated to measure the statistical heterogeneity among the studies. Trial sequential analysis suggested that the number of patients included in the analysis was adequate. Based on the qualitative and quantitative analyses, we concluded that the infiltration technique, either as a primary injection or as a supplementary injection, given after the failure of primary IANB, increases the overall anesthetic efficacy.

Keywords: Irreversible Pulpitis; Local Anesthesia; Mandibular Molars.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Pain-free dental treatment, a myth or a reality? Amidst all the technological advances in achieving predictable

success in endodontic therapy, the search for profound anesthesia in patients with irreversible pulpitis continues to be the Achilles heel for many clinicians today. Complete or near-complete control of pulpal pain reduces the patient’s fear of endodontic therapy [1-3]. The

Received: March 23, 2021 • Revised: May 26, 2021 • Accepted: June 5, 2021

Corresponding Author: Vivek Aggarwal, Department of Conservative Dentistry and Endodontics, Faculty of Dentistry, Jamia Millia Islamia, New Delhi, India

E-mail: drvivekaggarwal@gmail.com

Copyright© 2021 Journal of Dental Anesthesia and Pain Medicine

primary objective of a clinician is to calm the patient suffering from acute pain, which may be attributed to either pulpal or periapical pain. Crystal-clear knowledge of various local anesthetic techniques is needed to battle the ongoing situation. Attaining profound anesthesia in teeth with irreversible pulpitis is a tedious task compared to achieving anesthesia in healthy mandibular molars [4]. Further studies have shown that block anesthesia is less successful in achieving anesthesia in mandibular molars [5,6]. Local anesthetic failure can be attributed to numerous mechanisms, including anatomic variations [7-10], decreased local pH [10,11], rapid desensitization of local anesthetic [10], and activation of nociceptors and associated central mechanisms [10,12,13]. Hence, it is imperative to assess an alternate method for achieving profound anesthesia in mandibular molars with irreversible pulpitis [14,15].

Many studies have suggested alternative methods to control the inconvenience and pain experienced by subjects during endodontic treatment, including the use of supplementary injections, such as buccal infiltrations (BI), intraligamentary (IL), intraosseous (IO), and intrapulpal injections [16,17]. Both IL and IO techniques provide anesthesia to the cancellous bone adjacent to the apex of the injected teeth [18,19].

The IL injection technique is a misleading term for an anesthetic solution that fans out through the outer surface of the cribriform plate, which ultimately diffuses inside the crestal marrow spaces and does not diffuse along the periodontal space [20,21]. It provides anesthesia for almost 20 minutes and may cause postoperative pain [22]. Many clinical studies have reported that the IO route increases pulpal anesthesia after the IANB injection in patients suffering from endodontic pain [10]. The IO route delivers higher doses but entails cortical perforation and requires specialized instruments [10]. Additionally, postoperative hyperocclusion and infection at the site of perforation, and cardiovascular problems may ensue [23-25]. Another problem with this technique is the friction that is generated during osseous perforation by the detachment of perforation drills off the plastic base.

In addition, necrosis of the bone and perforation of the roots would occur if the drills were rotated at a persistent motion in the bone, leading to an increase in heat generation [26]. Intrapulpal injection is used as the last choice because it involves restricted access cavity opening, which allows local anesthetic solution to backflow [10].

Although infiltration alone suffices the purpose in the maxillary quadrants, the same is not possible for the mandibular region because of the thick cortical plates [27]. However, it can serve as a supplemental mode of anesthesia in IANB [28-32]. The anesthetic success of endodontic treatment for mandibular molars in patients with irreversible pulpitis has been reviewed. A recent systematic review (Seema Yadav 2015 - add in reference) evaluated the anesthetic success rates of IANB alone or along with supplemental infiltration and found that the success rates of infiltration techniques were clinically evident. However, no meta-analysis has been conducted to confirm the results. Another systematic review (Zanjir 2019) used a network meta-analysis to study several pulpal anesthesia strategies, including different anesthetic solutions and injection techniques along with premedications. This systematic review and meta-analysis focuses on the success of primary and supplementary infiltration injection techniques (both buccal and lingual) in patients with irreversible pulpitis in human mandibular molars.

METHODS

The protocol of the systematic review was framed using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist [33]. The prepared protocol was registered with the International Prospective Register of Ongoing Systematic Reviews (PROSPERO) with the number CRD42020210896.

1. Eligibility criteria

The review question was formulated using the PICO

Table 1. Lists of a combination of keywords for electronic database search strategy

Database	Search strategy (2019)	n
Pubmed	("buccal infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" [All fields] OR "buccal infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" AND "human mandibular molars" [All fields] OR "lingual infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" OR "lingual infiltration" AND "irreversible pulpitis" OR "lingual infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "irreversible pulpitis" AND "human mandibular molars" OR "buccal infiltration" AND "irreversible pulpitis" AND "systematic review" OR "lingual infiltration" AND "irreversible pulpitis" AND "systematic review")	80
EbscoHost	("buccal infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" [All fields] OR "buccal infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" AND "human mandibular molars" [All fields] OR "lingual infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" OR "lingual infiltration" AND "irreversible pulpitis" OR "lingual infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "irreversible pulpitis" AND "human mandibular molars" OR "buccal infiltration" AND "irreversible pulpitis" AND "systematic review" OR "lingual infiltration" AND "irreversible pulpitis" AND "systematic review")	58
Scopus	("buccal infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" [All fields] OR "buccal infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" [All fields] OR "buccal infiltration" AND "irreversible pulpitis" AND "human mandibular molars" [All fields] OR "lingual infiltration" AND "symptomatic irreversible pulpitis" [All fields] OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" OR "lingual infiltration" AND "irreversible pulpitis" OR "lingual infiltration" AND "symptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "asymptomatic irreversible pulpitis" AND "human mandibular molars" OR "lingual infiltration" AND "irreversible pulpitis" AND "human mandibular molars" OR "buccal infiltration" AND "irreversible pulpitis" AND "systematic review" OR "lingual infiltration" AND "irreversible pulpitis" AND "systematic review")	57

framework of systematic review, defining the population, intervention, comparison, and outcome. The formulated PICO was "What will be the success of primary and supplemental infiltration injection technique during endodontic treatment in patients with irreversible pulpitis in human mandibular molars?"

2. Information sources and literature search strategy

A comprehensive literature search was performed using the following databases: Medline/Pubmed, Scopus, and Ebsco host, along with an extensive manual search. We used a combination of the following keywords: The systematic search was performed separately by three examiners (AS, NM, and VA) for relevant studies published up to November 2019. Boolean operators "AND" and "OR" were used to build a search string of keywords displayed in Table 1. The collected data were manually searched to identify and exclude duplicates.

3. Inclusion criteria

Three investigators (AS, NM, VA) evaluated the articles with their titles and abstracts that were discovered during the electronic and manual searches. Articles meeting the inclusion criteria were extracted. All the remaining articles were obtained and fully screened independently by the three reviewers to reach a consensus. We included studies published from 1960 to November 2019, clinical studies that were published in English with adult human patients undergoing non-surgical root canal treatment, and studies including patients with irreversible pulpitis in human mandibular molars and utilizing infiltration injection as part of the primary IANB injection or as a supplemental injection given after the failure of primary IANB. The included studies reported on the success of primary and supplemental infiltration injection, either in the form of percentages or numbers.




Fig. 1. PRISMA Flow Diagram of the included studies in systematic review and meta-analysis. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analysis.

4. Exclusion criteria

Studies that evaluated the success of anesthesia utilizing other supplemental techniques, such as IL and IO, and any study reporting the presence of any systemic disease or any effect of medication on anesthetic success, were excluded.

5. Data extraction

From the total database search, the relevant articles were manually searched by three authors (AS, AG, VA). The bibliographies of relevant papers and review articles were also screened to calculate the relevant data. Finally, the data were framed in the form of a PRISMA flowchart, as presented in Figure 1.

Two examiners (AG and VA) extracted the data and analyzed each study using the following parameters: author (year)/country, journal, language, age group, total

patient size, sex, tooth type, sample size, injection type, case/study type, pain evaluation scale, and success rate with the statistical result. Any disagreements between the two authors (AG and VA) were resolved by the third author (AS).

RESULTS

Using the search strategy, both through electronic and manual searches, we identified 195 relevant articles. After excluding duplicates, 128 articles were screened for abstract evaluation, out of which 54 were selected for full-text analysis. Considering the strict inclusion and exclusion criteria, after full-text analysis, only 26 articles were included in the systematic review. Of these, 13 studies were included in the meta-analysis and 28 articles were excluded after full-text reading.

Table 2. Summary of included studies

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
Dianat O, et al. [34]	2020, Clin Oral Invest	Mandibular molars Total patients = 90 Age group = 18-65 years Group 1 - Males = 16 Females = 14 1 st molars - 20 2 nd molars - 10 Group 2 - Males = 14 Females = 16 1 st molars - 18 2 nd molars - 12 Group 3 - Males = 17 Females = 13 1 st molars - 21 2 nd molars - 9	Group 1 - IANB - 1.7 ml of 2% lidocaine with 1:100,000 epinephrine Group 2 IANB + BI - 1.7 ml of 4% articaine with 1:100,000 epinephrine Group 3 IANB + IS + BI - 0.85 ml of 4% articaine with 1:100,000 epinephrine in mesial and distal aspect of dental papilla.	Symptomatic irreversible pulpitis Randomised clinical study	HP VAS EPT	Group 1 = 0.3% Group 2 = 66.6% Group 3 = 80% Group 3 results in significantly higher success rate than group 2 and group 1
Shapiro MR, et al. [35]	2018, J Endod	Total patients = 75 Age group - 39 ± 15 years Mandibular 1 st molars = 36 Males = 51 Females = 47 Mandibular 2 nd molars = 39 Males = 46 Females = 55	1° = IANB (1.7 ml of 4% articaine with 1: 100,000 epinephrine) 2° Group 1 = BI (first molars) - 1) Articaine - 1.7 ml 4% articaine 2) Lidocaine- 1.7 ml 2% lidocaine Group 2 = BI - (second molars) 1) Articaine-1.7 ml 4% articaine 2) Lidocaine-1.7 ml 2% lidocaine.	Irreversible pulpitis Randomised double-blind clinical trial	HP VAS	Group 1 - Articaine = 61% Lidocaine = 66% Group 2 - Articaine = 63% Lidocaine = 32% No significant difference among Group 1 but articaine was significantly more successful for second molars.
Kanaa MD, et al. [28]	2012, J Endod	Mandibular molars, premolars and anteriors Total patients = 182 Age group - 18-66 years Males = 133, Females = 49 Molars = 162, Premolars = 18, Anteriors = 2	1° - IANB (2 ml of 2% lidocaine with 1: 80000 epinephrine) Failure of 1° 2° - Group 1 = PDL injection Group 2 = repeat IANB Group 3 = BI (2ml of 2% articaine HCl with epinephrine 1:100,000) Group 4 = IO	Irreversible pulpitis Randomised clinical trial	EPT	Group 1 = 48% Group 2 = 32% Group 3 = 84% Group 4 = 68% Significant difference among group 1 & 2 and group 3 & 4
Dou L, et al. [36]	2013, Int Endod J	Mandibular molars Total patients = 80 Age group - 27-49 years Group 1 - 30 Males = 13, Females = 17 Group 2- 30 Males = 18 Females = 12	Group 1 IANB + BI - 4 ml of 2% lidocaine with 1: 100,000 epinephrine+ 0.9 ml of 4% articaine with 1: 100,000 epinephrine Group 2 IANB + BLI- 4 ml of 2% lidocaine with 1: 100,000 epinephrine + 0.9 ml of 4% articaine with 1:100,000 epinephrine	Symptomatic irreversible pulpitis Randomised double-blind clinical study	HP VAS	Group 1 - 70% Group 2 - 62.5% No significant difference among groups.

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
Poomi S, et al. [5]	2011, J Endod	Mandibular molars Total patients = 156 Age group - 18-30 years Males = 90, Females = 66	Group 1 = IANB - 1.8 ml with 4% articaine with 1:100,000 adrenaline Group 2 = BI - 1.8 ml with 4% articaine with 1:100,000 adrenaline Group 3 = IANB (control) - 1.8 ml with 2% lidocaine with 1:100,000 adrenaline	Irreversible pulpitis Randomized double-blind clinical trial	HP VAS	Access preparation Group 1 = 75% Group 2 = 69.2% Group 3 = 69.2% Pulp extirpation Group 1 = 69.2% Group 2 = 65.4% Group 3 = 65.4% No statistical significant difference among the study groups.
Matthews R, et al. [32]	2009, J Endod	Mandibular molars or premolars Total patients = 55 Age group - 18-71 years Males = 29 Females = 26 1 st molars - 26 2 nd molars - 23 1 st premolar - 3 2 nd premolar - 3	1st - IANB - 2% lidocaine with 1:100,000 epinephrine Failure of 1st 2nd - BI - 1 cartridge of 4% articaine with 1:100,000 epinephrine	Irreversible pulpitis Prospective study	HP VAS	1 st molars = 58% success 2 nd molars = 48% 2 nd premolar = 100% 1 st premolar = 100% Total anesthetic success with BI - 58%
Parirokh M, et al. [37]	2014, Int Endod J	Mandibular 1 st molars Total patients = 69 Age group- 18-52 years Control group - 36 Females = 21, Males = 15 Test group - 36 Females = 17, Males = 16	Group 1 - IANB (control) 3.6 ml of 2% lidocaine with 1:80000 epinephrine Group 2 - IANB + BI + IL 1.8 ml of 2% lidocaine with 1/80,000 epinephrine + 1.8 ml lidocaine with 1/80,000 epinephrine + IL	Asymptomatic irreversible pulpitis Randomised controlled trial	HP VAS	Group 1 - IANB = 22% Group 2 - IANB + BI + IL = 58% Statistical significant difference among the groups.
Shahi S, et al. [38]	2018, J Endod	Mandibular 1 st molars Total patients = 96 Age group - 18-65 years Males = 42 Females = 54	Group 1 = IANB 1.8 ml 4% articaine with 1:100,000 epinephrine Group 2 = IANB + BI 0.5 ml 4% articaine with 1:100,000 epinephrine Group 3 = IANB + BI + PDL	Symptomatic irreversible pulpitis Consort randomised double-blind clinical study	VAS	Group 1 = 28.1% Group 2 = 65.6% Group 3 = 75% No significant difference
Aggarwal V, et al. [39]	2011, J Endod	Mandibular molars (1 st and 2 nd molars) Total patients = 94 Group 1 - Age group - 24-34 years Males = 12 Females = 11 Group 2 -	Group 1 = IANB (control) 1.8 ml of 2% lidocaine with 1:200,000 epinephrine Group 2 - IANB + BI - 1.8 ml of 2% lidocaine with 1:200,000 epinephrine + 4% articaine with 100,000 epinephrine Group 3 - IANB + BI - 1.8 ml of 2% lidocaine with 1:200,000	Irreversible pulpitis Randomised double-blind study	HP VAS	Group 1 = 39% Group 2 = 54% Group 3 = 62% Group 4 = 45% No significant difference

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
		Age group - 26-35 years Males = 11, Females = 13 Group 3 - Age group - 24-34 years Males = 11 Females = 13 Group 4 - Age group - 25-36 years Males = 11 Females = 12	epinephrine + 1 ml / 30 mg of ketorolac tromethamine Group 4 - IANB + BI - 1.8 ml of 2% lidocaine with 1:200,000 epinephrine + 1 ml / 4 mg of dexamethasone			
Rosenberg PA, et al. [40]	2007, J Endod	Maxillary and Mandibular teeth Total = 48 Age group - 22-58 years Maxillary - 22 Mandibular - 26 Males = 23, Females = 25	Group 1 - (maxillary)- BI A) lidocaine B) articaine Group 2 - (mandibular) 1° IANB of 3.6cc of 2% lidocaine with 1:100,000 epinephrine. Failure 2° BI- a) 1.8cc of 2% lidocaine with 1: 100,000 epinephrine b) 1.8cc of 4% articaine with 100,000 epinephrine	Irreversible pulpitis Randomised double-blind trial.	VAS	Mean score for - 4% articaine = 70.5% For lidocaine = 62.2% No significant difference
Aggarwal V, et al. [30]	2009, J Endod	Mandibular posterior teeth (1 st and 2 nd molars) Total patients = 84 Group 1 - 24-37 years Males = 14 Females = 10 Group 2 - 23-36 years Males = 16 Females = 14 Group 3 - 26-37 years Males = 15 Females = 15	Group 1 - IANB (control) 2% lidocaine with 1:200,000 epinephrine Group 2 - IANB + BI + LI - 2% lidocaine with 1:200,000 epinephrine + 1.7 ml of 2% articaine with 1:200,000 epinephrine + 1.7 ml of 2% articaine with 1:200,000 epinephrine Group 3 - IANB + BI + LI - 2% lidocaine with 1:200,000	Irreversible pulpitis Randomized double-blind study	HP VAS	Group 1 - 33% Group 2 - 67% Group 3 - 47% No significant difference
Fan S, et al. [41]	2009, Oral Surg Med Oral Pathol Oral Radiol Endod	Mandibular first molars Total = 57 Age group - 18-50 years Males = 34, Females = 23	Group 1 - IANB + PDL Group 2 - IANB + BI - 1.7 ml of 4% articaine/hcl with epinephrine 1:100,000 + 0.4 ml of 4% articaine/hcl with epinephrine 1:100,000	Irreversible pulpitis	HP VAS	Group 1 - 83.3% Group 2 - 81.4% No significant difference
Pairokh M, et al. [29]	2010, Oral Surg Oral Med Oral Pathol Oral Radiol Endod	Mandibular 1 st molar teeth Total = 84 Age groups - 18-35 years	Group 1 - IANB 1.8 ml of 2% lidocaine with 1/80,000 epinephrine Group 2 - IANB	Symptomatic irreversible pulpitis Randomized double-blind study	HP VAS	Success Group 1 - 14.8% Group 2 - 39.3% Group 3 - 65.4%

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
		Group 1 - Male = 10 Females = 17 Group 2 - Males = 8 Females = 20 Group 3 - Males = 7 Females = 20	3.6 ml of 2% lidocaine with 1/80,000 epinephrine Group 3 - IANB + BI 1.8 ml of lidocaine with 1/80,000 epinephrine			Group 3 is more significant than group 1
Ashraf H, et al. [42]	2013, J Endod	1 st or 2 nd Mandibular molars Age group – above 20 years Total = 102 Males = 47 Females = 55	1° - IANB 1.8 ml of 2% lidocaine with 100,000 epinephrine or 4% articaine with 100,000 epinephrine respectively to the groups Failure 2° Group 1 - BI - 1.8 ml of 2% lignocaine with 100,000 epinephrine Group 2 - BI - 1.8 ml of 4% articaine with 100,000 epinephrine	Irreversible pulpitis Randomized double- blind Study	HP VAS	Success Group 1 - 29% Group 2 - 71% No significant difference
Ahmad ZH, et al. [43]	2014, J Contemp Dent Pract	Mandibular molars Age groups - 18-40 years Total = 45 Group 1 - Without BI - 6 With BI - 9 Failure with BI - 3 Group 2 - Without BI - 9 With BI - 6 Failure with BI - 2 Group 3 - Without BI - 13 With BI - 2 Failure with BI - 0	1° - IANB 2% lidocaine with 1:100,000 epinephrine Failure 2° Group 1 (1 st molars) BI = 4% articaine with 1:100,000 epinephrine Group 2 (2 nd molars) BI Group 3 (premolars) BI	Symptomatic irreversible pulpitis	HP VAS	Group 1 - 42% Group 2 - 48% Group 3 - 73% No significant difference
Fowler S, et al. [44]	2016, J Endod	Mandibular molars (1 st molars, 2 nd molars, premolars) Total = 221 Age groups - 18-65 years Males - 97 Females - 124	1° - IANB 2% lidocaine with 1:100,000 epinephrine Failure 2° Group 1 (1 st molars) BI = 4% articaine with 1:100,000 epinephrine Group 2 (2 nd molars) - BI	Symptomatic irreversible pulpitis	HP VAS	Group 1 - 42% Group 2 - 48% Group 3 - 73% No significant difference

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
Group 3 (premolars) - BI						
Singla M, et al. [45]	2015, Int Endod J	Mandibular 1 st and 2 nd molars Total = 147	1^o - IANB 1.8 ml of 4% articaine with 1:100,000 epinephrine Failure 2^o Group 1 - BI 1.8 ml of 4% articaine with 1:100,000 epinephrine Group 2 - BI 3.6 ml of 4% articaine with 100,000 epinephrine	Symptomatic irreversible pulpitis Randomized double-blind trial	HP VAS	Group 1 - 62% Group 2 - 64% No significant difference
Rogers BS, et al. [46]	2014, J Endod	Mandibular molars Total - 74 Group 1 1 st molars = 18 2 nd molars = 21 3 rd molars = - Males = 17 Females = 22 Group 2 1 st molars = 17 2 nd molars = 17 3 rd molars = 1 Males = 12 Females = 23	1^o - IANB 1.7 ml of 4% articaine with 1:100,000 epinephrine. Failure 2^o Group 1 - BI 1.7 ml of 4% articaine Group 2 - BI 1.7 ml of 2% lidocaine	Irreversible pulpitis Randomized, double-blind study	HP VAS	Group 1 - 62% Group 2 - 37% Group 1 is more effective than group 2
Yadav M, et al. [47]	2015, J Endod	1 st or 2 nd mandibular molars Total = 150 Group 1 - 75 (each group - 25) Group 2 - 75 (each group - 25)	Group 1 - 4% articaine with 1:100,000 epinephrine Group 1 - a) 0.9 ml BI + 0.9 ml LI Group 1 - b) pre-operative oral medication of ketorolac Group 1 - c) pre-operative oral medication of ketorolac (10 mg) + 0.9 ml (BI + LI) with either articaine or lidocaine Group 2 - 2% lidocaine with 1:80,000 epinephrine Group 2 - a) 0.9 ml BI + 0.9 ml LI Group 2 - b) pre-operative oral medication of ketorolac Group 2 - c) pre-operative oral medication of ketorolac (10 mg) + 0.9 ml (BI+LI) with either articaine or lidocaine	Irreversible pulpitis Randomized controlled, double-blind study	HP-VAS	Group 1 = 64% Group 2 = 32% Group 1c) is more significant with success rate of 76%
Saatchi M, et al. [48]	2016, J Endod	First mandibular molars Total = 100 Age group - 18-53 years Group 1 Males = 13	Group 1 BI- 0.7 ml of 8.4% sodium bicarbonate with 0.3 ml 2% lidocaine containing 1:80,000 epinephrine Group 2 - BI -	Symptomatic irreversible pulpitis Randomized double-blind study	HP VAS	Group 1 = 78% Group 2 = 44% Group 1 is significant than group 2

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
		Females = 37 Group 2 Males = 14 Females = 36	0.7 ml sterile distilled water with 0.3 ml of 2% lidocaine with 1:80,000 epinephrine. - After 15 minutes, all the patients received conventional IANB injection using 3.6 ml 2% lidocaine with 1:80,000 epinephrine			
Monteiro MR, et al. [49]	2015, Int Endod J	Mandibular molars (1 st and 2 nd molars) Total = 50 Age group - above 18 years Group 1 - 1 st molars - 17 2 nd molars - 13 Males = 5 Females = 25 Group 2 1 st molars - 13 2 nd molars - 7 Male = 4 Female = 16	Group 1 - BI 1.8 ml of 4% articaine with 1:100,000 epinephrine + LI of 0.6 ml of articaine solution. Group 2 - IANB 1.8 ml of 2% lidocaine with 1:100,000 epinephrine	Irreversible pulpitis Randomized clinical study	HP-VAS	Group 1 = 40% Group 2 = 10% No significant difference
Ghoddusi J, et al. [50]	2018, Iran Endod J	Mandibular molars Age group - 18-50 years Total = 45 IANB + BLI - 24 GG + BLI- 21	Group 1 - IANB 3.6 ml 2% lidocaine with 100,000 epinephrine Failure 2^o 1) BI 2) LI 1.8 ml of 2% lidocaine with 100,000 epinephrine Group 2 - Gow-Gates 3.6 ml 2% lidocaine with 100,000 epinephrine Failure 2^o 1) BI 2) LI 1.8 ml of 2% lidocaine with 100,000 epinephrine	Symptomatic irreversible pulpitis Randomized double-blind clinical trial	VAS	Group 1 - 42.5% Group 2 - 50% No significant difference
Zain M, et al. [51]	2016, J Coll Physicians Surg Pak	1 st mandibular molars Age group - 18-60 years Total: 156 Group 1 Males - 46 Females - 32 Group 2 Males = 46 Females = 32	Group 1 IANB 1.8 ml of 2% lidocaine with 1:100,000 epinephrine Group 2 BI 4% articaine with 1:100,000 epinephrine	Symptomatic irreversible pulpitis Randomized control trial	HP VAS	Group 1 - 62.8% Group 2 - 76.9% No significant difference
Saberi EA, et al. [52]	2014, Zahedan J Res Med Sci	Mandibular molars Total = 75 Age group - 20-50 years Males = 39	Group 1 - BI 1.8 ml of 4% articaine with 1:100,000 epinephrine Group 2 - BI	Symptomatic irreversible pulpitis Randomized double-blind clinical trial	VAS	Success Group 1 = 68% Group 2 = 52% Group 3 = 64%

(continued)

Author	Year and Journal of publication (language)	Tooth type, No. of subjects and Group wise division	Primary or supplemental injection type	Case / Study type	Pain evaluation	Success rate with statistical result
		Females = 36	1.8 ml of 4% articaine + 0.1 ml of morphine with 1:100,000 epinephrine Group 3 - IANB (control) 1.8 ml of 2% lidocaine with 1:100,000 epinephrine			No statistically difference
Akhlaghi NM, et al. [53]	2016, J Endod	Mandibular molars Total = 40 Age group = 18-65 years Group 1 - Males = 9 Females = 11 Group 2 - Males = 7 Females = 13	1° IANB+BI 1.8 ml of 4% articaine with 1:100,000 epinephrine + 0.9 ml articaine Group 1 - BI 30 mg/ml ketorolac tromethamine Group 2 - BI (control) Normal saline	Acute irreversible pulpitis double-blind randomized clinical trial	HP VAS	Group 1 = 40% Group 2 = 15% Significant difference.
Rajput F, et al. [54]	2015, Pak Oral Dental J	Mandibular first molar Total = 60 Age group = 18-65 years	Group 1 - BI 1.7 ml of 4% articaine with 1:100,000 epinephrine Group 2 - IANB 1.2 ml of 2% lidocaine with 1:200,000 epinephrine.	Symptomatic irreversible pulpitis Randomized clinical trial	HP VAS	Group 1 = 52.4% Group 2 = 47.6% Statistically not significant

BI, buccal infiltrations; EPT, electric pulp tester; HP VAS, Heft Parker Visual Analog Scale; IANB, inferior alveolar nerve block; IL, intraligamentary; IO, intraosseous.

A thorough qualitative and quantitative analysis was carried out for the included studies.

1. Qualitative analysis

1) Study characteristics

In the final interpretation, twenty-six studies [5,28-30, 32,34-54] were involved, which were related to the success of infiltration injection as a primary or supplemental technique in patients with irreversible pulpitis in mandibular molars.

The individual characteristics of each included study are described in Table 2.

2. Outcomes

Based on the individual study criteria, the evaluated results showed successful anesthesia. Either the Heft Parker Visual Analog Scale (HP VAS) or Visual analog scale (VAS) was used to define success in studies when

patients reported no pain, mild/bearable pain, or discomfort during access cavity preparation, except in one study conducted by Kanaa et al. [28], who defined profound anesthesia as 'no' response to the electric pulp tester (EPT).

Dianat et al. [34] used both HP VAS scores and EPT to define successful anesthesia.

3. Risk of bias assessment

The Cochrane Collaboration "Risk of Bias" tool was used to evaluate the methodological quality of the studies involved [55]. The risk of bias was assessed using RevMan 5.3 software. The risks of bias domains evaluated were selection bias (random sequence generation and allocation concealment), performance bias (blinding of participants and personnel), detection bias (blinding of outcome assessment), attrition bias (incomplete outcome data), reporting bias (selective reporting), and other potential sources of bias. Risk of

bias was indicated as low, high, or unclear. Thirteen studies [28-30,32,34,37-39,42-46] that were involved in the meta-analysis were part of the risk of bias assessment. The first analysis involved studies comparing a primary IANB versus a secondary infiltration injection given after the failure of the first injection [28,32,42-46]. The other analysis involved studies comparing the infiltration injections given along a primary IANB versus IANB given alone [29,30,34,37-39].

The evaluation criteria in both analyses were success rates. The remaining 13 studies did not meet the analysis criteria [5,35,36,40,41-54]. The evaluation was performed by two reviewers separately, and any disparity among them was resolved by consensus.

4. Interpretation of studies

A randomized clinical trial by Dianat et al. (2020) [34] compared the anesthetic success of three types of techniques in patients diagnosed with symptomatic irreversible pulpitis in their mandibular molars. Ninety patients were randomly divided into three groups according to anesthetic techniques. The first group received 2% lidocaine IANB, the second group received 4% articaine IANB plus BI, and the third group received IANB plus BI, along with articaine injected intra-septally in both the proximal papillae. Pain was recorded using HP VAS and EPT before and after the anesthetic injection, as well as during access preparation. No or mild pain (pain score of ≤ 54 mm on the HP VAS) on pulpal access was designated as successful. The success rates for groups I, II, and III were 30.33%, 66.66%, and 80.00%, respectively. The mean VAS score after injection was significantly lower in group II and group III than in group I, whereas the EPT score after injection was significantly higher in groups II and III than in group I ($P = 0.016$).

Shapiro et al. (2018) [35] comparatively evaluated the anesthetic efficiency of 4% articaine with 2% lidocaine for supplemental BI in mandibular first molars and second molars with irreversible pulpitis after the failure of IANB. A total of 199 patients diagnosed with irreversible pulpitis

in the mandibular molar were injected with an IANB using 4% articaine. If the patients had a positive response to cold or pain upon initiation of endodontic treatment, the subjects were randomly injected with supplemental 4% articaine or 2% lidocaine. A score of ≤ 54 mm on the HP VAS was considered successful anesthesia. IANB with 4% articaine showed anesthetic success in 25% of the subjects. The success rate for articaine supplemental injection in first molars was 61%, and that for second molars was 63% ($P > 0.05$). Lidocaine resulted in successful anesthesia in 66% of first molars and 32% of second molars ($P = 0.004$). The difference was significant only for the second molars and the results for the first molars were similar for both solutions.

Kanaa et al. (2012) [28], through a randomized clinical study, compared the efficiency of the supplemental repeat inferior alveolar nerve block (rIANB) injected with 2% lidocaine and epinephrine, BI with 4% articaine and epinephrine, and IL or IO injection (both using 2% lidocaine and epinephrine) after the failure of primary IANB to achieve complete analgesia during endodontic treatment of irreversible pulpitis in mandibular teeth. A total of 182 patients were injected with 2 mL of 2% lidocaine with 1:80,000 epinephrine as the primary IANB. One hundred out of 182 patients complained of pain during treatment and were then randomly divided into four groups according to four supplementary injection techniques: rIANB with lidocaine, articaine buccal infiltration (ABI), lidocaine IL, and orlidocaine IO injection. The treatment was initiated only after receiving a negative response to the maximum reading of the electric pulp tester. No pain was regarded as successful anesthesia. ABI and IO exhibited a higher success rate of painless treatment (84% and 68%, respectively) than rIANB or IL supplementary techniques (32% and 48%, respectively). The difference was statistically significant ($P = 0.001$).

Dou et al. (2013) [36] investigated the effect of supplemental lingual infiltration (LI) in mandibular molars after the failure of primary IANB and subsequent BI injection in patients with irreversible pulpitis. Eighty patients were given the standard IANB injection using

4 mL of 2% lidocaine with 1:100,000 epinephrine, after which, patients with marked lower lip numbness were further randomly divided into two groups. In one group, 40 subjects were given supplemental BI (0.9 mL) of 4% articaine with 1:100,000 epinephrine and in the second group of buccal LI (BLI), 40 patients received supplemental BI (0.9 mL) of 4% articaine with 1:100,000 epinephrine followed by LI using the same dose of the solution. Treatment was initiated 15 min after the injection. HP VAS scoring was used to record pain during treatment. No or mild pain on endodontic access and initial instrumentation were considered successful anesthesia. The success rate for BI was found to be 70%, while that of BLI was 62.5%. There was no statistically significant difference between the two groups ($P = 0.478$).

Poomi et al. (2011) [5] studied the anesthetic efficiency of 4% articaine with 1:100,000 epinephrine injected as IANB along with infiltration techniques to anesthetize mandibular molars with irreversible pulpitis. The study design included three groups (two tests and one control). The patients in the test groups were injected with either conventional IANB or 4% articaine with 1:100,000 epinephrine, whereas only conventional IANB injection of 2% lidocaine with 1:100,000 epinephrine was administered to subjects in the control group. The HP VAS was used to assess pain after injection during pulpal access and extirpation. The three groups showed significantly different results, although BI and IANB with 4% articaine exhibited equal efficacy.

Matthews et al. (2009) [32] determined the anesthetic success of secondary BI injection with 4% articaine, with 1:100,000 epinephrine in mandibular posteriors diagnosed with irreversible pulpitis after failed IANB. Fifty-five patients with irreversible pulpitis in a mandibular posterior tooth were injected with an IANB, and the block was considered to have failed if the patient complained of moderate-to-severe pain on pulpal access. Subsequent injections of BI were performed using 4% articaine with 1:100,000 epinephrine. A painless-access biomechanical preparation was designated as successful anesthesia. The resulting anesthetic success was achieved in 58% of the

mandibular posterior teeth. The anesthetic success rate for BI of articaine in the first molar was 58%, the second molar was 48%, the second premolar was 100%, and the first premolar was 100%.

Parirokh et al. (2014) [37] studied the anesthetic action of primary IANB injection for mandibular first molar teeth with irreversible pulpitis, with or without supplemental BI and IL injection. 82 patients with symptomatic irreversible pulpitis were injected with either a combined injection of IL + BI + IANB or standard IANB injection in mandibular first molars. HP VAS pain scores were recorded in response to the cold test at three stages: before the treatment began, during access preparation, and at instrumentation. Anesthetic success was defined as no or mild pain in any of the three stages. In the final stage of treatment, only 69 subjects were included in the study. Traditional IANB was successful in 22% of the patients, while the combination of injections was successful in 58% of the patients. The results were statistically significant ($P = 0.003$).

Shahi et al. (2018) [38] studied the anesthetic efficacy of three types of injection techniques using articaine in mandibular first molars with symptomatic irreversible pulpitis. The three methods were IANB, IANB + IL, and IANB + BI injected before beginning the endodontic treatment. Ninety-six patients were selected and randomly assigned to three groups. Treatment was initiated 20 min after the injection. Each injection used articaine 4% with 1:100,000 epinephrine. Based on the VAS, no or mild pain upon access cavity preparation or initial instrumentation was considered successful. The success rate for IANBs + IL injection was 75%, and that for IANBs + BI was 65.6%. For the standard IANB injected alone, the success rate was only 28.1% ($P < 0.05$).

Aggarwal et al. (2011) [39] evaluated the effects of ketorolac and dexamethasone infiltration injected with conventional IANB. Ninety-four subjects who were actively experiencing pain were injected with a traditional IANB of 2% lidocaine with 1:200,000 epinephrine. Twenty-four patients were set as controls and were not

given any supplemental injections. Another 24 patients were given supplemental BI of 4% articaine with 1:100,000 epinephrine, and another set of 24 patients was given supplemental BI of 1 mL/4 mg of dexamethasone. Twenty-six patients received supplemental BI of 1 mL/30 mg of ketorolac tromethamine, but because the first 2 patients complained of severe pain after ketorolac infiltration and were excluded, the rest of the subjects were given an infiltration of 0.9 mL of 4% articaine before ketorolac. Pulpal access was started 15 minutes after the primary IANB. HP VAS scoring was performed to record the pain. Anesthesia was considered successful if the patient had no pain or mild pain; 39% of the patients in the control group showed successful anesthesia. BI of articaine and articaine plus ketorolac showed significantly high success rates of 54% and 62%, respectively ($P < 0.05$). Supplemental dexamethasone infiltration showed 45% success, which was not significant when compared with the results in the control group.

Rosenberg et al. (2007) [40] studied the anesthetic success of 4% articaine with 1:100,000 epinephrine compared with 2% lidocaine with 1:100,000 epinephrine as a solution for supplemental injections on maxillary and mandibular teeth with irreversible pulpitis. Forty-eight patients required supplemental BI for root canal treatment and were administered either 4% articaine or 2% lidocaine (both with 1:100,000 epinephrine as a vasoconstrictor). VAS scores were evaluated to record pain severity after the injection, and the mean scores were 15.28 for articaine and 19.70 for lidocaine. The mean percentage change in the VAS score was 70.5% and 62.2% for articaine and lidocaine, respectively. However, these differences were not significant in the statistical analyses.

Aggarwal et al. (2009) [30] studied the anesthetic efficiency of the supplemental infiltration technique in 84 subjects with active pain, receiving a standard IANB of 2% lidocaine with 1:200,000 epinephrine. Twenty-four patients were included in the control group and were not injected with supplemental infiltration. Thirty patients received supplemental BI + LI of 2% articaine with

1:200,000 epinephrine and another 30 patients received BI + LI of 2% lidocaine with 1:200,000 epinephrine. After 15 minutes of initial IANB, pulpal access was initiated, and the pain scores were recorded on the HP VAS. Patients who reported 'none' or 'mild' pain were considered to have achieved successful anesthesia. Supplemental BI + LI with lidocaine or articaine improved the success rate from 33% (of the primary IANB alone) to 47% and 67%, respectively. Additionally, the success rate with 4% articaine was significantly higher than that with 2% lidocaine ($P < 0.05$).

Fan et al. (2009) [41] evaluated the anesthetic efficiency of IANB + BI and IANB + PDL articaine injections in patients with irreversible pulpitis in the mandibular first molars. Fifty-seven subjects randomly received traditional IANB injections of 1.7 mL 4% articaine/HCl with 1:100,000 epinephrine with either BI or PDL injections containing 0.4 mL articaine/HCl with 1:100,000 epinephrine. The patients reported the severity of the pain experienced during injections and access cavity preparation on the HP VAS. Anesthetic success was seen in 81.48% of IANB + BI cases compared with 83.33% for IANB + PDL. The observed differences between the two groups was significant ($P < 0.05$).

Parirokh et al. (2010) [29] evaluated the success of IANB combined with BI for mandibular molars with irreversible pulpitis. Eighty-four patients were randomly divided into three groups ($n = 28$ in each group). Lidocaine 2% with 1:80,000 epinephrine was used as the solution of choice. Patients received 1.8 mL of anesthesia as IANB in first group, 3.6 mL as IANB in the second group, and 1.8 mL of IANB and 1.8 mL of BI in the third group. VAS was used to record pain intensity before anesthesia and discomfort experienced before and during root canal access. Groups I, II, and III showed success rates of 14.8%, 39.3%, and 65.4%, respectively, with the third group showing significantly better anesthetic success ($P < 0.05$).

Ashraf et al. (2013) [42] conducted a comparative analysis of the anesthetic efficacy of BI as a supplemental injection using articaine or lidocaine with an IANB. A

total of 125 mandibular molars with irreversible pulpitis were injected with an IANB using either 2% lidocaine or 4% articaine (both with 1:100,000 epinephrine). A total of 102 patients complained of moderate-to-severe pain on pulpal access and were given supplemental BI injections using the same anesthetic solution as that of primary IANB. No pain or minimal pain during instrumentation was deemed successful. Lidocaine was successful in only 29% of the teeth, whereas articaine was successful in 71% ($P < 0.001$), and this difference was significant in the statistical analysis.

Ahmad et al. (2014) [43] investigated the effect of IANB using 4% articaine and 2% lidocaine (with different concentrations of vasoconstrictor). Supplemental BI was provided if the IANB failed. Forty-five patients with irreversible pulpitis of a mandibular posterior tooth were divided into three groups according to the anesthetic injection: Group I received 2% lidocaine with 1:200,000 epinephrine, Group II received 2% lidocaine with 1:80,000 epinephrine, and group III was injected with 4% articaine with 1:100,000 epinephrine. Supplemental BI was administered to patients who complained of pain during root canal therapy. The percentage of successful anesthesia and failed anesthesia were calculated and tabulated using the VAS. Most subjects (87%) injected with articaine reported satisfactory anesthesia with IANB alone. Only 2 (13%) subjects were given an additional infiltration injection, and none of the patients failed to obtain complete anesthesia with articaine. On the other hand, 40% of the subjects receiving lidocaine injections obtained satisfactory results with 1:200,000 epinephrine and 60% were satisfied with 1:80,000 epinephrine.

Fowler et al. (2016) [44] investigated the anesthetic effect of primary IANB and supplemental articaine BI injected after the failure of primary injection in the posterior teeth of patients presenting with symptomatic irreversible pulpitis. A total of 375 subjects were selected and injected with 2% lidocaine with 1:100,000 epinephrine as IANB. After confirming lip numbness, treatment was initiated. A total of 204 patients complained of moderate-to-severe pain, following which,

a supplementary BI injection was administered using 4% articaine with 1:100,000 epinephrine. Successful anesthesia was confirmed on no or mild pain on access preparation and instrumentation (VAS rating of 0 or ≤ 54). Twenty five percent of first and second molars and 39% of premolars achieved successful anesthesia. For the supplemental articaine infiltration, 42% of first molars, 48% of second molars, and 73% of premolars were successfully anesthetized. No significant differences were found between the molars ($P = 0.3450$), but there was a significant difference between the percentages of premolars and molars, that is, second molars with premolars ($P = 0.0411$) and first molars with premolars ($P = 0.0132$).

Singla et al. (2015) [45] comparatively studied the anesthetic success of two volumes (1.8 mL versus 3.6 mL) of 4% articaine with 1:100,000 epinephrine solution injected as a supplemental BI after the failure of primary IANB in teeth with symptomatic irreversible pulpitis. A total of 234 subjects received IANB with 1.8 mL of 4% articaine with 1:100,000 epinephrine. Pain scoring was performed using the HP VAS, and no or weak/mild pain during the root canal procedures was deemed satisfactory (HP VAS score < 55 mm). Total 147 patients experienced 'moderate-to-severe' pain (HP VAS score ≥ 55 mm), so they were further randomized into 2 groups and received BI with either 1.8mL or 3.6mL of the same anaesthetic solution. Primary articaine IANB was successful in 37% of the cases and supplementary BI with 1.8 mL and 3.6 mL volumes succeeded in 62% and 64% cases, respectively. However, these differences were not statistically significant.

Rogers et al. (2014) [46] evaluated and compared 4% articaine with 2% lidocaine for their anesthetic effect when injected as supplemental BI injections after failed IANB in mandibular molars with irreversible pulpitis. One hundred emergency subjects were selected and injected with 1.7 mL of 4% articaine with 1:100,000 epinephrine as an IANB. Seventy-four patients were randomly injected with 4% articaine or 2% lidocaine as supplemental BI after failed IANB. No pain or mild pain

according to the VAS score during the procedure was considered successful. IANB was successful in 26% of the cases. The success rate of articaine BI (62%) was significantly higher than that of lidocaine BI (37%).

Yadav et al. (2015) [47] conducted a randomized study on 150 emergency patients with mandibular first and/or second molars diagnosed with irreversible pulpitis. The patients were divided into two groups: Group 1 was injected with 4% articaine and 1:100,000 epinephrine and group II with 2% lidocaine and 1:80,000 epinephrine. Furthermore, the two groups had three sub-groups each, with 25 patients in each sub-group: BI and LI with articaine and lidocaine, respectively; preoperative oral medication of ketorolac and preoperative oral medication of ketorolac followed by BI and LI with articaine and lidocaine, respectively. All patients with profound lip numbness were considered to have successful anesthesia. No pain or mild pain during root canal procedures was deemed successful analgesia. Articaine IANB with infiltration and oral ketorolac premedication had a significantly high success rate of 76%. Articaine and lidocaine IANB plus infiltration injection without premedication showed success rates of 64% and 32%, respectively ($P < 0.05$).

Saatchi et al. (2016) [48] evaluated the anesthetic effect of BI using sodium bicarbonate for mandibular first molars in patients with symptomatic irreversible pulpitis. A total of 100 patients were randomly administered a BI injection. In group 1, 0.7 mL of 8.4% sodium bicarbonate with 0.3 mL of 2% lidocaine containing 1:80,000 epinephrine was injected, and in group 2, 0.7 mL sterile distilled water was injected with the same dose of lidocaine as in the other group. Following the BI injection, all patients were administered standard IANB using 3.6 mL of the same lidocaine solution. Anesthetic success was determined only when lip numbness was present. HP VAS scores of no or mild pain during access cavity preparation or initial instrumentation were deemed to have a satisfactory anesthetic effect. The BI of sodium bicarbonate was successful in 78% of the cases, whereas for BI without bicarbonate, only 44% of the cases were

considered successful ($P < 0.001$).

Monteiro et al. (2015) [49] investigated the anesthesia achieved by IANB of 1.8 mL 2% lidocaine compared to a BI of 1.8 mL 4% articaine, both with 1:100,000 epinephrine, in patients with symptomatic irreversible pulpitis. Fifty subjects were randomly divided into 30 patients receiving articaine injections and 20 receiving lidocaine injections. No pain after a single injection (IANB or BI) or one supplemental injection needed for emergency root canal treatment was considered successful anesthesia. Articaine showed a higher success rate of 40% compared to only 10% with lidocaine. No significant differences were observed between the groups. Ghoddsi et al. (2018) [50] compared the anesthetic effect of IANB and the Gow-Gates technique in mandibular molars with symptomatic irreversible pulpitis. Eighty patients randomly received either IANB or Gow-Gates injection using 2% lidocaine with 1:100,000 epinephrine. Afterward, pain during caries removal and pulpal access were recorded, following which patients were randomly injected with either BI or LI. A total of 45 patients reported pain in both groups. The VAS was used to evaluate pain. The success rates of the Gow-Gates and IANB techniques were 50% and 42.5%, respectively, with no significant difference ($P = 0.562$). There was a significant reduction in pain severity in all subgroups, including supplementary infiltrations ($P < 0.05$). A significantly greater reduction in pain severity was observed with LI given after IANB, as compared to LI after Gow-Gates injection ($P < 0.05$).

Zain et al. (2016) [51] conducted a randomized control trial to study the pulpal anesthetic effect in mandibular first molars by injecting either 4% articaine BI or 2% lidocaine IANB. A total of 156 patients diagnosed with irreversible pulpitis in the first molar were randomly divided into two groups. Group 1 patients were injected with 4% articaine BI and group 2 patients were administered IANB of 2% lidocaine. HP VAS scoring was performed to report pain during the root canal treatment procedures. In 76.9% of the cases, 4% Articaine BI was successful, whereas lidocaine IANB was

successful in 62.8%. The differences were not statistically significant.

Saberi et al. (2014) [52] conducted a randomized double-blind clinical trial to identify and compare the anesthetic effects of articaine and articaine plus morphine for BI in mandibular posterior teeth diagnosed with irreversible pulpitis. Seventy-five patients were divided into 3 groups and randomly injected with either a BI of 4% articaine or articaine morphine (both with 1:100,000 epinephrine) or traditional IANB of 2% lidocaine with 1:800,000 epinephrine. Pain scoring on the VAS was performed before and after the anesthetic injection and during endodontic access. The success rates of articaine, articaine-morphine, and lidocaine were 68%, 52%, and 64%, respectively. No statistically significant difference was found among the groups.

Akhlaghi et al. (2016) [53] investigated the effect of ketorolac BI on the success of IANB injection in patients with acute irreversible pulpitis. The HP VAS was used to evaluate pain. Forty subjects were injected with a standard IANB injection and a BI of 4% articaine with 1:100,000 epinephrine. The patients were then divided into two groups, with the first test group receiving BI of 30 mg/mL ketorolac, and the second control group receiving BI of normal saline. After observing a negative response to the EPT twice, treatment was initiated. HP VAS scoring of pain was performed. Successful anesthesia was defined as no or minimal pain during the procedure, without any need for supplemental injections. Adding BI significantly increased the success rate from 15% in the control group to 40% in the test group ($P < 0.05$).

Rajput et al. (2015) [54] comparatively studied the anesthetic efficacy of 4% articaine with 1:100,000 epinephrine for BI and 2% lidocaine with 1:200,000 epinephrine for IANB in patients with irreversible pulpitis in mandibular first molars. Sixty patients were randomly divided into two groups. Group 1 received a BI of 1.7 ml of 4% articaine with 1:100,000 epinephrine and group 2 patients were given standard IANB with 1.8 ml of 2% lidocaine with 1:200,000 epinephrine. Anesthetic success

was defined as no or minimal pain on endodontic access, according to the VAS score. Twenty two patients (52.4%) achieved successful anesthesia with 4% articaine, while 20 patients (47.6%) had pain reduction in the 2% lignocaine group. However, the observed results were not statistically significant ($P = 0.220$).

5. Quality assessment


Figures 2 and 3 show the assessment for risk of bias categories. The categories analyzed included biases such as selection bias, performance bias, detection bias, attrition bias, reporting bias, and assessment of randomization and blinding followed in each individual study. Of all the included studies, six depicted a low risk of bias in almost every domain [29,30,39,42-46]. One study reported a high risk of bias in the selection [34]. Another study showed a high risk of performance and detection bias and a low risk in every bias domain [28]. Two studies were in the low risk of bias category, except for a few domains that were under unclear risk, as randomization and blinding was not possible in these two studies [32,44]. Two other studies had a low risk of bias, except that the performance bias of both studies is unclear [37,38]. All individual studies were assessed qualitatively for each category.

6. Quantitative analysis

- 1) Meta-analysis
- 2) Forest plot calculations

The meta-analysis was performed using RevMan 5.3, and two different meta-analyses were conducted for the included studies. The first analysis involved studies comparing a primary IANB versus a secondary infiltration injection (given after the failure of the first injection) [28,34,42-46].


The other analyses involved studies comparing the infiltration injections given along a primary IANB versus IANB given alone [29,30,34,37-39]. The effect size in both analyses was the success rate. The remaining studies were excluded from meta-analysis. Figure 2 shows the



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Fig. 2. Forest plot representation of meta-analysis performed on the studies evaluating the secondary infiltration injections.



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Fig. 3. Forest plot representation of meta-analysis performed on the studies evaluating primary IANB + infiltration versus primary IANB. IANB, inferior alveolar nerve block.

forest plot representation of the random-effect meta-analysis performed on studies evaluating secondary infiltration injections. The combined risk ratio (RR) was 1.88 (95% CI: 1.49, 2.37) in favor of the secondary infiltration injections. The statistical heterogeneity among the studies was 77%. The high statistical heterogeneity can be attributed to the varying confidence intervals (CIs) of different studies. In contrast, the forest plot analysis (Fig. 3) of studies comparing primary IANB + infiltration

versus primary IANB alone had a low heterogeneity (0%). The included studies had similar RRs and the combined RR was 1.84 (95% CI: 1.44, 2.34). The results of both forest plots suggest that supplemental infiltration injections, given along with a primary IANB, provide better success rates. Moreover, when the primary IANB fails, supplementary infiltration injections improve success rates.




Fig. 4. Publication bias quantified using Egger regression test for studies evaluating the secondary infiltration injections.




Fig. 5. Publication bias quantified using Egger regression test for studies evaluating primary IANB + infiltration versus primary IANB. IANB, inferior alveolar nerve block.




Fig. 6. L'Abbe plots for studies comparing secondary infiltrations after failed IANB. IANB, inferior alveolar nerve block.




Fig. 7. L'Abbe plots for studies comparing primary IANB + infiltration versus primary IANB. IANB, inferior alveolar nerve block.

7. Funnel plots

To assess publication bias, funnel plots were created. The extent of publication bias was based on visual examination of funnel plot symmetry. The funnel plot of studies evaluating secondary infiltration injections had a low publication bias. The extent of publication bias was

quantified using the Egger regression test (Fig. 4), which showed a low publication bias ($P = 0.281$). Studies comparing primary IANB + infiltration with primary IANB alone also had low publication bias. The Egger regression test (Fig. 5) showed a low extent of publication bias ($P = 0.6044$).




Fig. 8. Required information size and the combined information size of the included studies evaluating secondary infiltration.

8. L'Abbe plots

L'Abbe plots were generated for both analyses, and are shown in Figures 6 and 7. The risks of success were plotted against each other. The risk of success for supplementary infiltration was plotted on the y-axis and the risk of success of IANB was plotted on the x-axis. The solid diagonal line represents the studies in which the risk of success did not differ between the groups. Studies above this line have a higher risk of success in the supplementary infiltration group than in the IANB group. The dashed line shows the estimated effects based on the analysis model. Figure 6 shows the L'Abbe plots for studies comparing the secondary infiltrations given after the failure of IANB. All the studies were above the solid line along the dashed line, showing that the risk of success was higher in the infiltration injection group. The scattered studies also represented the level of heterogeneity among the studies. The L'Abbe plots (Fig. 7) of studies comparing primary IANB + infiltration versus primary IANB also showed that the estimated effect was in favor of the IANB + infiltration group. Both

studies were near the dashed line, suggesting minimal statistical heterogeneity among them.

9. Trial sequential analysis (TSA)

TSA calculations were performed and graphs were plotted with a type I error of 5%. The required information size (RIS) was calculated using a random-effects model to evaluate RRs with a power of 80%. The incidence in the control group was maintained at 30%. The incidence in the intervention arm was maintained at 60% (relative risk reduction of -100%). For the group evaluating secondary infiltrations given after failure of a primary IANB, the RIS was calculated using a heterogeneity correction of 79%, as suggested by the model variance. The high statistical heterogeneity among the included studies increased the RIS to 587. However, the combined information size of the included studies surpassed that of the RIS, with cumulative $Z > 1.96$ (Fig. 8). TSA for the group comparing primary IANB + infiltration versus primary IANB alone has a low heterogeneity and the heterogeneity correction was maintained at 0%, as suggested by the model variance. The RIS was calculated




Fig. 9. Required information size and the combined information size of the included studies evaluating primary IANB + infiltration versus primary IANB. IANB, inferior alveolar nerve block.

using a random model with an incidence of 30% and 55% in the control and intervention groups, respectively (relative reduction of -83.33%). The RIS for this group was estimated to be 123 subjects (Fig. 9). The information size was greater than that of the RIS, with $Z > 1.96$, confirming the beneficial effect of primary IANB + infiltration. It should be noted that an increase in heterogeneity led to an increased RIS in the first analysis.

DISCUSSION

Complete analgesia in the mandibular posterior region during dental procedures is often regarded as difficult to achieve. IANB is the most standardized and frequently used injection technique for inducing anesthesia in this region. However, the reduced efficacy of IANB can be attributed to factors such as the thickness of the cortical bone, overlying soft tissue, and accessory innervations [28,56,57]. Nerve block success in an un-inflamed pulp was reported to be 70%, as compared to only 30% in

patients diagnosed with irreversible pulpitis [11,14, 58-63]. Reports suggest that the failure rate of local anesthesia in patients with irreversible pulpitis is eight times higher than the failure rate in normal patients.

The current review summarizes the anesthetic value of infiltration injection techniques, either applied as a primary or supplemental injection in patients diagnosed with irreversible pulpitis in human mandibular molars.

Other techniques, such as IO and IL, are often applied to raise anesthetic efficacy in difficult situations. However, the infiltration technique is a simple, easy, and comfortable technique for achieving desirable anesthetic levels [42].

Recently, the BI technique has been used in the mandible in various studies. Some of the authors have shown that the use of only BI or BLI can produce profound anesthesia in subjects with 32% to 67% when administered with lidocaine, and 57% to 92% when administered with articaine, even when the IANB is not administered [31,64-66]. A recent meta-analysis (involving studies with healthy and inflamed pulp) revealed that

articaine was 3.8 times more potent in an infiltration injection than lidocaine [67].

In the mandible, a BLI anesthetic injection containing articaine was suggested for sufficient anesthesia in the anterior tooth pulps [68]. This BLI technique showed significant improvement in the anesthetic action in healthy mandibular incisor pulps than when labial infiltration used solo [68]. However, in mandibular molars, a part of the supplementary LI remains debatable. It has been observed that in the first mandibular molars, LI was less efficacious than BI in achieving anesthesia in healthy subjects [69]. Meechan et al. [16] observed no difference in anesthetic efficacy for healthy mandibular molars between BI alone and BLIs. The effectiveness of a supplementary LI in mandibular molars when IANB and BI have already been administered, especially in irreversible pulpitis cases, where it has not been examined. Only a few studies have searched supplemental LI with IANB + BI and have not found any significant results in the success rates of anesthesia [30,36]. More such studies are required to determine an appropriate and reliable outcome of LI as a supplemental injection technique to primary IANB + BI.

Yadav [70] compared and evaluated the anesthetic success rates of the IANB injection technique alone or injected along with supplemental infiltration technique to achieve profound anesthesia in mandibular posteriors with irreversible pulpitis. This review included studies published up to 2014, and no meta-analysis was conducted for the included studies.

Infiltration injection has been tested in the field of endodontics, both as a primary or supplemental technique. Various clinical studies have focused on the success of infiltration techniques (BI/LI), either as a primary or supplemental injection in subjects with mandibular posteriors suffering from irreversible pulpitis. Studies showing the anesthetic success of supplemental infiltration injected after the failure of primary IANB in teeth with irreversible pulpitis exhibited a high success rate of 40% to 73% [34,42,43,45,46], apart from the study conducted by Kanaa et al. [28], which gave an even

higher success rate of 84%. This high success rate could be attributed to the injection of a larger volume of anesthetic solution with BI. Lastly, studies in which the IANB plus infiltration technique was given as a primary injection in subjects suffering from irreversible pulpitis showed a success rate of 45% to 67% [29,30,34,37-39].

The current review only focused on the success of pulpal anesthetic injection techniques, regardless of the different anesthetic solutions and concentrations used in the included studies. Therefore, only the success of the infiltration technique on anesthetic efficacy was considered.

Risk of bias assessment was performed for the included studies in the current review, and it was found to be low in every category that was tested. The qualitative results of the review were quantified through two types of meta-analyses based on specific characteristics of the study. The first analysis involved a comparison of the anesthetic success rates of primary IANB versus secondary infiltration injections administered after failure of primary IANB. This meta-analysis, when represented as a forest plot, revealed 87% statistical heterogeneity among the studies that further increased the calculated RIS to 778. However, the calculated RIS was still less than the combined information size, which reaffirmed the reliability of our meta-analysis, which could be considered sufficient to prove that secondary infiltration injection is better than primary IANB administered alone. In contrast, the second meta-analysis involved a comparison of success rates of the infiltration injections given as part of the primary injection alongside IANB versus IANB administered alone. The forest plot revealed a low heterogeneity of 0%, and eventually, the calculation of RIS was performed at 123, which was again less than the combined information size. One limitation of the second meta-analysis quantification is that one of the studies included by Parirokh et al. [37] used a combined success rate of the IL that was injected along with IANB and BI, which could have influenced the final result of the analysis to certain limits. Overall, these results suggest better success rates of infiltration techniques, given along

with a primary IANB.

The qualitative and quantitative results of the current review suggest the incorporation of a simple and easy infiltration injection technique as a primary injection along with IANB in routine clinical practice.

CONCLUSION

Based on the qualitative and quantitative analyses, we can conclude that the infiltration technique, either as a primary injection or as a supplementary injection, when given after the failure of primary IANB increases the overall anesthetic efficacy.

AUTHOR ORCID*s*

Alpa Gupta: <https://orcid.org/0000-0003-2834-0608>

Aarushi Sahai: <https://orcid.org/0000-0002-7519-9339>

Vivek Aggarwal: <https://orcid.org/0000-0002-9792-4554>

Namrata Mehta: <https://orcid.org/0000-0002-4836-524X>

Dax Abraham: <https://orcid.org/0000-0002-6870-6689>

Sucheta Jala: <https://orcid.org/0000-0001-9680-3147>

Arundeeep Singh: <https://orcid.org/0000-0003-2342-9486>

AUTHOR CONTRIBUTIONS

Alpa Gupta: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing - original draft, Writing - review & editing

Aarushi Sahai: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resources, Software, Supervision, Validation, Visualization, Writing - original draft

Vivek Aggarwal: Data curation, Formal analysis, Methodology, Project administration, Software, Supervision, Validation, Visualization, Writing - review & editing

Namrata Mehta: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Software, Validation, Visualization, Writing - original draft, Writing - review & editing

Dax Abraham: Conceptualization, Formal analysis, Investigation, Project administration, Resources, Supervision, Validation, Writing - review & editing

Sucheta Jala: Data curation, Formal analysis, Software, Supervision, Validation, Writing - review & editing

Arundeeep Singh: Formal analysis, Project administration, Resources, Supervision, Validation, Visualization

DECLARATIONS OF INTERESTS: The authors have no conflicts of interest to declare.

REFERENCES

1. Brännström M, Johnson G, Nordenvall KJ. Transmission and control of dentinal pain: resin impregnation for the desensitization of dentin. *J Am Dent Assoc* 1979; 99: 612-8.
2. Milgrom P, Coldwell SE, Getz T, Weinstein P, Ramsay DS. Four dimensions of fear of dental injections. *J Am Dent Assoc* 1997; 128: 756-62.
3. Glickman GN. AAE consensus conference on diagnostic terminology: background and perspectives. *J Endod* 2009; 35: 1619-20.
4. Modaresi J, Dianat O, Mozayeni MA. The efficacy comparison of ibuprofen, acetaminophen-codeine, and placebo premedication therapy on the depth of anaesthesia during treatment of inflamed teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006; 102: 399-403.
5. Poomi S, Veniashok B, Senthilkumar AD, Indira R, Ramachandran S. Anaesthetic efficacy of four percent articaine for pulpal anaesthesia by using inferior alveolar nerve block and buccal infiltration techniques in patients with irreversible pulpitis: a prospective randomized double-blind clinical trial. *J Endod* 2011; 37: 1603-7.
6. Malamed SF. *Handbook of local anaesthesia* 4th ed. St Louis, Mosby. 1997, pp 223-5.
7. Potočník I, Bajrović F. Failure of inferior alveolar nerve block in endodontics. *Endod Dent Traumatol* 1999; 15: 247-51.
8. Rood JP. Some anatomical and physiological causes of failure to achieve mandibular analgesia. *Br J Oral Surg* 1977; 15: 75-82.
9. Parente SA, Anderson RW, Herman WW, Kimbrough WF, Weller RN. Anaesthetic efficacy of the supplemental intraosseous injection for teeth with irreversible pulpitis. *J Endod* 1998; 24: 826-8.
10. Hargreaves KM, Keiser K. Local anaesthetic failure in endodontics: mechanisms and management. *Endod Topics*

- 2002; 1: 26-39.
11. Tortamano IP, Siviero M, Costa CG, Buscariolo IA, Armonia PL. A comparison of the anaesthetic efficacy of articaine and lidocaine in patients with irreversible pulpitis. *J Endod* 2009; 35: 165-8.
 12. Chaudhary P, Martenson ME, Baumann TK. Vanilloid receptor expression and capsaicin excitation of rat dental primary afferent neurons. *J Dent Res* 2001; 80: 1518-23.
 13. Stenholm E, Bongehiellm U, Ahlquist M, Fried K. VR1-and VRL-1-like immunoreactivity in normal and injured trigeminal dental primary sensory neurons of the rat. *Acta Odontol Scand* 2002; 60: 72-9.
 14. Claffey E, Reader A, Nusstein J, Beck M, Weaver J. Anaesthetic efficacy of articaine for inferior alveolar nerve blocks in patients with irreversible pulpitis. *J Endod* 2004; 30: 568-71.
 15. Kaufman E, Weinstein P, Milgrom P. Difficulties in achieving local anaesthesia. *J Am Dent Assoc* 1984; 108: 205-8.
 16. Meechan JG. Supplementary routes to local anaesthesia. *Int Endod J* 2002; 35: 885-96.
 17. Moore PA, Cuddy MA, Cooke MR, Sokolowski CJ. Periodontal ligament and intraosseous anaesthetic injection techniques: alternatives to mandibular nerve blocks. *J Am Dent Assoc* 2011; 142: 13S-8S.
 18. Jastak JT, Yagiela JA, Donaldson D. Local anaesthesia of the oral cavity. 2nd ed. Vancouver, WB Saunders Company. 1995.
 19. Smith GN, Walton RE. Periodontal ligament injection: distribution of injected solutions. *Oral Surg Oral Med Oral Pathol* 1983; 55: 232-8.
 20. Walton RE, Abbott BJ. Periodontal ligament injection: a clinical evaluation. *J Am Dent Assoc* 1981; 103: 571-5.
 21. Tagger M, Tagger E, Sarnat H. Periodontal ligament injection: spread of the solution in the dog. *J Endod* 1994; 20: 283-7.
 22. McLean ME, Wayman BE, Mayhew RB. Duration of anaesthesia using the periodontal ligament injection: a comparison of bupivacaine to lidocaine. *Anesth Pain Control Dent* 1992; 1: 207-13.
 23. Dunbar D, Reader A, Nist R, Beck M, Meyers WJ. Anaesthetic efficacy of the intraosseous injection after an inferior alveolar nerve block. *J Endod* 1996; 22: 481-6.
 24. Guglielmo A, Reader A, Nist R, Beck M, Weaver J. Anaesthetic efficacy and heart rate effects of the supplemental intraosseous injection of 2% mepivacaine with 1: 20,000 levonordefrin. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999; 87: 284-93.
 25. Coggins R, Reader A, Nist R, Beck M, Meyers WJ. Anaesthetic efficacy of the intraosseous injection in maxillary and mandibular teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1996; 81: 634-41.
 26. Woodmansey KF, White RK, He J. Osteonecrosis related to intraosseous anaesthesia: report of a case. *J Endod* 2009; 35: 288-91.
 27. Haas DA, Harper DG, Saso MA, Young ER. Comparison of articaine and prilocaine anaesthesia by infiltration in maxillary and mandibular arches. *Anesth Prog* 1990; 37: 230-7.
 28. Kanaa MD, Whitworth JM, Meechan JG. A prospective randomized trial of different supplementary local anaesthetic techniques after failure of inferior alveolar nerve block in patients with irreversible pulpitis in mandibular teeth. *J Endod* 2012; 38: 421-5.
 29. Parirokh M, Satvati SA, Sharifi R, Rekabi AR, Gorjestani H, Nakhaee N, et al. Efficacy of combining a buccal infiltration with an inferior alveolar nerve block for mandibular molars with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 109: 468-73.
 30. Aggarwal V, Jain A, Kabi D. Anaesthetic efficacy of supplemental buccal and lingual infiltrations of articaine and lidocaine after an inferior alveolar nerve block in patients with irreversible pulpitis. *J Endod* 2009; 35: 925-9.
 31. Jung IY, Kim JH, Kim ES, Lee CY, Lee SJ. An evaluation of buccal infiltrations and inferior alveolar nerve blocks in pulpal anaesthesia for mandibular first molars. *J Endod* 2008; 34: 11-3.
 32. Matthews R, Drum M, Reader A, Nusstein J, Beck M. Articaine for supplemental buccal mandibular infiltration anaesthesia in patients with irreversible pulpitis when the inferior alveolar nerve block fails. *J Endod* 2009; 35: 343-6.
 33. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*

- 2009; 6: e1000097.
34. Dianat O, Mozayeni MA, Layeghnejad MK, Shojaeian S. The efficacy of supplemental intraseptal and buccal infiltration anaesthesia in mandibular molars of patients with symptomatic irreversible pulpitis. *Clin Oral Investig* 2020; 24: 1281-6.
 35. Shapiro MR, McDonald NJ, Gardner RJ, Peters MC, Botero TM. Efficacy of articaine versus lidocaine in supplemental infiltration for mandibular first versus second molars with irreversible pulpitis: a prospective, randomized, double-blind clinical trial. *J Endod* 2018; 44: 523-8.
 36. Dou L, Luo J, Yang D. Anaesthetic efficacy of supplemental lingual infiltration of mandibular molars after inferior alveolar nerve block plus buccal infiltration in patients with irreversible pulpitis. *Int Endod J* 2013; 46: 660-5.
 37. Parirokh M, Sadr S, Nakhaee N, Abbott PV, Askarifard S. Efficacy of supplementary buccal infiltrations and intraligamentary injections to inferior alveolar nerve blocks in mandibular first molars with asymptomatic irreversible pulpitis: a randomized controlled trial. *Int Endod J* 2014; 47: 926-33.
 38. Shahi S, Rahimi S, Yavari HR, Ghasemi N, Ahmadi F. Success rate of 3 injection methods with articaine for mandibular first molars with symptomatic irreversible pulpitis: a CONSORT randomized double-blind clinical trial. *J Endod* 2018; 44: 1462-6.
 39. Aggarwal V, Singla M, Rizvi A, Miglani S. Comparative evaluation of local infiltration of articaine, articaine plus ketorolac, and dexamethasone on anaesthetic efficacy of inferior alveolar nerve block with lidocaine in patients with irreversible pulpitis. *J Endod* 2011; 37: 445-9.
 40. Rosenberg PA, Amin KG, Zibari Y, Lin LM. Comparison of 4% articaine with 1: 100,000 epinephrine and 2% lidocaine with 1: 100,000 epinephrine when used as a supplemental anaesthetic. *J Endod* 2007; 33: 403-5.
 41. Fan S, Chen WL, Pan CB, Huang ZQ, Xian MQ, Yang ZH, et al. Anaesthetic efficacy of inferior alveolar nerve block plus buccal infiltration or periodontal ligament injections with articaine in patients with irreversible pulpitis in the mandibular first molar. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009; 108: e89-93.
 42. Ashraf H, Kazem M, Dianat O, Noghrehkar F. Efficacy of articaine versus lidocaine in block and infiltration anaesthesia administered in teeth with irreversible pulpitis: a prospective, randomized, double-blind study. *J Endod* 2013; 39: 6-10.
 43. Ahmad ZH, Ravikumar H, Karale R, Preethanath RS, Sukumaran A. Study of the anaesthetic efficacy of inferior alveolar nerve block using articaine in irreversible pulpitis. *J Contemp Dent Pract* 2014; 15: 71-4.
 44. Fowler S, Drum M, Reader A, Beck M. Anaesthetic success of an inferior alveolar nerve block and supplemental articaine buccal infiltration for molars and premolars in patients with symptomatic irreversible pulpitis. *J Endod* 2016; 42: 390-2.
 45. Singla M, Subbiya A, Aggarwal V, Vivekanandhan P, Yadav S, Yadav H, et al. Comparison of the anaesthetic efficacy of different volumes of 4% articaine (1.8 and 3.6 mL) as supplemental buccal infiltration after failed inferior alveolar nerve block. *Int Endod J* 2015; 48: 103-8.
 46. Rogers BS, Botero TM, McDonald NJ, Gardner RJ, Peters MC. Efficacy of articaine versus lidocaine as a supplemental buccal infiltration in mandibular molars with irreversible pulpitis: a prospective, randomized, double-blind study. *J Endod* 2014; 40: 753-8.
 47. Yadav M, Grewal MS, Grewal S, Deshwal P. Comparison of preoperative oral ketorolac on anaesthetic efficacy of inferior alveolar nerve block and buccal and lingual infiltration with articaine and lidocaine in patients with irreversible pulpitis: a prospective, randomized, controlled, double-blind study. *J Endod* 2015; 41: 1773-7.
 48. Saatchi M, Farhad AR, Shenasa N, Haghghi SK. Effect of sodium bicarbonate buccal infiltration on the success of inferior alveolar nerve block in mandibular first molars with symptomatic irreversible pulpitis: a prospective, randomized double-blind study. *J Endod* 2016; 42: 1458-61.
 49. Monteiro MR, Groppo FC, Haiter-Neto F, Volpato MC, Almeida JF. 4% articaine buccal infiltration versus 2% lidocaine inferior alveolar nerve block for emergency root canal treatment in mandibular molars with irreversible pulpitis: a randomized clinical study. *Int Endod J* 2015; 48: 145-52.
 50. Ghoddsu J, Zarrabi MH, Daneshvar F, Naghavi N.

- Efficacy of IANB and Gow-Gates techniques in mandibular molars with symptomatic irreversible pulpitis: a prospective randomized double-blind clinical study. *Iran Endod J* 2018; 13: 143-8.
51. Zain M, Rehman Khattak SU, Sikandar H, Shah SA, Fayyaz. Comparison of anaesthetic efficacy of 4% articaine primary buccal infiltration versus 2% lidocaine inferior alveolar nerve block in symptomatic mandibular first molar teeth. *J Coll Physicians Surg Pak* 2016; 26: 4-8.
 52. Saberi EA, Farhadmollashahi N, Aliabadi T. A comparison of the anaesthesia efficacy of articaine and articaine plus morphine for buccal infiltration in mandibular posterior teeth with irreversible pulpitis. *Zahedan J Res Med Sci* 2014; 16: 26-30.
 53. Akhlaghi NM, Hormozi B, Abbott PV, Khalilak Z. Efficacy of ketorolac buccal infiltrations and inferior alveolar nerve blocks in patients with irreversible pulpitis: a prospective, double-blind, randomized clinical trial. *J Endod* 2016; 42: 691-5.
 54. Rajput F, Katpar S, Shaikh MI, Khatoon S. Evaluation of anaesthetic efficacy of 4% articaine as buccal infiltration vs 2% lidocaine as ianb in the mandibular 1st molar with irreversible pulpitis. *Pak Oral Dent J* 2015; 35: 125-8.
 55. Deeks JJ, Higgins J, Altman DG, Green S. *Cochrane handbook for systematic reviews of interventions version 5.1.0 (updated March 2011)*. The Cochrane Collaboration. 2011.
 56. Ogle OE, Mahjoubi G. Local anaesthesia: agents, techniques, and complications. *Dent Clin North Am* 2012; 56: 133-48.
 57. Malamed SF. Is the mandibular nerve block passé? *J Am Dent Assoc* 2011; 142: 3S-7S.
 58. Cohen HP, Cha BY, Spångberg LS. Endodontic anaesthesia in mandibular molars: a clinical study. *J Endod* 1993; 19: 370-3.
 59. Goldberg S, Reader A, Drum M, Nusstein J, Beck M. Comparison of the anaesthetic efficacy of the conventional inferior alveolar, Gow-Gates, and Vazirani-Akinosi techniques. *J Endod* 2008; 34: 1306-11.
 60. Childers M, Reader A, Nist R, Beck M, Meyers WJ. Anaesthetic efficacy of the periodontal ligament injection after an inferior alveolar nerve block. *J Endod* 1996; 22: 317-20.
 61. Dagher FB, Yared GM, Machtou P. An evaluation of 2% lidocaine with different concentrations of epinephrine for inferior alveolar nerve block. *J Endod* 1997; 23: 178-80.
 62. Vreeland DL, Reader A, Beck M, Meyers W, Weaver J. An evaluation of volumes and concentrations of lidocaine in human inferior alveolar nerve block. *J Endod* 1989; 15: 6-12.
 63. Mikesell P, Nusstein J, Reader A, Beck M, Weaver J. A comparison of articaine and lidocaine for inferior alveolar nerve blocks. *J Endod* 2005; 31: 265-70.
 64. Meechan JG, Kanaa MD, Corbett IP, Steen IN, Whitworth JM. Pulpal anaesthesia for mandibular permanent first molar teeth: a double-blind randomized cross-over trial comparing buccal and buccal plus lingual infiltration injections in volunteers. *Int Endod J* 2006; 39: 764-9.
 65. Kanaa MD, Whitworth JM, Corbett IP, Meechan JG. Articaine and lidocaine mandibular buccal infiltration anaesthesia: a prospective randomized double-blind cross-over study. *J Endod* 2006; 32: 296-8.
 66. Robertson D, Nusstein J, Reader A, Beck M, McCartney M. The anaesthetic efficacy of articaine in buccal infiltration of mandibular posterior teeth. *J Am Dent Assoc* 2007; 138: 1104-12.
 67. Brandt RG, Anderson PF, McDonald NJ, Sohn W, Peters MC. The pulpal anaesthetic efficacy of articaine versus lidocaine in dentistry: a meta-analysis. *J Am Dent Assoc* 2011; 142: 493-504.
 68. Nuzum FM, Drum M, Nusstein J, Reader A, Beck M. Anaesthetic efficacy of articaine for combination labial plus lingual infiltrations versus labial infiltration in the mandibular lateral incisor. *J Endod* 2010; 36: 952-6.
 69. Meechan JG, Jaber AA, Corbett IP, Whitworth JM. Buccal versus lingual articaine infiltration for mandibular tooth anaesthesia: a randomized controlled trial. *Int Endod J* 2011; 44: 676-81.
 70. Yadav S. Anaesthetic success of supplemental infiltration in mandibular molars with irreversible pulpitis: a systematic review. *J Conserv Dent* 2015; 18: 182-6.