

Angiotensin II and Renal Tubular Ion Transport

Patricia Valles¹, Jan Wysocki², and Daniel Batlle^{2,*}

¹Area de Fisiopatología, Departamento de Patología, Facultad de Ciencias Médicas, Universidad Nacional de Cuyo, Mendoza, Argentina; ²Division of Nephrology and Hypertension, Department of Medicine, Northwestern University, The Feinberg School of Medicine, Chicago

E-mail: d-batlle@northwestern.edu

Received June 30, 2005; Revised August 22, 2005; Accepted August 23, 2005; Published August 29, 2005

Angiotensin II, a potent vasoconstrictor, also participates in the regulation of renal sodium and water excretion, not only via a myriad of effects on renal hemodynamics, glomerular filtration rate, and regulation of aldosterone secretion, but also via direct effects on renal tubule transport. In addition, angiotensin II stimulates H⁺ secretion and HCO₃⁻ reabsorption in both proximal and distal tubules and regulates H⁺-ATPase activity in intercalated cells of the collecting tubule.

Different results regarding the effect of angiotensin II on bicarbonate reabsorption and proton secretion have been reported at the functional level, depending on the angiotensin II concentration and tubule segment studied. It is likely that interstitial angiotensin II is more important in regulating hemodynamic and transport functions than circulating angiotensin II. In proximal tubules, stimulation of bicarbonate reabsorption, Na⁺/H⁺-exchange, and Na⁺/HCO₃⁻ cotransport has been found using low concentrations (<10⁻⁹ M), while inhibition of bicarbonate reabsorption has been documented using concentrations higher than 10⁻⁸ M.

Evidence for the regulation of H⁺-ATPase activity *in vivo* and *in vitro* by trafficking/exocytosis has been provided. An additional level of H⁺-ATPase regulation via protein synthesis may be important as well. Recently, we have shown that both aldosterone and angiotensin II provide such a mechanism of regulation *in vivo* at the level of the medullary collecting tubule. Interestingly, in this part of the nephron, the effects of aldosterone and angiotensin II are not sodium dependent, whereas in the cortical collecting duct, both aldosterone and angiotensin II, by contrast, affect H⁺ secretion by sodium-dependent mechanisms.

KEYWORDS: Angiotensin II -Bicarbonate transport- Vacuolar H⁺-ATPase- Proximal convoluted tubule (PCT) Cortical collecting ducts (CCDs)- Medullary collecting ducts (MCDs)- ATP6V₁ B1- ATP6V₀A4.

INTRODUCTION

A main function of angiotensin II is the maintenance of extracellular Na^+ and volume homeostasis by controlling renal Na^+ reabsorption through the stimulation of Na^+ -reabsorptive transport via the Na^+/H^+ antiporter, the epithelial Na^+ -channel, and Na^+/K^+ -ATPases[1,2,3,4]. Angiotensin II is a potent vasoconstrictor and, in addition to its effects on Na^+ -homeostasis and blood pressure regulation, plays a role in the regulation of acid-base balance. Volume depletion leads to activation of angiotensin II, often resulting in metabolic alkalosis, suggesting a close association between the regulation of renal Na^+ and HCO_3^- handling. Some of these effects are direct and others indirect via activation of aldosterone secretion by the adrenal gland. In the kidney, there is increasing evidence of a regulation of bicarbonate reabsorption and vacuolar H^+ -ATPase activity by angiotensin II, as will be reviewed below.

ANGIOTENSIN II

Angiotensin II is mainly formed by renin-mediated activation of circulating angiotensinogen, originating from the liver and after activation of angiotensin I by angiotensin converting enzyme (ACE). Intrarenal angiotensin II is regulated by several complex processes, involving formation of both systemically delivered and intrarenally formed substrate, as well as receptor-mediated internalization. Angiotensinogen, renin, and ACEs are also present along the nephron. Angiotensinogen is secreted into the lumen and activated there, forming a paracrine renin-angiotensin II system along the nephron[5,6]. The complex and extensive actions of angiotensin II on renal function are due to the widespread distribution of angiotensin II in nephron segments as well as in the vasculature and interstitium. Angiotensin II signals through two main receptor subtypes, AT_1 and AT_2 [1,7,8]. Most actions of angiotensin II are primarily attributed to the AT_1 receptor because of its multiple vascular and transport effects. The AT_1 receptors have been localized in glomerular podocyte cells, proximal tubule brush border and basolateral membranes, thick ascending limb epithelia, distal tubule collecting ducts, macula densa cells, and interstitial cells of the renal medulla[9,10]. Due to the extensive localization of AT_1 receptors in luminal as well as basolateral membranes of proximal and distal nephron segments, interest is growing on the effects of angiotensin II at the tubular level.

Stimulation of the AT_1 receptor leads to increased Na^+ , HCO_3^- , and fluid reabsorption along the nephron, which together with the induced vasoconstriction, result in a rise of blood pressure[1,11,12,13]. For these short-term effects, concentrations of angiotensin II in the picomolar range are needed, which is in contrast to higher nanomolar concentrations needed to activate the AT_2 receptor[1,14]. AT_2 receptor immunostaining in adult kidneys has been found in proximal tubules, collecting ducts, and renal vasculature. The AT_2 receptor seems to counterbalance the effects of the AT_1 receptor on electrolyte transport and blood pressure[14]. The very high abundance of AT_1 receptors in the kidney is associated with tissue angiotensin II levels that are much higher than plasma levels[15,16].

The intrarenal content of angiotensin II is not distributed in a homogenous manner, but is compartmentalized in both a regional and segmental manner[7]. Angiotensin II levels in the deep medulla are much higher than in the cortex and may be involved in the regulation of medullary hemodynamics[17,18]. Moreover, receptor binding studies have demonstrated that angiotensin II receptor density is much higher in the medulla than in the cortex[19]. Within the cortex, there is a substantial compartmentalization of intrarenal angiotensin II, with levels in the renal interstitial fluid and in proximal tubule fluid being much higher than that which can be accounted for by the circulating levels[20]. It is likely that interstitial angiotensin II is more important in regulating hemodynamic and transport functions than circulating angiotensin II. Angiotensin II in interstitial fluid might be responsible for maintaining tone of the pre- and postglomerular vessels and influencing tubular transport function by acting on angiotensin II receptors on the basolateral membranes of the tubular cells[21].

The finding that angiotensin II concentrations in perfused tubules were similar to those measured in nonperfused tubules indicates that the tubular fluid angiotensin II concentration is not derived from glomerular filtration[22]. Moreover, it has also been demonstrated that extracellular fluid volume

expansion reduced plasma angiotensin II levels, but did not reduce intratubular angiotensin II concentrations[23].

Because angiotensin I and angiotensinogen are present in the proximal tubule fluid, intratubular angiotensin II could be formed from precursors secreted into the tubular lumen. The localization of intrarenal angiotensinogen mRNA and protein in proximal tubule cells indicates that proximal tubule cells provide the substrate for intratubular and interstitial angiotensin I and angiotensin II, and may not be dependent on circulating angiotensin II[24]. The angiotensinogen produced in proximal tubules is secreted into the tubular lumen. Angiotensin I may be formed in the tubular lumen by renin or renin-like enzymes[25]. In addition, cultured proximal cells have been found to produce renin and contain renin mRNA, thus suggesting that constitutive renin secretion occurs in proximal tubule cells[26]. When angiotensin I is formed, the conversion to angiotensin II occurs by the action of ACE in the proximal tubule brush border[27]. A novel ACE homologue, ACE 2, is also abundantly present in the proximal tubule and may play a counter-regulatory role by preventing angiotensin II accumulation[28]. ACE has been measured in proximal and distal tubular fluid, but is more abundant in proximal tubule fluid[29]. It seems that angiotensin II would have to be continuously produced or added to the proximal tubule fluid in view of the abundance of angiotensinases found in the brush border of proximal tubules[30]. It remains uncertain how much of the peptides are formed intracellularly and how much are formed in the tubular fluid.

The augmentation of intrarenal angiotensin II that occurs during physiologic increases in angiotensin levels includes a component of AT_1 receptor-mediated accumulation of angiotensin II into an intracellular compartment. Angiotensin peptides, ACE, and angiotensin II are present in renal endosomes[31]. The role of the internalized angiotensin II remains unclear. Angiotensin II might exert cytosolic actions. In addition, it has been suggested that angiotensin II migrates to the nucleus to exert transcriptional effects[32,33]. It has been shown that endocytosis of the angiotensin II/ AT_1 receptor coupled to the activation of signal transduction pathways leads to enhanced sodium transport[34].

Previous studies support a role for luminal angiotensin II in regulating reabsorptive function in distal nephron and collecting ducts as well as in proximal tubule segments[12,35,36]. A direct relationship between urinary angiotensinogen excretion rates and renal angiotensin II has been demonstrated[37].

REGULATION OF BICARBONATE REABSORPTION BY ANGIOTENSIN II IN PROXIMAL TUBULES

Several studies have shown, either in animal or cell culture models, that angiotensin II is a potent activator of bicarbonate reabsorption and proton secretory mechanisms along the nephron, thus stimulating overall bicarbonate reabsorption. The target transport mechanisms include the Na^+/HCO_3^- cotransporter and Na^+/H^+ exchanger (at least NHE-3) in the proximal tubule, and the vacuolar H^+ -ATPase in the proximal tubule, the distal convoluted tubule, and intercalated cells of the cortical collecting duct[12,38,39,40,41].

The proximal tubule is the primary site of HCO_3^- reabsorption in the mammalian kidney. Angiotensin II plays an important role in modulating proximal tubule transport. Systemic infusion of angiotensin II, at doses that do not affect glomerular hemodynamics or blood pressure, result in a significant increase in proximal tubule transport. Both *in vivo* and *in vitro* microperfusion studies demonstrate that physiological concentrations of peritubular angiotensin II increase proximal tubule transport[42].

Recent studies have provided evidence for the local production and luminal secretion of angiotensin II by the proximal convoluted tubule (PCT), a mechanism that maintains a luminal concentration of angiotensin II along the proximal tubule higher than that measured in the systemic circulation[22,43]. The effect of luminal angiotensin II at low concentrations ($<10^{-9}$ M) increases HCO_3^- reabsorption at 1.5-fold in the early segment of the proximal tubule[39]. This effect of angiotensin II is, at least in part, due to the stimulation of luminal Na^+/H^+ exchanger[12]. In this regard, angiotensin II, at the same low concentration (10^{-9} M), has been shown to stimulate both an intracellular acid extruder (luminal Na^+/H^+ exchange) and

an intracellular acid loader (basolateral $\text{Na}^+/\text{HCO}_3^-$ cotransport), thus greatly increasing transepithelial HCO_3^- reabsorption in proximal tubules, with only a modest change in pHi [12].

Conversely, inhibition of bicarbonate reabsorption has been shown with angiotensin II concentrations higher than 10^{-8} M[44]. Stimulation of bicarbonate reabsorption, observed at low angiotensin II concentrations, has been attributable to inhibition of the production of cAMP via the G-protein G_i [45,46]. On the other hand, intracellular Ca^{2+} serves as a second messenger in the inhibitory effect of high concentrations of angiotensin II on bicarbonate reabsorption in the proximal tubule of the kidney[47].

Angiotensin II has potent effects on ammonia production and secretion rates by the proximal tubule. It has been demonstrated that isolated perfused mouse S3 proximal tubule segments produce and secrete ammonia. Angiotensin II added to the luminal fluid, enhanced renal ammonia production and secretion in S3 segments from acidotic mice[48].

REGULATION OF BICARBONATE REABSORPTION BY ANGIOTENSIN II IN DISTAL TUBULE

Because relatively high concentrations of angiotensin II are present in the tubule fluid, it appears likely that this hormone might also have distinct effects on transport beyond the proximal tubule[13,40,49]. In distal tubules, the role of angiotensin II on acid secretion and bicarbonate reabsorption has been less extensively studied. Most of the previous studies evaluating electrolyte transport in the cortical distal tubule (DT) were based on either free-flow micropuncture or by perfusion of the whole distal tubule[50]. Indeed, *in vivo* microperfusion studies of the distal tubule have shown that an intravenous infusion of angiotensin II enhances distal tubule net bicarbonate reabsorption in the normal rat[40]. Since the distal tubule of cortical nephrons consists of several segments with distinct cell types and transport modes[51], angiotensin II transport modulation was examined separately on microperfused early and late distal tubule segments. By separate perfusion of early and late segments of cortical distal tubule *in vivo* (early segments corresponding to distal convoluted tubule and late segments to connecting segment and initial collecting duct), it has been shown that angiotensin II at concentration of 10^{-11} M stimulated early distal bicarbonate reabsorption by stimulating the Na^+/H^+ exchanger. The late distal effect of angiotensin II (10^{-11} M) was mostly dependent on amiloride-sensitive Na^+ reabsorption, i.e., on sodium channels[36]. In luminal perfusion studies, it has been shown that angiotensin II stimulates Na^+/H^+ exchange in both early and late distal tubule via activation of AT_1 receptors, and the vacuolar H^+ -ATPase in the late distal tubule. In all situations where the pharmacological action was evaluated, stimulation occurred through AT_1 receptors[35,49,52]. Consistent with these observations, Geibel et al. recently found that angiotensin II (10^{-8} M) activates H^+ -ATPase in intercalated cells via an AT_1 receptor which stimulates trafficking of the H^+ -ATPase into the membrane[38]. In addition to acute effects of angiotensin II on vacuolar H^+ -ATPase activity, it has been shown that chronic AT_1 blockade of *in vivo* microperfused distal tubules from remaining nephrons in 2/3 nephrectomized rats reduced bicarbonate reabsorption in close association with the reduction of synthesis and insertion of apical H^+ -ATPase[53,54].

AT_1 receptors couple intracellularly mainly to phospholipase C, Ca^{2+} , and the PKC pathway[55]. Thus, it could be speculated that most stimulatory effects of angiotensin II on the vacuolar H^+ -ATPase should be mediated via these signal pathways, but also through activation of tyrosine kinases[56]. In a subclone of MDCK cells exhibiting some characteristics of intercalated cells, angiotensin II was found to increase intracellular Ca^{2+} and stimulate vacuolar H^+ -ATPase activity. Moreover, chelating intracellular Ca^{2+} abolished the stimulatory effect of angiotensin II[57].

In contrast to the studies of H^+ -ATPase activation, one study using permeabilized CCD segments showed a specific dose-dependent inhibitory effect on the vacuolar H^+ -ATPase in the presence of angiotensin II[58]. Weiner et al. demonstrated that angiotensin II acts through a basolateral AT_1 receptor to stimulate outer CCD luminal alkalization in β -intercalated cells[59]. Moreover, angiotensin II has been found to inhibit HCO_3^- reabsorption to a similar extent in the absence and presence of luminal furosemide, which indicates that inhibition occurs independent from the effects on net NaCl absorption and is likely mediated through regulation of apical membrane Na^+/H^+ exchange in medullary thick

ascending limb (MTAL). Thus, the inhibition of Na^+/H^+ exchange activity by angiotensin II in MTAL is an effect opposite to its physiological action to stimulate apical Na^+/H^+ exchange and HCO_3^- reabsorption in the proximal tubule and early distal tubule. Angiotensin II inhibited HCO_3^- reabsorption in MTAL in a concentration of 10^{-8} M, similar to angiotensin II levels measured in the renal medulla *in vivo*. This inhibition is mediated by the cytochrome P-450–dependent signaling pathway. Tyrosine kinase pathways also appear to play a role in the angiotensin II–induced transport inhibition[60].

A reduction in bicarbonate reabsorption after application of angiotensin II was also found in perfused rat outer medullary collecting ducts[61]. This effect of angiotensin II was not observed in the presence of the AT_1 receptor blocker, candesartan. Although angiotensin II upregulates H^+ secretion in proximal and distal tubules, angiotensin II may reduce H^+ secretion in other segments when perfused *in vitro* depending on angiotensin II concentration and tubule segments studied. An inhibitory effect on distal H^+ excretion, however, would be unexpected in light of the stimulatory effect of angiotensin II on vacuolar H^+ -ATPase (see below).

VACUOLAR H^+ -ATPASE

Vacuolar H^+ -ATPases are ubiquitous multisubunit complexes mediating ATP-driven vectorial transport of protons across membranes. The vacuolar H^+ -ATPases are hetero-oligomeric complexes composed of 13 polypeptides arranged into a soluble cytoplasmic V_1 domain composed of 8 subunit types, which is the site of ATPase activity, and a membrane-associated V_0 domain composed of 5 subunit types that includes the proton translocation pathway[62,63,64,65]. Little is known about the expression patterns of these subunits in the kidney with the exception of the B1 and “a4” subunits. The B1 isoform is a part of the peripheral V_1 domain[62]. Two isoforms of the B subunit exist: the B2 isoform (ATP6V1B2) is ubiquitously expressed[63,64], whereas, in the kidney, the B1 subunit is present in intercalated cells of the late distal tubule, connecting segment, and cortical and medullary collecting ducts[65]. The B1 subunit is also expressed in the cochlea and endolymphatic sac[66]. The “a4” subunit (ATP6V0A4) is one of four isoforms of the “a” subunit which form part of the membrane bound V_0 sector. All four “a” subunits are expressed in the kidney as detected by Northern blot and RT-PCR analysis[67,68,69]. The “a4” isoform is expressed in the kidney (proximal tubule, loop of Henle, and all subtypes of intercalated cells along the late distal tubule, connecting segment, and entire collecting duct)[70].

Mutations in either the B1 or the 116-kDa subunit (a4 isoform) cause inherited forms of distal renal tubular acidosis (dRTA). Whereas mutations in the B1 subunit (ATP6B1) cause dRTA with sensorineural deafness[66], mutations in the 116-kD accessory subunit (ATP6N1B) affect predominantly the kidney[71,72].

ANGIOTENSIN II REGULATION OF VACUOLAR H^+ -ATPASE ACTIVITY BY TRAFFICKING/EXOCYTOSIS

Regulation of H^+ -ATPase activity includes mechanisms such as trafficking from submembraneous pools, regulation by cytosolic activator and inhibitor proteins, or gene expression and protein expression[73]. There is evidence for regulation of H^+ -ATPase activity *in vivo* and *in vitro* by trafficking/exocytosis.

The B1 subunit interacts directly with filamentous actin, which could allow for the trafficking[74]. Recently, in cell culture, evidence has been found for the involvement of the vesicular SNARE docking proteins in the fusion of H^+ -ATPase containing vesicles with the plasma membrane[75,76]. Moreover, the B1 subunit of the kidney H^+ -V-ATPase has been identified as a PDZ domain-binding protein that colocalized with NHE-RF in renal type B-intercalated cells, providing a target for regulation. NHE-RF has been shown to link other transport proteins to the cytoskeleton and may have a role in the regulation of H^+ -ATPase localization or trafficking in B intercalated cells[77].

Regulation of a4 isoform subunit by trafficking, but not protein expression, with a marked redistribution of the subunit in intercalated cells in response to acid-base changes and dietary potassium

intake, has recently been reported[70]. These findings are in good agreement with earlier reports showing trafficking of other H^+ -ATPase subunits in response to metabolic acidosis or alkalosis or dietary electrolyte intake [78]. In intercalated cells of isolated mouse cortical collecting ducts, angiotensin II (10^{-8} M) stimulation of vacuolar H^+ -ATPase has been shown to be prevented by colchicine, an agent that disrupts the microtubular network. This suggests that vacuolar H^+ -ATPases, some of their subunits, or other stimulatory proteins may be trafficked to the membrane[38].

The electrogenic vacuolar H^+ -ATPase, by the translocation of H^+ ions across membranes, renders the cell interior negative with respect to the exterior. This process generates both a transmembrane potential and a chemical gradient, resulting in a self-limiting activity of the vacuolar H^+ -ATPase[79]. In many intracellular organelles and at cell plasma membranes, a parallel Cl^- conductance provides an electric shunt, compensating for the positive charge transferred by the pump and, thus, dissipating the electrical gradient[80,81,82,83].

The presence of this parallel Cl^- conductance has been found in most intracellular organelles that are acidified by vacuolar H^+ -ATPases, such as lysosomes[84] and endosomes[80,81,85]. In the kidney, it is clear that vacuolar H^+ -ATPase-dependent acidification is Cl^- dependent in endosomal fractions[80,81,85,86,87,88] and in brush border membrane vesicles[89,90]. The chloride dependence on plasma membrane H^+ -ATPase, however, is a complex issue because of the expression of numerous other conductances or electrogenic transporters in apical and basolateral membrane domains[79]. In isolated rat proximal tubules, vacuolar H^+ -ATPase-dependent H^+ extrusion was reduced after preincubation in Cl^- -free media[91]. In another study, Na^+ -independent, H^+ -ion secretion was stimulated by physiological concentration of angiotensin II. This angiotensin II stimulation of H^+ -ATPase-dependent proton extrusion was demonstrated to occur via a process involving a Cl^- -dependent insertion of vesicles into the brush border membrane[41]. In addition, in mouse proximal tubules, the insertion of vacuolar H^+ -ATPase containing vesicles was delayed in the absence of chloride[92]. Notwithstanding, there has not been convincing evidence of the role of chloride in apical H^+ -ATPase-dependent proton secretion in intercalated cells. The *in vivo* microperfusion of late distal tubule, however, showed a reduction of electrogenic H^+ secretion by the Cl^- channel blocker, NPPB, indicating coupling between H^+ -ATPase and Cl^- transport[93].

Bicarbonate reabsorption is independent of the presence of luminal chloride in isolated rabbit medullary collecting duct, although it is completely abolished in the absence of basolateral chloride, presumably due to impairment of peritubular bicarbonate exit in exchange for Cl^- via the Cl^-/HCO_3^- exchanger AE1[94].

ANGIOTENSIN II AND EXPRESSION OF VACUOLAR H^+ -ATPase SUBUNITS *IN VIVO*

It has been generally believed that H^+ -ATPase is regulated to only a minor extent by transcriptional and translational mechanisms and very little is known about the *in vivo* regulation of the various subunits of the H^+ -ATPase. Recently, we examined the expression of H^+ -ATPase subunits, B1 and a4, in an animal model of selective aldosterone deficiency (adrenalectomized [ADX] rats maintained on glucocorticoid replacement). The effect of exogenous angiotensin II on the *in vivo* expression of the B1 H^+ -ATPase subunit was also studied[95].

Our data showed an increase in the relative protein abundance of B1 subunit isoform of the H^+ -ATPase after chronic infusion of angiotensin II in the renal medulla of ADX rats. This increase in H^+ -ATPase expression by angiotensin II was limited to intercalated cells in the medullary collecting tubule. The antibody to the kidney isoform of the B1 subunit stained intensely on apical membrane of intercalated cells from outer medullary collecting tubules and initial inner medullary collecting tubules in ADX infused with angiotensin II. By contrast, we did not find significant differences in B1 subunit of H^+ -ATPase protein in cortical membrane preparations. The lack of difference in B1 protein expression renal cortex could reflect, in part, the predominance of renal proximal tubules, which do not express B1. In contrast to the B1 subunit, the expression of the a4 subunit was not altered significantly by either ADX or

exogenous angiotensin II addition. Interestingly, the $\alpha 4$ subunit is found in both the early segments of proximal tubules and the intercalated cells in the kidney. The B1 subunit of the kidney H^+ -V-ATPase, however, is expressed only in the intercalated cells of the kidney collecting tubules. Thus, one could speculate that regulation of the B1 subunit could be a mechanism by which hormones, such as aldosterone and angiotensin II, specifically control H^+ -ATPase synthesis in intercalated cells of the distal nephron.

Related to the different effect of angiotensin II and ADX on medullary and cortical collecting tubules, it has been previously demonstrated that medullary angiotensin II levels are higher than the cortical levels in normal rats and increase further in angiotensin II-infused hypertensive rats[96]. In addition, increased uptake and trafficking of angiotensin II into renal endosomes is mediated by AT_1 receptors in renal cortex after chronic angiotensin II infusion[97]. This could suggest that in renal cortex, B1 regulation is under control of a mechanism(s) other than protein synthesis. Regulation of $\alpha 4$ and B1 subunits by trafficking, but not protein expression, has been demonstrated under conditions of acid base and electrolyte changes[38,41,67]. It should be noted that different results regarding the effect of angiotensin II on H^+ secretion have been reported at the functional level depending on the angiotensin II concentration and tubule segments studied[12,54].

On a functional level, aldosterone stimulates electrogenic Na^+ reabsorption through the luminal epithelial Na^+ channel resulting in lumen-negative potential that leads to increase the electrical driving force for H^+ secretion in the cortical collecting duct[98,99]. A direct action of aldosterone on H^+ secretion has been shown in rat medullary collecting duct. The effect of aldosterone was independent of Na^+ in medullary collecting ducts perfused *in vitro* pointing also to mechanisms in addition to changes in the electrical driving force[100]. Different sensitivities of vacuolar H^+ -ATPase enzymatic activity in distinct collecting duct segments have been described, even though aldosterone stimulated H^+ -ATPase-dependent bicarbonate reabsorption in all collecting duct segments[101,102].

Our findings in ADX rats suggest that the removal of aldosterone results in a decrease in the relative abundance and expression of B1 H^+ -ATPase in the medullary, but not the cortical, collecting tubule. These results suggest that aldosterone affects vacuolar H^+ -ATPase activity in the collecting duct through different mechanisms that may be specific for tubular segment and cell type. Our data suggest that some of aldosterone actions may require protein synthesis of H^+ -ATPase subunits.

We also have found that aldosterone increases $\alpha 4$ H^+ -ATPase synthesis in a cell line of the collecting duct[103]. Importantly, in microdissected cortical and medullary collecting duct segments from ADX rats, a pharmacological dose of aldosterone increased enzyme H^+ -ATPase activity irrespective of potassium level[104]. A recent study showed nongenomic targeting of the $\alpha 4$ subunit of vacuolar H^+ -ATPase to the apical membrane by aldosterone[105].

An additional level of regulation via protein synthesis, however, may be important as well. Our results suggest that both aldosterone and angiotensin II provide such a mechanism *in vivo* at the level of the medullary collecting tubule. Interestingly, in this part of the nephron, the effects of aldosterone and angiotensin II are not sodium dependent whereas in the cortical collecting duct both aldosterone and angiotensin II, by contrast, affect H^+ secretion by sodium-dependent mechanisms[98,100].

In summary, angiotensin II and ADX affect H^+ -ATPase B1-subunit protein expression in medullary collecting ducts[95]. Selective aldosterone deficiency created by adrenalectomy with glucocorticoid replacement resulted in the downregulation of the expression of H^+ -ATPase B1 subunit in medullary collecting ducts. Angiotensin II increased the expression of the B1 subunit of H^+ -ATPase in the medullary collecting duct and thus may upregulate H^+ secretion in this tubule segment. Such an action may be responsible, in part, for an increase in H^+ secretion, independent of aldosterone[95]. The expression of the $\alpha 4$ subunit, by contrast, was not altered by either ADX or exogenous angiotensin II[95]. Angiotensin II exerts significant direct effects on acid-base balance, separate from those related to its stimulatory effect on aldosterone secretion. The renal vacuolar H^+ -ATPase is an important target for angiotensin II action particularly in renal medullary collecting tubules.

REFERENCES

1. Allen, A.M., Zhuo, J., and Mendelsohn, F.A. (2000) Localization and function of angiotensin AT1 receptors. *Am. J. Hypertens.* **13**, 31S–38S.
2. Inagami, T. (1999) Molecular biology and signaling of angiotensin receptors: an overview. *J. Am. Soc. Nephrol.* **10**, S2–7.
3. Verrey, F. (1999) Early aldosterone action: toward filling the gap between transcription and transport. *Am. J. Physiol.* **277**, F319–327.
4. Verrey, F., Hummler, E., Schild, L., and Rossier, B. (2000) Control of Na⁺ transport by aldosterone. In *The Kidney: Physiology and Pathophysiology*. 3rd ed. Seldin, D.W. and Giebisch, G., Eds. Lippincott Williams & Wilkins, Philadelphia. pp. 1441–1471.
5. Quan, A. and Baum, M. (1996) Endogenous production of angiotensin II modulates rat proximal tubule transport. *J. Clin. Invest.* **97**, 2878–2882.
6. Rohrwasser, A., Morgan, T., Dillon, H.F., Zhao, L., Callaway, C.W., Hillas, E., Zhang, S., Cheng, T., Inagami, T., Ward, K., Terreros, D.A., and Lalouel, J.M. (1999) Elements of a paracrine tubular renin-angiotensin system along the entire nephron. *Hypertension* **34**, 1265–1274.
7. Ardaillou, R. (1999) Angiotensin II receptors. *J. Am. Soc. Nephrol.* **10**, S30–39.
8. Paxton, W.G., Runge, M., Horaist, C., Cohen, C., Alexander, R.W., and Bernstein, K.E. (1993) Immunohistochemical localization of rat angiotensin II AT₁ receptor. *Am. J. Physiol.* **264**, F989–F995.
9. Harrison-Bernard, L.M., Navar, L.G., Ho, M.M., Vinson, G.P., and el-Dahr, S.S. (1997) Immunohistochemical localization of angiotensin II AT1 receptor in adult rat kidney using a monoclonal antibody. *Am. J. Physiol. Renal Physiol.* **273**, F170–177.
10. Wang, Z.Q., Millatt, L.J., Heiderstadt, N.T., Siragy, H.M., Johns, R.A., and Carey, R.M. (1999) Differential regulation of renal angiotensin subtype AT1A and AT2 receptor protein in rats with angiotensin-dependent hypertension. *Hypertension* **33**(1), 96–101.
11. Baum, M., Quigley, R., and Quan, A. (1997) Effect of luminal angiotensin II on rabbit proximal convoluted tubule bicarbonate absorption. *Am. J. Physiol.* **273**, F595–600.
12. Geibel, J., Giebisch, G., and Boron, W.F. (1990) Angiotensin II stimulates both Na⁺-H⁺ exchange and Na⁺/HCO₃⁻ cotransport in the rabbit proximal tubule. *Proc. Natl. Acad. Sci. U. S. A.* **87**, 7917–7920.
13. Levine, D.Z., Iacovitti, M., Buckman, S., and Burns, K.D. (1996) Role of angiotensin II in dietary modulation of rat late distal tubule bicarbonate flux in vivo. *J. Clin. Invest.* **97**, 120–125.
14. Haithcock, D., Jiao, H., Cui, X.L., Hopfer, U., and Douglas, J.G. (1999) Renal proximal tubular AT₂ receptor: signaling and transport. *J. Am. Soc. Nephrol.* **10**, S69–74.
15. Navar, L.G., Harrison-Bernard, L.M., Wang, C.T., Cervenka, L., and Mitchell, K.D. (1999) Concentrations and actions of intraluminal angiotensin II. *J. Am. Soc. Nephrol.* **10**, S189–195.
16. Navar, L.G., Lewis, L., Hymel, A., Braam, B., and Mitchell, K.D. (1994) Tubular fluid concentrations and kidney contents of angiotensins I and II in anesthetized rats. *J. Am. Soc. Nephrol.* **5**, 1153–1158.
17. Pallone, T.L., Robertson, C.R., and Jamison, R.L. (1990) Renal medullary microcirculation. *Physiol. Rev.* **79**, 885–920.
18. Chou, S.Y., Porush, J.G., and Faubert, P.F. (1990) Renal medullary circulation: hormonal control. *Kidney Int.* **37**, 1–13.
19. Zhuo, J., Alcon, D., Allen, A.M., and Mendelsohn, F.A.O. (1992) High resolution localization of angiotensin II receptors in rat renal medulla. *Kidney Int.* **42**, 1372–1380.
20. Nishiyama, A., Seth, D.M., and Navar, L.G. (2002) Renal interstitial fluid concentrations of angiotensin I and II in anesthetized rats. *Hypertension* **39**, 129–134.
21. Navar, L.G. (2004) The intrarenal renin-angiotensin system in hypertension. *Kidney Int.* **65**, 1522–1532.
22. Braam, B., Mitchell, K.D., Fox, J., and Navar, L.G. (1993) Proximal tubular secretion of angiotensin II in rats. *Am. J. Physiol. Renal Physiol.* **264**, F891–898.
23. Boer, W.H., Braam, B., Fransen, R., Boer, P., and Koomans, H.A. (1997) Effects of reduced renal perfusion pressure and acute volume expansion on proximal tubule and whole kidney angiotensin II content in the rat. *Kidney Int.* **51**, 44–49.
24. Kobori, H., Harrison-Bernard, L.M., and Navar, L.G. (2001) Expression of angiotensinogen mRNA and protein in angiotensin II-dependent hypertension. *J. Am. Soc. Nephrol.* **12**, 431–439.
25. Moe, O.W., Ujiie, K., Star, R.A., Miller, R.T., Widel, J., Alperm, R.J., and Henrich, W.L. (1993) Renin expression in renal proximal tubule. *J. Clin. Invest.* **91**, 774–779.
26. Yamagawa, N., Capparelli, A.W., Jo, O.D., Friedal, A., Barrett, J.D., and Eggena, P. (1991) Production of angiotensinogen and renin-like activity by rabbit proximal tubular cells in culture. *Kidney Int.* **39**, 938–941.
27. Sibony, M., Gasc, J.M., Soubrier, F., Alhenc-Gelas, F., and Corvol, P. (1993) Gene expression and tissue localization of the two isoforms of angiotensin I converting enzyme. *Hypertension* **21**, 827–835.
28. Ye, M., Wysocki, J., Naaz, P., Salabat, R., LaPointe, M.S., and Batlle, D. (2004) Increased ACE2 and decreased ACE protein in renal tubules from diabetic mice: a renoprotective combination? *Hypertension* **43**(5), 1120–1125.
29. Casarini, D.E., Boim, M.A., Stella, R.C., Krieger-Azzolini, M.H., Krieger, J.E., and Schor, N. (1997) Angiotensin I-converting enzyme activity in tubular fluid along the rat nephron. *Am. J. Physiol.* **272**(3 Pt 2), F405–409.

30. Goodfriend, T.L. (1993) Angiotensinases. In *Renin-Angiotensin System*. Robertson, J.I.S. and Nichols, M.G., Eds. Gower, London.
31. Imig, J.D., Navar, G.L., Zou, L.X., O'Reilly, K.C., Allen, P.L., Kaysen, J.H., Hammond, T.G., and Navar, L.G. (1999) Renal endosomes contain angiotensin peptides, converting enzyme, and AT₁ receptors. *Am. J. Physiol. Renal Physiol.* **277**, F303–311.
32. Anderson, K.M. and Peach, M.J. (1994) Receptor binding and internalization of a unique biologically active angiotensin II-colloidal gold conjugate. Morphological analysis of angiotensin II processing in isolated vascular strips. *J. Vasc. Res.* **31**, 10–17.
33. Chen, R., Mukhim, Y.V., Garnovskaya, M.N., Thielen, T.E., Iijima, Y., Huang, C., Raymond, J.R., Ullian, M.E., and Paul, R.V. (2000) A functional angiotensin II receptor GFP fusion protein: evidence for agonist-dependent nuclear translocation. *Am. J. Physiol. Renal Physiol.* **279**, F440–448.
34. Schelling, J.R. and Linas, S.L. (1004) Angiotensin II-dependent proximal tubule sodium transport requires receptor-mediated endocytosis. *Am. J. Physiol. Cell Physiol.* **266**, C669–675.
35. Barreto-Chaves, M.L. and Mello-Aires, M. (1996) Effect of luminal angiotensin II and ANP on early and late cortical distal tubule HCO₃⁻ reabsorption. *Am. J. Physiol. Renal Physiol.* **271**, F977–984.
36. Wang, T. and Giebisch, G. (1996) Effects of angiotensin II on electrolyte transport in the early and late distal tubule in rat kidney. *Am. J. Physiol.* **271**, F143–149.
37. Kobori, H., Nyshiyama, A., Harrison-Bernard, L.M., and Navar, L.G. (2003) Urinary excretion of angiotensinogen as an indicator of intrarenal angiotensin status in hypertension. *Hypertension* **41**, 42–49.
38. Wagner, C.A., Giebisch, G., and Geibel, J.P. (2000) Stimulation of H⁺-ATPase in intercalated cells from isolated mouse cortical collecting ducts by angiotensin II. *J. Am. Soc. Nephrol.* **11**(Abstr), A0054.
39. Liu, F.Y. and Cogan, M.G. (1987) Angiotensin II: a potent regulator of acidification in the rat early proximal convoluted tubule. *J. Clin. Invest.* **80**, 272–275.
40. Levine, D.Z., Iacovitti, M., Buckman, S., and Harrison, V. (1994) In vivo modulation of rat distal tubule net HCO₃⁻ flux by VIP, isoproterenol, angiotensin II, and ADH. *Am. J. Physiol.* **266**, F878–883.
41. Wagner, C.A., Giebisch, G., Lang, F., and Geibel, J.P. (1998) Angiotensin II stimulates vesicular H⁺-ATPase in rat proximal tubular cells. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 9665–9668.
42. Li, L., Wang, Y.P., Capparelli, A.W., Jo, O.D., and Yanagawa, N. (1994) Effect of luminal angiotensin II on proximal tubule fluid transport: role of apical phospholipase A₂. *Am. J. Physiol.* **266**(2 Pt 2), F202–209.
43. Seikaly, M.G., Arant, B.S., and Seney, F.S. (1990) Endogenous angiotensin concentrations in specific intrarenal fluid compartments in the rat. *J. Clin. Invest.* **86**, 1352–1357.
44. Chatsudthipong, V. and Chan, Y.L. (1991) Inhibitory effect of angiotensin II on renal tubular transport. *Am. J. Physiol.* **260**, F340–346.
45. Le Goas, F., Amiel, C., and Friedlander, G. (1991) Protein kinase C modulates cAMP content in proximal tubular cells: role of phosphodiesterase inhibition. *Am. J. Physiol.* **261**, F587–592.
46. Liu, F.Y. and Cogan, M.G. (1989) Angiotensin II stimulates early proximal bicarbonate absorption in the rat by decreasing cyclic adenosine monophosphate. *J. Clin. Invest.* **84**, 83–91.
47. Dominguez, J.H., Snowdowne, K.W., Freudenrich, C.C., Brown, T., and Borle, A.B. (1987) Intracellular messenger for action of angiotensin II on fluid transport in rabbit proximal tubule. *Am. J. Physiol.* **252**, F423–428.
48. Nagami, G.T. (2004) Ammonia production and secretion by S3 proximal tubule segments from acidotic mice: role of ANG II. *Am. J. Physiol.* **287**, F707–712.
49. Coppola, S. and Frömter, E. (1994) An electrophysiological study of angiotensin II regulation of Na-CO₃H cotransporter and K conductance in renal proximal tubules. Effect of picomolar concentrations. *Pflügers Arch.* **427**, 143–50.
50. Kunau, R.T., Jr. and Walker, K.A. (1987) Total CO₂ absorption in the distal tubule of the rat. *Am. J. Physiol.* **252**(3 Pt 2), F468–473.
51. Kriz, W. and Kaissling, B. (1992) Structural organization of the mammalian kidney. In *The Kidney: Physiology and Pathophysiology*. Seldin, D.W. and Giebisch, G., Eds. Raven, New York. pp. 707–778.
52. De Mello –Aires, M. and Malnic, G. (2002) Distal tubule bicarbonate transport. *J. Nephrol.* **15**(suppl 5), S97–S111.
53. Levine, D.Z., Iacovitti, M., Buckman, S., Hincke, M.T., Luck, B., and Fryer, J.N. (1997) ANG II-dependent HCO₃⁻ reabsorption in surviving rat distal tubules: expression/activation of H⁺-ATPase. *Am. J. Physiol.* **272**, F799–808.
54. Levine, D.Z., Iacovitti, M., Luck, B., Hincke, M.T., Burns, K.D., and Fryer, J.N. (2000) Surviving rat distal tubule bicarbonate reabsorption: effects of chronic AT₁ blockade. *Am. J. Physiol. Renal Physiol.* **278**, F476–483.
55. Bouby, N., Hus-Citharel, A., Marchetti, J., Bankir, L., Corvol, P., and Llorens-Cortes, C. (1997) Expression of type 1 angiotensin II receptor subtypes and angiotensin II-induced calcium mobilization along the rat nephron. *J. Am. Soc. Nephrol.* **8**, 1658–1667.
56. Doan, T.N., Ali, M.S., and Bernstein, K.E. (2001) Tyrosine kinase activation by the angiotensin II receptor in the absence of calcium signaling. *J. Biol. Chem.* **276**, 20954–20958.
57. Oliveira-Souza, M., and De Mello-Aires, M. (2000) Interaction of angiotensin II and atrial natriuretic peptide on pH_i regulation in MDCK cells. *Am. J. Physiol. Renal Physiol.* **279**, F944–953.
58. Tojo, A., Tisher, C.C., and Madsen, K.M. (1994) Angiotensin II regulates H⁺-ATPase activity in rat cortical collecting duct. *Am. J. Physiol.* **267**, F1045–1051.
59. Weiner, I.D., New, A.R., Milton, A.E., and Tisher, C.C. (1995) Regulation of luminal alkalization and acidification

- in the cortical collecting duct by angiotensin II. *Am. J. Physiol.* **269**, F730–F738.
60. Good, D.W., Thampy, G., and Wang, D.H. (1999) Angiotensin II inhibits HCO₃ absorption via a cytochrome P-450-dependent pathway in MTAL. *Am. J. Physiol. Renal Physiol.* **276**, F726–F736.
61. Wall, S.M., Fischer, M.P., Glapion, D.M., and De La Calzada, M. (2003) ANG II reduces net acid secretion in rat outer medullary collecting duct. *Am. J. Physiol. Renal Physiol.* **285**, F930–F937.
62. Nelson, N. and Harvey, W.R. (1999) Vacuolar and plasma membrane proton-adenosinetriphosphatases. *Physiol. Rev.* **79**, 361–385.
63. Bernasconi, P., Rausch, T., Struve, I., Morgan, L., and Taiz, L. (1990) An mRNA from human brain encodes an isoform of the B subunit of the vacuolar H⁺-ATPase. *J. Biol. Chem.* **265**, 17428–17431.
64. van Hille, B., Richener, H., Schmid, P., Puettner, I., Green, J.R., and Bilbe, G. (1994) Heterogeneity of vacuolar H⁺-ATPase: differential expression of two human subunit B isoforms. *Biochem. J.* **303**, 191–198.
65. Nelson, R.D., Guo, X.L., Masood, K., Brown, D., Kalkbrenner, M., and Gluck, S. (1992) Selectively amplified expression of an isoform of the vacuolar H⁺-ATPase 56-kilodalton subunit in renal intercalated cells. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 3541–3545.
66. Karet, F.E., Finberg, K.E., Nelson, R.D., Nayir, A., Mocan, H., Sanjad, S.A., Rodriguez-Soriano, J., Santos, F., Cremers, C.W., Di Pietro, A., Hoffbrand, B.I., Winiarski, J., Bakkaloglu, A., Ozen, S., Dusunsal, R., Goodyer, P., Hulton, S.A., Wu, D.K., Skvorak, A.B., Morton, C.C., Cunningham, M.J., Jha, V., and Lifton, R.P. (1999) Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat. Genet.* **21**, 84–90.
67. Oka, T., Murata, Y., Namba, M., Yoshimizu, T., Toyomura, T., Yamamoto, A., Sun-Wada, G.H., Hamasaki, N., Wada, Y., and Futai, M. (2001) a4, a unique kidney-specific isoform of mouse vacuolar H⁺-ATPase subunit a. *J. Biol. Chem.* **276**, 40050–40054.
68. Peng, S.B., Li, X., Crider, B.P., Zhou, Z., Andersen, P., Tsai, S.J., Xie, X.S., and Stone, D.K. (1999) Identification and reconstitution of an isoform of the 116-kDa subunit of the vacuolar proton translocating ATPase. *J. Biol. Chem.* **274**, 2549–2555.
69. Toyomura, T., Oka, T., Yamaguchi, C., Wada, Y., and Futai, M. (2000) Three subunit a isoforms of mouse vacuolar H⁺-ATPase. Preferential expression of the a3 isoform during osteoclast differentiation. *J. Biol. Chem.* **275**, 8760–8765.
70. Stehberger, P., Schulz, N., Finberg, K.E., Karet, F.E., Giebisch, G., Lifton, R.P., Geibel, J.P., and Wagner, C.A. (2003) Localization and regulation of the ATP6V0A4 (a4) vacuolar H⁺-ATPase subunit defective in an inherited form of distal renal tubular acidosis. *J. Am. Soc. Nephrol.* **14**, 3027–3038.
71. Smith, A.N., Finberg, K.E., Wagner, C.A., Lifton, R.P., Devonald, M.A., Su, Y., and Karet, F.E. (2001) Molecular cloning and characterization of ATP6N1b: a novel fourth murine vacuolar H⁺-ATPase a-subunit gene. *J. Biol. Chem.* **276**, 42382–42388.
72. Smith, A.N., Skaug, J., Choate, K.A., Nayir, A., Bakkaloglu, A., Ozen, S., Hulton, S.A., Sanjad, S.A., Al-Sabban, E.A., Lifton, R.P., Scherer, S.W., and Karet, F.E. (2000) Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. *Nat. Genet.* **26**, 71–75.
73. Wagner, C.A. and Geibel, J.P. (2002) Acid-base transport in the collecting duct. *J. Nephrol.* **15**, S112–S127.
74. Holliday, L.S., Lu, M., Lee, B.S., Nelson, R.D., Solivan, S., Zhang, L., and Gluck, S.L. (1999) The amino-terminal domain of the B subunit of vacuolar H⁺-ATPase contains a filamentous actin binding site. *J. Biol. Chem.* **275**, 32331–32337.
75. Banerjee, A., Li, G., Alexander, E.A., and Schwartz, J.H. (2001) Role of SNAP-23 in trafficking of H⁺-ATPase in cultured inner medullary collecting duct cells. *Am. J. Physiol. Cell Physiol.* **280**, 775–781.
76. Banerjee, A., Shih, T., Alexander, E.A., and Schwartz, J.H. (1999) SNARE proteins regulate H⁺-ATPase redistribution to the apical membrane in rat renal inner medullary collecting duct cells. *J. Biol. Chem.* **274**, 26518–26522.
77. Breton, S., Wiederhold, T., Marshansky, V., Nsumu, N.N., Ramesh, V., and Brown, D. (2000) The B1 subunit of the H⁺ ATPase is a PDZ domain-binding protein. Colocalization with NHE-RF in renal B-intercalated cells. *J. Biol. Chem.* **275**, 18219–18224.
78. Brown, D. and Breton, S. Structure, function and cellular distribution of the vacuolar H⁺-ATPase (H⁺ V-ATPase/proton pump). In: *The Kidney: Physiology and Pathophysiology*. 3rd Ed., edited by Seldin DW, Giebisch G, Philadelphia: Lippincott Williams and Wilkins, 2000, 171–191.
79. Wagner, C.A., Finberg, K.E., Breton, S., Marshansky, V., Brown, D., and Geibel, J.P. (2004) Renal vacuolar H⁺-ATPase. *Physiol. Rev.* **84**(4), 1263–1314. Review.
80. Marshansky, V., Fleiser, A., Noel, J., Bourgoin, S., and Vinay, P. (1996) Isolation of heavy endosomes from dog proximal tubules in suspension. *J. Membr. Biol.* **153**, 59–73.
81. Marshansky, V. and Vinay, P. (1996) Proton gradient formation in early endosomes from proximal tubules. *Biochim. Biophys. Acta* **1284**, 171–180.
82. Nelson, N., Perzov, N., Cohen, A., Hagai, K., Padler, V., and Nelson, H. (2000) The cellular biology of proton-motive force generation by V-ATPases. *J. Exp. Biol.* **203**, 89–95.
83. Nishi, T. and Forgac, M. (2002) The vacuolar (H⁺)-ATPases—nature's most versatile proton pumps. *Nat. Rev. Mol. Cell. Biol.* **3**, 94–103.

84. Ohkuma, S., Moriyama, Y., and Takano, T. (1982) Identification and characterization of a proton pump on lysosomes by fluorescein-isothiocyanate-dextran fluorescence. *Proc. Natl. Acad. Sci. U. S. A.* **79**, 2758–2762.
85. Bae, H.R. and Verkman, A.S. (1990) Protein kinase A regulates chloride conductance in endocytic vesicles from proximal tubule. *Nature* **348**, 637–639.
86. Hilden, S.A., Johns, C.A., and Madias, N.E. (1988) Cl⁻-dependent ATP-driven H⁺ transport in rabbit renal cortical endosomes. *Am. J. Physiol.* **255**, F885–897.
87. Reenstra, W.W., Sabolic, I., Bae, H.R., and Verkman, A.S. (1992) Protein kinase A dependent membrane protein phosphorylation and chloride conductance in endosomal vesicles from kidney cortex. *Biochemistry* **31**, 175–181.
88. Sabolic, I. and Burckhardt, G. (1986) Characteristics of the proton pump in rat renal cortical endocytotic vesicles. *Am. J. Physiol.* **250**, F817–826.
89. Kaunitz, J.D., Gunther, R.D., and Sachs, G. (1985) Characterization of an electrogenic ATP and chloride-dependent proton translocating pump from rat renal medulla. *J. Biol. Chem.* **260**, 11567–11573.
90. Krick, W., Dolle, A., Hagos, Y., and Burckhardt, G. (1998) Characterization of the chloride conductance in porcine renal brush-border membrane vesicles. *Pflugers Arch.* **435**, 415–421.
91. Zimolo, Z., Montrose, M.H., and Murer, H. (1992) H⁺ extrusion by an apical vacuolar-type H⁺-ATPase in rat renal proximal tubules. *J. Membr. Biol.* **126**, 19–26.
92. Malnic, G. and Geibel, J.P. (2000) Cell pH and H⁺ secretion by S3 segment of mammalian kidney: role of H⁺-ATPase and Cl⁻. *J. Membr. Biol.* **178**, 115–125.
93. Fernandez, R., Bosqueiro, J.R., Cassola, A.C., and Malnic, G. (1997) Role of Cl⁻ in electrogenic H⁺ secretion by cortical distal tubule. *J. Membr. Biol.* **157**, 193–201.
94. Stone, D.K., Seldin, D.W., Kokko, J.P., and Jacobson, H.R. (1983) Anion dependence of rabbit medullary collecting duct acidification. *J. Clin. Invest.* **71**, 1505–1508.
95. Vallés, P., Wysocki, J., Salabat, M.R., Cokic, I., Ye, M., Lapointe, M.S., and Batlle, D. (2005) Angiotensin II increases H⁺-ATPase B1 subunit expression in medullary collecting ducts. *Hypertension* **45**(4), 818–823.
96. Von Thun, A.M., Vari, R.C., el-Dahr, S.S., and Navar, L.G. (1994) Augmentation of intrarenal angiotensin II levels by chronic angiotensin II infusion. *Am. J. Physiol.* **266**(1 Pt 2), F120–128.
97. Zhuo, J.L., Imig, J.D., Hammond, T.G., Orenco, S., Benes, E., and Navar, L.G. (2002) Ang II accumulation in rat renal endosomes during Ang II-induced hypertension: role of AT(1) receptor. *Hypertension* **39**, 116–121.
98. Hamm, L.L. and Alpern, R.J. (2000) In *The Kidney: Physiology and Pathophysiology*. 3rd ed. Seldin, D. and Giebisch, G., Eds. Lippincott Williams & Wilkins, Philadelphia. pp. 1935–1979.
99. Koeppen, B.M. and Helman, S.I. (1982) Acidification of luminal fluid by the rabbit cortical collecting tubule perfused in vitro. *Am. J. Physiol.* **242**, F521–531.
100. Stone, D.K., Seldin, D.W., Kokko, J.P., and Jacobson, H.R. (1983) Mineralocorticoid modulation of rabbit medullary collecting duct acidification. A sodium-independent effect. *J. Clin. Invest.* **72**, 77–83.
101. Garg, L.C. and Narang, N. (1988) Effects of aldosterone on NEM-sensitive ATPase in rabbit nephron segments. *Kidney Int.* **34**, 13–17.
102. Mujais, S.K. (1987) Effect of aldosterone on rat collecting tubule NEM sensitive ATPase. *J. Lab. Clin. Med.* **109**, 34–39.
103. Moorthi, K., Wysocki, J., Salabat, R., Wang, M., Cokic, I., Ikramuddin, P., LaPointe, M., and Batlle, D. (2003) Aldosterone increases the synthesis and cell surface expression of α4, a kidney specific subunit of H⁺-ATPase in a mouse renal collecting tubule cell line. *J. Am. Soc. Nephrol.* **14**, 68^a.
104. Eiam-Ong, S., Kurtzman, N.A., and Sabatini, S. (1993) Regulation of collecting tubule adenosine triphosphates by aldosterone and potassium. *J. Clin. Invest.* **91**, 2385–2392.
105. Winter, C., Schulz, N., Giebisch, G., Geibel, J.P., and Wagner, C.A. (2004) Nongenomic stimulation of vacuolar H⁺-ATPases in intercalated renal tubule cells by aldosterone. *Proc. Natl. Acad. Sci. U. S. A.* **101**(8), 2636–2641.

This article should be referenced as follows:

Valles, P., Wysocki, J., and Batlle, D. (2005) Angiotensin II and renal tubular ion transport. *TheScientificWorldJOURNAL* **5**, 680–690

Handling Editor:

Rajiv Agarwal, Editorial Board Member for *Nephrology* — a domain of *TheScientificWorldJOURNAL*.

