

## REVIEW

# Animals and People in the Netherlands' Past: >50 Years of Archaeozoology in the Netherlands

Canan Çakırlar\*, Youri van den Hurk\*, Inge van der Jagt†, Yvonne van Amerongen‡, Jan Bakker§, Rianne Breider\*, Joyce van Dijk||, Kinie Esser||, Maaïke Groot¶, Theo de Jong\*\*, Lisette Kootker††, Frits Steenhuizen\*, Jørn Zeiler‡‡, Thijs van Kolfschoten§§, Wietske Prummel\* and Roel Lauwerier†

More than fifty years ago, Anneke T. Clason published the first English-language archaeozoological study on Dutch faunal assemblages. Inspired by the anniversary of this landmark publication, this paper presents a status overview of Dutch archaeozoology organized in twelve themes (e.g. rituals, Mesolithic-Neolithic transition, medieval period). The paper also discusses the common methods applied in Dutch archaeozoology, and includes extensive supplementary material that summarizes data from gray literature in Dutch. Our aim is to provide a guide to archaeozoological questions pertaining to the Netherlands and open a window for researchers working outside the Netherlands to the highly active world of Dutch archaeozoology.

**Keywords:** Zooarchaeology; Archaeozoology; Dutch; the Netherlands; Anneke Clason

## Introduction

It has been more than 50 years since the late Prof. Anneke T. Clason published her dissertation, *Animal and Man in Holland's Past* (Clason 1967). After that publication, Clason became the leading archaeozoologist in the Netherlands, as well as its international face. Her dissertation was arguably the best-known Dutch archaeozoological study in the English-speaking scholarly world until at least the early 1990s. Since the late 1960s, her colleagues and students have built a strong tradition of archaeozoology in the Netherlands, producing, interpreting and publishing an enormous amount of data. While some of these data were—and continue to be—collected in the context of academic research and higher education (IJzereef 1981; Prummel 1983; Lauwerier 1988; Brinkhuizen 1989b; Schelvis 1992; Gehasse 1995; Zeiler 1997b; Groot 2008; van Amerongen 2016), the great majority have been

produced in the context of development-led archaeology. Archaeozoologists working in the commercial archaeology sector regularly record and publish massive amounts of data, in compliance with the regulations put into effect in 2007, through the Valletta Convention (Lauwerier 2017).

However, most data remain difficult to access and/or are located in the Dutch grey literature. Syntheses are scarce, and some are in Dutch and thus not readily accessible to non-Dutch speakers. The scarcity of overviews hinders efforts to train, strategize, and internationalize. Inspired by the 50<sup>th</sup> anniversary of Clason's career launch, this review article is an attempt to partly redress this deficiency by producing a widely accessible introduction for beginners in and outsiders to Dutch archaeozoology, one that documents the state of the art.

First, we provide a very brief summary of the Quaternary geographical setting of the Netherlands. Next, we explain the common methods of Dutch archaeozoology, as we consider this a fundamental prerequisite for digging further into the data. Then, we provide summaries of the 12 themes that have received the most attention from us and our predecessors over the past 50 years. The themes vary from regional (e.g. archaeozoology of terps) to chronological (e.g. archaeozoology of the metal ages) to overarching overviews (e.g. expositions and introductions). Each thematic section draws data from the most thoroughly studied, interesting and/or best-known sites and assemblages. Maps with relevant sites are provided for several sections. Site names are provided in Supplementary Table 21. The methods used vary from theme to theme and are

\* University of Groningen, NL

† Cultural Heritage Agency of the Netherlands, NL

‡ Archol, NL

§ University of Amsterdam, NL

|| Archeoplan Eco, NL

¶ Freie Universität Berlin, DE

\*\* Heritage Centre, Municipality of Eindhoven, NL

†† Vrije Universiteit Amsterdam, NL

‡‡ Archaeobone, NL

§§ Leiden University, NL

Corresponding author: Canan Çakırlar ([c.cakirlar@rug.nl](mailto:c.cakirlar@rug.nl))

provided in Supplementary Document 1. To summarize, we provide a comprehensive but concise discussion of the state of the art of archaeozoology in the Netherlands, from a *longue durée* perspective.

### A Note on the Quaternary Geography of the Netherlands

Having a basic knowledge of the Quaternary evolution of the country's dynamic landscapes is crucial to understanding the Dutch archaeozoological record. Luckily, the Dutch landscape is arguably one of the best and most systematically investigated landscapes in the world. In the Late Pleistocene, most of the Netherlands was part of Doggerland, a vast area of land connecting Britain and continental Europe. In the early Holocene, the coastline lay farther west than it does at present, and around 9000 BC, the relative sea level of the southern North Sea was 26 m below the present level. As sea levels continued to rise, at an average speed of ~0.80 m per century, peat was formed in the pericoastal zone due to the wet conditions that prevailed there, and a major part of the Netherlands became part of the tidal basins of various river systems. In the west, the coastline reached its easternmost location around 3850 BC. In the north, the dune barriers of the Wadden Sea Islands (with the exception of the island of Texel) have been in place since the Iron Age and the Roman period (Vos 2015). Around 500 BC, the sea-level rise had decreased to about 0.10–0.05 m per century and, due to a sediment surplus in the coastal zone of the western part of the Netherlands, the dune systems were shifting westwards again. In the northern part of the Netherlands, salt-marsh areas were growing, and from the Early Iron Age onwards, dwelling mounds (known in Dutch as *terpen* or *wierden*) were created to protect people against floods. Artificial ditches and channels made peat areas habitable. This was the beginning of centuries of reclaiming land from the sea that the Netherlands is renowned for. Works to manage the landscape continued under the Romans, who incorporated the southern part of the country into their empire.

The meandering river channels shifted continuously, with new watercourses being developed through the deltas until the Middle Ages, when humans started to embank the rivers to control the watercourses (Vos 2015). Anthropogenic impact on the landscape increased from the High Middle Ages onwards (since the 12<sup>th</sup> century). Dykes were constructed along the salt marshes and floodplains of coastal and river regions, and the peat landscape almost disappeared (Vos 2015). Estuary systems (including those of the Rhine and the Meuse) still dominate, creating a complex and dynamic landscape. Continuous and extensive waterworks coupled with a completely tamed 'nature' define the Dutch landscape today.

### Methods in Dutch Archaeozoology

Dutch archaeozoology has been both innovative and practical in terms of methodology.

Methodological innovations born out of the necessity to interpret Dutch assemblages have become fundamental methods worldwide. Examples include Lauwerier

on recording butchery marks (1988: 181–212) and pig (*Sus domesticus*) seasonality (1983); Prummel and Frisch (1986) on the distinction of sheep (*Ovis aries*) and goat (*Capra hircus*); Prummel (1987b, 1987a, 1988, 1989) on the identification of foetal skeletal elements of cattle (*Bos taurus*), horse (*Equus caballus*), sheep and pig; Schelvis (1992) on mites (Arachnida); and Jans (2005) and Cuijpers (2009) on the application of histology.

Practical methods can be categorized into on-site, recording and interpreting, publishing and post-analysis storage methods.

At prehistoric sites, finds are usually collected through trowelling (known as hand collecting) and sieving. For later periods, a combination of hand collecting and sieving is the norm. Precise taxonomic identifications are prioritized in faunal analysis. Researchers both in public and in commercial institutions are keen on using and improving reference collections across the country. Archaeozoological reference collections are located in Amsterdam (Department of Archaeology, University of Amsterdam), Leiden (Faculty of Archaeology, Leiden University), Groningen (Groningen Institute of Archaeology, University of Groningen) and Amersfoort (Rijksdienst voor het Cultureel Erfgoed (RCE), the Dutch government's cultural heritage agency).

The methods used in recording and reporting are largely in line with methods used elsewhere in Europe, especially in continental Europe, and thus can be considered standardized. Standardization is further achieved through frequent contact among the small and well-organized archaeozoological community (through a yearly workshop meeting of archaeozoologists organized by the RCE and such organizations as the Biologisch Archeologisch Platform (the biological archaeology platform), and a recent Dutch textbook/manual on archaeozoology (Groot 2010)). While not dictated, best practices are recommended by the RCE (Lauwerier 2010). The Dutch archaeology quality standard (*Kwaliteitsnorm Nederlandse Archeologie*), based on heritage legislation and self-regulation, sets minimum standards for specialist work in archaeology. Since 2017, quality has been ensured through a certification-based system.

Tooth wear and eruption are generally recorded following Grant (1982), in combination with Hambleton (1999) or Habermehl (1975). Fusion is recorded following Habermehl (1975), Silver (1969), Zietschmann and Krölling (1955) and Johansson and Hüster (1987), as visualized by the Archäologisch-Zoologische-Arbeitsgruppe Schleswig-Kiel (Reichstein (1991: 21–22)). Number of fragments (*Aantal Fragmenten*) is the most common unit of quantification. Although it is sometimes equivalent to NISP (Number of Identified Specimens), it usually refers to the number of fragments rather than specimens. Bone weight frequently accompanies fragment counts in reports. Other quantification methods or derived types of data used to estimate relative abundance or importance (e.g. MNI = Minimum Number of Individuals), etc., are not in general use.

Most standardized biometric measurements are omitted, especially in development-led research, with the exception of the greatest length of complete long bones (Von den Driesch 1976). These are then converted to

withers heights following von den Driesch & Boessneck (1974) and Matolcsi (1970) for cattle; May (1985) for horse; Harcourt (1974) for dog (*Canis familiaris*); and Teichert (1969, 1975) for pig, sheep and goat. In general, pathologies, butchery, burning and gnawing marks are recorded and briefly described in reports. Occasionally, the publication by Lauwerier (1988) is used to describe butchery marks. A description of the preservation of the bone material is usually given, often referring to Behrensmeyer (1978) or Huisman *et al.* (2009), and sometimes a description of the fragmentation is given, as percentages of the elements present (Groot 2010: 99, Table 7.2).

Archaeozoological reports produced through development-led research are, by default, uploaded to Archis (a database managed by the RCE containing information on rescue excavations in the Netherlands) and often also to the electronic archives (known as e-depot DANS) administered by the Royal Netherlands Academy of Arts and Sciences (KNAW) (2018). BoneInfo, a system with archaeozoological meta-data operated by the (RCE) is the best starting point to obtain regional and chronological absence/presence information about taxonomic groups (Lauwerier & de Vries 2004). The post-analysis storage of archaeofaunal assemblages is the responsibility of provincial and local authorities. Assemblages are submitted to provincial heritage depots along with meta-data (Supplementary Table 1) (Royal Dutch Academy of Sciences (KNAW) 2018, RCE 2018).

### The Archaeozoology of the Pleistocene

Pleistocene/Palaeolithic faunal remains are investigated to address questions about the date ranges for the occurrence of Pleistocene species, the role of animals in hominin/human subsistence, bone tool manufacture, and hominin/human environment.

Hotspots that yield Pleistocene fauna are the North Sea area (both the sea and the coast) and the river valleys. The fossil record is, however, rather fragmented (van Kolfschoten 2001). Even boreholes, such as the ones at Zuurland, with an extremely rich, stratified small-mammal record show large hiatuses in time (van Kolfschoten 1998; van Kolfschoten & Tesakov 1998; van Kolfschoten, Tesakov & Bell 2018). Rich faunal assemblages collected *in situ* from stratigraphically well-embedded deposits are limited: the localities Tegelen and Tegelen-Maalbeek yielded fossil vertebrates from the onset of the Early Pleistocene, and Maastricht-Belvédère yielded faunal remains from the later part of the Middle Pleistocene and the late Pleistocene. However, a number of localities (e.g. Bavel, Dorst, Neede, Wageningen-Fransche Kamp, Rhenen, and Orvelte) yielded smaller *in situ* assemblages dated to the Pleistocene.

Most of the Pleistocene faunal remains are from stratigraphically disturbed contexts; they were collected, for example, from sediments dredged from sand and gravel pits throughout the country (e.g. Woerden, Rhederlaag) or from beaches, such as Maasvlakte 1, Maasvlakte 2, Zandmotor and Hoek van Holland, where sediments from the bottom of the North Sea have been redeposited to create artificial land or sea defences (van Kolfschoten

& Vervoort-Kerkhoff 1999). The Brown Bank and the Eurogeul appear to be very rich in fossil remains. While the majority of the North Sea fossil assemblages date from the late Pleistocene, both early Pleistocene and early Holocene fossils have been identified. A similar situation exists more inland, in the southern Netherlands, where fossil vertebrate remains are found in mixed Pleistocene and Holocene deposits (de Jong 2012).


The Pleistocene (2.6 Ma to 10,000 BP) deposits in the Netherlands are rich in faunal remains, including a large variety of small and large mammal species (**Figure 1**). The large mammal record includes carnivores (e.g. lion (*Panthera* sp.), sabretooth cat (*Homotherium latidens*), hyena (*Crocuta* sp.), bear (*Ursus* sp.), wolf (*Canis* sp.), and wolverine (*Gulo* sp.)), at least five different elephants (assigned to the genera *Mammut*, *Mammuthus* and *Palaeoloxodon*), different horses (Equidae), at least six different rhino species (Rhinocerotidae), tapir (*Tapirus arvernensis*), wild boar (*Sus scrofa*), hippopotamus (*Hippopotamus* sp.), a large number of deer species (Cervidae), and a variety of bovids (Bovidae) (van Kolfschoten 2001).

The role of reindeer (*Rangifer tarandus*) in Late Palaeolithic cultures, a persistent issue in Palaeolithic research in Europe, has also been tackled in Dutch contexts. Campsites of Late Pleistocene hunter-gatherers in northern Germany and Denmark have yielded reindeer bones radiocarbon-dated to the Hamburg (13,000–15,500 BC) and Ahrensburg (11,000–9800 BC) cultures (Lanting & van der Plicht 1996). Characteristic for these cultures are flint arrow heads which were most probably used for reindeer hunting. These types of artefacts have also been found in the Netherlands, but not in association with reindeer remains. In order to get a better idea of the occurrence and distribution of the reindeer in the Netherlands during the Pleistocene, mammalian fossils have been radiocarbon dated (van Kolfschoten *et al.* 2011). This research showed that the reindeer remains from the Netherlands and the North Sea are older than ca. 30,000 years (Kuitens 2007; van Kolfschoten *et al.* 2011; Lauwerier & Deebe 2011; Lauwerier, Prummel & van Kolfschoten 2016; Lanting & van der Plicht 1996) (**Figure 2**). The mammalian fauna from the late Pleistocene (Younger Dryas and Preboreal, 11,500 to 9,000 BP) is dominated by red deer (*Cervus elaphus*), aurochs (*Bos primigenius*) and beaver (*Castor fiber*), while reindeer seems to be absent (Snijders & Broertjes 2016: 32–33; De Jong 2016). This absence of reindeer has led to the hypothesis that the so-called reindeer hunters had no reindeer to hunt in the Netherlands (Lauwerier, Prummel & van Kolfschoten 2016). The Late Pleistocene biotope in the Netherlands was apparently no longer suitable for reindeer herds, whereas that in northern Germany and Denmark still was. More radiocarbon dates are needed to test this hypothesis and further investigate the subsistence strategies of the late Pleistocene and early Holocene hunter-gatherers.

### The Mesolithic–Neolithic Transition

Animal remains are central to the study of Mesolithic–Neolithic transitions. This is also the case for the Netherlands and for north-western Europe in general (Raemaekers





**Figure 1:** Map with Pleistocene localities discussed in text.

2003; Rowley-Conwy 2014). Although there is clear consensus that ceramics arrived in the Netherlands before plant and animal husbandry, the mechanisms, sequence and timing of the adoption of plant and animal husbandry are still under investigation (Cappers & Raemaekers 2008; Rowley-Conwy 2011; Rowley-Conwy 2014). Sites with Mesolithic–Middle Neolithic A (8400–3400 cal. BC) deposits are scarce (Supplementary Table 2; **Figure 3**), and assemblages are often small (Supplementary Table 3). Evidence is missing altogether from the sandy areas that cover most of the eastern and southern parts of the Netherlands, where organic remains are not preserved (Vos & de Vries 2013: maps 5500 and 3850 BC).

The Mesolithic–Neolithic transition research focuses on the period between 5600–4000 BC. The higher parts of the southern Netherlands, on the other hand, as exemplified by the site Elsloo-Koolweg (Lauwerier & Laarman 2015) (5500–4900 BC), are already settled by farmers associated with the Linearbandkeramik culture, who husbanded sheep, goat, cattle and pig. In the Dutch wetlands, ample remains of pike (*Esox lucius*), cyprinids (Cyprinidae), perch (*Perca fluviatilis*), wels catfish (*Silurus*

*glanis*), beaver, otter (*Lutra lutra*), ducks (Anatidae), swans (*Cygnus* sp.) and geese (Anatidae), wild boar and red deer, all associated with freshwater, demonstrate the significance of freshwater faunal resources in the life of humans throughout the period (van Wijngaarden-Bakker et al. 2001; Oversteegen et al. 2001; Clason & Brinkhuizen 1978; Zeiler 1997b; Prummel et al. 2009b; Gehasse 1995; Kranenburg & Prummel in press), and nitrogen and carbon isotopic ratios in human remains confirm this (Smits et al. 2010; Smits & van der Plicht 2009).

There is no evidence for overexploitation of frequently hunted species. For example, in several sites, beavers are abundant, with butchery marks indicating their intensive use as a source of food and fur, but they are almost always adult individuals, indicating that the beaver population remained stable despite human pressure. Mesolithic kill sites in the north, however, such as Jardinga and Balkweg, with assemblages dominated by aurochs and red deer, deviate from this general pattern of focus on aquatic resource exploitation (Prummel et al. 2002; Prummel & Niekus 2005; Prummel et al. 2009b; Prummel & Niekus 2011). While properties of the landscape and availability




Figure 2: Radio-carbon-dated Pleistocene faunal remains from the southern Netherlands (De Jong 2016).




Figure 3: Map with Mesolithic and Neolithic sites discussed in text.



of resources explain some patterns in the faunal exploitation during the Mesolithic, other patterns cannot be explained by optimal exploitation of whatever is available in the environment. For example, even at sites near the North Sea coast, e.g. Yangtze Harbour, remains of marine fish are rare (Zeiler & Brinkhuizen 2015). Wild horse (*Equus ferus*) seems to have been part of the fauna in the early Holocene as well, but in very small numbers (Laarman 2001; Zeiler 1997b; Zeiler & Brinkhuizen 2016).


Wetland foraging remains an important form of subsistence between 5400 and 4000 BC. The first few remains of morphologically domestic animals apart from dog in the Dutch wetland sites come from deposits dated after 4400 BC and represent cattle, sheep, goat and pig (Oversteegen et al. 2001; Louwe Kooijmans 2001). The identified mammal assemblages from the wetlands that date between 4400 and 3400 BC contain, on average, one third morphologically domestic mammals (mainly cattle and pig and a few sheep) and two thirds wild mammals (in terms of the number of fragments) (Gehasse 1995; Zeiler 1997b; Prummel et al. 2009a; Kranenburg & Prummel in press). Direct radiocarbon dates on a few caprine remains from key transitional sites indicate this taxon's presence in the

period 4400–4100 BC (but note that there is a plateau in the radiocarbon curve at the end of the 5<sup>th</sup> millennium BC), but sheep/goat remains do not appear in the Dutch wetlands again until 2500 cal. BC (Çakırlar et al. in press). MtDNA does not show a presence of Near Eastern lineage in any of the suids in the period 4400–4000 BC (Larson et al. 2007; Krause-Kyora 2011; Kranenburg & Prummel in press).

The changing relationships between humans and animals during this crucial period need to be explored in greater detail, with new methods and from multi-disciplinary perspectives.

### The Subsistence Economy of the Metal Ages

The metal ages (the Bronze Age (2000–800 BC) and the Iron Age (800–12 BC)) have been important research foci in the Netherlands. The landscape was very suitable for animal husbandry, with large areas ideal for grazing. In general, the metal ages are characterized by small farmsteads where people kept cattle, sheep, and goat, followed in terms of proportion of fragment counts by pigs, horses and dogs (**Figure 4**), but within this general pattern substantial changes must have occurred, relating to social and




**Figure 4:** Map with Bronze Age and Iron Age sites discussed in text.

technological innovations. Although the relative proportions of domestic animals vary among Bronze Age assemblages, the general pattern that emerges is an emphasis on cattle breeding (**Figure 5**) (van Amerongen 2016).


Most archaeozoological research focusing on the metal ages of the Netherlands investigates diachronic change, testing old hypotheses. Research suggests, for example, that with the emergence of a crop husbandry system known as Celtic fields in the Late Bronze Age, the interdependence between crop and animal husbandry became more pronounced, with crops providing staple food and animals providing meat and secondary products (van den Broeke 2005; Kooistra & Maas 2008). Indeed, preserved trackways at the present island of Texel (Woltering 2001) indicate that the landscape was also husbanded to

facilitate overland connectivity. Isotopic studies suggest that these routes may have been used to transport live animals (e.g. Brusgaard 2014; Brusgaard, Fokkens & Kootker 2019).

A claim that there was an emphasis on milk production (van Wijngaarden-Bakker 1988) was tested by recent research, which showed no clear economic specialisation in a particular use of livestock in the metal ages (van Amerongen 2016; van Dijk 2015). This probably also holds true for the Iron Age, because the farmstead settlement pattern continues. However, the significant changes in culling patterns that are evident towards the end of the Late Iron Age may point at adjustments that pre-date Roman rule (**Figure 6**). Further investigation is necessary to test this hypothesis (van Dijk 2015). The intensive



**Figure 5:** Use potential, birth rate, and herd growth rate of cattle from several Dutch Bronze Age sites (modified after van Amerongen 2016, Figure 9.14). Use potential is a relative value to evaluate the production potential of a herd for meat or milk (Cribb 1985).



**Figure 6:** Mortality profile of cattle from Iron Age sites in the wetlands in the western Netherlands (after van Dijk 2015, figure 6).

animal husbandry practices in the metal ages led to inbreeding abnormalities and perhaps also to smaller-sized cattle (van Amerongen 2016: 138, 139; Manning et al. 2015). Whether the small size was an unwanted outcome or the result of a conscious strategy on the part of the farmers remains unclear.

The impact of farmers and their livestock on the environment was substantial, placing pressure on populations of wild animals (van Amerongen 2016: 259). Hunting and fishing as subsistence strategies were becoming less important. However, the wide range of wild animal species present at Bronze Age sites indicate that the intricate knowledge needed for hunting was maintained. The contribution of wild animals to the seasonal diet remained important (van Amerongen 2016: 96–104), and a possible taboo on fish, as has been suggested for Belgium and England during the Iron Age, for example (Dobney & Ervynck 2007), was not present in the Netherlands.


Although the Roman period influenced many things in the south, the type of animal husbandry system established in the Bronze Age most likely continued in rural areas until the medieval period.

### The Roman Period

The Roman period in the Netherlands started in the last decades BC, when a legionary camp was constructed in Nijmegen. Between the 1<sup>st</sup> and 4<sup>th</sup> centuries AD, a series of forts were built along the south bank of the Rhine, and urban settlements developed in Voorburg and Nijmegen. Only the southern half of the Netherlands formed part of the Roman Empire (**Figure 7**). Archaeozoological work focusing on the Roman period mainly investigates the scale and extent of the Roman presence on the production and consumption of animal products in the 'Dutch part' of the Roman Empire.

An increase in agrarian production resulting from the economic demands of the army and towns was also observed in the Dutch record. Beef was the main type of meat consumed at all types of sites, but local farms probably supplied much of it (e.g. Groot, 2008; Groot et al. 2009). According to culling profiles, cattle were just as important as providers of labour and manure (Groot 2016; Groot & Deschler-Erb 2015, 2017).

A typical Roman phenomenon is the large-scale processing of cattle carcasses in towns, including in



**Figure 7:** Map with Roman sites discussed in text.



Nijmegen (Filean 2006). Cattle also provided leather, horn, grease and bones, and the working of these materials took on an industrial character. The increased use of the meat cleaver in rural sites is reflected in an increase in chop marks on bone (Groot 2016). Meat was also preserved. Cattle shoulder blades often show butchery marks that indicate the smoking of beef shoulders, such as perforations caused by a hook. Concentrations of such shoulder blades have been found in Nijmegen (Lauwerier 1988), but smoked meat was also consumed in the countryside (Groot 2016, Lauwerier 1988: 156). The find of a shoulder blade of a horse displaying similar butchery marks, from the vicus or castellum of Kesteren (Zeiler 2005), is unexpected because horse consumption is not a typically Roman practice. Nevertheless, there is some other evidence for horse consumption in Roman contexts (e.g. Esser 2013), perhaps indicating local tastes or meat fraud.

Domestic chickens were introduced in this period. Finds in sites north of the Rhine (Lauwerier & Laarman 1999; Knol 1983; Prummel 2013; Esser & Zeiler 2013) show that chicken was also accessible to people living outside the Roman Empire. While oysters and other sea molluscs were already being exploited before the Roman period, they were now transported far inland (Groot 2016).

Occasionally, animal bones demonstrate the luxurious side of the Roman diet. A pot with preserved song thrush (*Turdus philomelos*) breasts, imported from the Ardennes, was found in the 1st-century fort on the Kops Plateau in Nijmegen (Lauwerier 1993). Preserved Spanish mackerels (*Scomber japonicus*) were found at several military and urban sites, such as Velsen and Nijmegen (Brinkhuizen 1989a; Lauwerier 1993, 2009: 162), and barracuda (*Sphyraena* sp.) was found in Nijmegen (Lauwerier 1988: 149). Edible snails (*Helix pomatia*) and garden dormice (*Eliomys quercinus*), found only at Valkenburg and the villa of Hoogeloon (Laarman 1987; Kuijper 1990, Groot 2013; Kooistra & Groot 2015), and quail (*Coturnix coturnix*), found in the Fortuna temple in Nijmegen and the villa at Maasbracht (Zeiler 1997a; Esser, Laarman & Rijkeljkhuizen 2017), can also be regarded as delicacies.

Despite these introduced and luxury foods, the diet retained much of its traditional Iron Age character, with meat consumption consisting mostly of beef and lamb. In military sites, pork is consumed more than lamb or mutton (Groot 2016; 2017). Hunting and fishing may have contributed to a varied diet, but their role was negligible in terms of calories.

### The Size and Shape of Cattle and Horse in the Roman Period

Although several thousand Dutch archaeozoological assemblages have been studied over the past 50 years, many of them yielding useful metric information, only a few regional or summarising studies have been published about animal size and shape, mainly focusing the Roman period. The chronological focus is not only a result of the overrepresentation of the historic periods in development-led research, and the relatively well-preserved state, and hence reliable measurability, of the Roman-period


material, but also of the interest in the debate concerning food production by farmers for the soldiers and inhabitants of towns in the 'Dutch' part of the lower Rhine delta (e.g. Groot et al. 2009; Kooistra et al. 2013). The sites considered in this section are provided in **Figure 8**.

Groot (2008, 2016) demonstrated that the average horse size in rural sites in the southern area increased during the Roman period, from 133 cm to 142 cm, indicating that local breeders tailored their production towards the military market's demand for big horses (**Figure 9**). In the Middle Iron Age the average withers height of horses in the rivers area was 131 cm (Van Dijk & Groot 2013; Groot & van Haasteren 2017). The withers height of horses increased in some places, but not everywhere, during the Roman period (Lauwerier 1988; Lauwerier & Robeerst 2001). Horses in rural settlements north of the Roman border were the smallest, with a mean withers height of 132 cm. At military sites within the Roman Empire, the average height was 142 cm, while at villa sites it was 144 cm.


In the Iron Age, cattle were usually no taller than 106 cm at the withers on average (e.g. Terpstra 1986; Knol 1983; Zeiler 2001; Groot 2005; Groot & van Haasteren 2017). In the Roman period, this did not change in the northern coastal area, which was beyond the border of the Empire (Knol 1983; Terpstra 1986; Halici 2002; Bazelmans et al. 2009). The west coast, however, saw a slight increase in average withers height, to around 110–115 cm, during the Roman period (e.g. Laarman 1983; Zeiler 1996; Lauwerier & Laarman 1999).

The trend is different in the rivers area. As far back as the Early Roman period, the withers height in the various settlements increased to a mean of 114 cm (Groot 2016). Large animals were particularly common at the consumer city of Nijmegen. This increase in average withers height was probably the combined effect of breeding for size, the importation of larger animals from other parts of the Empire, and the maintaining of old local breeds. The difference between the small animals in the north and the much larger ones in the rivers area (i.e. south of the Roman border) may reflect a different system of farming. While cattle were probably kept mainly for stock breeding in the north, in the rivers area they were kept above all to provide traction in the mixed agrarian system, and for transporting military supplies and trade goods (Lauwerier 2015).

Morphological attributes of livestock other than size have been subject to research as well. Research showed that polledness occurred mainly in the Roman period and disappeared again in the early Middle Ages in the Netherlands, unlike elsewhere in Europe, where hornless cattle had been around since the Neolithic (Schaffberg & Swalve 2015; Lauwerier 2015). Polled cattle were particularly common in the coastal area beyond the borders of the Roman Empire, while they were rare south of there (**Figure 10**). There is little evidence to explain the differences on the basis of natural selection or functional considerations; selection therefore was most likely based on aesthetic considerations or differences in perception about the appearance of a 'good' cow.



**Figure 8:** Map with sites discussed in Size and Shape section.



**Figure 9:** Withers height of cattle and horse in the Dutch rivers area. n-Values are number of specimens on which the average was based for cattle and horse, respectively.



**Figure 10:** Polled cow with light scurs from the Roman period at Achlum (Hullegie & Prummel 2015; photo W. Prummel).

### The Early Medieval Period

Little is known about the early Middle Ages (450–525 AD) from an archaeozoological point of view, partly because Roman and medieval remains are often mixed in the stratigraphy and because the transition between the two periods occurred gradually, while, at the same time, the population declined. In the northern part of the Netherlands, the Roman and early medieval periods are often separated by an occupation hiatus, which is followed by a deviating settlement pattern (**Figure 11**). Which animal-related socio-economic trends are visible in the archaeological record in the Netherlands during the early medieval period?

During the Merovingian period (525–725 AD), clear regional differences in the representation of cattle, sheep, and pig arise (Supplementary table 4). Along the coast, from the salt marsh areas in Friesland in the north, to the marsh areas in the southwestern parts of the Netherlands, the percentage of sheep remains is high and sometimes exceeds that of cattle, which are much better represented in other regions. Changes in the archaeozoological record in the north are seemingly less pronounced, but – at least in the province of Friesland – a gradual shift from cattle to sheep emerges as well (Nieuwhof 2006; Prummel, Esser & Zeiler 2013).

In the Carolingian period (725–900 AD), the differences in the proportions of sheep and cattle are even more pronounced. With an average of 17% sheep remains, the emporium of Dorestad is an outlier in the rivers area; in the rural settlements in this area, sheep represent barely 6% on average. Striking is the dominance of pig at the Carolingian monastery of St. Salvator, in Susteren, representing almost two thirds of the animal remains (Esser & van Hees in press).

Other archaeozoological characteristics for this period are the appearance of dog and horse burials in cemeteries (Prummel 1992, 1993; van der Jagt et al. 2014). Another, unique feature is the enormous number of remains of wading birds found at Wijnaldum, a central terp mound

in the north (Prummel, Esser & Zeiler 2013). Elsewhere, the avian assemblages consist mainly of chickens and of aquatic birds, such as ducks, geese, mute swan (*Cygnus olor*), and cranes (*Grus* sp.).

Fish consumption concerned mainly freshwater fish and migratory fish. Sea fish are found only in settlements along and near the coast, such as The Hague (Magendans & Waasdorp 1989), Wijnaldum (Prummel, Esser & Zeiler 2013) and Oegstgeest (Cavallo 2006, Esser 2011). Inland, fish remains are scarce and seem to have a connection with exchange and trade contacts. Remains of herring (*Clupea harengus*) and plaice (*Pleuronectes platessa*) are known from settlements near the major rivers: Leidsche Rijn (Esser 2009), Stenen Kamer/Linge (Esser & van Dijk 2001) and Dorestad (e.g. Prummel 1983; Esser, Beerenhout & Rijkelijhuizen 2012). Imported garfish (*Belone belone*) was found in Deventer (Beerenhout 2011a).


Archaeozoological data on the succeeding Ottonian period (900–1050 AD) are too scarce at the moment to enable an overview.

### Animals in Castles and Monasteries

From the beginning of the 11<sup>th</sup> century, a feudal society developed in Europe, based on a philosophical theory of three sharply divided social orders: nobility, clergy and peasants. This society continued to exist until the end of the 18<sup>th</sup> century (Ervynck 2004). We know from the historical record that each of these three groups had its own function, activities and differences in diet. Archaeozoological research concerning this period often aims at testing these differences by comparing castles (known in Dutch as *borg*, *stins*, *versterkt huis*, *havezathe* or *ridderhofstad*) and monasteries in the Netherlands (**Figure 12**) (Supplementary Tables 5 and 6). The current overview is restricted to vertebrates, although molluscs were also consumed.

At all castles, meat, poultry and fish were consumed. Cattle remains dominate the assemblages except at some early castles, where pigs are more abundant





**Figure 11:** Map with early medieval sites discussed in text.


(Beerenhout 2011b; Esser, Kootker & van der Sluis 2014; Rompelman & Eeltink 2011; Cavallo 2011). The nobility distinguished themselves from the other orders by their hunting privileges. Large game hunting is visible in the presence of wild boar and several deer species at some of the castle sites (Supplementary Table 7). But, generally, remains of large game are scarce and those of small game, such as hare (*Lepus europaeus*) and, from the 13<sup>th</sup> century onwards, rabbit (*Oryctolagus cuniculus*), dominate. Hunters made use of dogs and trained birds of prey, and remains of the latter are found occasionally. A typical example of noble falconry is found at the Valkhof (*valk* being the Dutch word for falcon) in The Hague (Pavlovic & Nieweg 2006).

Typical noble dishes were copious and consisted of diverse and often rare luxury products. The species diversity, especially that of birds, is enormous (Supplementary Table 8). Some of these animals are rare, including turkey (*Meleagris gallopavo*) and Eurasian bittern (*Botaurus stellaris*). Banquets were decorated with swans, herons, storks, cranes and other large birds. Sea fish found on

inland sites can also be considered high-status food (Supplementary Table 9). Even more rare are finds of large marine mammals, such as killer whale (*Orcinus orca*) at Velsen (Zeiler & Kompanje 2010) and North Atlantic right whale (*Eubalaena glacialis*) (Esser, Kootker & van der Sluis 2014; de Jong 2010) at both coastal and inland sites.

A variety of meat, poultry and fish was eaten at monasteries as well. Cattle is again dominant, except for the Carolingian and Ottonian period monastery in Susteren, where pigs dominate (Esser & van Hees in press). Clergy probably rarely engaged in hunting. Large wild mammals (both terrestrial and aquatic) are rare, and the few accounts of deer consist (with some exceptions) of the meatless parts. Common however, was the consumption of hare and rabbit (Supplementary Table 10).

A remarkable hunt-related find group is the hunting birds and (hunted?) crows at Ursulaklooster in Delft (van der Jagt 2017a). The diversity in domestic birds and wild ducks, geese and swans equals that at the castle sites, but that of other bird groups is lower (Supplementary



**Figure 12:** Map with castle and monastery sites discussed in the text and tables.

Table 11). Fish consumption is often related to monasteries; historical documents suggest that fish was frequently consumed during fasting. The number of fish taxa in monasteries is surprisingly similar to that found in castles (Supplementary Table 12), but the amount of fish consumed at monasteries is unknown.

#### Animals in Ritual: A Focus on Special Deposits

This section gives a chronological overview of known animal burials and other special animal deposits in the Netherlands, often referred to as ritual deposits. For the purposes of this discussion, we will keep the criteria for recognizing rituals broad and inclusive (see Groot 2008: 115–117).

Between the Late Palaeolithic and the Roman period, ritually deposited remains of diverse species are found in peat soils and stream valleys (Supplementary Table 13; **Figure 3, 7 and 11**). These off-site finds highlight the special significance of the landscape to prehistoric societies.

Special animal deposits from other contexts are scarcer dating from the Palaeolithic to the Bronze Age than they

are in the later periods. Three ‘ritually deposited’ aurochs skulls from Almere, 5300–4200 BC (Peeters & Hogestijn 2001) and antlers and bones of red deer deposited with wooden sticks in pits at Hardinxveld-Giessendam, 5000–4600 BC, are among the better-known of such prehistoric finds (Louwe Kooijmans & Nökkert 2001). There are only a few accounts of animals in human graves: a complete cattle skeleton next to Late Neolithic graves at Garderen (2600–2500 BC), a cattle skull in a grave from Emst (2600–2500 BC), a presumable cattle skull from a Corded Ware culture grave from Zeijen (2850–2450 BC) (Behrens 1964, Louwe Kooijmans 1974: 323), and two cremated dogs in urns in a Middle Bronze Age burial mound at Borger (Prummel 2006a).

Animals played an important role in the burial tradition of the Iron Age and Roman period in the Netherlands. Meat portions were placed in inhumation graves (Lauwerier 2013) or burned with the body on the funeral pyre. Sometimes unburned portions were added to cremations. Studies of animal remains from cemeteries show a clear selection (Supplementary Table 14). Pig and sheep



or goat are most commonly found in burials, followed by chicken, which is only found in Roman cemeteries. Apart from animal remains included within graves, separate animal burials (three horses and one calf) were found in cemeteries in Valkenburg, Tiel and Zaltbommel, respectively (Verhagen 1987, Groot 2008: 178, Esser, van Dijk & Groot 2010). Animals were also part of rituals in Roman temples (Supplementary Table 15) (for a summary, see Groot 2016: 167–174) and were used in rituals that took place within settlements or fields. A range of deposits, from skulls and lower limbs to burials of complete animals, mostly dating to the Roman period, were found (see Groot 2009 for details). These comprise cattle, sheep, horse, dog, piglet, red deer and crow, and were found in relation to farmhouses, in or very close to enclosure ditches, in ditches of field systems, and in wells (van Giffen 1963: 246–248; van Londen 2006; Groot, 2009; van Haasteren & Groot 2013). These animals were probably offered for a variety of purposes.

Special depositions of animals or animal remains in the early Middle Ages are mainly found in extramural cemeteries or in burials situated within settlements. In a departure from previous periods, these concern mostly horses and dogs (Supplementary Table 16) (van der Jagt et al. 2014). The connection with animals in the early medieval burial rituals is also evident from the many finds of animal remains as additional gifts in the form of objects, such as pendants made of teeth of bears, beaver, and deer, and food (Supplementary Table 17) (van der Jagt 2017b). Other special depositions are not well documented, with the exception of a few from the northern part of the Netherlands that were described by Thilderkvist (2013). There is not much written about animals in rituals in the late Middle Ages. Special depositions of animals, especially burials of partial or whole animals, however, are plentiful. They are mainly found near farmsteads and are often interpreted as non-ritual. Establishing to what extent this interpretation is correct will require further study.


## The Persistence of Fishing and Hunting through the Ages

Hunting and fishing continued to be practiced in some regions and periods even after the introduction of animal husbandry. How important were wild resources for subsistence or for other reasons, for example to express social differentiation or as a pastime? These questions are discussed for some large game species, birds, fish and cetaceans for the Mesolithic to the Middle Ages.

### 1. Prehistory

As mentioned above, hunting and fishing remained a key activity throughout the Neolithic, with Ewijk as the only exception dating prior to 4000 BC (Bakels & Zeiler 2005). In the central rivers area and along the west coast, wild boar and red deer, as well as otter, beaver, ducks, geese and swans, were hunted most frequently during both the Mesolithic and the Neolithic. Locally, where the landscape was more open (or became so as a result of human activities), roe deer was an important hunted species. Other large game species, such as aurochs, horse and elk (*Alces alces*), were also hunted, but in far lower numbers, most probably because their populations were small. The same goes for brown bear (*Ursus arctos*), marten (*Martes* sp.), polecat (*Mustela putorius*) and other fur-bearing animals (Prummel 1987c; Gehasse 1995; Zeiler 1997b; Zeiler 2006). In the Late Neolithic, wild mammals seem to have been hunted less frequently. On just one occasion, at the site of Mienakker, have large numbers of seal bones been unearthed (Zeiler & Brinkhuizen 2013).

The shift towards a primarily agricultural subsistence at the start of the Bronze Age, when ‘...people started [...] living with their backs to nature...’ (Louwe Kooijmans 1993: 80) will have caused wildlife exploitation to become increasingly less important (see **Figures 13–16** for selected taxa). This shift is demonstrated by the decreasing number of sites yielding wild boar, beaver and bird remains compared with the increasing number of sites with pigs and dogs (**Figure 13**). However, recent research at several



**Figure 13:** Percentages of archaeological sites in the Netherlands where remains of Eurasian beaver, wild boar, domesticated pig, and dog have been found, based on the online database ‘BoneInfo’ (RCE 2018).



West Frisian Bronze Age sites has shown that a wide range of habitats were still exploited, but likely through a more passive and concentrated form of hunting and fishing, e.g. by using traps during specific times of the year (van Amerongen 2014).

## 2. The Roman and medieval periods


Despite the increasing symbolic importance of wild mammals and birds in the Roman period, hunted mammals and birds were of minor importance as a source of meat (Lauwerier 1988). In her study of the Dutch rivers area, Groot (2016) shows this to be true at both producer (rural) and consumer (urban and military) sites. Military sites show a higher percentage of wild mammals than urban sites (Groot 2016: 178). The increase in wild mammals mainly occurred in the Late Roman period (Lauwerier 1988; Groot 2008). This may be related to a decline in population and a regeneration of woodland (Bakels 1996; Lauwerier 1988).

During the (late) medieval period, hunting became a pastime of the aristocracy, who owned the land and allocated


the hunting rights (den Hartog 2005). This way, hunting became unavailable to the largest part of society. Fishing, however, became significantly more important during the late medieval period. The rise of in particular marine fish exploitation in this era (Figure 14) can be attributed to rapid developments in maritime technology and fish conservation techniques (e.g. Hoffmann 2005), together with the increasing demand for fish that accompanied the re-establishment of urban centres and associated population growth from the 12<sup>th</sup> century onwards (van Houtte 1977).

## 3. Birds

In the coastal areas, fowling was practiced on a larger scale than at inland sites (Zeiler & Clason 1993). This may simply be a consequence of the ecological potential of these habitats: in tidal areas, birds tend to gather in large numbers because of the presence of rich food sources. Fowling was of great interest during the Mesolithic and Neolithic, but of little interest during the Bronze and Iron Ages. It became more important again in the Roman and the medieval periods (Figure 15). Ducks, geese and



**Figure 14:** Percentages of archaeological sites in the Netherlands where remains of selected freshwater, marine and migratory fish taxa have been found, based on the online database 'BoneInfo' (RCE 2018).



**Figure 15:** Percentages of archaeological sites in the Netherlands where remains of bird (Aves) or selected (groups of) bird species have been found, based on the online database 'BoneInfo' (RCE 2018).

swans were the most often fowled birds in all periods. It must be noted, however, that, by the medieval period, these three taxonomic groups include the remains of domestic ducks (*Anas platyrhynchos domesticus*) and geese (*Anser anser domesticus*), which could not always be distinguished from those of their wild relatives. In order to illustrate the medieval upsurge of domestic fowl, the proportions of sites that include the remains of chicken (introduced in the Netherlands during the Roman period) are shown as well. White-tailed eagle (*Haliaeetus albicilla*) and crane (*Grus grus*) were hunted quite often during the Mesolithic, the Neolithic and the Roman period. Goshawk (*Accipiter gentilis*) and sparrowhawk (*Accipiter nisus*) were hunted during the Mesolithic and Neolithic in small numbers. Their bones and those of peregrine (*Falco peregrinus*) and common kestrel (*Falco tinnunculus*) are rather common at late medieval elite sites, suggesting that they belonged to hawks, which were trained birds of prey. The increase in the proportions of waders and corvids during the medieval period can be partly explained by the fact that they (and ducks, geese and many other species) are among the quarry of hawks (Prummel 2013; Prummel 2018).


#### 4. Cetaceans

Cetacean remains are common in Late Neolithic, Roman and medieval sites (Figure 16) (van den Hurk 2014). Medieval sources mentioning cetacean exploitation for the Netherlands are rare, but sources from Flanders, France and England suggest that hunting was occasionally undertaken, that the meat was highly prized, and that especially porpoise meat was seen as a delicacy (De Smet 1981). Hunting

for cetaceans may therefore also have occurred in the Netherlands during the medieval period, as their remains are frequently encountered in ecclesiastical and high-status sites. The majority of the exploited cetaceans, however, probably derive from stranded individuals (Speller et al. 2016).

#### The Use of Isotopes in Dutch Archaeozoological Research

During the past decade, the use of isotope geochemistry in Dutch human and faunal osteoarchaeological research has shifted from sporadic application to systematic integration. In particular strontium (Sr) and oxygen (O) isotopes have proven their value in enabling researchers to deduce the geographical catchment area where inhabitants and animals of the archaeological settlements originated, as specific  $^{87}\text{Sr}/^{86}\text{Sr}$  and  $\delta^{18}\text{O}$  values refer to specific geological and geographical locations, respectively. Moreover, carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotopes are often applied to monitor the individual dietary aspects of human and animals, which can also assist as indicators for mobility and to gain insight into animal husbandry. Hence, multidisciplinary studies of sites from all archaeological periods provide the opportunity to study the provenance of humans and animals, the age at which mobility took place, the dietary aspects of life, and, most importantly, how they were related, in order to provide a better understanding of the social-economic influences of migration, dietary change, trade, exchange and animal husbandry. Although most isotopic work in the Netherlands has focussed on human individuals (see Kootker & Davies 2017 for an overview), isotopic analysis has become a staple component of archaeozoological research as well.



**Figure 16:** Number of sites with cetacean remains present within their archaeozoological assemblages per 100 year period ( $n = 60$ ), based on the online database 'BoneInfo' (RCE 2018). Identified species are harbour porpoise (*Phocoena phocoena*), common bottlenose dolphin (*Tursiops truncatus*), killer whale (*Orcinus orca*), Atlantic right whale (*Eubalaena glacialis*), and sperm whale (*Physeter macrocephalus*). Most cetacean bones remain unidentified below the taxonomic level of family, a major problem in cetacean research in archaeozoology (Speller et al. 2016).

The first bioavailable strontium isotope distribution map of the Netherlands is solely based on archaeological rodents and carefully selected remains of archaeological medium-size mammals (such as foxes (*Vulpes vulpes*); Kootker et al. 2016). The map provides a first insight into the distribution of the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios in the Netherlands and has become an essential component for the interpretation of archaeological strontium isotopic data. During the past few years, strontium isotope research has been conducted on a vast variety of animal species from archaeological contexts, including cattle, horse, sheep/goat, pig, wolf (*Canis lupus*), lynx (*Lynx lynx*), cat, and dog, dating to multiple archaeological periods. The obtained isotopic data show that cattle, horses, sheep/goat and pigs were subject to trade, exchange and long-distance mobility from as early as the Bronze Age (2000–800 BC) onwards (see e.g. Brusgaard 2014 (Bronze Age); Kootker et al. 2018 (Iron Age); van der Jagt et al. 2012 (Early Medieval); Esser, Kootker & van der Sluis 2014 (Modern era)).

In addition, baseline data for carbon and nitrogen isotope research heavily depends on the analysis of archaeological animal remains as well. Archaeological background species from three biological classes, mammals (e.g., cattle, pigs, sheep/goat), birds (e.g., duck, chicken), and fish (e.g., haddock (*Melanogrammus aeglefinus*), cod (*Gadus morhua*)), have been analysed. Research executed to date provides invaluable information about animal husbandry techniques. Pig isotopic data from Esser et al. (2014), for instance, are suggestive of landscape- and possibly household-dependent management, resulting in a variety of diets, ranging from purely herbivorous to omnivorous foods. Similar research carried out on medieval sheep/goat and cattle remains from the Groningen area in northern Netherlands points toward a grazing system in which the more salt-tolerant sheep were kept on the saline thrifts and cattle were kept on grasslands, farther


away from the coast; a system that remained in use for at least a few hundred years (late medieval to modern era, Kootker et al. 2016).

In conclusion, the application of several isotope systems in Dutch archaeozoological research has proven its potential. We are able to collect more knowledge about provenance and mobility patterns of animals and gain valuable information about animal husbandry practises. The incorporation of isotope geochemistry in commercial faunal osteoarchaeological research, however, is still subject to improvement. Future isotopic work will undoubtedly contribute significantly to our understanding of the past human–animal relationships.

### Extirpations and introductions

The Dutch Holocene vertebrate faunal record shows many changes due to human-mediated extirpations and introductions of other species (Figure 17). Knowing which species became extirpated, which species were introduced, as well as how and when, is crucial to inform present-day conservation management. While some of the extirpations and introductions have similar timelines and causes as those in the rest of northwestern Europe, others are specific for the Netherlands. Habitat change, mostly induced by the introduction of domestic fauna and arable farming, is the most important cause of the extirpation of wild fauna.

Wild horse seems to have been present in small numbers in the Netherlands until at least the 5<sup>th</sup> millennium BC, but probably even later into the Neolithic (Laarman 2001; Zeiler 1997b; Zeiler & Brinkhuizen 2015). Aurochs survived until about AD 600 on the northern salt marshes, despite having been surrounded by large populations of domestic cattle (Prummel & Olivier 2008; Thilderkvist 2013: 58–59). Elk survived in the central part of the country until the 11<sup>th</sup>–13<sup>th</sup> centuries AD (Esser 2000;



**Figure 17:** Overview of introductions and extinctions of some mammal and bird species in the Netherlands during the Holocene, based on the online database 'BoneInfo' (RCE 2018) and references mentioned in the section 'Extirpations and introductions'. Figure adapted from (Baker & Worley 2019, Fig. 1.2).



Buitenhuis & Brinkhuizen 2003). The brown bear disappeared from the Netherlands ca. AD 1000. A partial brown bear skeleton found in the dunes near Noordwijk in 2016 is dated 880–970 cal AD (Kuijper et al. 2016). Little is known about the presence of lynx in the Netherlands, and hardly any lynx bones have been found. Vagrant lynxes have, however, occasionally been encountered in the southern-most part of the Netherlands since AD 2000 (Mulder 2016a; Bakker 2018).

Other extirpations, especially of aquatic fauna, are directly related to overexploitation and habitat destruction. Beaver had become extinct by 1826; the last was killed at Zalk, along the River IJssel. This species was successfully reintroduced from the River Elbe in 1988 (Sluiter 2003; Dijkstra 2016). The otter had a similar history, with, in this case, a 20<sup>th</sup> century extirpation and a reintroduction in 2002–2008 (Lammertsma et al. 2008; Lammertsma & Niewold 2016). Waterworks and pollution have had a negative effect on Dutch fish populations and have heavily affected the occurrence of species, such as the Atlantic salmon (*Salmo salar*) (Lenders et al. 2016).

The increased human habitation of the wetlands had a negative effect on the bird population. White-tailed eagle, crane, and the grey-lag goose (*Anser anser*) ceased breeding in the Netherlands in the post-medieval period. Thanks to protection and habitat recovery since the last decades of the 20<sup>th</sup> century, these species have once again become breeding birds of the Netherlands (van Straalen 2018; Feenstra 2018; Voslamber & Koffijberg 2018).

A number of species are difficult to trace in the archaeozoological record, either because humans did not hunt the species or because they did not bring the skeletons to their settlement. Wolf is a good example. The few archaeozoological records of wolves shed no light on the population dynamics of this species in the past. A small number of wolves have been observed in the southern part of the Netherlands in the 19<sup>th</sup> century. Several incidental appearances of wolves originating from lowland Germany have been recorded since 2015. They made wildlife managers and the public question whether the wolf is returning to the Netherlands. The latest information indeed suggests that this is the case (Lelieveld et al. 2016; La Haye & Verboom 2017). The wild cat (*Felis silvestris*) disappeared from the country during the Roman period, but returned to the south-eastern part of the country in 2012 (Mulder 2016b; Kuipers 2017).

Many introduced species are domesticated. Domestic cat and chicken, which were introduced elsewhere in Europe during the Iron Age (but remained rare until the Roman period), are absent in the Dutch archaeological record of the Iron Age. Domestic cat was introduced in the Netherlands in the Early (12 BC–AD 70) or Middle Roman period (AD 70–270). Domestic chicken, which was introduced to Europe as an 'exotic' species during the Iron Age and was mainly used in burials during this period, assumed economic importance much later in the Roman period (Sykes 2012). Chicken appears in the Dutch record in the Early Roman period (the first finds are from the Roman Castellum Velsen 1, AD 15–30 Prummel 1987d).

It is interesting that some of the domestic animals that are associated elsewhere with Romanisation were not introduced to the Netherlands until early Modern times. Although an implement made of a fallow deer (*Dama dama*) metatarsus has been found in the Roman Castellum Valkenburg (Prummel 1977), there is no indication of any fallow deer population in the Netherlands during the Roman period. The first historical record of fallow deer in deer parks of an aristocratic class dates to AD 1516 (Litjens & Pelzers 1988). This is in contrast with Britain, where enclosed fallow deer populations are known to have been established by the Romans (Sykes et al. 2011).

A peacock (*Pavo cristatus*) bone discovered in the Roman Castellum Velsen 1 (Prummel 1987) may come from a single import of peacock meat. The peacock bones found at some late medieval elite sites definitely come from peacocks living at those sites. The earliest late medieval archaeozoological peacock find is from Helmond (AD 1170–1400, de Jong 1992).

Other introductions in the late Middle Ages include rabbit, pheasant and common carp (*Cyprinus carpio*). Rabbits appear in the Dutch historical record in AD 1297, when they were introduced as game, at first within enclosures. After they escaped, they had devastating effects on the landscape (van Dam 2001, Lauwerier & Zeiler 2001). The pheasant was introduced as game; the first archaeozoological record is from Maastricht, AD 1250–1350 (BoneInfo Cluster 259) (Hiddingh 1983, Lauwerier 1997). The first carp was introduced as lent food; the first archaeozoological record is from Zutphen, AD 1125–1175 (Beerenhout 2011b).

Domestic duck and goose seem to have been introduced during the late Middle Ages, a conclusion that is based on the dramatic increase in duck and goose bones in the archaeozoological record of the Netherlands and the increase in the prevalence of large individuals. See the section on the persistence of fishing and hunting. Whether and how much the Dutch wild populations contributed to their domestication remains a question. After all, the word decoy derives from the Dutch word *eendenkooi*, referring to the combination of a pond, screens, nets a dog and domestic ducks to capture wild ducks.


Introductions from the Americas are also reflected in the archaeological record. Turkey (*Meleagris gallopavo*) was introduced soon after AD 1492; the first records are from Breda (AD 1530–1540, De Jong, Carmiggelt & van den Eynde 1997) and Alkmaar (AD 1475–1550 and AD 1450–1600, van Haaster, Zeiler & Brinkhuizen 2012). Exotic species like the raccoon (*Procyon lotor*), raccoon dog (*Nyctereutes procyonoides*), muskrat (*Ondatra zibethicus*), and American mink (*Neovison vison*) have made their way into the Netherlands even more recently, often bringing harm to the populations of native species.

### The Archaeozoology of the Terp Area


Between 2000 and 650 BC, tidal flats and salt marshes developed between the Pleistocene sandy soils in the north of the Netherlands and the islands along the coast (in the Provinces of Friesland and Groningen) (**Figure 18**). Settlers came to live here ca. 600 BC and built dwelling mounds (terps) on the salt marsh to be safe at high water

(Vos & Knol 2015: 211). Fresh clay was deposited during flood inundations, which made the salt marsh very fertile. It is commonly accepted that bones and shells preserve exceptionally well in terp soil. The rich archaeozoological record of the unique terp phenomenon has fascinated archaeozoologists since the time of van Giffen, the founder of Dutch archaeology. Subsistence in the unem-


banked salt marsh and tidal flat areas in this part of the Netherlands must have been quite different than elsewhere in the Netherlands or Europe in general, at least until the first dykes were built around AD 1200. The first dwellers, of the Iron Age, had a strong preference for cattle husbandry (Supplementary Table 18, **Figure 19**). Sheep husbandry became more important




**Figure 18:** Map with terp sites discussed in text.



**Figure 19:** Percentages of cattle, sheep, horse, pig and dog remains in selected Iron Age terp sites. Site, time period, (NISP), (G): Groningen, (F): Friesland.



**Figure 20:** Percentages of cattle, sheep, horse, pig and dog remains in selected Roman period terp sites. Site, time period, (NISP), (G): Groningen, (F): Friesland.



**Figure 21:** Percentages of cattle, sheep, horse, pig, dog and cat remains in selected early medieval terp sites. Site, time period, (NISP), (G): Groningen, (F): Friesland.

during the Roman period (Supplementary Table 19, **Figure 20**) and even more during the early Middle Ages (Supplementary Table 20, **Figure 21**). The increase in sheep husbandry was probably related to wool becoming a trading commodity. The Roman army had a large demand for wool. The area was part of an extensive trading system along the North Sea during the early Middle Ages, with woollen textile as an important commodity (Prummel 2014).

Sheep husbandry seems to have been more important at terps in Friesland than at those in Groningen, probably thanks to the influence of marine conditions in Friesland creating suitable meadows for sheep herds (Prummel 2006b: 42–45, Prummel, Esser & Zeiler 2013; Prummel 2014, Hullege & Prummel 2015: 151–153).

Hunting was practiced on a limited scale on the terps. Fowling and fishing were hardly practiced during the Iron Age, but they were of some importance during the Roman



and early medieval periods, especially at terps near the coast and at what are argued to be elite terps (Prummel, Esser & Zeiler 2013; Zeiler 2014; Timmerman 2012). Ducks, geese and wading birds were the most important game, and flatfishes and eel (*Anguilla anguilla*) the most captured fish species. Marine molluscs were consumed at most terps. Bones of stranded large whales were sometimes collected to use the oil or the bones (Prummel, van Gent & Kompanje 2012).

Between AD 1200 and 1300, most of the salt marshes were embanked, bringing an end to this unique way of life in the northern parts of the Netherlands (Vos & Knol 2015: 215).

## Conclusions

Thousands of archaeozoological assemblages have been analysed in the Netherlands since Clason published her dissertation in the 1960s, immensely enriching our understanding of past human–animal interactions. The number of archaeozoologists and the amount of archaeozoological data in the Netherlands are growing steadily. An increase in the employment of molecular techniques (stable isotope analysis, palaeogenetics, and ZooMS (zooarchaeology by mass spectrometry)) has begun to add a much-welcomed layer to the information that can be gained from conventional faunal analysis.

With the incorporation of data from molecular analyses, the data are becoming more solid and research questions are becoming more refined.

These are great developments in which we all participate and advocate for in governmental, academic and commercial institutions. However, assemblage-based research, regardless of the methods used to conduct it, continues to lead to knowledge fragmentation. The paper is also the result of an exercise in the re-use of (digital) archaeozoological data and in crowd-sourcing among archaeozoologists who operate primarily in the heritage sector. The BoneInfo database proved to be an excellent open-source starting point. However, quantitative meta-analysis had to depend for the most part on published reports, which was, as expected, time consuming. For more detailed and reproducible overviews and more interesting narratives of *Animals and People in the Netherlands*, interoperable data-sharing platforms are desirable.

## Additional Files

The additional files for this article can be found as follows:

- **Supplementary Table 1.** Archaeological periods used in Dutch archaeology. The table gives the period codes used in the online database 'BoneInfo', the name of the period and the period in years; ages in 'C14' are uncalibrated dates BP; ages in 'BC' and 'AD' are calibrated dates. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 2.** Sites and assemblages used in Figure 3. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 3.** Zooarchaeological and site data used in Figure 3. DOI: <https://doi.org/10.5334/oq.61.s1>

- **Supplementary Table 4.** Ratio of cattle, sheep/goat and pig for Merovingian and Carolingian sites. Only sites with a NISP  $\geq 100$  of cattle, sheep/goat or pig (excluding associated bone groups) were used. \* burnt remains. \*\* two exceptional pits filled with pig mandibles have been excluded from the NISP. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 5.** Overview of the excavated and archaeozoologically studied Dutch castle sites, including references. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 6.** Overview of the excavated and archaeozoologically studied Dutch monasteries, including references. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 7.** Mammal species found at castle sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 8.** Bird species found at castle sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 9.** Fish species found at castle sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 10.** Mammal species found at monastery sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 11.** Bird species found at monastery sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 12.** Fish species found at monastery sites in the Netherlands. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 13.** Overview of animal species found in peat soils and stream valleys in the northern and southern Netherlands. 'X' denotes the presence of the species. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 14.** Number of bone fragments for temple sites for the pre-temple period (Late Iron Age/Early Roman period). Additional data from the Fortuna temple: four fragments of quail, one of herring (*Clupea harengus*), one of smelt (*Osmerus eperlanus*) and one of Cyprinidae. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 15.** Animal species found in Late Iron Age and Roman cremation cemeteries in the Netherlands. Only cemeteries with  $\geq 5$  graves with identified animal bones have been included. x: burned; +: unburned. LIA: Late Iron Age; ROM: Roman period. DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 16.** Animal burials found in early medieval cemeteries. The numbers in parentheses are individuals of that are described as possible animal burials. (inh: inhumation; cr: cremation). DOI: <https://doi.org/10.5334/oq.61.s1>

**Supplementary Table 17.** Overview of animal remains in early medieval cemeteries, excluding animal burials and objects. Some of these remains are clearly food gifts, but others may not be. In Elst, the inhumation graves contained burnt animal remains. 'X' denotes the presence of the species, '(x)' denotes that bone

was found but it is not certain whether it is animal or human. DOI: <https://doi.org/10.5334/oq.61.s1>

- **Supplementary Table 18.** Percentages of cattle, sheep, horse, pig and dog remains from selected Iron Age terp sites. (h: hand collected, s: sieved). DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 19.** Percentages of cattle, sheep, horse, pig and dog remains from selected Roman period terp sites. (h: hand collected, s: sieved). DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 20.** Percentages of cattle, sheep, horse, pig, dog and cat remains from selected early medieval terp sites. (h: hand collected, s: sieved). DOI: <https://doi.org/10.5334/oq.61.s1>
- **Supplementary Table 21.** Site information for all the sites considered as part of this study. DOI: <https://doi.org/10.5334/oq.61.s1>
- **References for the Supplementary Tables.** References for the Supplementary Tables. DOI: <https://doi.org/10.5334/oq.61.s2>
- **Methods.** Methods used for the sections. DOI: <https://doi.org/10.5334/oq.61.s3>

### Acknowledgements

We would like to thank the Stichting Nederlands Museum voor Anthropologie en Praehistorie (SNMAP) for their financial support to deliver this paper. We additionally would like to thank Polydora Baker and two anonymous reviewers for their time and constructive feedback, which helped us to improve our paper immensely. RCE provided financial support for language and copy editing.

### Competing Interests

The authors have no competing interests to declare.

### Author Contributions

The archaeozoology of the Pleistocene: Thijs van Kolfschoten, Theo de Jong; The Mesolithic–Neolithic transition: Canan Çakırlar, Jörn Zeiler, Rianne Breider, Wietske Prummel; The subsistence economy of the metal ages: Joyce van Dijk, Yvonne van Amerongen; The Roman period: Maaïke Groot, Roel Lauwerier; The size and shape of cattle and horse in the Roman period: Roel Lauwerier, Kinie Esser, Theo de Jong, Joyce van Dijk, Canan Çakırlar, Maaïke Groot; The early medieval period: Kinie Esser; Animals in castles and monasteries: Inge van der Jagt, Kinie Esser; Animals in ritual: A focus on special deposits: Inge van der Jagt, Maaïke Groot, Wietske Prummel; The persistence of fishing and hunting through the ages: Jörn Zeiler, Wietske Prummel, Joyce van Dijk, Jan Bakker, Youri van den Hurk; The use of isotopes in Dutch archaeozoological research: Lisette Kootker; Extirpations and introductions: Wietske Prummel, Thijs van Kolfschoten, Kinie Esser; The archaeozoology of the terp area: Wietske Prummel. Frits Steenhuisen and Youri van den Hurk created the maps. Canan Çakırlar, Youri van den Hurk, Inge van der Jagt, Thijs van Kolfschoten, Wietske Prummel, Roel Lauwerier designed the study, organized the work, wrote and edited the manuscript.

### References

- Bakels, CC.** 1996. The Pollendiagram Voerendaal-7. In: Kooistra, LI (ed.), *Borderland Farming: Possibilities and limitations of Farming in the Roman Period and Early Middle Ages between the Rhine and Meuse*, 139–146. Assen: Van Gorcum.
- Bakels, C and Zeiler, J.** 2005. The fruits of the land: Neolithic subsistence (5300–2000 BC). In: Louwe Kooijmans, LP, van Gijn, AL, Fokkens, H and van den Broeke, P (eds.), *The Prehistory of the Netherlands*, 1: 311–336. Amsterdam: Amsterdam University Press.
- Baker, P and Worley, F.** 2019. *Animal bones and archaeology: Recovery to archives*. Swindon: Historic England.
- Bakker, H.** 2018. Rechter mandibulafragment van de Europese Lynx, *Lynx lynx* (Linnaeus, 1758) van het Noordzeestrand van Texel. *Cranium*, 35(1): 9–25.
- Bazelmans, J, Groenendijk, H, de Langen, G, Nicolay, J and Nieuwhof, A.** 2009. *De late prehistorie en protohistorie van Holoceen Noord-Nederland: Onderzoeksagenda, versie 2*. Leeuwarden: Waddenacademie, KNAW.
- Beerenhout, B.** 2011a. *Visresten van het Burseplein in Deventer (9e & 19e eeuw)*. Intern rapport Archeo-Zoo.
- Beerenhout, B.** 2011b. Visspectrum uit afvalkuilen van de grafelijke palts te Zutphen uit de late 10e t/m 12e eeuw. In: Groothedde, M (ed.), *De vorstelijke palts van Zutphen, De uitwerking van de opgravingen van 1946 en 1993 t/m 1999 op en rond het plein 's-Gravenhof te Zutphen (=Zutphense Archeologische Publicaties 66)*. Zutphen. Digital appendix.
- Behrens, H.** 1964. *Die neolithisch-fruhmetallzeitlichen Tierskelettfunde der Alten Welt: Studien zu ihrer Wesensdeutung und historischen Problematik (Veröffentlichungen des Landesmuseums für Vorgeschichte in Halle 19)*. Ost-Berlin: Deutscher Verlag der Wissenschaften.
- Behrensmeyer, AK.** 1978. Taphonomic and ecologic information from bone weathering. *Paleobiology*, 4(2): 150–162. DOI: <https://doi.org/10.1017/S0094837300005820>
- Brinkhuizen, D.** 1989a. Een wervel van een zeldzame vissoort uit de Romeinse haven van Velsen 1 (N.-H.). *Paleo-Aktueel*, 1: 69–72.
- Brinkhuizen, DC.** 1989b. Ichthyo-archeologisch onderzoek: methoden en toepassing aan de hand van Romeins vismateriaal uit Velsen (Nederland). Rijksuniversiteit Groningen.
- Brusgaard, N.** 2014. The social significance of cattle in Bronze Age northwestern Europe. Unpublished thesis (Masters), Leiden University.
- Brusgaard, NØ, Fokkens, H and Kootker, LM.** 2019. An isotopic perspective on socio-economic significance of livestock in Bronze Age West-Frisia, the Netherlands (2000–800 BCE). *Journal of Archaeological Science: Reports*, 24: 101944. DOI: <https://doi.org/10.1016/j.jasrep.2019.101944>
- Buitenhuis, H and Brinkhuizen, DC.** 2003. Faunaresten. In: Hielkema, JB (ed.), *Aanvullend Archeologisch Onderzoek op terrein 9 te Houten-Loerik, gemeente Houten (U) (ARC-Publicatie 69)*, 35–46. Groningen.

- Cakırlar, C, Breider, R, Koolstra, F and Raemaekers, DCM.** in press. Dealing with domestic animals in fifth millennium cal BC Dutch wetlands: new insights from old Swifterbant assemblages. In: Rowley-Conwy, P, Sorensen, L and Gron, K (eds.), *Farmers at the frontier: a Pan-European Perspective on Neolithisation*. Oxford: Oxbow.
- Cappers, RTJ and Raemaekers, DCM.** 2008. Cereal cultivation at Swifterbant? Neolithic wetland farming on the North European plain. *Current anthropology*, 49(3): 385–402. DOI: <https://doi.org/10.1086/588494>
- Cavallo, C.** 2006. De dierlijke resten. In: Hemminga, M and Hamburg, T (eds.), *Een Merovingische nederzetting op de oever van de Oude Rijn (Archol Rapport 69)*, 73–81. Leiden.
- Cavallo, C.** 2011. Adellijke eetgewoonten. De vorstelijke palts van Zutphen. Zooarchaeologisch onderzoek naar de resten uit Huize van de Kastele, Zutphen (1994–95). ZAP, 67.
- Clason, A.** 1967. Animal and man in Holland's past: an investigation of the animal world surrounding man in prehistoric and early historic times in the provinces of North and South Holland. Unpublished thesis (PhD), Rijksuniversiteit Groningen.
- Clason, AT and Brinkhuizen, DC.** 1978. Swifterbant, mammals, birds, fishes. A preliminary report (Swifterbant contribution 8). *Helinium*, 18: 69–82.
- Cribb, R.** 1985. The analysis of ancient herding systems: an application of computer simulation in faunal studies. *Beyond domestication in prehistoric Europe*, 75–106.
- Cuijpers, AGFM.** 2009. The application of bone histology for species identification in archaeology; with a photo catalogue. *Geoarchaeological and bioarchaeological studies*, 12.
- de Jong, T.** 1992. Van dagelijkse kost tot Vorstelijke maaltijd. *De Vlasbloem. Helmonds Historisch Jaarboek*, 12: 27–58.
- de Jong, T.** 2010. Dierlijk bot. In: Peters, S (ed.), *Sint-Oedenrode Kerkstraat. Archeologisch onderzoek, 's Hertogenbosch/Deventer (BAAC Rapport A-05.0339)*, 112–133.
- de Jong, T.** 2016. Brabantse beekdalen: prehistorische fauna. In: Amkreutz, L, Brounen, F, Deeben, J, Machiels, R, van Oorsouw, MF and Smit, B (eds.), *Vuursteen verzameld. Over het zoeken en onderzoeken van steentijd-vondsten en -vindplaatsen*, 284–289. Amersfoort.
- de Jong, TP.** 2012. Holocene Fauna from Brook Valleys in the Southern Netherlands. *A Bouquet of Archaeological Studies. Essays in Honour of Wietske Prummel*, 47–59. DOI: <https://doi.org/10.2307/j.ctt227285m.7>
- de Jong, T, Carmiggelt, A and van den Eynde, G.** 1997. Met de Nassaus aan tafel. Dierlijk botmateriaal uit het kasteel van Breda onderzocht. *Brabants Heem*, 49(4): 121–129.
- De Smet, WMA.** 1981. Evidence of Whaling in the North Sea and English Channel during the Middle Ages. *Mammals in the seas: Reports*, 5(3): 301–309.
- Den Hartog, E.** 2005. *Dieren in en rond de kastelen Teylingen en Brederode*. Haarlem: Kastelenstichting Holland en Zeeland.
- Dijkstra, V.** 2016. Bever Castor fiber. In: Broekhuizen, S, Spoelstra, K, Thissen, JBM, Canters, KJ and Buys, JC (eds.), *Atlas van de Nederlandse Zoogdieren (Natuur van Nederland 12)*, 108–110. Leiden.
- Dobney, K and Ervynck, A.** 2007. To fish or not to fish? Evidence for the possible avoidance of fish consumption during the Iron Age around the North Sea. In: Haselgrove, C (ed.) *The Later Iron Age in Britain and beyond*, 403–418. Oxford: Oxbow Books. DOI: <https://doi.org/10.2307/j.ctvh1dsh9.25>
- Ervynck, A.** 2004. Orant, pugniant, laborant. The diet of the three orders in the feudal society of medieval north-western Europe. In: Jones O'Day, S, van Neer, W and Ervynck, A (eds.), *Behaviour behind bones. The zooarchaeology of ritual, religion, status and identity (Proceedings of the 9th ICAZ Conference, Durham 2002)*, 215–223. Oxford: Oxbow Books.
- Esser, E.** 2000. Archeozoölogie. In: Oudhof, JWM, Dijkstra, J and Verhoeven, AAA (eds.), *Archeologie in de Betuweroute, 'Huis Malburg' van spoor tot spoor. Een middeleeuwse nederzetting in Kerk-Avezaath*, 199–277. Amersfoort.
- Esser, E.** 2009. Archeozoölogie – zoogdieren en vogels. In: Nokkert, M, Aarts, AC and Wynia, HL (eds.), *Vroeg-middeleeuwse bewoning langs de A2. Een nederzetting uit de zevende en achtste eeuw in Leidsche Rijn. (Basisrapportage archeologie 26)*. Utrecht.
- Esser, E.** 2011. Archeozoölogisch onderzoek. In: Jezeer, W (ed.), *Een Merovingische nederzetting aan de monding van de Rijn. Een archeologische opgraving te Oegstgeest Nieuw Rhijngeest-Zuid (ADC rapport 2054)*, 95–105. Amersfoort.
- Esser, E.** 2013. Archeozoölogisch onderzoek. In: Dielemans, L (ed.), *Wacht aan het water. VLEN3-00: archeologisch onderzoek naar sporen en vondstassemblages uit de Romeinse tijd in Vleuterweide. (Basisrapportage Archeologie, 52)*, 83–98. Utrecht.
- Esser, E, Beerenhout, B and Rijkkelijkhuizen, MJ.** 2012. Dierlijke resten van het Veilingterrein. In: Dijkstra, J (ed.), *Het domein van de boer en de ambachtsman. Een opgraving op het terrein van de voormalige fruitveiling te Wijk bij Duurstede: een deel van Dorestad en de villa Wijk archeologisch onderzocht (ADC monografie 12)*, 479–558. Amersfoort.
- Esser, E, Kootker, LM and van der Sluis, L.** 2014. *Dineren in de burcht van Rode. Archeozoölogisch en isotopeenonderzoek naar de samenstelling, productie en distributie van voedsel uit de burcht van Sint-Oedenrode. Special editie Ossiculum 1*. Delft.
- Esser, E, Laarman, FL and Rijkkelijkhuizen, MJ.** 2017. Animal remains and bone artefacts. In: Vos, WK, Bakels, CC and Goosens, TA (eds.), *The roman villa at Maasbracht. The archaeology and history of a Roman settlement on the banks of the river Meuse (province of Limburg, The Netherlands)*, 129–149. Leiden.



- Esser, E and van Dijk, J.** 2001. Archeozoölogie. In: Verhoeven, AAA and Brinkkemper, O (eds.), *Archeologie in de Betuweroute: Twaalf eeuwen bewoning langs de Linde bij De Stenen Kamer in Kerk-Avezaath (Rapportage Archeologische Monumentenzorg 58)*, 363–484. Amersfoort.
- Esser, E, van Dijk, J and Groot, M.** 2010. Archeozoölogisch onderzoek. In: Veldman, HAP and Blom, E (eds.), *Onder de zoden van Zaltbommel. (ADC Monografie 8)*, 201–231. Amersfoort.
- Esser, E and van Hees, L.** in press. De voedselvoorziening in het klooster – zoogdieren en vogels. In: Stoeper, H. (ed.), *Het klooster van Susteren. Archeologisch onderzoek van een Karolingisch abdij en een 11<sup>e</sup> – tot 18<sup>e</sup>-eeuws kanunnikessenstift, deel 1: Vondsten*.
- Feenstra, H.** 2018. Kraanvogel *Grus grus*. In: *SOVON Vogelonderzoek Nederland 2018, Vogelatlas van Nederland*, 232–233. Utrecht/Antwerpen: Kosmos Uitgevers.
- Filean, EP.** 2006. Domestic cattle and economic change in the Roman period lower Rhineland: the Civitas Batavorum. Unpublished thesis (PhD), University of Iowa.
- Gehasse, EF.** 1995. Ecologisch-archeologisch onderzoek van het Neolithicum en de Vroege Bronstijd in de Noordoostpolder met de nadruk op vindplaats P14 gevolgd door een overzicht van de bewoningsgeschiedenis en de bestaans economie binnen de Holocene Delta. Unpublished thesis (PhD), Universiteit van Amsterdam.
- Grant, A.** 1982. The use of tooth wear as a guide to the age of domestic animals. *Ageing and sexing animal bones from archaeological sites*, 91–108.
- Groot, M.** 2005. Dierlijk en menselijk botmateriaal. In: Hiddink, HA and de Boer, E (ed.), *Fossiele beekbeddingen met vondsten uit de Late IJzertijd bij Neerbeek*, 29–46. Amsterdam: Archeologisch Centrum Vrije Universiteit (Zuidnederlandse Archeologische Rapporten 21).
- Groot, M.** 2008. *Animals in ritual and economy in a Roman frontier community: Excavations in Tiel-Passewaaij*. Amsterdam: Amsterdam University Press. DOI: <https://doi.org/10.5117/9789089640222>
- Groot, M.** 2009. Searching for patterns among special animal deposits in the Dutch River Area during the Roman period. *Journal of Archaeology in the Low Countries*, 1(2).
- Groot, M.** 2010. *Handboek zoöarcheologie*. Amsterdam: ACVU-HBS.
- Groot, M.** 2013. Een Romeinse eikelmuis uit Hoogeloon. *Cranium*, 30(2): 8–12.
- Groot, M.** 2016. *Livestock for Sale: Animal Husbandry in a Roman Frontier Zone: the Case Study of the Civitas Batavorum*. Amsterdam: Amsterdam University Press.
- Groot, M.** 2017. Developments in Animal Husbandry and Food Supply in Roman Germania Inferior. *European Journal of Archaeology*, 20(3): 451–471. DOI: <https://doi.org/10.1017/ea.2016.31>
- Groot, M and Deschler-Erb, S.** 2015. Market strategies in the Roman provinces: different animal husbandry systems explored by a comparative regional approach. *Journal of Archaeological Science: Reports*, 4: 447–460. DOI: <https://doi.org/10.1016/j.jasrep.2015.10.007>
- Groot, M and Deschler-Erb, S.** 2017. Carnem et circenses—Consumption of Animals and their Products in Roman Urban and Military Sites in two Regions in the Northwestern Provinces. *Environmental Archaeology*, 22(1): 96–112. DOI: <https://doi.org/10.1179/1749631415Y.0000000027>
- Groot, M, Heeren, S, Kooistra, LI and Vos, WK.** 2009. Surplus production for the market? The agrarian economy in the non-villa landscapes of Germania Inferior. *Journal of Roman Archaeology*, 22: 231–252. DOI: <https://doi.org/10.1017/S1047759400020687>
- Groot, M and van Haasteren, M.** 2017. Dierlijk bot. In: van Renswoude, J and Kabermehl, D (eds.), *Opgravingen te Houten-Castellum. Bewoning langs een restgeul in de IJzertijd, Romeinse tijd en Vroege Middeleeuwen (Zuidnederlandse Archeologische Rapporten 65)*, 687–734.
- Habermehl, K.** 1975. Die Altersbestimmung beim Hauswiederkäuer. *Die Altersbestimmung bei Haus- und Labortieren, Berlin, Paul Parey*, 62–63.
- Halici, H.** 2002. Dierlijk botmateriaal. In: Koopstra, CG (ed.), *Archeologisch onderzoek in de Bullepolder, gemeente Leeuwarden. (ARC-Publicaties 52)*, 35–49. Groningen: ARC.
- Hambleton, E.** 1999. *Animal husbandry regimes in Iron Age Britain: a comparative study of faunal assemblages from British Iron Age sites*. British Archaeological Reports Limited.
- Harcourt, RA.** 1974. The dog in prehistoric and early historic Britain. *Journal of archaeological science*, 1(2): 151–175. DOI: [https://doi.org/10.1016/0305-4403\(74\)90040-5](https://doi.org/10.1016/0305-4403(74)90040-5)
- Hiddingh, H.** 1983. Geveugelde maaltijden in de middeleeuwen; Een onderzoek naar vogelresten uit de late middeleeuwen van vier Nederlandse vindplaatsen. Unpublished thesis (PhD), University of Groningen.
- Hoffmann, RC.** 2005. A brief history of aquatic resource use in medieval Europe. *Helgoland Marine Research*, 59(1): 22–30. DOI: <https://doi.org/10.1007/s10152-004-0203-5>
- Huisman, DJ, Lauwerier, RCGM, Jans, MME, Cuijpers, AGFM and Laarman, FJ.** 2009. Bone. In: Huisman, DJ (ed.), *Degradation of archaeological remains*, 33–54. Den Haag: SDU Uitgevers.
- Hullegie, AGJ and Prummel, W.** 2015. Dieren op en rond de Achlumer terp. In: Nicolay, JAW and de Langen, GJ (eds.), *Graven aan de voet van de Achlumer dorpsterp. Archeologische sporen rondom een terpnederzetting. (Jaarverslagen van de Vereniging voor Terpenonderzoek 97)*, 134–159. Groningen: Vereniging voor Terpenonderzoek.

- IJzereef, GF. 1981. *Bronze Age animal bones from Bovenkarspel: the excavation at Het Valkje*. Amersfoort: ROB.
- Jans, MME. 2005. Histological characterisation of diagenetic alteration of archaeological bone. Unpublished thesis (PhD), Institute for Geo and Bioarchaeology, Vrije Universiteit Amsterdam.
- Johansson, F and Hüster, H. 1987. *Untersuchungen an Skelettresten von Katzen aus Haithabu: (Ausgrabung 1966–1969)*. Wachholtz.
- Knol, E. 1983. Farming on the banks of the river Aa. The faunal remains and bone objects of Paddepoel 200 BC–250 AD. *Palaeohistoria*, 25: 154–182.
- Kooistra, LI and Groot, M. 2015. The agricultural basis of the Hoogeloon villa and the wider region. In: Roymans, N, Derks, T and Hiddink, H (eds.), *The Roman Villa of Hoogeloon and the Archaeology of the Periphery*, 141–162. Amsterdam: Amsterdam University Press.
- Kooistra, LI, van Dinter, M, Dütting, MK, van Rijn, P and Cavallo, C. 2013. Could the local population of the Lower Rhine delta supply the Roman army? Part 1: The archaeological and historical framework. *Journal of Archaeology in the Low Countries*, 4(2): 5–23.
- Kooistra, M and Maas, G. 2008. The widespread occurrence of Celtic field systems in the central part of the Netherlands. *Journal of Archaeological Science*, 35(8): 2318–2328. DOI: <https://doi.org/10.1016/j.jas.2008.03.007>
- Kootker, LM and Davies, GR. 2017. Bones, teeth, and invisible tracers. The current state of human bioarchaeological isotope geochemical research in the Netherlands. In: Kluiving, SJ, Kootker, LM and Hermans, RAE (eds.), *Interdisciplinarity between humanities and science. A Festschrift in honour of Prof. Dr. Henk Kars*, 53–71.
- Kootker, LM, Geerdink, C, van den Broeke, PW, Davies, GR and Kars, H. 2018. Breaking traditions: an isotopic study on the changing funerary practices in the Dutch Iron Age (800–12 BC). *Archaeometry*, 60: 594–611. DOI: <https://doi.org/10.1111/arcm.12333>
- Kootker, LM, Van Lanen, RJ, Kars, H and Davies, GR. 2016. Strontium isoscapes in the Netherlands. Spatial variations in  $87\text{Sr}/88\text{Sr}$  as proxy for palaeomobility. *Journal of Archaeological Science: Reports*, 6: 1–13. DOI: <https://doi.org/10.1016/j.jasrep.2016.01.015>
- Kranenburg, H and Prummel, W. in press. The use of domestic and wild animals on S4. In: Raemaekers, DCM and de Roever, JP (eds.), *Swifterbant S4. Occupation and exploitation of a Neolithic levee site (Groningen Archaeological Series)*. Groningen.
- Krause-Kyora, B. 2011. Molekulargenetische und archäologische Untersuchungen zur Domestikation und Züchtung des Schweins (*Sus scrofa*). Christian-Albrechts-Universität.
- Kuijper, W. 1990. Schelpdieren in Romeins Valkenburg (Z.H.). In: Bult, EJ and Hallewas, DP (eds.), *Graven bij Valkenburg; het archeologisch onderzoek in 1987 en 1988*, 134–141.
- Kuijper, WJ, Verheijen, IKA, Ramcharan, A, van der Plicht, H and van Kolfshoten, T. 2016. One of the last wild brown bears (*Ursus arctos*) in the Netherlands (Noordwijk). *Lutra*, 59: 49–64.
- Kuipers, L. 2017. De wilde kat anno 2017. Ontwikkeling van de wilde katpopulatie in Zuid-Limburg. *Rapport in opdracht van Ark*.
- Kuitema, M. 2007. De stabiele isotopen  $^{14}\text{C}$  en  $^{15}\text{N}$  in gedateerd botmateriaal. *Inventarisatie en analyse van de stabiele isotopen uit de database van het CIO te Groningen*. Unpublished thesis (Bachelor), University of Leiden.
- Laarman, F. 1983. Onderzoek van faunaresten uit opgravingen in de Assendelver polder 1988. Unpublished thesis (MSc), University of Amsterdam.
- Laarman, F. 1987. In de put zitten.... (I). Dierlijke overblijfselen uit een Romeinse waterput. In: Bult, EJ and Hallewas, DP (eds.), *Graven bij Valkenburg II; het archeologisch onderzoek in 1986*, 51–55. Delft.
- Laarman, F. 2001. Deel 16. Archeozoölogie: aard en betekenis van dierlijke resten. *De mesolithische en vroeg-neolithische vindplaats Hoge Vaart-A27 (Flevoland), Rapportage Archeologische Monumentenzorg*, 79.
- La Haye, M and Verboom, B. 2017. Year of the serotine or year of the wolf? *Lutra*, 60: 69–71.
- Lammertsma, DR, Kuiters, AT, Niewold, FJJ, Jansman, HAH, Koelewijn, HP, Perez-Haro, MI, Boerwinkel, MC and Bovenschen, J. 2008. Het gaat goed met de otter. *Zoogdier*, 19(2): 3–5.
- Lammertsma, DR and Niewold, FJJ. 2016. Otter *Lutra*. In: Broekhuizen, S, Spoelstra, K, Thissen, JBM, Canters, KJ and Buys, JC (eds.), *Atlas van de Nederlandse Zoogdieren (Natuur van Nederland 12)*, 247–249. Leiden.
- Lanting, J and van der Plicht, J. 1996. De  $^{14}\text{C}$ -chronologie van de Nederlandse Pre-en Protohistorie I.: Laat-Paleolithicum. *Palaeohistoria*, 37(38): 71–125.
- Larson, G, Albarella, U, Dobney, K, Rowley-Conwy, P, Schibler, J, Tresset, A, Vigne, J-D, Edwards, CJ, Schlumbaum, A, Dinu, A, Balacescu, A, Dolman, G, Tagliacozzo, A, Manaseryan, N, Miracle, P, Van Wijngaarden-Bakker, L, Masseti, M, Bradley, DG and Cooper, A. 2007. Ancient DNA, pig domestication, and the spread of the Neolithic into Europe. *Proceedings of the National Academy of Sciences*, 104(39): 15276–15281. DOI: <https://doi.org/10.1073/pnas.0703411104>
- Lauwerier, R. 2013. Food as gravegifts. (Nederlandse Oudheden 17). In: Steures, DC (ed.), *The Late Roman Cemeteries of Nijmegen. Stray Finds and Excavations 1947–1983 (Nederlandse Oudheden 17)*, 255–265. Amersfoort: Rijksdienst voor het Cultureel Erfgoed.
- Lauwerier, RCGM. 1983. Pigs, piglets and determining the season of slaughtering. *Journal of Archaeological Science*, 10(5): 483–488. DOI: [https://doi.org/10.1016/0305-4403\(83\)90063-8](https://doi.org/10.1016/0305-4403(83)90063-8)

- Lauwerier, RCGM.** 1988. *Animals in Roman times in the Dutch eastern river area. (Nederlandse Oudheden 12).* Amersfoort: Rijksdienst voor het Oudheidkundig Bodemonderzoek.
- Lauwerier, RCGM.** 1993. Twenty-eight bird brisquets in a pot; roman preserved food from Nijmegen. *Archaeofauna*, 2: 15–19.
- Lauwerier, RCGM.** 1997. Faunal remains from Dutch Medieval towns. A survey. *Anthropozoologica*, 25–26: 479–486.
- Lauwerier, RCGM.** 2009. Animal husbandry and fishing. In *Ulpia-Noviomagus-Roman Nijmegen: The Batavian Capital at the Imperial Frontier. Journal of Roman Archaeology*, 158–164. Portsmouth, RI: Journal of Roman Archaeology.
- Lauwerier, RCGM.** 2010. *Archeologie en resten van dieren: leidraad archeozoölogie.* Den Haag: Sdu Uitgevers.
- Lauwerier, RCGM.** 2015. Polled cattle in the Roman Netherlands. *Livestock Science*, 179: 71–79. DOI: <https://doi.org/10.1016/j.livsci.2015.05.018>
- Lauwerier, RCGM.** 2017. Knowledge for informed choices. Tools for decision making in archaeological heritage management in the Netherlands. In: Lauwerier, RCGM, Eerden, MC, Groenewoudt, BJ, et al (eds.), *Knowledge for informed Choices. Tools for more effective and efficient selection of valuable archaeology in the Netherlands.* Nederlandse Archeologische Rapporten 55), 11–24. Amersfoort: Rijksdienst voor het Cultureel Erfgoed.
- Lauwerier, RCGM and de Vries, LS.** 2004. Lifting the Iceberg – BoneInfo and the Battle to Save Archaeological Information. In: Lauwerier, RCGM and Plug, I (eds.), *The Future from the Past – Archaeozoology in wildlife conservation and heritage management*, 167–175. Oxford: Oxbow Books.
- Lauwerier, RCGM and Deeben, J.** 2011. Burnt animal remains from Federmesser sites in the Netherlands. *Archäologisches Korrespondenzblatt*, 41(1): 1–20.
- Lauwerier, R and Laarman, F.** 2015. Informatiewaarde van verbrand bot van de bandkeramische nederzetting te Elsloo. In: van Wijk, IM and Porreij-Lyklema, A (eds.), *Opgravingen in de bandkeramische nederzetting van Elsloo-Koolweg. Uitwerking van drie opgravingscampagnes aan de Joannes Rivijsstraat en Paulus Potterstraat te Elsloo. (Archol Rapport 252)*, 169–174. Leiden: Archol.
- Lauwerier, RCGM and Laarman, FJ.** 1999. Dierlijk botmateriaal. In: Hagers, J-KH and Sier, MM (eds.), *Castricum Oosterbuurt, bewoningssporen uit de Romeinse tijd en de Middeleeuwen (=ROB Rapportage Archeologische Monumentenzorg 53)*, 129–151.
- Lauwerier, R, Prummel, W and van Kolfschoten, T.** 2016. Dieren en de steentijd-mens. In: Amkreutz, L, Brounen, F, Deeben, J, Machiels, R, van Oorsouw, MF and Smit, B (eds.), *Vuursteen verzameld. Over het zoeken en onderzoeken van steentijdvondsten en -vindplaatsen.* (Nederlandse Archeologische Rapporten 50), 273–276. Amersfoort.
- Lauwerier, RCGM and Robeerst, JMM.** 2001. Horses in Roman times in the Netherlands. In: Buitenhuis, H and Prummel, W (eds.), *Animals and Man in the Past. Essays in Honour of Dr. A.T. Clason*, 275–289. ARC: ARC-Publicaties.
- Lauwerier, RCGM and Zeiler, JT.** 2001. Wishful thinking and the introduction of the rabbit to the Low Countries. *Environmental Archaeology*, 6: 87–90. DOI: <https://doi.org/10.1179/env.2001.6.1.87>
- Lelieveld, G, Beekers, B, Kamp, J, Klees, D, Linnartz, L, Polman, E and Vermeulen, R.** 2016. The first proof of the recent presence of wolves in the Netherlands. *Lutra*, 59: 23–31.
- Lenders, HJR, Chamuleau, TPM, Hendriks, AJ, Lauwerier, RCGM, Leuven, RSEW and Verberk, WCEP.** 2016. Historical rise of waterpower initiated the collapse of salmon stocks. *Scientific Reports*, 6. DOI: <https://doi.org/10.1038/srep29269>
- Litjens, BEJ and Pelzers, E.** 1988. The damhert Dama dama in Nederland. *Lutra*, 31: 132–144.
- Louwe Kooijmans, L.** 2001. Synthese. *Hardinxveld-De Bruin: Een kampplaats uit het Laat-Mesolithicum en het begin van de Swifterbant-cultuur (5500–4450 v.Chr.)*, 499–528.
- Louwe Kooijmans, L and Nokkert, M.** 2001. Sporen en structuren. *Archeologie in de Betuweroute. Hardinxveld-Giessendam De Bruin. Een kampplaats uit het Laat-Mesolithicum en het begin van de Swifterbant-cultuur (5500–4450 v.Chr.)*, Amersfoort (Rapportage Archeologische Monumentenzorg 88), 75–115.
- Louwe Kooijmans, LP.** 1974. *The Rhine/Meuse Delta; four studies on its prehistoric occupation and Holocene geology (Analecta Praehistorica Leidensia 7).* Leiden.
- Louwe Kooijmans, LP.** 1993. Wetland exploitation and upland relations of prehistoric communities in the Netherlands. In: Gardiner, J (ed.), *Flatland and Wetlands: Current Themes in East Anglian Archaeology*, 71–116. Norwich: Scole Archaeological Committee for East Anglia.
- Magdanas, JR and Waasdorp, JA.** 1989. *Franken aan de Frankenslag; een vroeg-middeleeuwse nederzetting in 's-Gravenhage (VOM-reeks 1089 nr 2).* Dan Haag: VOM.
- Manning, K, Timpson, A, Shennan, S and Crema, E.** 2015. Size reduction in early European domestic cattle relates to intensification of Neolithic herding strategies. *PloS one*, 10(12): e0141873. DOI: <https://doi.org/10.1371/journal.pone.0141873>
- Matolcsi, J.** 1970. Historische Erforschung der Körpergröße des Rindes auf Grund von ungarischem Knochenmaterial. *Zeitschrift für Tierzüchtung und Züchtungsbiologie*, 87(1–4): 89–137. DOI: <https://doi.org/10.1111/j.1439-0388.1970.tb01330.x>
- May, E.** 1985. Widerristhöhe und Langknochenmasse bei Pferden—ein immer noch aktuelles Problem. *Zeitschrift für Säugetierkunde*, 50(6): 368–381.
- Mulder, JL.** 2016a. *Lynx Lynx Lynx.* In: Broekhuizen, S, Spoelstra, K, Thissen, JBM, Canters, KJ and Buys, JC (eds.), *Atlas van de Nederlandse Zoogdieren (Natuur van Nederland 12)*, 234–235. Leiden.
- Mulder, JL.** 2016b. Wilde kat *Felis silvestris*. In: Broekhuizen, S, Spoelstra, K, Thissen, JBM, Canters,



- KJ and Buys, JC (eds.), *Atlas van de Nederlandse Zoogdieren (Natuur van Nederland 12)*, 232–233. Leiden.
- Nieuwhof, A.** 2006. *De wierde Wierum (provincie Groningen). Een archeologisch steilkantonderzoek (Groningen Archaeological Studies 3)*. Groningen: Barkhuis Publishing & Groningen University Library.
- Oversteegen, JFS, van Wijngaarden-Bakker, LH, Maliepaard, R and van Kofschoten, T.** 2001. Zoogdieren, vogels en reptielen. In: Kooijmans, LP (ed.), *Archeologie in de Betuweroute. Hardinxveld-Giesendam Polderweg. Een Mesolithisch jachtkamp in het rivierengebied (5500–5000 v. Ch.) (Rapportages Archeologisch Monumentenzorg 88)*, 209–297. Amersfoort: Rijksdienst voor het Oudheidkundig Bodemonderzoek.
- Pavlovic, A and Nieweg, DC.** 2006. *Archeologisch onderzoek Vijverhof (Valkhuis), gemeente Den Haag. (rapport 0605 Afdeling Archeologie, Dienst Stadsbeheer, gemeente Den Haag)*. Den Haag.
- Peeters, H and Hogestijn, JWH.** 2001. *De mesolithische en vroeg-neolithische vindplaats Hoge Vaart-A27 (Flevoland): Rapportage Archeologische Monumentenzorg 79, deel 20*. Amersfoort: Rijksdienst voor het Oudheidkundig Bodemonderzoek.
- Prummel, W.** 1977. Een bewerkt damhertbeen uit het Romeinse castellum te Valkenburg (Z.H.). *Westerheem*, 26: 4–14.
- Prummel, W.** 1983. *Excavations at Dorestad 2. Early medieval Dorestad, an archaeozoological study, Nederlandse Oudheden 11*. Amersfoort: ROB.
- Prummel, W.** 1987a. Atlas for identification of foetal skeletal elements of Cattle, Horse, Sheep and Pig. *Part*, 2: 11–41.
- Prummel, W.** 1987b. Atlas for the identification of foetal skeletal elements of cattle, horse, sheep and pig, part I. *Archaeozoologia*, 1(1): 23–30.
- Prummel, W.** 1987c. The faunal remains from the neolithic site of Hekelingen III. *Helinium*, 27(2): 190–258.
- Prummel, W.** 1987d. Poultry and fowling at the Roman castellum Velsen 1. *Palaeohistoria*, 29: 183–201.
- Prummel, W.** 1988. Atlas for identification of foetal skeletal elements of cattle, horse, sheep and pig. *Part*. 3. *Archaeozoologia II* (1–2), 13–26.
- Prummel, W.** 1989. Appendix to atlas for identification of foetal skeletal elements of cattle, horse, sheep and pig. *Part*, 2: 71–78.
- Prummel, W.** 1992. Early medieval dog burials among the Germanic tribes. *Helinium*, 32(1–2): 132–194.
- Prummel, W.** 1993. Paarden en honden uit vroeg-middeleeuwse grafvelden. In: Drenth, E, Hessing, WAM and Knol, E (eds.), *Het tweede leven van onze doden*, 53–60. Amersfoort: NAR.
- Prummel, W.** 2006a. Bronze age dogs from graves in Borger (Netherlands) and Dimini (Greece). In: Snyder, LM and Moore, EA (eds.), *Dogs and People in Social, Working, Economic or Symbolic Interaction*, 67–76. Oxford: Oxbow Books.
- Prummel, W.** 2006b. Dierlijk bot. In: Nieuwhof, A (ed.), *De wierde Wierum (provincie Groningen). Een archeologisch steilkantonderzoek (Groningen Archaeological Studies, 3)*, 31–45. Groningen: Barkhuis.
- Prummel, W.** 2013. Falconry in continental settlements as reflected by animal bones from the 6th to 12th centuries AD. In: Grimm, O and Schmölcke, U (eds.), *Hunting in northern Europe until 1500 AD. Old traditions and regional developments, continental influences. Schriften des archäologischen landesmuseums, Ergänzungsreihe 7*, 357–377. Neumünster: Wacholtz.
- Prummel, W.** 2014. Schapenteelt in terpengebied belangrijker dan gedacht. *Van Wierden en Terpen, Mededelingen van de Vereniging voor Terpenonderzoek*, 19: 19–21.
- Prummel, W.** 2018. The archaeological-archaeozoological identification of falconry – methodological remarks and some Dutch examples. In: Gersmann, K and Grimm, O (eds.), *Raptor and human – falconry and bird symbolism throughout the millennia on a global scale (Advanced studies of archaeology and history of hunting, vol. 1.1–1.4)*, 467–478. Kiel/Hamburg: Wacholz Verlag, Murmann Publishers.
- Prummel, W, Esser, E and Zeiler, J.** 2013. The animals on the terp at Wijnaldum-Tjitsma (The Netherlands)–reflections on the landscape, economy and social status. *Settlement and Coastal Research in the Southern North Sea Region*, 36: 87–98.
- Prummel, W and Frisch, H.** 1986. A guide for the distinction of species, sex and body side in bones of sheep and goat. *Journal of archaeological Science*, 13(6): 567–577. DOI: [https://doi.org/10.1016/0305-4403\(86\)90041-5](https://doi.org/10.1016/0305-4403(86)90041-5)
- Prummel, W and Niekus, MJLT.** 2011. Late Mesolithic hunting of a small female aurochs in the valley of the River Tjonger (the Netherlands) in the light of Mesolithic aurochs hunting in NW Europe. *Journal of Archaeological Science*, 38(7): 1456–1467. DOI: <https://doi.org/10.1016/j.jas.2011.02.009>
- Prummel, W, Niekus, MJT, van Gijn, AL and Cappers, RT.** 2002. A Late Mesolithic kill site of aurochs at Jardinga, Netherlands. *Antiquity*, 76(292): 413–424. DOI: <https://doi.org/10.1017/S0003598X00090529>
- Prummel, W and Niekus, MT.** 2005. De laatmesolithische vindplaats Jardinga: de opgravingen in 2002 en 2003. *Paleo-aktueel*, 14(15): 31–37.
- Prummel, W, Niekus, MT, Van der Meulen, S and Fens, R.** 2009b. Mesolithische botten uit het dal van de Tjonger. *Paleo-aktueel*, 20: 25–31.
- Prummel, W and Olivier, L.** 2008. Twee bijzondere terpvondsten uit Holwerd. *Van Wierden en Terpen, Mededelingen van de Vereniging voor Terpenonderzoek*, 5: 19–21.
- Prummel, W, Raemaekers, DCM, Beckerman, SM, Bottema, N, Mac Gillavry, JN, Cappers, RTJ, Cleveringa, P, Devriendt, I, de Wolf, H and Zeiler, JT.** 2009a. Terug naar Swifterbant. Een kleinschalige

- opgraving te Swifterbant-S2 (gemeente Dronten). *Archeologie*, 13: 17–47.
- Prummel, W, van Gent, JT and Kompanje, EJO.** 2012. Walvisbotten uit Friese en Groninger terpen. *Paleo-aktueel*, 23: 41–48.
- Raemaekers, D.** 2003. Cutting a long story short? The process of neolithization in the Dutch delta re-examined. *Antiquity*, 77(298): 740–748. DOI: <https://doi.org/10.1017/S0003598X00061688>
- RCE.** 2018. BoneInfo. Available at <https://archeologieinederland.nl/boneinfo> [Last accessed 18 November 2018].
- Reichstein, H.** 1991. *Die Fauna des germanischen Dorfes Feddersen Wierde*. Steiner.
- Rompelman, E and Eeltink, N.** 2011. Zutphen-Stadhuis 1997, vondstnummer 862. Een zoölogische quickscan met aanvullende gegevens. *Aestimatica Publicatie*, 27.
- Rowley-Conwy, P.** 2011. Westward Ho! The spread of agriculture from Central Europe to the Atlantic. *Current anthropology*, 52(S4): S431–S451. DOI: <https://doi.org/10.1086/658368>
- Rowley-Conwy, P.** 2014. Foragers and farmers in Mesolithic/Neolithic Europe, 5500–3900 cal BC: beyond the anthropological comfort zone. In: *Wild things: recent advances in palaeolithic and mesolithic research*, 185–201. Oxford: Oxbow Books. DOI: <https://doi.org/10.2307/j.ctvh1dhm5.19>
- Royal Dutch Academy of Sciences (KNAW).** 2018. DANS EASY. Available at <https://easy.dans.knaw.nl/ui/home> [Last accessed 18 November 2018].
- Schaftberg, R and Swalve, HH.** 2015. The history of breeding for polled cattle. *Livestock science*, 179: 54–70. DOI: <https://doi.org/10.1016/j.livsci.2015.05.017>
- Schelvis, J.** 1992. Mites and Archaeozoology. Unpublished thesis (PhD), Rijksuniversiteit Groningen.
- Silver, IE.** 1969. The ageing of domestic animals. In: Brothwell, DR and Higgs, ES (eds.), *Science and Archaeology. A survey of progress and research*, 250–268. Thames and Hudson, London.
- Sluiter, H.** 2003. The reintroduction and the present status of the beaver (*Castor fiber*) in the Netherlands: an overview. *Lutra*, 46(429): 133.
- Smits, E, Millard, AR, Nowell, G and Pearson, DG.** 2010. Isotopic investigation of diet and residential mobility in the Neolithic of the Lower Rhine Basin. *European Journal of Archaeology*, 13(1): 5–31. DOI: <https://doi.org/10.1177/1461957109355040>
- Smits, I and van der Plicht, H.** 2009. Mesolithic and Neolithic human remains in the Netherlands: physical anthropological and stable isotope investigations. *Journal of Archaeology in the Low Countries*, 1(1): 55–85.
- Snijders, F and Broertjes, J.** 2016. *Bivak aan de Beerze, deel 2. Laatpaleolitische en mesolithische vindplaatsen te Westelbeers. Uitgave van de Voormalige Vereniging Archeologische Werkgroep 't Oude Slot*. Veldhoven.
- Speller, C, van den Hurk, Y, Charpentier, A, Rodrigues, A, Gardeisen, A, Wilkens, B, McGrath, K, Rowsell, K, Spindler, L, Collins, M and Hofreiter, M.** 2016. Barcoding the largest animals on Earth: ongoing challenges and molecular solutions in the taxonomic identification of ancient cetaceans. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 371(1702). DOI: <https://doi.org/10.1098/rstb.2015.0332>
- Sykes, N.** 2012. A social perspective on the introduction of exotic animals: The case of the chicken. *World Archaeology*, 44(1): 156–165. DOI: <https://doi.org/10.1080/00438243.2012.646104>
- Sykes, NJ, Baker, KH, Carden, RF, Higham, TF, Hoelzel, AR and Stevens, RE.** 2011. New evidence for the establishment and management of the European fallow deer (*Dama dama*) in Roman Britain. *Journal of Archaeological Science*, 38(1): 156–165. DOI: <https://doi.org/10.1016/j.jas.2010.08.024>
- Teichert, M.** 1969. Osteometrische Untersuchungen zur Berechnung der Widerristhöhe bei vor- und frühgeschichtlichen Schweinen. *Kühn Archiv*, 237–292.
- Teichert, M.** 1975. Osteometrische Untersuchungen zur Berechnung der Widerristhöhe bei schafen. *Archaeozoological studies*, 51–69.
- Terpstra, H.** 1986. Runderen van Ezinge. Unpublished thesis (MSc), Rijksuniversiteit Groningen.
- Thilderkvist, JGM.** 2013. Ritual bones or common waste. Unpublished thesis (PhD), Rijksuniversiteit Groningen. DOI: <https://doi.org/10.2307/j.ctt2204s6c>
- Timmerman, D.** 2012. De rol en het gebruik van dieren en het landschap in het terpengebied in de vroege middeleeuwen aan de hand van de case-study Firdgum. Unpublished thesis (Master), University of Groningen.
- van Amerongen, Y.** 2014. Het wilde West-Friesland: jacht en visserij in de bronstijd. In: Theunissen, EM and Arnoldussen, S (eds.), *Metaaltijden 1. Bijdragen in de studie van de metaaltijden*, 81–96. Leiden: Sidestone Press.
- van Amerongen, Y.** 2016. Wild west Frisia. The role of domestic and wild resource exploitation in Bronze Age subsistence. (Archaeological Studies Leiden University 33). Unpublished thesis (PhD), Rijksuniversiteit Groningen.
- van Dam, PJEM.** 2001. De rol van de warande. Geschiedenis van de inburgering met het konijn. *Jaarboek voor Ecologische Geschiedenis 2000 (Vreemde delingen in de natuur)*, 2000: 59–85.
- van den Broeke, PW.** 2005. Toenemende verscheidenheid: synthese. In: Kooijmans, L, van den Broeke, PW, Fokkens, H and van Gijn, A (eds.), *Nederland in de prehistorie*, 683–694. Amsterdam.
- van den Hurk, Y.** 2014. A Whale of a Question: History and Social Implications of Cetacean Exploitation in the North Sea Area through Space and Time. Unpublished thesis (Masters), University of Nottingham.
- van der Jagt, I.** 2017a. Sperwer-*Accipiter nisus*. In: Esser, E and Lauwerier, R (eds.), *Archeozoologisch Vogel-ABC*, 79–82. Oostvoorne/Amersfoort.
- van der Jagt, IMM.** 2017b. *Dierlijk botmateriaal grafveld Lent. Intern rapport RCE*. Amersfoort: RCE.

- van der Jagt, IMM, Kootker, LM, van Kolfschoten, T, Kars, H and Davies, G.** 2012. An insight into animal exchange in Early Medieval Oegstgeest: a combined archaeozoological and isotopic approach. In: Raemaekers, DCM, Esser, E, Lauwerier, RCGM and Zeiler, JT (eds.), *A bouquet of archaeological studies. Essays in honour of Wietske Prummel*. GAS 21, 141–151.
- van der Jagt, IMM, Laarman, FJ, Kuijper, W, Nieman, AM, van Os, BJH and Zwaan, JC.** 2014. Dierlijk materiaal. In: Lauwerier, RCGM and de Kort, JE (eds.), *Merovingers in een villa 2. Romeinse villa en Merovingisch grafveld Borgharen – Pasestraat, onderzoek 2012 (RAM 222)*, 157–190. Amersfoort.
- van Dijk, J.** 2015. Iron Age animal husbandry in the wetlands of the western Netherlands. *Environmental Archaeology*, 21(1): 45–58. DOI: <https://doi.org/10.1179/1749631414Y.00000000060>
- van Dijk, J and Groot, M.** 2013. The Late Iron Age-Roman transformation from subsistence to surplus production in animal husbandry in the Central and Western parts of the Netherlands. In: Groot, M, Lentjes, DM and Zeiler, J (eds.), *Barely surviving or more than enough? The environmental archaeology of subsistence, specialisation and surplus food production*, 175–200. Leiden: Sidestone.
- van Giffen, A.** 1963. Het bouwoffer uit de oudste hoeve te Ezinge (Gr.). *Helinium*, 3: 246–253.
- van Haaster, H, Zeiler, JT and Brinkhuizen, DC.** 2012. *De voedingsgewoonten van (post) middeleeuws Alkmaar. Resultaten van het archeobotanisch en archeozoologisch onderzoek*. BIAxiaal 453. Zaandam.
- van Haasteren, M and Groot, M.** 2013. The biography of wells: a functional and ritual life history. *Journal of Archaeology in the Low Countries*, 4(2): 25–51. DOI: <https://doi.org/10.1111/gend.2013.25.issue-3>
- van Houtte, JA.** 1977. *An Economic History of the Low Countries (800–1800)*. London: Weidenfeld and Nicolson Ltd.
- van Kolfschoten, T.** 1998. The *Allophaiomys* record from Zuurland, the Netherlands. *Paludicola*, 2(1): 110–115.
- van Kolfschoten, T.** 2001. Pleistocene mammals from the Netherlands. *Bollettino-Societa Paleontologica Italiana*, 40(2): 209–216.
- Van Kolfschoten, T and Tesakov, AS.** 1998. The late Pliocene *Mimomys hordijki* sp. nov. from the Zuurland borehole (the Netherlands). *Mededelingen Nederlands Instituut voor Toegepaste Geowetenschappen TNO*, 60: 187–192.
- van Kolfschoten, T, Tesakov, AS and Bell, CJ.** 2018. The first record of *Phenacomys* (Mammalia, Rodentia, Cricetidae) in Europe (Early Pleistocene, Zuurland, The Netherlands). *Quaternary Science Reviews*, 192: 274–281. DOI: <https://doi.org/10.1016/j.quascirev.2018.06.005>
- van Kolfschoten, T, van der Jagt, I, Beeren, Z, Argiti, V, van der Leije, J, van Essen, H, Busschers, FS, Stoel, P and van der Plicht, H.** 2011. A remarkable collection of Late Pleistocene reindeer (*Rangifer tarandus*) remains from Woerden (The Netherlands). *Quaternary international*, 238(1–2): 4–11. DOI: <https://doi.org/10.1016/j.quaint.2010.12.033>
- van Kolfschoten, T and Vervoort-Kerkhoff, Y.** 1999. The Pleistocene and Holocene Mammalian assemblages from the Maasvlakte near Rotterdam (the Netherlands): with special reference to the *Ovibovini Soergelia minor* and *Praeovibos cf. priscus*/. In: Reumer, JWF and de Vos, J (eds.), *Elephants have a snorkel! Papers in honour of Paul Y. Sondaar – DEINSEA 7*, 373–386.
- van Londen, H.** 2006. Midden-Delfland: the roman native landscape past and present. Unpublished thesis (PhD), University of Amsterdam.
- van Straalen, D.** 2018. Zeearend *Haliaeetus albicilla*. In: *SOVON Vogelonderzoek Nederland 2018, Vogelatlas van Nederland*, 214–215. Utrecht/Antwerpen: Kosmos Uitgevers.
- van Wijngaarden-Bakker, LH.** 1988. Zoöarcheologisch onderzoek in de west-Nederlandse delta 1983–1987. *Archeologie en oecologie van Holland tussen Rijn en Vlie*, 154–185.
- van Wijngaarden-Bakker, LH, Cavallo, C, van Kolfschoten, T, Maliepaard, CH and Oversteegen, HFS.** 2001. Zoogdieren, vogels, reptielen. In: Kooijmans, LP (ed.), *Archeologie in de Betuweroute. Hardinxveld-Giessendam Polderweg. Een Mesolithisch jachtkamp in het rivierengebied (5500–5000 v. Ch.) (Rapportages Archeologisch Monumentenzorg 83)*, 181–242. Amersfoort: Rijksdienst voor het Oudheidkundig Bodemonderzoek.
- Verhagen, M.** 1987. Een Romeins paardengraf uit het grafveld. In: Bult, EJ and Hallewas, DP (eds.), *Graven bij Valkenburg 2. Het archeologisch onderzoek in 1986*, 92–98. Delft.
- Von den Driesch, A.** 1976. *A guide to the measurement of animal bones from archaeological sites: as developed by the Institut für Palaeoanatomie, Domestikationsforschung und Geschichte der Tiermedizin of the University of Munich*. Peabody Museum Press.
- Von den Driesch, A and Boessneck, J.** 1974. Kritische Anmerkungen zur Widerristhöhenberechnung aus Längenmassen vor- und frühgeschichtlicher Tierknochen. *Säugetierkundliche Mitteilungen*, 22(4): 325–348.
- Vos, P and de Vries, S.** 2013. *2e generatie palaeogeografische kaarten van Nederland (versie 2.0)*. Utrecht: Deltares.
- Vos, PC and Knol, E.** 2015. Holocene landscape reconstruction of the Wadden Sea area between Marsdiep and Weser. Explanation of the coastal evolution and visualisation of the landscape development of the Northern Netherlands and Niedersachsen in five palaeogeographical maps from 500 BC to present. In: Vos, PC (ed.), *Origin of the Dutch Coastal landscape. Long-term landscape evolution of the Netherlands during the Holocene, described and visualized in national, regional and local palaeogeographical map*



- series, 202–229. Groningen: Barkhuis. DOI: <https://doi.org/10.1017/njg.2015.4>
- Voslamber, B** and **Koffijberg, K.** 2018. Grauwe gans *Anser anser*. In: *SOVON vogelonderzoek Nederland 2018, Vogelatlas van Nederland*, 68–69. Utrecht/Antwerpen: Kosmos Uitgevers.
- Woltering, P.** 2001. Occupation history of Texel, IV. Middle Bronze Age-Late Iron Age. *Berichten van de Rijksdienst voor het Oudheidkundig Bodemonderzoek*, 44: 9–396.
- Zeiler, J.** 1996. De faunaresten van Schagen Witte Paal III (1e–3e eeuw na Chr). *ArchaeoBone Rapport*, 8.
- Zeiler, J.** 1997a. Offers en slachtoffers. Faunaresten uit de Fortunatempel te Nijmegen (2e eeuw n. Chr.).
- Zeiler, J.** 2001. Archeozoölogie. In: Sier, MM and Koot, CW (eds.), *Archeologie in de Betuweroute 'Kesteren-De Woerd'. Bewoningssporen uit de IJzertijd en de Romeinse Tijd. (Rapportage Archeologische Monumentenzorg 82)*, 217–293. Amersfoort: Rijksdienst voor het Oudheidkundig Bodemonderzoek.
- Zeiler, J.** 2005. Paardenrookvlees uit Kesteren (Gld.). *Paleo-aktueel*, 14(15): 143–145.
- Zeiler, JT.** 1997b. *Hunting, fowling and stock-breeding at Neolithic sites in the western and central Netherlands*. Proefschrift Rijksuniversiteit Groningen. edn. Groningen: ArchaeoBone.
- Zeiler, JT.** 2006. Mammals. In: Louwe Kooijmans, LP and Jongste, PFB (eds.), *Schipluiden. A Neolithic settlement on the Dutch North Sea coast c. 3500 cal BC. Leiden (Analecta Praehistorica Leidensia 37/38)*, 375–420. (Analecta Praehistorica Leidensia 37/38) edn. Leiden University.
- Zeiler, JT.** 2014. Birds for the Elite? Fowling in het Northern Netherlands in the Roman Period and the Early Middle Ages. *International Journal of Osteoarchaeology*, 24: 378–383. DOI: <https://doi.org/10.1002/oa.2370>
- Zeiler, JT** and **Brinkhuizen, DC.** 2013. Faunal Remains. In: Kleijne, JP, Brinkkemper, O, Lauwerier, RCGM, Smit, BI and Theunissen, EM (eds.), *A matter of life and death at Mienakker (The Netherlands). Late Neolithic behavioural variability in a dynamic landscape*, 155–173. Amersfoort: Cultural heritage agency of the Netherlands.
- Zeiler, JT** and **Brinkhuizen, DC.** 2015. Fauna. In: Moree, JM and Sier, MM (eds.), *Interdisciplinary archaeological research programme Maasvlakte 2, Rotterdam (=BOORrapporten 566). Part 1 twenty metres deep! The Mesolithic period at the Yangtze Harbour Site – Rotterdam Maasvlakte, the Netherlands. Early Holocene landscape development and habitation*, 201–221. Rotterdam: Bureau Oudheidkundig Onderzoek.
- Zeiler, JT** and **Brinkhuizen, DC.** 2016. *Terug naar 't Klumke. Archeozoologisch onderzoek van dierlijke resten van de vindplaats 't Klumke te Oosterhout, gemeente Nijmegen (Neolithicum, Late Bronstijd/Vroege IJzertijd en Vroege Middeleeuwen). (ArchaeoBone rapport nr. 132)*. Nijmegen: Gemeente Nijmegen.
- Zeiler, JT** and **Clason, AT.** 1993. Fowling in the Dutch Neolithic at inland and coastal sites. *Archaeofauna*, 2: 67–74.
- Zeiler, JT** and **Kompanje, EJ.** 2010. A killer whale (*Orcinus orca*) in the castle: first find of the species in a Dutch archaeological context. *The bat, the weevil and other 'species'*, 101.
- Zietzschmann, O** and **Krollig, O.** 1955. *Lehrbuch der Entwicklungsgeschichte der Haustiere*. Hamburg/Berlin: P. Parey.

**How to cite this article:** Çakırlar, C, van den Hurk, Y, van der Jagt, I, van Amerongen, Y, Bakker, J, Breider, R, van Dijk, J, Esser, K, Groot, M, de Jong, T, Kootker, L, Steenhuisen, F, Zeiler, J, van Kolfschoten, T, Prummel, W, and Lauwerier, R. 2019. Animals and People in the Netherlands' Past: >50 Years of Archaeozoology in the Netherlands. *Open Quaternary*, 5: 13, pp. 1–30. DOI: <https://doi.org/10.5334/oq.61>

**Submitted:** 20 March 2019

**Accepted:** 28 September 2019

**Published:** 19 November 2019

**Copyright:** © 2019 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.