Another advanced test of theory of mind: evidence from very high functioning adults with autism or Asperger Syndrome

Simon Baron-Cohen+, Therese Jolliffe, Catherine Mortimore*, and Mary Robertson*

Departments of Experimental Psychology and Psychiatry,

University of Cambridge, Downing St, Cambridge, CB2 3EB, UK.

*Raymond Way Neuropsychiatry Group, Institute of Neurology,

University of London, Queen Square, London, WC1, UK.

Abstract

Previous studies have found a subgroup of people with autism or Asperger Syndrome (AS) who pass second-order tests of theory of mind. However, such tests have a ceiling in developmental terms corresponding to a mental age of about 6 years old. It is therefore impossible to say if such individuals are intact or impaired in their theory of mind skills. We report the performance of very high functioning adults with autism or AS on an <u>adult</u> test of theory of mind ability. The task involved inferring the mental state of a person just from the information in photographs of a person's eyes. Relative to age-matched normal controls and a clinical control group (adults with Tourette Syndrome), the group with autism and AS were significantly impaired on this task. The autism and AS sample was also impaired on Happé's strange stories tasks. In contrast, they were unimpaired on two control tasks: recognizing gender from the eye region of the face, and recognizing basic emotions from the whole face. This provides evidence for subtle mindreading deficits in very high functioning individuals on the autistic continuum.

There is considerable evidence that the majority of children with autism have impairments in the development of a theory of mind (see Baron-Cohen, 1993,1995, for reviews). Such a deficit may underlie the social, communicative, and imaginative abnormalities that are diagnostic of the condition, since a theory of mind is necessary for normal development in each of these three areas. The theory of mind deficit appears to be expressed very early, from at least the end of the first year of life, as joint attention deficits (Sigman, Mundy, Ungerer, & Sherman, 1986; Baron-Cohen, 1989a; Baron-Cohen, Allen, & Gillberg, 1992; Baron-Cohen, Cox, Baird, et al, 1996a).

However, there is some evidence that appears to contradict the notion that a theory of mind deficit is a core cognitive deficit in autism. First, Bowler (1992) found that adults with Asperger Syndrome (AS) - who share the social and communicative symptoms of autism but who had no history of language delay - pass second-order theory of mind tests. Second-order theory of mind tests involve the subject reasoning about what one person thinks about another person's thoughts. Ozonoff and her colleagues (Ozonoff, Pennington, & Rogers, 1991; Ozonoff, Rogers, & Pennington, 1991) also found some adults with "high-functioning autism" or AS who passed second-order theory of mind tests. Both studies contradict the earlier finding from Baron-Cohen (1989b) that in autism there is an impairment in theory of mind ability in autism. In the latter study, no subjects with autism passed the second-order test of theory of mind.

However, these studies cannot be taken as conclusive evidence for an <u>intact</u> theory of mind in such individuals with autism or AS, because such second-order tests used can

easily produce ceiling effects if used with subjects with a mental age above 6 years old. This is because children with normal intelligence pass second-order theory of mind tasks at about 6 years old (Perner & Wimmer, 1985). It is unfortunate that many workers in this field have thought of second-order tests as "complex" or high-level tests of theory of mind. Certainly, they are more complex than first-order tests (in which the subject simply has to infer the thoughts of another person) [Wimmer & Perner, 1983; Baron-Cohen, Leslie, & Frith, 1985] - but recall that normal 4 year old children pass first-order tests.

In short, neither first or second-order tests are complex tests of theory of mind. They are simply probes for 4 or 6 year old level skills in this domain, respectively. Perhaps if they had been labelled as such, noone would have ever thought of them as suitable tests of whether an <u>adult</u> (with autism, AS, or any other condition) has a fully functional theory of mind. Finding a 30 year old individual with autism, of normal intelligence, who can pass a theory of mind test at the level of a normal 6 year old does not lead to the conclusion that they are necessarily normal in this domain. All we can conclude is that they have intact theory of mind skills at the level of a 6 year old.

Happé's (1994a) study is the only one to take this issue seriously. She tested adults with autism or AS on an "advanced" theory of mind task. This involved story comprehension, where the key questions in the task either concerned a character's mental states (the experimental condition) or physical events (the control condition). Happe's task was pitched at the level of a normal 8-9 year old, and in this respect, it is certainly more advanced than previous tests of theory of mind. She found that adults with autism or AS

had more difficulty with the mental state stories than matched controls, and that they used fewer appropriate mental state terms in their justifications of why characters behaved as they did.

In the present study we used a new, <u>adult</u> test of theory of mind competence, as another advanced test, with which to test high-functioning adults with autism or AS. This extends Happé's line of research. The task involves looking at photographs of the eye region of the faces, and making a forced choice between which of two words best describes what the person (in the photograph) might be thinking or feeling. The task is called the 'Reading the Mind in the Eyes' Task, or the Eyes Task for short.

The Eyes Task involves theory of mind skills in the sense that the subject has to understand mental state terms and match them to faces (or parts of faces, in this case). The choice is always between two mental state terms, some of which are "basic", in Ekman's (1992) sense, (such as happy, sad, angry, or afraid), and others of which are more "complex", (such as reflective, arrogant, scheming, planning, etc.). The forced choice method for interpreting faces in terms of mental states has been used successfully before (Baron-Cohen et al, 1996b). The task could equally be called a test of 'mindreading' in that nothing in the test addresses whether the subject is using a 'theory' or not. Here, as elsewhere, we use the terms 'theory of mind' and 'mindreading' interchangeably. An earlier study using this task found that parents of children with AS perform significantly worse than matched controls, whilst performing significantly better on the Embedded Figures Task (Baron-Cohen, & Hammer, in press a).

In the study reported below, 3 groups of subjects were compared on the Eyes Task: adults with autism or AS, normal adults, and a clinical control group, adults with Tourette Syndrome (TS). The latter was chosen because of the following similarities between autism, AS, and TS: (i) they all had intelligence in the normal range; (ii) they had all suffered from a developmental disorder since childhood; (iii) these disorders all cause disruptions to both normal schooling, and normal peer relations; (iv) these disorders are also all postulated to involve frontal abnormalities (Bishop, 1992; Baron-Cohen, Robertson, & Moriarty, 1994; Baron-Cohen et al, 1994); (v) and all have a sizeable genetic aetiology (Robertson, 1994; Bolton & Rutter, 1990). (vi) These disorders also all affect males more than females.

Naturally, there are many differences between autism and AS on the one hand, and TS on the other (e.g., different frontal abnormalities are implicated in these disorders), but the similarities they share serve to control for the presence of an organic, childhood-

onset psychiatric disorder. We predicted that despite their psychiatric history, patients with Tourette Syndrome would be unimpaired on this advanced theory of mind test, whilst the subjects with autism or AS would show a significant impairment on this test.

We make the assumption that the Eyes Task is a theory of mind task, because of the mental state attribution component. However, this assumption warrants direct testing. This was possible because subjects in the present study using the Eyes Task also took

part in a separate study using Happé's (1994) Strange Stories Task (Jolliffe, 1997). We predicted that if both the Eyes Task and the Strange Stories Task were both indexes of a relatively advanced theory of mind, then if subjects had difficulties with one of these tasks, they should also have difficulties with the other.

The Eyes Task also involves some other process, namely basic aspects of emotion recognition and face perception. In order to test if difficulties on the Eyes Task were specific to the mental state attribution component or were due to these other processes, we included two control tasks: a basic emotion recognition task (adapted from Ekman, 1992), and a gender recognition task.

Finally, if the Eyes Task was validated as an adult test of theory of mind, this afforded us the opportunity to test for a subtle sex difference in the normal group. Our folk psychology would lead us to expect that normal females may be superior to normal males in the domain of social sensitivity or empathy, but most previous theory of mind research has not used sufficiently subtle tests to evaluate if there is any basis to this. (An exception is a study by Hall, 1977). We therefore analysed larger numbers of normal male and female subjects separately, to examine this possibility.

The Experiment

Subjects

3 groups of subjects were tested:

<u>Group 1</u> comprised 16 subjects with high functioning autism (HFA) or AS (4 with highfunctioning autism and 12 with AS). The sex ratio was 13:3 (m:f). The HFA Group all showed a history of 'classical' autism (i.e. autism accompanied by language delay) and fulfilled established diagnostic criteria (American Psychiatric Association, DSM-IV, 1994). Note that because they were high-functioning adults, they would be considered "residual" cases. The AS Group all met the same criteria for autism, but without any clinically significant language delay (i.e., they had single words by 2, or phrase speech by 3, as reported by their parent). They thus met criteria for AS as defined in ICD-10 (World Health Organisation, 1994). They were all of normal intelligence. As such, they are relatively rare. They can be considered as cases of "pure" autism or AS, unconfounded by mental handicap. They were recruited from a variety of clinical sources, as well as an advert in the National Autistic Society magazine, <u>Communication</u>.

<u>Group 2</u> comprised 50 normal age matched adults (25 male and 25 female), drawn from the general population of Cambridge (excluding members of the University), all of whom had no history of any psychiatric condition (as established by self-report). They were selected randomly from the subject panel held in the University Department. IQ

information was not collected on subjects in this group, but they were assumed to all have intelligence in the normal range.

<u>Group 3</u> comprised 10 adult patients with Tourette Syndrome (TS), also age matched with Groups 1 and 2. The sex ratio was 8:2 (m:f), thus mirroring the ratio in Group 1.They were attending a tertiary referral centre in London, and had all been diagnosed by a leading expert in the field of TS (Dr Mary Robertson), on the basis of meeting DSM-IV criteria for TS.

The subjects with autism or AS were selected for being of at least normal intelligence (i.e., scoring >85) on the Wechsler Adult Intelligence Test - Revised Edition [WAIS-R] (Wechsler, 1981: full scale, performance, and verbal IQ). The WAIS-R was used because of previous work showing discrepancies between performance and verbal IQ in these groups (Frith, 1989, Happé, 1994b). We therefore ensured that these subjects had an IQ >85 on both verbal and performance IQ.

The subjects with TS were also selected for being in the normal IQ range, prorated from 4 subtests of the WAIS-R (Block Design, Object Assembly, Vocabulary, and Comprehension). This short version of the WAIS-R was administered to the subjects with TS because of its brevity, since these subjects were only available for limited

testing in this study. It correlates (0.91) with full-scale WAIS-R IQ (Crawford et al, 1992).

Finally, subjects in the two clinical groups were selected for being able to pass two 1st order false belief tasks (Baron-Cohen, Leslie, & Frith, 1985; Perner, Frith, Leslie, & Leekam, 1989), and a 2nd order false belief task (Baron-Cohen, 1989b). This was so that if any deficits were found on the Eyes Task, this could be attributed to mindreading problems beyond a 6 year old level. In fact, this did not lead to any of the clinical subjects being excluded.

Table 1 gives the subject characteristics in terms of chronological age (CA), and WAIS-R IQ. ANOVAs revealed no significant differences between 3 groups on age (p>.05), or between the clinical groups in terms of WAIS-R.

insert Table 1 here

Method and Design

The Eyes Task, The Strange Stories Task, and the two Control taks were presented in random order, to all subjects. The subjects were tested individually in a quiet room either in their own home, in our clinic, or in our lab at the University.

The Eyes Task

Items from the Eyes Task were first described by Baron-Cohen (1995) as an adult test of the "language of the eyes". This extends an earlier idea of Numenmaa's (1964) that there

might be a "language of the face", for expressing mental states. The test comprises photographs of the eye region of 25 different faces (male and female). The faces were taken from magazine photos. All faces were standardized to one size (15x10 cms), all black and white, with the same region of the face selected for each photo - from midway along the nose, to just above the eye-brow. Figure 1 shows a selection of the photographs used. (Figure 1 also shows the relevant mental state terms for each photo). Each picture was shown for 3 seconds, with a forced choice between two mental state terms printed under each picture. The Experimenter says to the subject "Which word best describes what this person is feeling or thinking?" The maximum score on this test is 25.

insert Figure 1 here

Construction of the Eyes Task

The target word to describe the mental state behind each pair of eyes was generated by 4 judges (2 male and 2 female) in open discussion. A foil word was selected which was the semantic opposite of the target word, in all cases. These were then tested on a panel of 8 judges (4 male and 4 female) who were all independent raters, blind to the hypotheses of the study. On the target words there was unanimous agreement by all 8 independent raters. The full set of mental state terms (and their foils) is shown in Table 2. Notice that the mental state terms include both basic and complex mental states.

insert Table 2 here

Validity of the Eyes Task

The Eyes Task is designed to be a 'pure' theory of mind test, at an advanced level. This is because, as far as we can see, the test itself involves no executive function component (no attention switching, inhibition, planning, etc., cf Russell, Mauthner, Sharpe, & Tidswell, 1991; Ozonoff et al, 1991) and no central coherence component (since there is little contextual information available, cf Frith, 1989). This is not to say that such processes may not play a role in the development of a theory of mind - only that the task itself has no planning or context component. As mentioned earlier, in order to validate the Eyes Task as a theory of mind task, subjects in the two clinical groups were also tested on Happé's (1994a) Strange Stories. In the case of the subjects with autism and AS, this was part of a separate study (Jolliffe, 1997). If the Eyes Task was indeed tapping theory of mind, then performance on the Eyes Task should correlate with performance on Happé's strange stories.

Control Tasks

Finally, in order to check whether deficits on the Eyes Task were due to other factors, we administered two control tasks to the subjects in Group 1.

(a) *Gender Recognition Task*: this involved looking at the same sets of eyes in the experimental task, but this time identifying the <u>gender</u> of person in each photograph.

This is a social judgement without involving mindreading, and allowed us to check if any deficits on the Eyes Task could be attributed to general deficits in face perception, perceptual discrimination, or social perception. This naturally had a maximum score of 25.

(b) *Basic Emotion Recognition Task*: (Emotion Task): this involved judging photographs of whole faces displaying the basic emotions (based on the Ekman categories). This was to check whether any deficits on the Eyes Task could be attributed to a deficit in basic emotion expression recognition. 6 faces were used, testing the following basic emotions: happy, sad, angry, afraid, disgust, and surprise. Examples of the faces used are shown in Figure 2. Note that the Basic Emotion Recognition Task differes from the Eyes Task in 2 ways: (1) The Emotion Task afforded information from the whole face, whereas the Eyes Task only had information from the eyes alone. (2) The Emotion Task tested just the basic (six) emotions, whereas the Eyes Task tested the full range of mental states.

insert Figure 2 here

Results

The results on the Eyes Task from the between-groups analyses are shown in Table 3, and the results from the between-sex analysis in the normal group are shown in Table 4. To test a priori predictions on the Eyes Task, independent t tests were used, with a significance level of p < .01

set to correct for carrying out multiple comparisons. The subjects with TS did not differ from normal subjects (combined male and female) on this task (t = 0.092, 58df, p > .93), but both control groups performed significantly better than the group with autism or AS (Autism/AS x Normal, t = -5.16, 64df, p = .0001; Autism/AS x TS, t = -3.98, 24df, p = .001). In the normal group, as predicted, female subjects performed significantly better than male subjects (t = -4.8, 48df, p = .0001). The autism/AS group difference remained significant even when the group with autism/AS was compared more conservatively to the normal male group (t = 2.93, 39df, p = .006). Inspection suggested there was no difference between subjects with autism and AS, though because of sample sizes this was not tested statistically.

It is of interest to look at individual performance. If one takes a cut-off of >15/25 on the Eyes Test, as above chance (Binomial Test), then only 8/16 of the Autism/AS group were above chance, versus 10/10 of the TS group, and 50/50 of the normal group. This is a highly significant difference (Chi Square, p < .01).

On the Gender and Emotion Control Tasks, there were no differences between the groups (Gender: F (3, 69) = 0.3, p > .1; Emotion: ceiling performance by all groups). Within the Autism/AS group, there was no significant correlation between IQ and performance on the Eyes task (r = -.08). Finally, on Happé's Strange Stories, no subjects with TS made any errors, but the subjects with autism or AS made errors on this task, and were significantly impaired on this task, relative to controls. Full details of this task, including coding and results, are reported separately (Jolliffe, 1997).

insert Tables 3 and 4 here

Discussion

This study tested several predictions. First, that adults with autism or Asperger Syndrome (AS), despite being of normal or above average IQ, would nevertheless be impaired on a subtle theory of mind test. This prediction was confirmed. Secondly, that within the normal population, females would be significantly better on this test of theory of mind than males. This was also confirmed. This extends earlier work (Hall, 1977).

Regarding the autism/AS impairment, this is not easily accounted for in terms of IQ, since these subjects were in the normal or above normal range, and since performance on the task by this group shows no significant correlation with IQ. It is also not easily accounted for in terms of being the result of having any developmental neuropsychiatric disability, since the subjects with Tourette Syndrome (TS) were unimpaired on this test. The impairment in the group with autism or AS is also not easily accounted for either in terms of what Frith (1989) calls "weak central coherence" (or a difficulty in using context), or in terms of "executive function". This is because the stimuli in this experiment are relatively 'pure' theory of mind items - they are devoid of contextual information (save the person's expression in the eyes) and there is no requirement to "disengage from the salience of reality" (Russell et al, 1991).

The good performance by the group with TS is interesting in the light of other studies suggesting frontal lobe, and executive-type, dysfunction in TS (Baron-Cohen & Moriarty, 1995; Baron-Cohen, Cross, Crowson, & Robertson, 1994). To the extent that theory of mind depends on specific frontal processes (Baron-Cohen, Ring, et al, 1994; Fletcher et al, 1995; Goel, Graffman, Sadato, & Hallet, 1995), these appear to be intact in patients with TS.

We interpret the results as providing experimental evidence for subtle theory of mind deficits in individuals with autism or AS, at later points in development and at higher points on the IQ continuum than has been previously demonstrated. The justification that the Eyes Task is indeed measuring theory of mind comes from four sources. First, the target words are mental state terms. Secondly, these are not just emotion terms, but include terms describing cognitive mental states. This is therefore more than just an emotion perception test. Thirdly, the pattern of results from the Eyes Task mirrored the pattern of performance on the Happé Strange Stories - an existing advanced theory of mind task. Finally, the deficit on the Eyes Task was not mirrored on the two control tasks, suggesting that the poor performance by subjects with autism or AS was not due to the stimuli being eyes, or to a deficit in extracting social information from minimal cues, or to a subtle perceptual deficit, or to basic emotion recognition. It should also be noted that some of the subjects with autism or AS in our sample have university degrees, yet scored poorly on the Eyes Task. This strongly suggests this aspect of social cognition is independent of general intelligence.

We wish to point out, however, that even this "very advanced test" of theory of mind is still simpler than the real demands of live social situations. For one thing, our stimuli are static, in a way that the real social world never is. A closer approximation to the real social world might be based around a task assessing comprehension of movies. Anecdotally, many of our subjects with autism or AS have told us that going to the movies is, for them, often a frustrating experience, a waste of their time. This is because the social action proceeds rapidly, and they find it hard to work out why a character did or said something (their intentions or motives), who knows what and who doesn't, and why the audience laughs at particular points in the film. It just all happens too fast. Movies, however, are not pure tests of theory of mind because they do involve central coherence and executive function. Hence our decision to use the Eyes Task here.

The finding of an impairment on the Eyes Task mirrors other difficulties that have been found in autism in relation to understanding the mentalistic significance of the eyes. For example, toddlers with autism have joint attention deficits, thought to reflect a failure to interpret gaze direction as a cue to the mental state of attention (see Baron-Cohen, 1994, for a review). Young children with autism have also been found to have difficulty in interpreting direction of gaze in terms of a person's goals or desires (Baron-Cohen, Campbell, Karmiloff-Smith, Grant, & Walker, 1995; Phillips, Baron-Cohen, & Rutter, 1992) and in terms of their intention to refer (Baron-Cohen, Baldwin, & Crowson, in press). In addition, and in contrast to normal preschool children, they are relatively blind to the significance of gaze direction as a cue to when someone is thinking (Baron-Cohen & Cross, 1992; Baron-Cohen et al, 1995).

Regarding the sex difference found in the normal group, this is of interest for several reasons. First, previous studies of theory of mind have not allowed for a subtle test of individual differences in this ability, being mostly all-or-none, pass-fail type tests. Sex differences might therefore simply have been missed. It may be that there are sex differences in the rate of development of theory of mind (and its developmental precursor, joint attention) in early childhood. This remains to be tested. Certainly, there are clear precedents for sex differences in cognitive development, the paradigm cases being female superiority in language development and male superiority in spatial skills (Halpern, 1992; Witelson, 1976; Kimura, 1992). Note that the female advantage on the mindreading task could be taken to reflect genetic or socialization factors (Baron-Cohen & Hammer, in press b). This remains to be investigated further.

References

Baron-Cohen, S, (1989a) Perceptual role-taking and protodeclarative pointing in autism. British Journal of Developmental Psychology, 7, 113-127.

Baron-Cohen, S, (1989b) The autistic child's theory of mind: a case of specific developmental delay. Journal of Child Psychology and Psychiatry, **30**, 285-298.

Baron-Cohen, S, (1991) The development of a theory of mind in autism: deviance and delay? <u>Psychiatric Clinics of North America</u>, **14**, 33-51.

Baron-Cohen, S, (1993) From attention-goal psychology to belief-desire psychology: the development of a theory of mind and its dysfunction. In Baron-Cohen, S, Tager-Flusberg, H, & Cohen, D.J. (eds) <u>Understanding other minds: perspectives from autism</u>. Oxford University Press.

Baron-Cohen, S, (1994) How to build a baby that can read minds: Cognitive mechanisms in mindreading. <u>Cahiers de Psychologie Cognitive/ Current Psychology of Cognition</u>, **13(5)**, 513-552.

Baron-Cohen, S, (1995) <u>Mindblindness: an essay on autism and theory of mind</u>. MIT Press/Bradford Books.

Baron-Cohen, S, Allen, J, & Gillberg, C, (1992) Can autism be detected at 18 months? The needle, the haystack, and the CHAT. <u>British Journal of Psychiatry</u>, **161**, 839-843.

Baron-Cohen, S, Baldwin, D, & Crowson, M, (in press) Do children with autism use the Speaker's Direction of Gaze (SDG) strategy to crack the code of language? <u>Child</u> <u>Development</u>

Baron-Cohen, S, Campbell, R, Karmiloff-Smith, A, Grant, J, & Walker, J, (1995) Are children with autism blind to the mentalistic significance of the eyes? <u>British</u> Journal of Developmental Psychology. **13**, 379-398.

Baron-Cohen, S, Cox, A, Baird, G, Swettenham, J, Drew, A, Nightingale, N, Morgan, K, & Charman, T, (1996) Psychological markers of autism at 18 months of age in a large population. <u>British Journal of Psychiatry</u>, **168**, 158-163.

Baron-Cohen, S, & Cross, P, (1992) Reading the eyes: Evidence for the role of perception in the development of a theory of mind. <u>Mind and Language</u>, **6**, 173-186.

Baron-Cohen, S, Cross, P, Crowson, M, & Robertson, M, (1994) Can children with Tourette's Syndrome edit their intentions? Psychological Medicine, **24**, 29-40.

Baron-Cohen, S, & Howlin, P, (1993) The theory of mind deficit in autism: some questions for teaching and diagnosis. In Baron-Cohen, S, Tager-Flusberg, H,

& Cohen, D.J. (eds) <u>Understanding other minds: perspectives from autism</u>. Oxford University Press.

Baron-Cohen, S, & Hammer, J, (in press a) Parents of children with Asperger Syndrome: What is the cognitive phenotype? <u>Journal of Cognitive Neuroscience</u>.

Baron-Cohen, S, & Hammer, (in press b) Is autism an extreme form of the male brain? In Infant Behaviour and Development, **11**.

Baron-Cohen, S, Leslie, A.M., & Frith, U, (1985) Does the autistic child have a `theory of mind'? <u>Cognition</u>, **21**, 37-46.

Baron-Cohen, S, Ring, H, Moriarty, J, Shmitz, P, Costa, D, & Ell, P, (1994) Recognition of mental state terms: a clinical study of autism, and a functional neuroimaging study of normal adults. <u>British Journal of Psychiatry</u>, **165**, 640-649.

Baron-Cohen, S, Robertson, M, & Moriarty, J, (1994) The development of the will: a neuropsychological analysis of Gilles de la Tourette's Syndrome. In Cicchetti, D, & Toth, S, (ed) <u>The Self and its Dysfunction: Proceedings of the 4th Rochester</u> <u>Symposium</u>. University of Rochester Press.

Baron-Cohen, S, Riviere, A, Cross, P, Fukushima, M, Bryant, C, Sotillo, M, Hadwin, J, & French, D, (1996b) Reading the mind in the face: a cross-cultural and developmental study. <u>Visual Cognition</u>, **3**, 39-59.

Bishop, D, (1983) Test of Perception of Grammer. University of Manchester.

Bolton, P, & Rutter, M, (1990) Genetic influences in autism. <u>International Review of</u> <u>Psychiatry</u>, **2**, 67-80.

Bowler, D.M. (1992) Theory of Mind in Asperger Syndrome. <u>Journal of Child</u> <u>Psychology and Psychiatry</u>, **33**, 877-893.

Crawford, J.R, Allan, K.M., & Jack, A.M. (1992) Short forms of the UK WAIS-R: Regression equations and their predictive validity in a general population sample. <u>British</u> <u>Journal of Clinical Psychology</u>, **31**, 191-202.

DSM-IV (1994) Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Washington DC: American Psychiatric Association.

Ekman, P, (1992) An argument for basic emotions. Cognition and Emotion, 6, 169-200.

Fletcher, P, Happé, F, Frith, U, Baker, S, Dolan, R, Frackowiak, R, & Frith, C,

(1995) Other minds in the brain: a functional imaging study of "theory of mind" in story comprehension. <u>Cognition</u>, **57**, 109-128.

Frith, U, (1989) Autism: explaining the enigma. Oxford: Basil Blackwell.

Goel, V, Grafman, J, Sadato, N, & Hallett, M, (1995) Modeling other minds. <u>Neuroreport</u>, **6**, 1741-1746.

Hall, J, (1977) Gender effects in decoding non-verbal cues. <u>Psychological Bulletin</u>, **85**, 845-857.

Halpern, D, (1992) Sex differences in cognitive ability. Second edition.

Happé, F, (1994a) An advanced test of theory of mind: Understanding of story characters' thoughts and feelings by able autistic, mentally handicapped, and normal children and adults. Journal of Autism and Developmental Disorders, **24**, 129-154.

Happé, F, (1994b) Autism. London: UCL Press.

Jolliffe, T, (1997) Central coherence in adults with high-functioning autism or Asperger Syndrome. Unpublished PhD Thesis, University of Cambridge.

Kimura, D, (1992) Sex differences in the brain. <u>Scientific American</u>, **September**, 119-125.

Nummenmaa, T, (1964) The language of the face. <u>Jyvaskyla studies in education</u>, <u>psychology</u>, and <u>social research</u>. Jyvaskyla, Finland.

Ozonoff, S, Pennington, B, & Rogers, S, (1991) Executive function deficits in highfunctioning autistic children: relationship to theory of mind. Journal of Child Psychology and Psychiatry, **32**, 1081-1106.

Ozonoff, S, Rogers, S, & Pennington, B, (1991) Asperger's Syndrome: evidence of an empirical distinction from high-functioning autism. Journal of Child Psychiatry and Psychology, **32**, 1107-1122.

Perner, J, Frith, U, Leslie, A.M, & Leekam, S, (1989) Exploration of the autistic child's theory of mind: knowledge, belief, and communication. <u>Child Development</u>, 60, 689-700.

Perner, J. & Wimmer, H. (1985) "John thinks that Mary thinks that..." Attribution of second-order beliefs by 5-10 year old children. Journal of Experimental Child Psychology, 39, 437-471.

Phillips, W, Baron-Cohen, S, & Rutter, M, (1992) The role of eye-contact in the detection of goals: evidence from normal toddlers, and children with autism or mental handicap. <u>Development and Psychopathology</u>, **4**, 375-383.

Robertson, M, (1994) Gilles de la Tourette Syndrome: an update. Journal of Child Psychology and Psychiatry, **35**,597-612. Russell, J, Mauthner, N, Sharpe, S, & Tidswell, T, (1991) The Windows Task as a measure of strategic deception in preschoolers and autistic children. <u>British Journal of</u> <u>Developmental Psychology</u>, **9**, 331-349.

Sigman, M, Mundy, P, Ungerer, J, & Sherman, T, (1986) Social interactions of autistic, mentally retarded, and normal children and their caregivers. Journal of Child Psychology and Psychiatry, **27**, 647-656.

Wechsler, D, (1981) <u>Wechsler Adult Intelligence Scale - Revised</u>. New York: The Physiological Corporation.

Wimmer, H. & Perner, J. (1983) Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's understanding of deception. <u>Cognition</u>, 13, 103-128.

Witelson, S, (1976) Sex and the single hemisphere: specialization of the right hemisphere for spatial processing. <u>Science</u>, **193**, 425-427.

Figure 1: Examples of stimuli from the Eyes Test.

Fig 1(a) CONCERNED vs Unconcerned

Fig 1(b) SERIOUS MESSAGE vs Playful Message

Fig 1(c) REFLECTIVE vs Unreflective

Fig 1(d) SYMPATHETIC vs Unsympathetic

Figure 2: Examples of the Basic Emotion Recognition Control Task

Fig 1(a) Happy vs Sad

Fig 1(b) Angry vs Afraid

<u>Acknowledgements</u>: The first author was supported by grants from the Medical Research Council (UK), the Wellcome Trust, and the Gatsby Foundation during the period of this work. We thank the following students for their assistance in data collection: Alison Pickett, Avram Gilbert, Jon Rohrer, Rikin Patel, Kannan Aranasalam, Parag Sharma and Jessica Hammer. The photographs used in Figure 1 are fragments of faces taken from a variety of commercial magazines, and unfortunately their sources can no longer be traced. Liz Tennent kindly agreed to be photographed for stimuli shown in Figure 2. +To whom correspondence should be addressed.

Table 1: Chronological age (CA) and IQ of the subjects in each of the 3 groups.

		СА	WAIS-R IQ
AUTISM/	mean	28.6	105.31
AS	sd	9.7	13.0
(n = 16)	range	18-49	86-133
		СА	-
	mean	30.0	-
NORMAL	sd	9.12	-
(n = 50)	range	18-48	-
	mean	27.77	103.5
TOURETTE	sd	7.81	10.0
(n = 10)	range	18-47	93-115

Stimulus No	Target term	Foil
1	Concerned	Unconcerned
2	Noticing you	Ignoring you
3	Attraction	Repulsion
4	Relaxed	Worried
5	Serious message	Playful message
6	Interested	Disinterested
7	Friendly	Hostile
8	Sad reflection	Happy reflection
9	Sad thought	Happy thought
10	Certain	Uncertain
11	Far away focus	Near focus
12	Reflective	Unreflective
13	Reflective	Unreflective
14	Cautious about something over there	Relaxed about something over there
15	Noticing someone else	Noticing you
16	Calm	Anxious
17	Dominant	Submissive
18	Fantasizing	Noticing
19	Observing	Daydreaming
20	Desire for you	Desire for someone else

Table 2: Target mental state terms, and their foil terms.

25	Decisive	Indecisive
24	Sympathetic	Unsympathetic
23	Flirtatious	Disinterested
22	Nervous about you	Interested in you
21	Ignoring you	Noticing you

Table 3 Performance on the Eyes Test

		Eyes Task	Gender
			Control
AUTISM/	mean	16.3*	24.1
AS	sd	2.9	0.7
(n = 16)	range	13-23	23-25
	mean	20.3	23.3
NORMAL	sd	2.63	1.1
(n = 50)	range	16-25	22-25
	mean	20.4	23.7
TOURETTE	sd	2.63	0.9
(n = 10)	range	16-25	23-25

* Autism/AS x Normal, t = -5.16, 64df, p = .0001; Autism/AS x TS, t = -3.98, 24df, p = .001.

Table 4: Performance by males and females in the normal group

		Eyes Task	Gender
			Control
	mean	18.8*	24.0
Males	sd	2.53	0.6
(n = 25)	range	16-22	23-25
	mean	21.8	23.8
Females	sd	1.78	0.6
(n = 25)	range	20-25	23-25

* t = -4.8, 48df, p = .0001