

[Orthopaedic Surgery]



Anterior Cruciate Ligament Reconstruction Rehabilitation: MOON Guidelines

Rick W. Wright, MD,^{*†} Amanda K. Haas, MA,[†] Joy Anderson, PT, ATC, CSCS,[‡] Gary Calabrese, PT,[§] John Cavanaugh, PT, ATC,^{||} Timothy E. Hewett, PhD,[¶] Dawn Loring, PT, SCS,[§] Christopher McKenzie, PT,[¶] Emily Preston, PT,[#] Glenn Williams, PhD, PT, ATC,^{**} and the MOON Group^{††}

Context: Anterior cruciate ligament (ACL) reconstruction rehabilitation has evolved over the past 20 years. This evolution has been driven by a variety of level 1 and level 2 studies.

Evidence Acquisition: The MOON Group is a collection of orthopaedic surgeons who have developed a prospective longitudinal cohort of the ACL reconstruction patients. To standardize the management of these patients, we developed, in conjunction with our physical therapy committee, an evidence-based rehabilitation guideline.

Study Design: Clinical review.

Level of Evidence: Level 2.

Results: This review was based on 2 systematic reviews of level 1 and level 2 studies. Recently, the guideline was updated by a new review. Continuous passive motion did not improve ultimate motion. Early weightbearing decreases patellofemoral pain. Postoperative rehabilitative bracing did not improve swelling, pain range of motion, or safety. Open chain quadriceps activity can begin at 6 weeks.

Conclusion: High-level evidence exists to determine appropriate ACL rehabilitation guidelines. Utilizing this protocol follows the best available evidence.

Keywords: anterior cruciate ligament; ACL reconstruction; rehabilitation

Anterior cruciate ligament (ACL) reconstruction is a frequently performed procedure in this country. While surgical aspects are critical to the success of the patient, rehabilitation following surgery is just as crucial. The best surgically reconstructed ACL can be ruined by inappropriate or ineffective rehabilitation. Our goal with this publication is to present the Multicenter Orthopaedic Outcomes Network (MOON) ACL rehabilitation guidelines (see Appendix, available at <http://sph.sagepub.com/content/suppl>) and the clinical evidence used to formulate the guidelines.

The MOON group began in 2001 as a 10-member group of orthopaedic sports medicine specialists at 7 sites that

assembled to study outcomes of lower extremity injuries and surgeries.^{14,15,39,66,72} It is now a National Institutes of Health-funded group of greater than 20 clinicians who have developed a longitudinal prospective cohort of more than 3000 ACL reconstructions. Shortly following formation of the group, it was acknowledged that standardized rehabilitation guidelines were important for analyzing and reporting our reconstruction outcomes. We undertook a groupwide systematic review to determine the best evidence to guide the formation of an ACL reconstruction rehabilitation protocol.^{74,75} Following a review of the evidence and an agreement on important milestones and approaches to rehabilitation, we engaged the physical

From [†]Washington University in Saint Louis, St Louis, Missouri, [‡]University of Colorado Denver, Denver, Colorado, [§]Cleveland Clinic, Cleveland, Ohio, ^{||}Hospital for Special Surgery, New York, New York, [¶]The Ohio State University, Columbus, Ohio, [#]Vanderbilt University, Nashville, Tennessee, and ^{**}University of Iowa, Iowa City, Iowa. ^{††}All members are listed in the Contributing Authors section at the end of this article.

*Address correspondence to Rick W. Wright, MD, Department of Orthopaedic Surgery, Washington University, 660 South Euclid Avenue, Campus Box 8233, St Louis, MO 63110 (e-mail: Rwwright1@aol.com).

The following authors declared potential conflicts of interest: Rick W. Wright, MD, received research grants from Smith & Nephew and the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health (NIH) and receives royalties from Wolters Kluwer and Lippincott Williams & Wilkins; and Timothy Hewett, PhD, is employed by The Ohio State University and received a grant from the NIH.

DOI: 10.1177/1941738113517855

© 2014 The Author(s)

therapists at our sites to develop practical ACL reconstruction rehabilitation guidelines. The guidelines have been developed to service the spectrum of ACL-injured people, from the nonathlete to the elite athlete. For this reason, sample exercises are provided instead of a highly structured rehabilitation guideline. Therefore, we recommend that attending rehabilitation specialists tailor the guidelines to each patient's specific needs.

The multicenter nature of the MOON group necessitates that the MOON ACL rehabilitation guidelines include only treatment methods that can be employed at all sites without purchasing expensive equipment. Consequently, some treatment methods with supporting evidence (eg, high-intensity electric stimulation, aquatic therapy) are not included in the guidelines because the expert panel believes that it is unreasonable to expect all sites to carry out such treatments.

Progression from one phase to the next is based on readiness by achieving functional criteria rather than the time elapsed since surgery. The time frames identified in parentheses after each phase are approximate times for the average patient, not guidelines for progression. Some patients will be ready to progress sooner than the time frame identified, whereas others will take longer.

This has now been in use for 10 years with high success. More than 3000 patients enrolled in MOON studies have utilized it, and most MOON surgeons use it for all their ACL reconstructions.

Recently, an update of this systematic review of level 1 and level 2 studies regarding ACL reconstruction rehabilitation was performed.³⁷ On the basis of this update, the physical therapist committee reconvened to address potential updates of the protocol. Minor changes were made to the protocol to reflect incremental progress in our understanding of ACL reconstruction rehabilitation that would influence the protocol for our patients. The goal of this report is to briefly review the available scientific evidence and to provide the guidelines in their current form.

REHABILITATION EVIDENCE

The initial systematic reviews covered level 1 and level 2 evidence, predominantly randomized controlled trials found through 2005.^{74,75} The updated systematic review covers the period from 2005 to 2011.³⁷ For many aspects of ACL rehabilitation, either there are no studies that qualify as "best evidence," or the number of studies is too few for conclusions to be drawn with confidence. These circumstances are identified.

CONTINUOUS PASSIVE MOTION

Six randomized trials evaluated continuous passive motion for the rehabilitation of the ACL-reconstructed knee.^{17,41,42,54,58,76} No long-term benefits were determined for continuous passive motion. There was no long-term improved range of

motion. These machines are frequently difficult to approve for insurance reasons, and thus, cost becomes an issue.

EARLY WEIGHTBEARING AND MOTION

Immediate weightbearing following ACL reconstruction was investigated by 1 study.⁶⁹ A significant decrease in patellofemoral pain from 35% to 8% was demonstrated by this study. Thus, immediate full weightbearing is initiated following ACL reconstruction in the MOON protocol. Immediate range of motion has not been studied by randomized trial but reflects a fundamental principle that most therapists ascribe to as important in modern ACL reconstruction rehabilitation and is also immediately begun during rehabilitation.

POSTOPERATIVE BRACING

The use of postoperative bracing utilizing rehabilitative braces has been evaluated by a variety of studies.⁷³ This is exclusive of the question of functional bracing for return to sport but involves the use of knee immobilizers or hinged knee braces in the immediate postoperative time frame. Eleven studies in the initial review evaluated this question, and no study demonstrated a clinically significant or relevant improvement in safety, range of motion including extension, or other outcome measures.^{7,15,19,25,27,31,36,43,47,50,57,67} Given these studies and the expense of postoperative bracing, we do not include bracing following ACL reconstruction as part of our protocol. This was reinforced by an additional 6 studies published since 2005.^{5,26,32,34,35,40} None of these demonstrated an advantage from bracing.

HOME-BASED REHABILITATION

Four studies evaluated home-based ACL rehabilitation.^{2,20,24,60} Each featured minimal official visits with a physical therapist. Two additional studies have been performed since 2005.^{23,53} Although many of these studies have biases and limitations, none have indicated that home-based ACL rehabilitation was deleterious when prescribed with motivated patients.

OPEN VERSUS CLOSED CHAIN EXERCISES

Few studies have adequately evaluated this question.^{9,33,45,48,49} Based on the available evidence, it appears that open chain activities after 6 weeks may improve strength without adversely affecting the graft and/or increasing graft laxity.^{22,30} There is currently insufficient evidence on the safety of open chain knee exercises before 6 weeks postsurgery. With this lack of evidence and a concern raised by strain values that have been demonstrated in the ACL graft with open chain activities, we have limited the open chain activities in the first 6 weeks to light-load, short-arc quadriceps exercises.^{22,30} Additional studies need to be performed in this area to provide more appropriate evidence regarding the safety of initiating open chain exercise sooner.

NEUROMUSCULAR ELECTRICAL STIMULATION

Many studies have evaluated neuromuscular electrical stimulation in the ACL-reconstructed population.^{1,11,13,18,21,38,51,52,59,62-65,71} The evidence suffers from a lack of standardization and homogeneity among the studies to allow comparison. Some handheld devices are too weak to provide stimulation that will result in improved outcomes. Rehabilitation specialists should ensure that portable stimulators are capable of adequately recruiting the target muscle, or they should use a clinical stimulator in an outpatient setting. Some patients who are lagging in muscle recruitment and redevelopment following surgery will benefit from this adjunctive treatment.^{11,12} The electrical parameters that best stimulate the muscle for improved outcome are uncertain and require further analysis. Given this, we do not make neuromuscular electrical stimulation a requirement for the MOON ACL reconstruction rehabilitation protocol, and we leave it to the discretion of the individual physical therapist.

ACCELERATED REHABILITATION

Two randomized trials have evaluated accelerated rehabilitation.^{4,16} Neither study had a group that dramatically lessened the rehabilitation below 6 months. The question remains whether patients can rehabilitate from this surgery and be ready to return to sports at less than 4 months. There is minimal or no scientific evidence to support this earlier return. The guidelines are based on a functional criterion that typically has patients ready at 5 to 6 months. Additional research will be necessary to prove that shorter time frames are safe for the graft, menisci, articular cartilage, and the patient in general.

NEUROMUSCULAR TRAINING

Since 2005, 9 randomized trials evaluated neuromuscular training—including “proprioceptive” and balance training, perturbation training, and vibratory stimulation as part of ACL reconstruction rehabilitation—and demonstrated safety with some efficacy in their use.^{3,8,10,28,29,46,55,56,70} Neuromuscular training has been suggested in most phases of the MOON ACL rehabilitation guidelines since inception and continues to be included.

MISCELLANEOUS

A variety of randomized trials evaluated several miscellaneous topics. These included the safety of aquatic training,⁶⁸ slide board safety,⁶ and stair-climber versus cycle exercise efficacy.⁴⁴ Aquatic training was deemed to be safe and may decrease effusions. Slide board work was safe and may improve quadriceps strength. The stair-climber was as efficacious and safe as stationary cycling. Early quadriceps strengthening with straight-leg raises has been evaluated and improved strength has been noted with no untoward effects.⁶¹

CONCLUSION

The MOON ACL reconstruction rehabilitation guidelines have been in use for 10 years. Rehabilitation effectiveness is largely based on exercise selection and dosing. Accordingly, it is important that exercise dosing is scientifically based as well.

CONTRIBUTING AUTHORS

Annunziato Amendola, MD (University of Iowa); Jack T. Andrich, MD (University of Iowa); Robert H. Brophy, MD (Washington University in St Louis); Charles L. Cox, MD, MPH (Vanderbilt University); Warren R. Dunn, MD, MPH (University of Wisconsin); David C. Flanigan, MD (The Ohio State University); Carolyn M. Hettrich, MD, MPH (University of Iowa); Laura J. Huston, MS (Vanderbilt University); Morgan H. Jones, MD (Cleveland Clinic); Christopher C. Kaeding, MD (The Ohio State University); Christian Lattermann, MD (University of Kentucky); Robert A. Magnussen, MD (The Ohio State University); Robert G. Marx, MD (Hospital for Special Surgery); Matthew J. Matava, MD (Washington University in St Louis); Eric C. McCarty, MD (University of Colorado); Richard D. Parker, MD (Cleveland Clinic); Emily K. Reinke, PhD (Vanderbilt University); Matthew V. Smith, MD (Washington University in St Louis); Kurt P. Spindler, MD (Vanderbilt University); Armando F. Vidal, MD (University of Colorado); Michelle L. Wolcott, MD (University of Colorado); Brian R. Wolf, MD (University of Iowa).

ACKNOWLEDGMENT

This project was partially funded by grant 5R01 AR053684 from the National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases.

REFERENCES

1. Arvidsson I, Arvidsson H, Eriksson E, Jansson E. Prevention of quadriceps wasting after immobilization: an evaluation of the effect of electrical stimulation. *Orthopedics*. 1986;9:1519-1528.
2. Beard DJ, Dodd CA. Home or supervised rehabilitation following anterior cruciate ligament reconstruction: a randomized controlled trial. *J Orthop Sports Phys Ther*. 1998;27:134-143.
3. Benazzo F, Zanon G, Pederzini L, et al. Effects of biophysical stimulation in patients undergoing arthroscopic reconstruction of anterior cruciate ligament: prospective, randomized and double blind study. *Knee Surg Sports Traumatol Arthrosc*. 2008;16:595-601.
4. Beynon BD, Uh BS, Johnson RJ, et al. Rehabilitation after anterior cruciate ligament reconstruction: a prospective, randomized, double-blind comparison of programs administered over 2 different time intervals. *Am J Sports Med*. 2005;33:347-359.
5. Birmingham TB, Bryant DM, Giffin JR, et al. A randomized controlled trial comparing the effectiveness of functional knee brace and neoprene sleeve use after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2008;36:648-655.
6. Blanpied P, Carroll R, Douglas T, Lyons M, Macalisang R, Pires L. Effectiveness of lateral slide exercise in an anterior cruciate ligament reconstruction rehabilitation home exercise program. *J Orthop Sports Phys Ther*. 2000;30:602-608.
7. Brandsson S, Faxen E, Kartus J, Eriksson BI, Karlsson J. Is a knee brace advantageous after anterior cruciate ligament surgery? A prospective, randomised study with a two-year follow-up. *Scand J Med Sci Sports*. 2001;11:110-114.

8. Brunetti O, Filippi GM, Lorenzini M, et al. Improvement of posture stability by vibratory stimulation following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2006;14:1180-1187.
9. Bynum EB, Barrack RL, Alexander AH. Open versus closed chain kinetic exercises after anterior cruciate ligament reconstruction: a prospective randomized study. *Am J Sports Med.* 1995;23:401-406.
10. Cooper RL, Taylor NF, Feller JA. A randomised controlled trial of proprioceptive and balance training after surgical reconstruction of the anterior cruciate ligament. *Res Sports Med.* 2005;13:217-230.
11. Delitto A, Rose SJ, McKowen JM, Lehman RC, Thomas JA, Shively RA. Electrical stimulation versus voluntary exercise in strengthening thigh musculature after anterior cruciate ligament surgery. *Phys Ther.* 1988; 68:660-663 [erratum 1988;68:1145].
12. Dolan MG, Mendel FC. Clinical application of electrotherapy. *Atbl Ther Today.* 2004;9:11-16.
13. Draper V, Ballard L. Electrical stimulation versus electromyographic biofeedback in the recovery of quadriceps femoris muscle function following anterior cruciate ligament surgery. *Phys Ther.* 1991;71:455-461.
14. Dunn WR, Spindler KP, MOON Consortium. Predictors of activity level 2 years after anterior cruciate ligament reconstruction (ACLR): a Multicenter Orthopaedic Outcomes Network (MOON) ACLR cohort study. *Am J Sports Med.* 2010;38:2040-2050.
15. Dunn WR, Spindler KP, Amendola A, et al. Which preoperative factors, including bone bruise, are associated with knee pain/symptoms at index anterior cruciate ligament reconstruction (ACLR)? A Multicenter Orthopaedic Outcomes Network (MOON) ACLR Cohort Study. *Am J Sports Med.* 2010;38:1778-1787.
16. Ekstrand J. Six versus eight months of rehabilitation after reconstruction of the anterior cruciate ligament: a prospective randomized study on soccer players. *Science Football.* 1990;3:31-36.
17. Engstrom B, Sperber A, Wredmark T. Continuous passive motion in rehabilitation after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 1995;3:18-20.
18. Eriksson E, Haggmark T. Comparison of isometric muscle training and electrical stimulation supplementing isometric muscle training in the recovery after major knee ligament surgery: a preliminary report. *Am J Sports Med.* 1979;7:169-171.
19. Feller J, Bartlett J, Chapman S, Delahunt M. Use of an extension-assisting brace following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 1997;5:6-9.
20. Fischer DA, Tewes DP, Boyd JL, Smith JP, Quick DC. Home based rehabilitation for anterior cruciate ligament reconstruction. *Clin Orthop Relat Res.* 1998;347:194-199.
21. Fitzgerald GK, Piva SR, Irrgang JJ. A modified neuromuscular electrical stimulation protocol for quadriceps strength training following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2003;33:492-501.
22. Fleming BC, Oksendahl H, Beynon BD. Open- or closed-kinetic chain exercises after anterior cruciate ligament reconstruction? *Exerc Sport Sci Rev.* 2005;33:134-140.
23. Grant JA, Mohtadi NG. Two- to 4-year follow-up to a comparison of home versus physical therapy-supervised rehabilitation programs after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2010;38:1389-1394.
24. Grant JA, Mohtadi NG, Maitland ME, Zernicke RF. Comparison of home versus physical therapy-supervised rehabilitation programs after anterior cruciate ligament reconstruction: a randomized clinical trial. *Am J Sports Med.* 2005;33:1288-1297.
25. Haggmark T, Eriksson E. Cylinder or mobile cast brace after knee ligament surgery: a clinical analysis and morphologic and enzymatic studies of changes in the quadriceps muscle. *Am J Sports Med.* 1979;7:48-56.
26. Harilainen A, Sandelin J. Post-operative use of knee brace in bone-tendon-bone patellar tendon anterior cruciate ligament reconstruction: 5-year follow-up results of a randomized prospective study. *Scand J Med Sci Sports.* 2006;16:14-18.
27. Harilainen A, Sandelin J, Vanhanen I, Kivinen A. Knee brace after bone-tendon-bone anterior cruciate ligament reconstruction: randomized, prospective study with 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc.* 1997;5:10-13.
28. Hartigan E, Axe MJ, Snyder-Mackler L. Perturbation training prior to ACL reconstruction improves gait asymmetries in non-copers. *J Orthop Res.* 2009;27:724-729.
29. Hartigan EH, Axe MJ, Snyder-Mackler L. Time line for noncopers to pass return-to-sports criteria after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2010;40:141-154.
30. Hejine A, Fleming BC, Renstrom PA, Peura GD, Beynon BD, Werner S. Strain on the anterior cruciate ligament during closed kinetic chain exercises. *Med Sci Sports Exerc.* 2004;36:935-941.
31. Henriksson M, Rockborn P, Good L. Range of motion training in brace vs. plaster immobilization after anterior cruciate ligament reconstruction: a prospective randomized comparison with a 2-year follow-up. *Scand J Med Sci Sports.* 2002;12(2):73-80.
32. Hiemstra LA, Heard SM, Sasyniuk TM, Buchko GL, Reed JG, Monteone BJ. Knee immobilization for pain control after a hamstring tendon anterior cruciate ligament reconstruction: a randomized clinical trial. *Am J Sports Med.* 2009;37:56-64.
33. Hooper DM, Morrissey MC, Drechsler W, Morrissey D, King J. Open and closed kinetic chain exercises in the early period after anterior cruciate ligament reconstruction. Improvements in level walking, stair ascent, and stair descent. *Am J Sports Med.* 2001;29:167-174.
34. Isberg J, Faxen E, Brandsson S, Eriksson BI, Karrholm J, Karlsson J. Early active extension after anterior cruciate ligament reconstruction does not result in increased laxity of the knee. *Knee Surg Sports Traumatol Arthrosc.* 2006;14:1108-1115.
35. Ito Y, Deie M, Adachi N, et al. A prospective study of 3-day versus 2-week immobilization period after anterior cruciate ligament reconstruction. *Knee.* 2007;14:34-38.
36. Kartus J, Stener S, Kohler K, Sernert N, Eriksson BI, Karlsson J. Is bracing after anterior cruciate ligament reconstruction necessary? A 2-year follow-up of 78 consecutive patients rehabilitated with or without a brace. *Knee Surg Sports Traumatol Arthrosc.* 1997;5:157-161.
37. Kruse LM, Gray B, Wright RW. Rehabilitation after anterior cruciate ligament reconstruction: a systematic review. *J Bone Joint Surg Am.* 2012;94:1737-1748.
38. Lieber RL, Silva PD, Daniel DM. Equal effectiveness of electrical and volitional strength training for quadriceps femoris muscles after anterior cruciate ligament surgery. *J Orthop Res.* 1996;14:131-138.
39. Magnusson RA, Granan LP, Dunn WR, et al. Cross-cultural comparison of patients undergoing ACL reconstruction in the United States and Norway. *Knee Surg Sports Traumatol Arthrosc.* 2010;18:98-105.
40. Mayr HO, Hochrein A, Hein W, Hube R, Bernstein A. Rehabilitation results following anterior cruciate ligament reconstruction using a hard brace compared to a fluid-filled soft brace. *Knee.* 2010;17:119-126.
41. McCarthy MR, Buxton BP, Yates CK. Effects of continuous passive motion on anterior laxity following ACL reconstruction with autogenous patellar tendon grafts. *J Sport Rehabil.* 1993;2:171-178.
42. McCarthy MR, Yates CK, Anderson MA, Yates-McCarthy JL. The effects of immediate continuous passive motion on pain during the inflammatory phase of soft tissue healing following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 1993;17:96-101.
43. Melegati G, Tornese D, Bandi M, Volpi P, Schonhuber H, Denti M. The role of the rehabilitation brace in restoring knee extension after anterior cruciate ligament reconstruction: a prospective controlled study. *Knee Surg Sports Traumatol Arthrosc.* 2003;11:322-326.
44. Meyers MC, Sterling JC, Marley RR. Efficacy of stairclimber versus cycle ergometry in postoperative anterior cruciate ligament rehabilitation. *Clin J Sport Med.* 2002;12:85-94.
45. Mikkelsen C, Werner S, Eriksson E. Closed kinetic chain alone compared to combined open and closed kinetic chain exercises for quadriceps strengthening after anterior cruciate ligament reconstruction with respect to return to sports: a prospective matched follow-up study. *Knee Surg Sports Traumatol Arthrosc.* 2000;8:337-342.
46. Moezy A, Olyaei G, Hadian M, Razi M, Faghihzadeh S. A comparative study of whole body vibration training and conventional training on knee proprioception and postural stability after anterior cruciate ligament reconstruction. *Br J Sports Med.* 2008;42:373-378.
47. Moller E, Forssblad M, Hansson L, Wange P, Weidenhielm L. Bracing versus nonbracing in rehabilitation after anterior cruciate ligament reconstruction: a randomized prospective study with 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2001;9:102-108.
48. Morrissey MC, Drechsler WI, Morrissey D, Knight PR, Armstrong PW, McAuliffe TB. Effects of distally fixated versus nondistally fixated leg extensor resistance training on knee pain in the early period after anterior cruciate ligament reconstruction. *Phys Ther.* 2002;82:35-43.
49. Morrissey MC, Hudson ZL, Drechsler WI, Coutts FJ, Knight PR, King JB. Effects of open versus closed kinetic chain training on knee laxity in the early period after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2000;8:343-348.

50. Muellner T, Alacamlıoglu Y, Nikolic A, Schabus R. No benefit of bracing on the early outcome after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 1998;6:88-92.
51. Paternostro-Sluga T, Fialka C, Alacamlıoglu Y, Saradeth T, Fialka-Moser V. Neuromuscular electrical stimulation after anterior cruciate ligament surgery. *Clin Orthop Relat Res.* 1999;368:166-175.
52. Rebai H, Barra V, Laborde A, Bonny JM, Poumarat G, Coudert J. Effects of two electrical stimulation frequencies in thigh muscle after knee surgery. *Int J Sports Med.* 2002;23:604-609.
53. Revenas A, Johansson A, Leppert J. A randomized study of two physiotherapeutic approaches after knee ligament reconstruction. *Adv Physiother.* 2009;11:30-41.
54. Richmond JC, Gladstone J, MacGillivray J. Continuous passive motion after arthroscopically assisted anterior cruciate ligament reconstruction: comparison of short- versus long-term use. *Arthroscopy.* 1991;7:39-44.
55. Risberg MA, Holm I. The long-term effect of 2 postoperative rehabilitation programs after anterior cruciate ligament reconstruction: a randomized controlled clinical trial with 2 years of follow-up. *Am J Sports Med.* 2009;37:1958-1966.
56. Risberg MA, Holm I, Myklebust G, Engebretsen L. Neuromuscular training versus strength training during first 6 months after anterior cruciate ligament reconstruction: a randomized clinical trial. *Phys Ther.* 2007;87:737-750.
57. Risberg MA, Holm I, Steen H, Eriksson J, Ekland A. The effect of knee bracing after anterior cruciate ligament reconstruction: a prospective, randomized study with two years' follow-up. *Am J Sports Med.* 1999;27:76-83.
58. Rosen MA, Jackson DW, Atwell EA. The efficacy of continuous passive motion in the rehabilitation of anterior cruciate ligament reconstructions. *Am J Sports Med.* 1992;20:122-127.
59. Ross M. The effect of neuromuscular electrical stimulation during closed kinetic chain exercise on lower extremity performance following anterior cruciate ligament reconstruction. *Sports Med Train Rehab.* 2000;9:239-251.
60. Schenck RC Jr, Blaschak MJ, Lance ED, Turturro TC, Holmes CF. A prospective outcome study of rehabilitation programs and anterior cruciate ligament reconstruction. *Arthroscopy.* 1997;13:285-290.
61. Shaw T, Williams MT, Chipchase LS. Do early quadriceps exercises affect the outcome of ACL reconstruction? A randomised controlled trial. *Aust J Physiother.* 2005;51:9-17.
62. Sisk TD, Stralka SW, Deering MB, Griffin JW. Effect of electrical stimulation on quadriceps strength after reconstructive surgery of the anterior cruciate ligament. *Am J Sports Med.* 1987;15:215-220.
63. Snyder-Mackler L, Delitto A, Bailey SL, Stralka SW. Strength of the quadriceps femoris muscle and functional recovery after reconstruction of the anterior cruciate ligament. A prospective, randomized clinical trial of electrical stimulation. *J Bone Joint Surg Am.* 1995;77:1166-1173.
64. Snyder-Mackler L, Delitto A, Stralka SW, Bailey SL. Use of electrical stimulation to enhance recovery of quadriceps femoris muscle force production in patients following anterior cruciate ligament reconstruction. *Phys Ther.* 1994;74:901-907.
65. Snyder-Mackler L, Ladin Z, Schepsis AA, Young JC. Electrical stimulation of the thigh muscles after reconstruction of the anterior cruciate ligament: effects of electrically elicited contraction of the quadriceps femoris and hamstring muscles on gait and on strength of the thigh muscles. *J Bone Joint Surg Am.* 1991;73:1025-1036.
66. Spindler KP, Huston IJ, Wright RW, et al. The prognosis and predictors of sports function and activity at minimum 6 years after anterior cruciate ligament reconstruction: a population cohort study. *Am J Sports Med.* 2011;39:348-359.
67. Timm KE. The clinical and cost-effectiveness of two different programs for rehabilitation following ACL reconstruction. *J Orthop Sports Phys Ther.* 1994;25:43-48.
68. Tovin BJ, Wolf SL, Greenfield BH, Crouse J, Woodfin BA. Comparison of the effects of exercise in water and on land on the rehabilitation of patients with intra-articular anterior cruciate ligament reconstructions. *Phys Ther.* 1994;74:710-719.
69. Tyler TF, McHugh MP, Gleim GW, Nicholas SJ. The effect of immediate weightbearing after anterior cruciate ligament reconstruction. *Clin Orthop Relat Res.* 1998;357:141-148.
70. Vathrakokilis K. Effects of a balance training protocol on knee joint proprioception after anterior cruciate ligament reconstruction. *J Bank Musculoskeletal Rehab.* 2008;21:233-237.
71. Wigerstad-Lossing I, Grimby G, Jonsson T, Morelli B, Peterson L, Renstrom P. Effects of electrical muscle stimulation combined with voluntary contractions after knee ligament surgery. *Med Sci Sports Exerc.* 1988;20:93-98.
72. Wright RW, Dunn WR, Amendola A, et al. Risk of tearing the intact anterior cruciate ligament in the contralateral knee and rupturing the anterior cruciate ligament graft during the first 2 years after anterior cruciate ligament reconstruction: a prospective MOON cohort study. *Am J Sports Med.* 2007;35:1131-1134.
73. Wright RW, Fetzter GB. Bracing after ACL reconstruction: a systematic review. *Clin Orthop Relat Res.* 2007;455:162-168.
74. Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part I. Continuous passive motion, early weight bearing, postoperative bracing, and home-based rehabilitation. *J Knee Surg.* 2008;21:217-224.
75. Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part II. Open versus closed kinetic chain exercises, neuromuscular electrical stimulation, accelerated rehabilitation, and miscellaneous topics. *J Knee Surg.* 2008;21:225-234.
76. Yates CK, McCarthy MR, Hirsch HS, Pascale MS. Effects of continuous passive motion following ACL reconstruction with autogenous patellar tendon grafts. *J Sport Rehabil.* 1992;1:121-131.