

Anthocyanins from *Vaccinium arctostaphylos* Berries

Bahman Nickavar¹ and Gholamreza Amin²

¹Department of Pharmacognosy, School of Pharmacy, Shaheed Beheshti University of Medical Sciences, Tehran, Iran;

²Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

Abstract

The systematic investigation of the berries of *Vaccinium arctostaphylos* L. (Ericaceae), a native medicinal plant growing in Iran, permitted the identification of three major anthocyanins. The compounds were isolated from the acidified extract of the berries using the repeated paper chromatography with different mobile phases and were characterized using chromatographic, chemical, and spectroscopic methods as delphinidin 3-*O*- β -glucoside, petunidin 3-*O*- β -glucoside, and malvidin 3-*O*- β -glucoside.

Keywords: Anthocyanin, Ericaceae, *Vaccinium arctostaphylos*.

Introduction

The genus *Vaccinium* (Ericaceae) comprises nearly 200 species, most of them found in the Northern Hemisphere (Morazzoni & Bombardelli, 1996). This genus is represented with only one taxon (*Vaccinium arctostaphylos* L.) in Iran (Wendelbo, 1965). *V. arctostaphylos* is a compact shrub about 1.5–2.5 m in height with many ascending branches and is found in the northern forests of the country between 1600–1800 m above sea level and known by the local names “Qaraqat” and/or “Cyah-gileh” (Ghahreman, 2001). The plant is therapeutically important in Iranian traditional medicine, and the decoction from the berries has been used as an antidiabetic and antihypertensive agent for a long time (Amin, 1991).

In regard to phytochemical profiles of *V. arctostaphylos*, previous studies have shown the presence of flavonol glycosides (Mzhavanadze, 1971a) and coumarins (Mzhavanadze et al., 1971b) in leaves and the presence phenolic acids and their derivatives in leaves (Mzhavanadze

et al., 1971b) and unripe fruits (Mzhavanadze et al., 1972). In addition, the essential oil composition of the flowering aerial parts has been determined (Nickavar et al., 2002).

In past decades, the anthocyanin content of the genus *Vaccinium*, especially of *V. myrtillus*, has received attention due to its relevant pharmacological properties (Francis et al., 1966; Ballinger et al., 1981, 1982; Baj et al., 1983; Andersen, 1985; Fuleki, 1986; Andersen, 1987a, 1987b, 1989; Gao & Mazza, 1994; Morazzoni & Bombardelli, 1996; Cabrita & Andersen, 1999) but, to the best of our knowledge, there is no previous report on the anthocyanin content of *V. arctostaphylos*.

The aim of this study was to determine the major anthocyanins in the berries of *V. arctostaphylos*. This investigation will be useful to identify the bioactive compounds, which may be responsible for the therapeutic properties of the plant berries.

Materials and Methods

Plant material

The ripe berries of *V. arctostaphylos* were collected from the forest region of Asalem in the north of Iran in August 2000. Voucher specimens were deposited in the Herbarium of Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran (no. 6520 THE).

Extraction and isolation of compounds

The pigments were extracted by maceration of 400 g of the berries with methanol:glacial acetic acid:water (70:2:28). The extract was filtrated, concentrated, and then partitioned

Accepted: July 2, 2004

Address correspondence to: Bahman Nickavar, Department of Pharmacognosy, School of Pharmacy, Shaheed Beheshti University of Medical Sciences, P.O. Box 14155-6153, Tehran, Iran. E-mail: bnickavar@yahoo.com

against ethyl acetate (18 g). The pigments were isolated from the final aqueous extract by repeated descending paper chromatography (PC) on Whatman No. 3 MM paper with *n*-butanol:glacial acetic acid:water (4:1:5, upper phase) and 15% aqueous acetic acid as developing solvents. The bands were separately eluted with methanol:glacial acetic acid:water (90:5:5), concentrated and monitored by PC on Whatman No. 1 paper (Glassgen et al., 1992).

Identification of compounds

The structures of isolated compounds were determined on the basis of chromatographic, chemical, and spectral studies and confirmed by comparison with literature (Francis et al., 1966; Cabrita & Andersen, 1999).

UV-Vis spectra were recorded on Shimadzu UV-160A spectrophotometer in 0.01 N HCl-CH₃OH; ¹H NMR spectra were taken on a Bruker-Spectrospin 500 spectrometer in CD₃OD-CF₃COOD (5:1), and chemical shifts were recorded in δ -values; positive fast atom bombardment (FAB) mass spectra were obtained in glycerol-HCl as the matrix by Finigan MAT TSQ 700 mass spectrometer.

The nature of the sugar moiety in isolated compounds was determined through acid hydrolysis, according to the standard methods and direct comparison of each sugar moiety with authentic carbohydrates by co-chromatography on Merck cellulose TLC plates (Francis et al., 1966; Skaltsa et al., 1999).

Results and Discussion


V. arctostaphylos berries were extracted by the acidified hydroalcoholic solvent. The extract was chromatographed successively on PC with different mobile phases. Finally, three major pigments, **1** (13 mg), **2** (10 mg), and **3** (16 mg), were obtained. The structures of the isolated anthocyanins are shown in Figure 1. R_f data observed in standard PC conditions used for this group of pigments are presented in Table 1. The spectroscopic data for the compounds are in agreement with the reported values for delphinidin 3-*O*- β -glucoside, petunidin 3-*O*- β -glucoside, and malvidin 3-*O*- β -glucoside (Francis et al., 1966; Cabrita & Andersen, 1999).

Delphinidin 3-*O*- β -glucoside (1)

UV-Vis λ_{\max} (in CH₃OH-HCl): 539.5, 273, 219; + AlCl₃ (3 drops of 5% AlCl₃ in CH₃OH): 557.5, 273, 219 nm; E₄₄₀/E_{vis,max} = 29%.

¹H-NMR (500 MHz, in CD₃OD-CF₃COOD): δ (aglycone moiety) 8.93 (1H, *s*, H-4), 7.74 (2H, *s*, H-2' and H-6'), 6.84 (1H, *d*, *J* = 1.5 Hz, H-8), 6.64 (1H, *d*, *J* = 1.5 Hz, H-6); δ (sugar moiety) 5.32 (1H, *d*, *J* = 7.6 Hz, H-1''), 3.96–3.51 (5H, *m*, H-2''-H-6'') ppm.

FAB-MS *m/z*: 465 for M⁺ = C₂₁H₂₁O₁₂⁺.



Delphinidin 3-*O*- β -glucoside (1) : R₁ = R₂ = H

Petunidin 3-*O*- β -glucoside (2) : R₁ = CH₃; R₂ = H

Malvidin 3-*O*- β -glucoside (3) : R₁ = R₂ = CH₃

Figure 1. Structures of the anthocyanins isolated from *Vaccinium arctostaphylos*.

Table 1. R_f data of the anthocyanins isolated from *Vaccinium arctostaphylos*.

Anthocyanin	PC (R _f × 100)	
	BAW ^a	15% HOAc ^b
Delphinidin 3- <i>O</i> - β -glucoside (1)	22	32
Petunidin 3- <i>O</i> - β -glucoside (2)	26	37
Malvidin 3- <i>O</i> - β -glucoside (3)	32	49

Solvent systems: ^aBAW, *n*-butanol:glacial acetic acid:water (4:1:5, upper phase); ^b15% HOAc, water:glacial acetic acid (85:15). PC, paper chromatography.

Petunidin 3-*O*- β -glucoside (2)

UV-Vis λ_{\max} (in CH₃OH-HCl): 536, 275, 218.5; + AlCl₃ (3 drops of 5% AlCl₃ in CH₃OH): 550, 275, 218.5 nm; E₄₄₀/E_{vis,max} = 28%.

¹H-NMR (500 MHz, in CD₃OD-CF₃COOD): δ (aglycone moiety) 8.98 (1H, *s*, H-4), 7.94 (1H, *d*, *J* = 1.7 Hz, H-2'), 7.77 (1H, *d*, *J* = 1.7 Hz, H-6'), 6.88 (1H, *d*, *J* = 1.6 Hz, H-8), 6.64 (1H, *d*, *J* = 1.6 Hz, H-6), 3.97 (3H, *s*, OCH₃-3'); δ (sugar moiety) 5.23 (1H, *d*, *J* = 7.6 Hz, H-1''), 3.84–3.26 (5H, *m*, H-2''-H-6'') ppm.

FAB-MS *m/z*: 479 for M⁺ = C₂₂H₂₃O₁₂⁺.

Malvidin 3-*O*- β -glucoside (3)

UV-Vis λ_{\max} (in CH₃OH-HCl): 539, 277.5, 215.5; + AlCl₃ (3 drops of 5% AlCl₃ in CH₃OH): 539, 277.5, 215.5 nm; E₄₄₀/E_{vis,max} = 23%.

¹H-NMR (500 MHz, in CD₃OD-CF₃COOD): δ (aglycone moiety) 9.03 (1H, *s*, H-4), 7.97 (2H, *s*, H-2' and H-6'), 6.94 (1H, *d*, *J* = 1.5 Hz, H-8), 6.66 (1H, *d*, *J* = 1.5 Hz, H-6), 3.99 (6H, *s*, OCH₃-3' and OCH₃-5'); δ (sugar moiety) 5.25 (1H, *d*, *J* = 7.6 Hz, H-1''), 3.84–3.30 (5H, *m*, H-2'', H-6'') ppm.

FAB-MS *m/z*: 493 for M⁺ = C₂₃H₂₅O₁₂⁺.

The previous studies on the berries of the genus *Vaccinium* revealed the presence of 3-*O*-monoglycosides in which the aglycones delphinidin, cyanidin, petunidin, malvidin, and peonodin are combined with the sugars galactose, arabinose, and glucose. The current study, the first to report the anthocyanin content in *V. arctostaphylos*, also shows a similar chemical profile. These anthocyanins are likely the active compounds responsible for some therapeutic effects that have been supposed for the berries of *V. arctostaphylos* in Iranian traditional medicine (Amin, 1991). On the other hand, these compounds seem to have potential as taxonomic markers in the genus *Vaccinium* (Cabrita & Andersen, 1999).

References

- Amin G (1991): *Popular Medicinal Plants of Iran*, Vol. 1. Tehran, Research Deputy of Health Ministry, p. 126.
- Andersen ØM (1985): Chromatographic separation of anthocyanins in cowberry (lingonberry), *Vaccinium vitis-idaea* L. *J Food Sci* 50: 1230–1232.
- Andersen ØM (1987a): Anthocyanins in fruits of *Vaccinium japonicum*. *Phytochemistry* 26: 1220–1221.
- Andersen ØM (1987b): Anthocyanins in fruits of *Vaccinium uliginosum* L. *J Food Sci* 52: 665–666.
- Andersen ØM (1989): Anthocyanins in fruits of *Vaccinium oxycoccus* L. *J Food Sci* 54: 383–384.
- Baj A, Bombardelli B, Gabetta B, Martinelli EM (1983): Qualitative and quantitative evaluation of *Vaccinium myrtillus* anthocyanins by high-resolution gas chromatography and high-performance liquid chromatography. *J Chromatogr* 279: 365–372.
- Ballinger WE, Maness EP, Ballington JR (1981): Anthocyanin and total flavonol content of *Vaccinium stamineum* L. fruit. *Sci Hortic* 15: 173–178.
- Ballinger WE, Maness EP, Ballington JR (1982): Anthocyanins in ripe fruit of the sparkleberry, *Vaccinium arboreum* Marsh. *Can J Plant Sci* 62: 683–687.
- Cabrita L, Andersen ØM (1999): Anthocyanins in blue berries of *Vaccinium padifolium*. *Phytochemistry* 52: 1693–1696.
- Francis FJ, Harborne JB, Barker WG (1966): Anthocyanins in the lowbush blueberry, *Vaccinium angustifolium*. *J Food Sci* 31: 583–587.
- Fuleki T (1986): Individual anthocyanin composition of American cranberry cultivars. *Bull Liaison-Groupe Polyphenols* 13: 374–379.
- Gao L, Mazza G (1994): Quantitation and distribution of simple and acylated anthocyanins and other phenolics in blueberries. *J Food Sci* 59: 1057–1059.
- Ghahreman A (2001): *Flora of Iran*, Vol. 22, no. 2749. Tehran, Research Institute of Forest and Rangeland.
- Glassgen WE, Seitz HU, Metzger JW (1992): High-performance liquid chromatography/electrospray mass spectrometry and tandem mass spectrometry of anthocyanins from plant tissues and cell cultures of *Daucus carota* L. *Biol Mass Spectrom* 21: 271–277.
- Morazzoni P, Bombardelli E (1996): *Vaccinium arctostaphylos* L. *Fitoterapia* 67: 3–29.
- Mzhavanadze VV (1971a): Kaempferol glycosides from the leaves of the caucasian bilberry, *Vaccinium arctostaphylos*. *Soobshch Akad Nauk Gruz SSR* 62: 445–447.
- Mzhavanadze VV, Targamadze IL, Dranik LI (1971b): Polyphenols from *Vaccinium arctostaphylos* leaves. *Khim Prir Soedin* 7: 546.
- Mzhavanadze VV, Targamadze IL, Dranik LI (1972): Phenolic compounds of unripe blueberry *Vaccinium arctostaphylos* fruits. *Soobshch Akad Nauk Gruz SSR* 68: 205–208.
- Nickavar B, Salehi-Sormagi MH, Amin G, Daneshtalab M (2002): Steam volatiles of *Vaccinium arctostaphylos*. *Pharm Biol* 40: 448–449.
- Skaltsa H, Tzakou O, Singh M (1999): Polyphenols of *Ocimum sanctum* from Suriname. *Pharm Biol* 37: 92–94.
- Wendelbo P (1965): Ericaceae. In: Rechinger KH, ed., *Flora Iranica*, Vol. 11. Graz, Akademische Druck-U. Verlagsanstalt, p. 2.

Copyright of Pharmaceutical Biology is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.