



Antimicrobial Activity of Necklace Orchids is Phylogenetically Clustered and can be Predicted With a Biological Response Method

Richa Kusuma Wati^{1,2*}, Esmée F. de Graaf³, Diego Bogarin^{1,4}, Reinout Heijungs^{5,6}, Rogier van Vugt⁷, Erik F. Smets^{1,8,9} and Barbara Gravendeel^{1,8,10}

¹Naturalis Biodiversity Center, Endless Forms Group, Leiden, Netherlands, ²Center for Plant Conservation, Bogor Botanic Garden, Indonesian Institute of Sciences (LIPI), Bogor, Indonesia, ³Science and Technology Faculty, University of Applied Sciences Leiden, Leiden, Netherlands, ⁴Lankester Botanical Garden, University of Costa Rica, Cartago, Costa Rica, ⁵Institute of Environmental Sciences, Leiden University, Leiden, Netherlands, ⁶Department of Econometrics and Operations Research, Vrije Universiteit Amsterdam, Amsterdam, Netherlands, ⁷Hortus botanicus, Leiden University, Leiden, Netherlands, ⁸Institute of Biology Leiden, Leiden University, Leiden, Netherlands, ⁹Ecology, Evolution and Biodiversity Conservation, KU Leuven, Heverlee, Belgium, ¹⁰Institute of Water and Wetland Research, Radboud University, Nijmegen, Netherlands

OPEN ACCESS

Edited by:

Prof Pulok Kumar Mukherjee,
Institute of Bio-Resources and
Sustainable Development (IBSD), India

Reviewed by:

Guo-Qiang Zhang,
Orchid Conservation and Research
Center, China
Christian Agyare,
Kwame Nkrumah University of
Science and Technology, Ghana

*Correspondence:

Richa Kusuma Wati
richa.kusumawati@naturalis.nl;
rich002@lpi.go.id

Specialty section:

This article was submitted to
Ethnopharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 23 July 2020

Accepted: 24 November 2020

Published: 12 March 2021

Citation:

Wati RK, de Graaf EF, Bogarin D,
Heijungs R, van Vugt R, Smets EF and
Gravendeel B (2021) Antimicrobial
Activity of Necklace Orchids is
Phylogenetically Clustered and can be
Predicted With a Biological
Response Method.
Front. Pharmacol. 11:586345.
doi: 10.3389/fphar.2020.586345

Necklace orchids (Coelogyinae, Epidendroideae) have been used in traditional medicine practices for centuries. Previous studies on a subset of unrelated orchid species utilized in these traditional practices revealed they possessed antimicrobial, anti-inflammatory, and anti-oxidant activity, providing experimental proof for their medicinal properties. To date however none of these species have been investigated ethno-botanically in a phylogenetic context. This study carried out comparative bioprospecting for a group of wild orchids using EBDCS (the Economic Botany Data Collection Standards) organ targeted and biological response methods. The traditional medicinal use of necklace orchids was recorded from books and journals published between 1984 and 2016. Two orchids, *Coelogyne cristata* and *Coelogyne fimbriata*, were selected, cultivated both indoors and outdoors, and the antimicrobial properties on extracts from their leaves and pseudobulbs tested against a selection of human pathogens. A molecular phylogeny of Coelogyinae based on nuclear ribosomal ITS and plastid *matK* DNA sequences obtained from 148 species was reconstructed with Maximum Likelihood (ML) using RAxML, Maximum Parsimony (MP) using PAUP and Bayesian Inference using MrBayes. Bioprospecting comparison of EBDCS and biological response was carried out using customized R scripts. Ethanolic extracts obtained from leaves of *C. fimbriata* inhibited growth of *Bacillus cereus*, *Staphylococcus aureus*, and *Yersinia enterocolitica*, confirming the antimicrobial properties of these extracts. Leaf extracts were found to have slightly stronger antimicrobial properties for plants cultivated outdoors than indoors. These differences were not found to be statistically significant though. Three hot nodes with high potency for antimicrobial activities were detected with the EBDCS organ targeted classification method, and eight hot nodes were detected with the biological response classification method. The biological response classification method is thus a more effective tool in finding hot nodes amongst clades of species with high medicinal potential.

Keywords: bio-assays, bioprospecting, Coelogyinae, herbal medicine, horticulture, hot nodes

INTRODUCTION

For millennia, products of nature have been an important source of traditional medicine (Cragg and Newman, 2013). Even today, between 70 and 95% of the world population in developing countries continues to use traditional medicines (Robinson and Zhang, 2011). Plant-based antibiotics form the basis of these traditional medicinal systems (Newman et al., 2000). There is an increasing interest in the study of these plant-based antibiotics as a source of novel antibiotics that human pathogens may not have developed resistance against, and because of the increasing extinction rate of wild plant species (Savoia, 2012; Cragg and Newman, 2013; Ernst et al., 2016).

To discover potential new plant species with antimicrobial properties, a time-efficient and systematic approach is needed. Bioprospecting is an approach combining phylogeny with ethnobotanical knowledge to identify potential sources of bioactive compounds. The underlying assumption is that phylogenies can predict the traditional medicinal use of natural products in a bioprospecting approach (Saslis-Lagoudakis et al., 2012; Leonti et al., 2013; Ernst et al., 2016). The hypothesis is that closely related species share similar biochemical pathways and that the same bioactive compounds are present in all descendants of a single ancestor rather than in species scattered over unrelated clades. This method has been used in different plant species (Douwes et al., 2008; Zhu et al., 2011; Saslis-Lagoudakis et al., 2012; Siqueira et al., 2012; Leonti et al., 2013) and animal groups (Smith and Wheeler, 2006). For bioprospecting, two different methods are mainly used. The first method is the Economic Botany Data Collection Standard (EBDCS) classification method. The EBDCS provides a system where cultural plant uses are described using standardized descriptors and terms, and attached to taxonomic data sets. This classification is based on the treatment of symptoms, i.e. a medicine against stomach pain (Cook, 1995). The other method is a classification based on the biological response, such as a medicine with antimicrobial effects (Ernst et al., 2016).

Pathogens cause an array of diseases in humans, and their identification is important in administering the correct treatment. (Washington, 1996). It is expected that bioprospecting based on biological responses will produce different results from the organ targeted EBDCS method, as biological responses are focused on a classification based on a single effect in the entire human body rather than a single organ (Ernst et al., 2016). A growing number of studies report on the bioprospecting of medicinal plants, including orchids (Beena, 2011; Purkayastha, 2016). We have not yet come across any study carried out on a group of wild orchids from a phylogenetic perspective.

The orchid family is historically well-known for its medicinal properties (Lawler, 1984; Singh and Singh, 2012). Medicinal orchids contain phytochemicals such as alkaloids, bibenzyl derivatives, flavonoids, phenanthrenes and terpenoids, which are present in leaves, roots, pseudobulbs (modified stem parts for water and nutrient storage), and flowers (Gutiérrez, 2010; Hsiao et al., 2011; Pant, 2014). Necklace orchids (Coelogyinae, Epidendroideae) comprise over 680 species, that are distributed throughout Southeast Asia (Pridgeon et al., 2005). *Bletilla*, *Coelogyne*,

Dendrochilum, *Otochilus*, *Pholidota*, *Pleione*, and *Thunia* are examples of necklace orchid genera with documented medicinal properties (Singh and Duggal, 2009; Subedi et al., 2011; Pant and Raskoti, 2013; Teoh, 2016) (see **Figure 1**).

In this study, we 1) compiled traditional medicinal uses of necklace orchids from the literature, 2) carried out bio-assays on six human pathogens with ethanol and hexane extracts of leaves and pseudobulbs from *C. cristata* and *C. fimbriata* plants grown both inside a glasshouse and outside to experimentally validate whether traditional growth methods impacted the orchid medicinal properties, and 3) investigated whether an organ-targeted EBDCS or biological response-based classification was most informative for predicting the biological activity of related species.

MATERIALS AND METHODS

Medicinal Uses of Necklace Orchids Recorded in the Literature

Information on the medicinal use of different species of necklace orchids was compiled from scientific journals and books through September 2019 (**Table 1**). We included all data from publications that stated the local names, latin names and the traditional uses for the orchid species. We excluded the publications where only the local name and/or genus were given. All records were compiled into a list and coded according to the Economic Botany Data Collection Standard (EBDCS) as recommended by the Biodiversity Information Standards of the Taxonomic Databases Working Group (TDWG) (Cook, 1995). The medicinal properties of the orchid species were categorized into EBDCS level 2 characters and into biological response characters. We used the antimicrobial response character as defined in MedlinePlus with three different states: no response, possible response or unknown response. This definition assumes that a plant species should be categorized as no response when it is applied for anything other than antimicrobial treatments, such as bone fracture treatments. A possible response was scored when the disease dictionary of MedlinePlus dictated this. Reducing fever was for instance scored as a possible antimicrobial effect since fever is a biological response to infection. Finally, an 'unknown response' was given if no records of plant use were available. Medicinal properties of all *Glomera* species were categorized as unknown, as to the best of our knowledge no ethnobotanical information for this genus has been published.

Antimicrobial Activity Plant Material

Fresh pseudobulbs and leaves of mature sterile plants of *C. cristata* and *C. fimbriata* (3–5 different individuals per species) grown in greenhouses were obtained from Orchideeën Wubben (Hollandsche Rading, Netherlands) and Claessen Orchideeën (Nederweert-Eind, Netherlands). The same species were subsequently grown outside where they were exposed to UV light and herbivorous snails and insects during a period of 5 months in the Hortus botanicus (Leiden, Netherlands). A second batch of fresh pseudobulbs and leaves was then




FIGURE 1 | Examples of medicinally used necklace orchids investigated in this study. **(A)** *Pholidota imbricata*. **(B)** *Coelogyne fimbriata*. **(C)** *Coelogyne cristata*. **(D)** *Coelogyne fuscescens*. The *Pholidota* species depicted under A can only be cultivated in a humid greenhouse. The *Coelogyne* species depicted under B-D can be cultivated outdoors during the summer and early fall in temperate regions to stimulate the production of secondary compounds. Photographs by Rogier van Vugt.

harvested from these species. All leaves and pseudobulbs were sterilized and freeze-dried in a VirTis Benchtop Pro Freeze Dryer at -104°C until they reached a constant weight. The dried pseudobulbs and leaves were ground into a fine powder and about 1 g of the powder was extracted with 70% ethanol and 100% hexane in a vacuum speed extractor E-916 (Buchi, Switzerland) (40°C , 100 bar). The extracts were stored in a freezer (-20°C) before further use.

Bacterial Strains

The antimicrobial properties of extracts from pseudobulbs and leaves of *C. cristata* and *C. fimbriata* were evaluated with five bacterial strains that are common causes of human gastrointestinal tract infections and are resistant against a range of synthetic antibiotics (Mutsaers et al., 2001). Two strains of Gram-positive *Staphylococcus aureus* (ATCC 12600) and *Bacillus cereus* (ATCC 14579) and three strains of Gram-negative *Escherichia coli* (ATCC 10798), *Klebsiella pneumoniae* (ATCC BAA-3079), and *Yersinia enterocolitica* (ATCC 9610) bacteria were used for the experiments. The bacterial strains were provided by the University of Applied Sciences Leiden, Netherlands. All strains were cultured on Columbia Agar with 5% Sheep Blood (COL-S (BDTM)) overnight at aerobic conditions at 37°C (except for *Y. enterocolitica*, which was typically cultured at 30°C), followed by storage at 4°C for up to 1 week.

Antimicrobial Activity of Plants Extracts

A disk diffusion method was used to evaluate the antimicrobial activity for each plant extract. Each bacterial strain was streaked onto a plate, grown overnight, and used to inoculate Mueller-Hinton cation-adjusted agar broth 2 (Sigma-Aldrich). The cultures were incubated overnight under aerobic conditions at 37°C (except for *Y. enterocolitica*, which was grown at 30°C) on a rotary shaker (180 rpm) until a McFarland Standard of 0.5 (10^7 CFU/ml) was reached. The cultures were subsequently used to make a confluent growth on COL-S agar plates. Sterile filter paper disks with (10 mm diameter, Sigma-Aldrich) were loaded with the different plant extracts with a total content of 55 mg/ml. The disks were then evaporated by air at room temperature inside a laminar air flow hood for 20 min before they were placed onto the top of the inoculated plates. Sterile filter paper disks loaded with 7.5 μg of Levofloxacin (Sigma-Aldrich) were used as positive control, and sterile paper disks loaded with 5% DMSO (Sigma-Aldrich) were used as the negative control. All the samples were then incubated at 37°C (except for *Y. enterocolitica* at 30°C) for 24 h. All tests were performed in triplicate and the zones of inhibition were measured with an automatic Vernier caliper. The scale of the inhibitory effect was scored as follows: high (diameter zone ≥ 17 mm), intermediate ($14 \leq$ diameter zone < 16 mm), low (diameter zone ≤ 13 mm) (CLSI, 2011).

TABLE 1 | Information on traditional medicinal use of necklace orchids (Coelogyninae) compiled from the literature.

| Species | Use | | References |
|---|---|------------------------------|--|
| | Symptoms | Plant organ(s) | |
| <i>Bletilla formosana</i> (Hayata) Schltr. | Strengthen the lungs, stop bleeding and reduce swellings. Used for treatment of tuberculous cough, bronchiectasis, bleeding peptic ulcers, nose-bleed and treat cracks on the heel | Stems | Teoh (2016) |
| <i>Bletilla ochracea</i> Schltr. | See <i>B. striata</i> | Tubers/ pseudobulbs | Teoh (2016) |
| <i>Bletilla foliosa</i> (King & Pantl.) Tang & F.T.Wang | See <i>B. striata</i> | Tubers/ pseudobulbs | Teoh (2016) |
| <i>Bletilla striata</i> (Thunb.) Rchb.f. | Benefit the lungs (effect on pulmonary diseases), liver and stomach meridians. Effects of the medicine are haemostatic, reduce swelling and promotes regeneration of muscles and other tissues. Also used to treat sores, pustules and dry, chapped and burned skin | Tubers/ pseudobulbs | Teoh (2016) |
| <i>Coelogyne barbata</i> Lindl. Ex Griff. | The whole plant is valued for its ability to counter 'heat', relieve thirst, stop coughs and lessen pain. It is used to treat sore throat, pain at hernias, swelling of the scrotum, chappy extremities, traumatic injuries and 'lung-heat' | Entire plant | Teoh (2016) |
| <i>Coelogyne corymbosa</i> Lindl. | Paste applied to the forehead to relieve headaches, fresh juice applied to burns and wounds as an analgesic. It treats fractures and is used as haemostatic and to relieve pain. Reduces heat and taken for coughs, flu, and bronchitis | Pseudobulbs/ entire plant | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000), Yonzzone et al. (2012) |
| <i>Coelogyne cristata</i> Lindl. | Are given for constipation as well as diarrhoea and dysentery. It is also used as an aphrodisiac. Freshly collected paste or juice consumed to relieve headaches, fever and for indigestion. Pulp applied to burnt skin. Juice also applied to wounds and skin boils. Gum is used for sores. Used for cooling & soothing | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000) |
| <i>Coelogyne fimbriata</i> Lindl. | Powder used in tonic preparation and used to reduce heat | Pseudobulbs | Subedi et al. (2011, 2013), Teoh (2016) |
| <i>Coelogyne flaccida</i> Lindl. | Paste applied externally or consumed to relieve frontal headaches, fever, and boils. Juice is taken for indigestion. The whole plant is also used to clear heat, counter dryness, promote the production of body fluids, clear phlegm and stop coughs | Pseudobulbs/ entire plant | Pant and Raskoti (2013), Teoh (2016) |
| <i>Coelogyne fuscescens</i> Lindl. | Paste applied externally or consumed to relieve headaches, fever, and stomach/abdominal ache. Treat burns and otitis media. Has sometimes an aphrodisiac function | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011), Yonzzone et al. (2012, 2013), Teoh (2016), Vaidya et al. (2000) |
| <i>Coelogyne nitida</i> (Wall. Ex D. Don) Lindl. | Juice consumed against headaches and fever and recommended for stomach ache. Paste applied externally on burns | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016) |
| <i>Coelogyne ovalis</i> Lindl. | Used as a tonic, aphrodisiac and to treat coughs, urine infections and eye disorders | Not specified | Pant and Raskoti (2013), Teoh (2016), Yonzzone et al. (2012, 2013) |
| <i>Coelogyne prolifera</i> Lindl. | Paste consumed against headaches and fever. Paste applied externally on burns, boils and to relieve backache | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016) |
| <i>Coelogyne punctulata</i> Lindl. | Used to treat wounds, burns, dry coughs. Relieves pain and helps to heal the wounds | Pseudobulbs | Teoh (2016), Yonzzone et al. (2012, 2013) |
| <i>Coelogyne stricta</i> (D.Don) Schltr. | Paste applied externally against headaches and fever. Healing of fractured bones | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000), Yonzzone et al. (2013) |
| <i>Coelogyne trinervis</i> Lindl. | Used to treat fractures and sprains | Tuber | Teoh (2016) |
| <i>Otochilus lancilabius</i> Seidenf. | Paste applied to fractured and dislocated bones | Entire plant | Subedi et al. (2011, 2013) |
| <i>Pholidota articulata</i> Lindl. | Paste applied on fractured bones and consumed as a tonic. Root powder is used to treat cancer. Juice berries are used to treat ulcers, skin eruptions, traumatic injuries, and sores. Removes gas and reduce swelling. Also used to treat coughs caused by body heat, headache, dizziness, irregular menses, and uterine prolapse | Entire plant | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000) |
| <i>Pholidota cantonensis</i> Rolfe | Used to treat high fever, eczema, and haemorrhoids | Entire plant | Teoh (2016) |
| <i>Pholidota chinensis</i> Lindl. | Used for cooling, moistens the lungs, promotes salivation. Used to treat tuberculosis-associated haemoptysis, acute or chronic bronchitis, dry cough, pharyngitis, tonsillitis, toothache, peptic ulcer, gastroenteritis, dizziness, headache, post-concussion syndrome, neurasthenia, osteomyelitis and trauma | Entire plant | Teoh (2016), Yonzzone et al. (2012), Wang et al. (2006) |

(Continued)

TABLE 1 | (Continued) Information on traditional medicinal use of necklace orchids (Coelogyninae) compiled from the literature.

| Species | Use | | References |
|--|---|----------------------|---|
| | Symptoms | Plant organ(s) | |
| <i>Pholidota imbricata</i> Hook. | Paste consumed to relieve fever and powder as a tonic. Juice is applied to relieve navel pain, abdominal pain, rheumatic pain, and headache. Applied to boils and to treat fractures | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000), Yonzone et al. (2012, 2013) |
| <i>Pholidota pallida</i> Lindl. | Paste used to relieve fever, powder to induce sleep and to cure abdominal pain, juice used for navel pain and rheumatic pain and sore throat | Rhizome, pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016), Vaidya et al. (2000), Yonzone et al. (2012, 2013) |
| <i>Pleione bulbocodioides</i> (Franch.) Rolfe | Treatment for wet sores, sore throat, rabies, tuberculosis, asthma, boils, and carbuncles. Clears phlegm. It reduces inflammation and fever. It removes extravasated blood swellings. It is also used as a detoxifier | Entire plant | Teoh (2016) |
| <i>Pleione hookeriana</i> (Lindl.) Rolisson | Are used to remove heat, toxins, abscesses and lymphatic tuberculosis | Pseudobulbs | Teoh (2016) |
| <i>Pleione humilis</i> (Sm.) D.Don | Paste applied on cuts and wounds. Powder used as a tonic | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016) |
| <i>Pleione maculata</i> (Lindl.) Lindl. & Paxton | Used for liver and stomach ailments | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2013), Teoh (2016), Vaidya et al. (2000), Yonzone et al. (2012, 2013) |
| <i>Pleione praecox</i> (Sm.) D.Don | Dried powder consumed (with milk) as tonic and energizer. Paste externally applied on cuts and wounds | Pseudobulbs | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016) |
| <i>Thunia alba</i> (Lindl.) Rchb.f. | Paste used on fractured and dislocated bones. Benefit the lungs, clear phlegm and stop cough, remove bruises and improve blood flow | Entire plant | Pant and Raskoti (2013), Subedi et al. (2011, 2013), Teoh (2016) |

Phylogenetic Reconstructions Plant Sampling and DNA Extraction

Previously generated DNA sequences for necklace orchids (Gravendeel et al., 2001; Subedi et al., 2011; Sulistyono et al., 2015; Pedersen et al., 2020) were downloaded from NCBI GenBank (see **Supplementary Table S2** for more details). In addition, new DNA sequences were generated from 77 specimens of the necklace orchid genus *Glomera*. From these, 14 specimens were collected in the field in Seram, Papua and Papua New Guinea (Indonesia). The first author also identified living orchid material from the Bogor Botanical Garden with identification keys and taxonomic descriptions from Schuiteman and de Vogel (2001), Wati et al. (2018) and the website of de Vogel et al. (2019). Additionally, a total of 42 specimens from the living orchid collection of the Hortus botanicus Leiden, Netherlands were analysed. Lastly, 21 dried herbarium specimens from the Herbarium Bogoriense, Indonesia and the herbarium of Naturalis Biodiversity Center, Leiden, Netherlands, were analysed (see **Supplementary Table S2** for more details). Total genomic DNA was extracted from 50 mg of leaf tissue from herbarium or silica-gel dried material using the 2x CTAB (Cetyltrimethylammonium bromide) method of Doyle and Doyle (1987), or with the Qiagen DNeasy Plant mini kit (Qiagen) following the manufacturer's protocol.

Amplification and Sanger Sequencing

The nuclear ribosomal ITS-5.8S-ITS2 (nrITS) region of silica-gel dried leaf material was amplified using primers 17SE (5'-ACG AATTCATGGTCCGGTGAAGTGTC-3') and 26SE (5'-TAG AATCCCCGGTTCGCTCGCCGTTAC-3') as described by Sulistyono et al. (2015). Subsequently, a M13 universal sequencing primer was added to the 5' end of the forward

(ACGAATTCATGGTCCGGTGAAGTGTC) and reverse (TAGAATCCCCGGTTCGCTCGCCGTTAC) primers to improve Sanger sequencing efficiency. Each PCR reaction was 25 µL and included the template DNA, CoralLoad PCR buffer (Qiagen), dNTPs, Taq DNA Polymerase (Qiagen), and both primers. All PCR reactions were done on a C1000 Touch Thermal Cycler (Bio-Rad) instrument. The thermal cycling protocol began with a 5 min initial denaturation at 96°C, followed by 35 amplification cycles, each with 30 s denaturation at 96°C, 30 s annealing at 50°C, and 1 min extension at 72°C, followed by a final 7 min final extension at 72°C.

The nrITS region of herbarium preserved leaf material was amplified using primer p3 (5'-GACTCYCGGCAATGGATATCTCG-3') and p4 (5'-CCGCTTATTGATATGCTTAAACTCRGC-3') as described by Cheng et al. (2016) and primer efgF1 (5'-CGAGTC TTTGAACGCAAGTTGCG-3') and efgR1 (5'-GGCCAACGAGAC GATAACCC-3') that were newly designed. Each PCR reaction consisted of 25 µL, containing the template DNA, 5x Phire PCR buffer (ThermoScientific), BSA, dNTPs, Phire Hot Start II DNA Polymerase (ThermoScientific), and both primers. The thermal cycling protocol began with a 1 min initial denaturation at 98°C, followed by 40 amplification cycles, each with 10 s denaturation at 98°C, 10 s annealing at 50°C, and 20 s extension at 72°C, followed by a 1 min final extension at 72°C.

The *matK* region of silica dried silica-gel dried leaf material was amplified using two primer sets: 731F (5'-TCTGGAGTC TTTCTTGAGCGA-3') and 2R (5'-AACTAGTCGGAGTAG-3'), and 19F (5'-CGTTCTGACCATATTGCACTATG-3') and 881R (5'-TMTTCATCAGAATAAGAGT-3') as described by Sulistyono et al. (2015). The PCR reaction setup was the same as for nrITS with fresh plant material, but with additional BSA. The

TABLE 2 | Prevalence of various categories of medicinal use of necklace orchids for the organ targeted EBDCS classification method.

| EBDCS classification method | N _{Total species included in the category} | Prevalence |
|--|---|------------|
| Abnormalities | 5 | 0.033 |
| Circulatory system disorders | 2 | 0.013 |
| Digestive system disorders | 14 | 0.094 |
| Genitourinary system disorders | 5 | 0.033 |
| Ill-defined symptoms | 2 | 0.013 |
| Infections/infestations | 13 | 0.087 |
| Inflammations | 3 | 0.020 |
| Injuries | 16 | 0.108 |
| Mental disorders | 1 | 0.006 |
| Metabolic system disorders | 9 | 0.060 |
| Muscular-skeletal system disorders | 13 | 0.087 |
| Nervous system disorders | 1 | 0.006 |
| Nutritional disorders | 7 | 0.047 |
| Pain | 12 | 0.081 |
| Poisonings | 2 | 0.013 |
| Respiratory system disorders | 13 | 0.087 |
| Sensory system disorders | 1 | 0.006 |
| Skin/subcutaneous cellular tissue disorder | 13 | 0.087 |
| Unknown | 120 | 0.810 |
| N _{Total} (total number of species) | 148 | |

thermal cycling protocol began with a 5 min initial denaturation at 94°C, followed by 35 amplification cycles, each with a 1 min denaturation at 94°C, 30 s annealing at 50°C, and 1 min extension at 72°C, followed by a 7 min final extension at 72°C.

Sanger sequencing of the amplification products were performed at Baseclear (<http://www.baseclear.com/>), using an ABI-3730XL DNA Sequencer (Applied Biosystems). All sequences were deposited in NCBI GenBank. Accession numbers of all sequences can be found in **Supplementary Table S2**.

Sequence Editing and Phylogenetic Analysis

Sanger sequences were assembled and edited in Geneious® R8 (Biomatters Ltd., Auckland, New Zealand) (Kearse et al., 2012). The ends of all data sets were trimmed to avoid character misinterpretation. Ambiguous bases were replaced with “N” in the data matrix. DNA sequences were aligned using the MAFFT platform (Multiple Alignment Fast Fourier Transform) (Katoh and Standley, 2013) as implemented in Geneious® R8 with subsequent manual adjustment. Missing data were replaced with “?”.

A phylogenetic analysis was carried out using Bayesian Inference (BI) with *Arundina graminifolia* as an outgroup based on earlier studies (Gravendeel et al., 2001; Pedersen et al., 2020) that showed this genus to be most closely related to the necklace orchids. The chosen nucleotide substitution model GTR+G was calculated using the Akaike Information Criterion (AIC) in jModelTest2 v.2.1.6 (Darriba et al., 2015). The analyses were run in the CIPRES Science Gateway v.3.1.1 (Miller et al., 2010). We performed Bayesian interference analyses with Mr. Bayes v.3.2.6 on XSEDE (Huelsenbeck et al., 2004) with the following parameters for the alignment dataset: number of runs (nrns = 2), number of chains to run (nchains = 4), number of generations (ngen = 5×10^7), temperature parameter (temp =

2) and sampling frequency of 2000 yielding 25,000 trees per run. The log files from MrBayes were inspected in Tracer v.1.6 to check for convergence of independent runs (i.e. with estimated sample size (ESS) > 200). Maximum Likelihood analyses were performed with RAxML-HPC2 on XSEDE (8.2.10) (Stamakis et al., 2008) choosing the GTRGAMMA model for bootstrapping and 1,000 bootstrap iterations. Parsimony analyses were performed with PAUPRat: Parsimony ratchet searches using PAUP* (Nixon, 1999; Sikes and Lewis, 2001; Swofford, 2002) with 1,000 ratchet repetitions, seed value = 0, 20% percent of characters to perturb (pct = 20), original weights 1 for all characters (wtmode = uniform) and a tree bisection-reconnection branch swapping algorithm (swap = TBR). The 50% majority rule consensus for MP was obtained with PAUP v4.0a152. and inspected in FigTree v.1.3.1. The statistical support of the clades was evaluated with the values of posterior probability (PP) for BI reconstruction, bootstrap for ML (MLB) and parsimony bootstrap for MP (MPB). The support values (PP) were added to the branches on the Bayesian 50% majority-rule consensus tree with additional support values shown for ML and MP when the same topology was retrieved.

Bioprospecting Analysis

A randomly selected subset of 1,000 trees within the 95% highest posterior density (HPD) interval was used for further analyses using the packages *caper*, *ape*, *plyr*, and (Paradis et al., 2004; Kembel et al., 2010; Wickham, 2011; Orme et al., 2013) scripts in the R programming language (R Core Team, 2018) under RStudio (Gandrud, 2015). The R bioprospecting script of Ernst et al. (2016) was used to assess evolutionary patterns of medicinal properties of the necklace orchids analysed. The strength of the phylogenetic signal of the EBDCS category and the antimicrobial biological response category were investigated using D statistics (Fritz and Purvis, 2010), that was calculated

TABLE 3 | Prevalence of various categories of medicinal use of necklace orchids for the antimicrobial response classification method.

| Antimicrobial response classification method | N _{Total} number of species included in the category | Prevalence |
|--|---|------------|
| No documented response | 111 | 0,75 |
| Possible response | 19 | 0,123 |
| Unknown process | 122 | 0,824 |
| N _{Total} (total number of species) | 148 | |

TABLE 4 | Antimicrobial activity of extracts of *Coelogyne cristata* and *C. fimbriata* as recorded in the bioassays carried out in this study of five plants per species grown in greenhouses.

| Extracts | Zone of inhibition (mm) | | | | |
|---|-------------------------|-------------------------|------------------------------|------------------------------|--------------------------------|
| | <i>Bacillus cereus</i> | <i>Escherichia coli</i> | <i>Klebsiella pneumoniae</i> | <i>Staphylococcus aureus</i> | <i>Yersinia enterocolitica</i> |
| Positive control (7.5 µg/ml levofloxacin) | 22.87 ± 1.0 | 13.12 ± 0.2 | 22 ± 2.2 | 14.6 ± 0.6 | 38 ± 1.5 |
| 70% EtOH pseudobulbs <i>C. cristata</i> | — | — | — | — | — |
| 70% EtOH leaves <i>C. cristata</i> | — | — | — | — | — |
| Hexane pseudobulbs <i>C. cristata</i> | — | — | — | — | — |
| Hexane leaves <i>C. cristata</i> | — | — | — | — | — |
| 70% EtOH pseudobulbs <i>C. fimbriata</i> | — | — | — | — | — |
| 70% EtOH leaves <i>C. fimbriata</i> | 15.55 ± 0.6 | 13.88 ± 0.7 | 18.55 ± 0.6 | 13.3 ± 1.0 | 21.7 ± 2.0 |
| Hexane pseudobulbs <i>C. fimbriata</i> | — | — | — | — | — |
| Hexane leaves <i>C. fimbriata</i> | — | — | — | — | — |

All experiments were carried out in triplicate. Absence of growth inhibition is indicated with —

with the *phylo.d* function implemented in the R package *capser* (Orme et al., 2013). A boxplot of the D values for each category of the two classification methods investigated was made using *ggplot*. If 95% of the 1,000 trees had a median value of $D > 1$, the medicinal properties were considered as randomly distributed; for $D < 1$, the phylogenetic signal was considered as strong (Ernst et al., 2016). $D > 0$ indicates that the medicinal properties of the orchids possess a significantly different distribution from the standard Brownian model, implying that they are clustered within the phylogeny. $D < 0$ indicates that the categories are extremely clustered. The prevalence of each category was measured by $N_{\text{total species included in the category}} / N_{\text{total number of species}}$. For a prevalence < 0.020 the category was considered as too biased, and omitted from further analyses.

We also tested the phylogenetic diversity (PD) of the EBDCS category and the antimicrobial biological response category with the function *pd* in the R package *picante* v.1.6-2 (Kembel et al., 2010). The percentage of the possible response category of the antimicrobial biological response was compared with the Infections/Infestations category of the EBDCS classification method. A higher PD percentage means that species in this category are more scattered throughout the phylogeny. As a consequence, more potential species with medicinal properties are present because the PD-values are based on the total branch length spanned by the species (Ernst et al., 2016).

A consensus BI tree with 10% burnin was used to visualize the distribution of the two categories over the necklace orchid species investigated. Using the *nodesigl* command in R with the system

PHYLOCOM v4.2 (Webb et al., 2008), so-called 'hot nodes' were calculated to visualize potential medicinal species.

RESULTS

Medicinal Uses of Necklace Orchids Recorded in the Literature

For 28 necklace orchid species, traditional medicinal uses were compiled to determine 19 organ-targeted categories and a single biological response (i.e., antimicrobial) category with three different character states (see Tables 1, 2, and 3 for an overview of all data obtained from the literature). The prevalence of the categories Mental Disorders, Nervous System Disorders and Sensory System Disorders in the EBDCS classification method showed the lowest value of 0.006 because only one species was used in these categories.

Bioassays

None of the 100% hexane leaf and pseudobulb extracts and 70% of the ethanol pseudobulb extracts showed any antimicrobial effect in the bio-assays conducted. On the contrary, the 70% ethanol leaf extracts inhibited the growth of several of the human pathogens investigated (Table 4). Extracts obtained from freshly harvested leaves of *C. cristata* and *C. fimbriata* were found to inhibit growth of *Y. enterocolitica*, *B. cereus* and *S. aureus* and confirmed the traditional medicine uses recorded in the literature (Pyakurel and Gurung, 2008; Subedi, 2002; Subedi et al., 2013).

TABLE 5 | Antimicrobial activity of extracts of *Coelogyne cristata* and *C. fimbriata* as recorded in the bioassays carried out in this study of five plants per species grown outside for 5 months in the Hortus botanicus Leiden, Netherlands .

| Extracts | Zone of inhibition (mm) | | | | |
|---|-------------------------|-------------------------|------------------------------|------------------------------|--------------------------------|
| | <i>Bacillus cereus</i> | <i>Escherichia coli</i> | <i>Klebsiella pneumoniae</i> | <i>Staphylococcus aureus</i> | <i>Yersinia enterocolitica</i> |
| Positive control (7.5 µg/ml levofloxacin) | 22.51 ± 0.8 | 18.77 ± 0.4 | 26 ± 1.2 | 20.15 ± 0.2 | 39 ± 2.0 |
| 70% EtOH pseudobulbs <i>C. cristata</i> | — | — | — | — | — |
| 70% EtOH leaves <i>C. cristata</i> | — | — | — | — | — |
| 70% EtOH pseudobulbs <i>C. fimbriata</i> | — | — | — | — | — |
| 70% EtOH leaves <i>C. fimbriata</i> | 16.44 ± 0.8 | 14.55 ± 1.4 | 17.22 ± 1.0 | 22.55 ± 1.4 | 20.44 ± 1.1 |

All experiments were carried out in triplicate. Absence of inhibition zone is indicated with —.

The highest effect was recorded for the 70% EtOH leaf extracts of *C. fimbriata* against *Y. enterocolitica* (19.6 ± 4.2 mm). Intermediate effects were recorded for leaf extracts of the same *Coelogyne* species against *B. cereus* (14.3 ± 1.4 mm) and *S. aureus* (13.6 ± 1.2 mm). Leaf extracts were found to have slightly stronger (but not significant) antimicrobial properties for plants cultivated outdoors than indoors (Table 5).

Bioprospecting of Necklace Orchids

The majority consensus reconstructed BI tree, which is based on combined nrITS and plastid *matK* sequences for 148 species of necklace orchid species, is depicted in Figure 2. The consensus tree of ML and MP shows relatively high support for (>70%) and was congruent with the topology of the majority consensus BI tree. The Infections/Infestation category of the organ targeted EBDCS (Figure 2A) classification method and the biological (i.e. antimicrobial response) method (Figure 2B) were plotted on the BI tree.

The boxplots of the D-statistics for the organ-based EBDCS classification method and the antimicrobial biological response classification method are shown in Figures 3 and 4. For the EBDCS classification method, 7 of the 19 categories showed a $D > 1$, indicating that a minority of these categories were randomly distributed. A total of 12 of the 19 categories were (extremely) clustered. For the antimicrobial response method, all the categories were found to be (extremely) clustered.

The median of the phylogenetic diversity (PD) was calculated to compare the phylogenetic distribution of medicinal species from the Unknown, Possible and No Antimicrobial Response categories with the 19 categories of the organ based EBDCS classification method. In Figure 5, these medians are depicted. The Possible Antimicrobial Response category of the biological classification method had a median of 18.83%, whereas the Infections/Infestations category of the organ based EBDCS classification method had a median of 13.32%.

To narrow down potential new species with antimicrobial activities, the Possible Antimicrobial Response state of the biological response classification method was compared with the Infections/Infestations category of the EBDCS classification method using the PHYLOCOM platform. Figure 2 depicts the recovered hot nodes. Figure 2A shows the three hot nodes detected for the category within the EBDCS classification method with high potency for antimicrobial activities. Figure 2B shows the eight hot


nodes detected for the biological response classification method.

DISCUSSION

When compiling data on medicinal use of necklace orchids from the scientific literature, we noted that information for specific species was not always provided. This was for instance the case for the genus *Dendrochilum*. We thus urge ethnobotanists to make vouchers so that more detailed information for a particular genus can be obtained to link medicinal uses to the species level. Detailed information on plant organs used for medicinal purposes was not always provided either. We therefore recommend ethnobotanists to ask more detailed questions about specific organs used when interviewing traditional plant healers working with orchids. When information about plant organs was mentioned, this was sometimes contradictory among different publications. Our bio-assays showed that antimicrobial effects for extracts of leaves from *C. fimbriata* were much higher than those for pseudobulbs, which is not fully in agreement with previous publications, where it was found that pseudobulbs were the main source of secondary metabolites (Tóth et al., 2018). Our results are however supported by the publication of Buyun et al. (2016), who found that leaf extracts from *Coelogyne ovalis* produced larger inhibition zones than pseudobulbs extracts. The ethanol extract of leaves for *Bulbophyllum neilgherrense* also showed the same result (Priya and Krishnaveni, 2005). A possible explanation might be that the metabolites present in the pseudobulbs are more diluted as the relative amount of water is usually higher in these organs than in the leaves.

The exact method used to obtain plant extracts was also not always provided. Extracts dissolved in 70% ethanol had a higher antimicrobial effect in our bio-assays than extracts dissolved in hexane. This difference might be explained by the fact that hexane is a non-polar solvent that mostly extracts large fatty acid methyl esters with hydrocarbons and terpenes, whereas known antimicrobial substances isolated from necklace orchids are mostly phenanthrenes (Majumder et al., 2001; Kovács et al., 2008; Yang et al., 2012; Pant, 2014; Qian et al., 2015), which dissolve more readily in polar solvents such as ethanol.

Whether medicinal orchids were collected in the wild or from gardens or greenhouses was not mentioned in any of the



publications that we screened. Our bio-assays show that antimicrobial effects of extracts of plants grown outside were higher (but not significantly so) than those of plants cultivated in greenhouses. A possible explanation for the difference in antimicrobial activity may be that plants naturally produce secondary metabolites that have a role in the defence against abiotic and biotic stresses (Dangle and Jones, 2001; Kim et al., 2009; Ramakrishna and Ravishankar, 2011). Recent studies by Isah (2019) also show that both stress and defense responses are involved in secondary metabolite production in plants. The insignificant differences found in our experiments when comparing indoor vs. outdoor cultivation methods might be a result of a too short exposure to UV light and herbivory, resulting in a too low level of secondary metabolites to create a significant difference between the cultivation methods. Plants grown in temperature controlled sterile greenhouses are generally exposed to less abiotic (UV light) and biotic (herbivores) stress and might therefore produce fewer secondary metabolites. Li et al. (1996) for instance reported that a longer exposure period to direct sunlight promoted higher ginsenoside production in American ginseng plants. Nevertheless, our results show that while exposure to UV light and herbivores may increase the antimicrobial activity of leaf extracts for necklace orchids, plants grown indoors also possess antimicrobial activity. This result contradicts the common folk belief that medicinal orchids can only be harvested from the wild to maintain their potency. We therefore encourage cultivation of medicinal necklace orchids

in order to prevent overexploitation and extinction of rare species in the wild.

The 70% ethanol leaf extracts of *C. fimbriata* showed *in vitro* antimicrobial activity against *S. aureus*, *B. cereus* and *Y. enterocolitica*, all known to cause gastrointestinal tract infections in humans. Activity was observed against both Gram-positive and Gram-negative bacteria, which indicates a broad spectrum of antimicrobial effects of leaf extracts of necklace orchids. The extracts were not able to inhibit growth of *E. coli* and *K. pneumoniae*. This can be explained by the fact that Gram-negative bacteria generally develop more resistance against synthetic antibiotics as compared with Gram-positive bacteria because they can more efficiently regulate genes involved in antibiotic drug resistance (Peleg and Hooper, 2010).

In contrast with the organ targeted EBDCS classification method, all the categories from the biological response (i.e. antimicrobial) classification method were found to be (highly) clustered. The biological response classification method can therefore be considered as more informative for bioprospecting. The biological response classification method also had a more scattered distribution of medicinal species on the phylogeny than the EBDCS classification method, covering a wider group of potential medicinal necklace orchid species by retrieving eight hot nodes as compared with the organ targeted EBDCS classification method, that only found three hot nodes. One of the eight hot nodes detected by the biological response classification method, but not by the organ targeted EBDCS classification method, encompasses species of the necklace




FIGURE 3 | Boxplot of the 19 categories of the organ targeted EBCS classification method (indicated with different colors) over which the data on medicinally used necklace orchid species that were analysed phylogenetically can be divided. The red lines indicate the D reference values 0 (on the left) and 1 (on the right). The box boundaries indicate the first and third quartile (Q1 and Q3), the line indicates the median, and the whiskers extend to either the extreme values or 1.5 times the interquartile range (Q3–Q1).





FIGURE 4 | Boxplot of the Antimicrobial biological response classification method of the three characters states over which the data on medicinally used necklace orchid species, that were analysed phylogenetically, could be divided. Color codes: no response (orange), possible response (green), unknown response (blue). The red lines indicate the D reference values 0 (on the left) and 1 (on the right). The box boundaries indicate the first and third quartile (Q1 and Q3), the line indicates the median, and the whiskers extend to either the extreme values or 1.5 times the interquartile range (Q3–Q1).



orchid genus *Bletilla*. Yang et al. (2012) successfully isolated bletilin A, bletilin B and other phenanthrenes from *Bletilla ochracea* tuber extracts that showed antibacterial activities against *S. aureus*, *S. epidermis* and *B. subtilis*. The fibrous roots and tubers from *Bletilla striata* contain biphenanthrenes and stilbenoids, which possess antibacterial activity (Kovács et al., 2008; Qian et al., 2015). Additionally, dihydrophenanthrenes, phenanthrene, flavonoids, bibenzyl and phenolic compounds were isolated from entire plants of *B. formosana* by Lin et al. (2005). These research findings support the results of our bioprospecting analyses and show that the biological response classification method is more effective in uncovering potential clades with high medicinal potential as compared with the EBDCS classification method.

Ethno-directed approaches to identifying plants traditionally used to treat specific diseases received significantly more attention over the past decade as this method shows a relatively high success rate compared to random plant screening programmes (Douwes et al., 2008; Siqueira et al., 2012). Plotting ethno-pharmacological data on a phylogenetic tree can be used as a time-efficient approach to discover potential new plant species with medicinal properties (Ernst et al., 2015), especially when a plant group is as large and diverse as the orchid family. We could only analyse 10% of all

necklace orchid species for their medicinal properties. The reason for this was that while for some species with recorded medicinal use no DNA sequences were available, other species with known DNA sequences had not yet been investigated for their medicinal uses. We encourage more work into the ethnobotany and pharmacology of necklace orchids to increase species sampling. Especially species of the genera *Bletilla*, *Coelogyne* sect. *Bicellae*, sect. *Brachypterae*, sect. *Coelogyne*, sect. *Elatae*, sect. *Flaccidae*, sect. *Fuscescentes*, sect. *Hologyne*, sect. *Lawrenceana*, sect. *Lentiginosae*, sect. *Longifoliae*, sect. *Ocellatae*, sect. *Proliferae*, sect. *Ptychogyne*, sect. *Speciosae*, *Neogyne*, *Otochilus* and *Pholidota* sect. *Articulatae*, sect. *Chinenses*, sect. *Crinonia*, sect. *Pholidota* and sect. *Repentes* seem very promising for further research as these were identified to belong to hot node clades with high potency of antimicrobial activity.

CONCLUSION

We successfully applied bioprospecting to discover new necklace orchid species with antimicrobial activity. The traditional antimicrobial use of necklace orchids could be confirmed with bio-assays for leaf extracts prepared with 70% ethanol. Additionally, outdoor cultivation may result in increased

antimicrobial activity, though this needs to be further explored. The biological response classification method was more effective in uncovering hot nodes leading to clades of species of necklace orchids with high antimicrobial potential as compared to the EBDCS classification method.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found in the article/**Supplementary Material**.

AUTHOR CONTRIBUTIONS

RKW and EFG performed the experiments and analyzed the data. DB assisted with R analysis, RH helped with statistics, RV helped with orchid cultivation. RKW wrote the article with assistance from BG and EFS. All authors have read and approved the manuscript.

FUNDING

LPDP funded the PhD project of RKW (0044737/SC/D/2/lpdp2014) and Naturalis Biodiversity Center funded the open access publication fees. This research was partially supported by a machine allocation on the Kabré supercomputer at the Costa Rica National High Technology Center (CNCA/CeNAT).

REFERENCES

- Beena, C. (2011). Bioprospecting on the endangered medicinal plant *Nervilia aragoana* gaud. *J. Progress. Agric.* 2, 75–78. doi:10.7324/JABB.2019.70215
- Buyun, L., Tkachenko, H., Kovalska, L., and Osadowski, Z. (2016). "Preliminary screening of *Coelogyne ovalis* Lindl. (Orchidaceae) for antimicrobial activity against *Staphylococcus aureus*", in *Agrobiodiversity for improving Nutrition, health and life quality 2017*, 43–49. doi:10.15414/agrobiodiversity.2017.2585-8246
- Cheng, T., Xu, C., Lei, L., Li, C., Zhang, Y., and Zhou, S. (2016). Barcoding the kingdom Plantae: New PCR primers for ITS regions of plants with improved universality and specificity. *Mol. Ecol. Resour.* 16, 138–149. doi:10.1111/1755-0998.12438
- Cook, F. E. M. (1995). *Economic Botany data collection standard*. Editors J. M. Lock and H. D. Prendergast (Kew: the Royal Botanic Gardens). doi:10.1111/j.1756-1051.1997.tb00317.x
- CLSI, (2011). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-First Informational Supplement. Wayne, PA: Clinical and Laboratory Standards Institute.
- Cragg, G. M., and Newman, D. J. (2013). Natural products: a continuing source of novel drug leads. *Biochim. Biophys. Acta* 1830, 3670–3695. doi:10.1016/j.bbagen.2013.02.008
- Dangle, J. L., and Jones, J. D. (2001). Plant pathogens and integrated defence responses to infection. *Nature* 411, 826–833. doi:10.1038/35081161
- Darriba, D., Taboada, G. L., Doallo, R., and Posada, D. (2015). jModelTest 2: more models, new heuristics and high-performance computing Europe PMC Funders Group. *Nat. Methods* 9, 772. doi:10.1038/nmeth.2109
- de Vogel, E. F., Vermeulen, J. J., and Schuiteman, A. (2019). Orchids of New Guinea <http://www.orchidsnewguinea.com/>. Accessed in 2019.
- Douwes, E., Crouch, N. R., Edwards, T. J., and Mulholland, D. A. (2008). Regression analyses of southern African ethnomedicinal plants: informing the targeted selection of bioprospecting and pharmacological screening subjects. *J. Ethnopharmacol.* 119, 356–364. doi:10.1016/j.jep.2008.07.040

ACKNOWLEDGMENTS

The authors would like to thank Lina Juswara, Aninda Wibowo, Bima Atmadja, Wenni Setyo Lestari and Nicolien Sol for granting permission to extract DNA from specimens from the Herbarium Bogoriense (BO), Bali Botanic Gardens and the herbarium of Naturalis Biodiversity Center (L). Lengguru Scientific Exploration of Papuan Karst, Koos Wubben and Rik Claessen are thanked for providing fresh orchid material. Henrie Korthout from Fytogoras helped us with finding the most effective extraction protocol for the bioassays. Rieneke Boeije and Kiki Fontein helped with optimising protocols for the bioassays. We also thank Mark Arentshorst from the Institute of Biology of Leiden University for granting us access to lab facilities certified for carrying out bioassays and Jolanda van Schie of the University of Applied Sciences Leiden for providing us with bacterial stocks. Marcella Orwick Rydmark is thanked for helping with language checking and Ivo Horn and Hugo de Boer for valuable suggestions for this manuscript. Last but not least we thank the Indonesia Endowment Fund for Education (LPDP) of the Ministry of Finance of Indonesia for funding the PhD project of RW.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://doi.org/10.6084/m9.figshare.13071893.v2>.

- Doyle, J. J., and Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19, 11–15.
- Ernst, M., Grace, O. M., Saslis-Lagoudakis, C. H., Nilsson, N., Simonsen, H. T., and Rønsted, N. (2015). Global medicinal uses of *Euphorbia* L. (Euphorbiaceae). *J. Ethnopharmacol.* 176, 90–101. doi:10.1016/j.jep.2015.10.025
- Ernst, M., Saslis-Lagoudakis, C., Grace, O. M., Nilsson, N., Simonsen, H. T., Horn, J. W., et al. (2016). Evolutionary prediction of medicinal properties in the genus *Euphorbia* L. *Sci. Rep.* 6, 30531. doi:10.1038/srep30531
- Fritz, S. A., and Purvis, A. (2010). Selectivity in mammalian extinction risk and threat types: a new measure of phylogenetic signal strength in binary traits. *Conserv. Biol.* 24, 1042–1051. doi:10.1111/j.1523-1739.2010.01455.x
- Gandrud, C. (2015). *Reproducible research with R and RStudio*. 2nd ed. Boca Raton, Florida: CRC Press. doi:10.1201/9781315382548
- Gravendeel, B., Chase, M. W., De Vogel, E. F., Roos, M. C., Mes, T. H., and Bachmann, K. (2001). Molecular phylogeny of *Coelogyne* (Epidendroideae; Orchidaceae) based on plastid RFLPS, matK, and nuclear ribosomal ITS sequences: evidence for polyphyly. *Am. J. Bot.* 88, 1915–1927. doi:10.2307/3558367
- Gutiérrez, R. M. P. (2010). Orchids: a review of uses in traditional medicine, its phytochemistry and pharmacology. *J. Med. Plants Res.* 4, 592–638. doi:10.5897/JMPR10.012
- Hsiao, Y. Y., Pan, Z. J., Hsu, C. C., Yang, Y. P., Hsu, Y. C., Chuang, Y. C., et al. (2011). Research on orchid biology and biotechnology. *Plant Cell Physiol.* 52, 1467–1486. doi:10.1093/pcp/pcr100
- Huelsenberg, J. P., Larget, B., and Alfaro, M. E. (2004). Bayesian phylogenetic model selection using reversible jump Markov chain Monte Carlo. *Mol. Biol. Evol.* 21, 1123–1133. doi:10.1093/molbev/msh123
- Isah, T. (2019). Stress and defense responses in plant secondary metabolites production. *Biol. Res.* 52, 39. doi:10.1186/s40659-019-0246-3
- Katoh, K., and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi:10.1093/molbev/mst010

- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., et al. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199
- Kemmel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., et al. (2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26, 1463–1464. doi:10.1093/bioinformatics/btq166
- Kim, J. Y., Park, S. C., Hwang, I., Cheong, H., Nah, J. W., Hahm, K. S., et al. (2009). Protease inhibitors from plants with antimicrobial activity. *Int. J. Mol. Sci.* 10, 2860–2872. doi:10.3390/ijms10062860
- Kovács, A., Vasas, A., and Hohmann, J. (2008). Natural phenanthrenes and their biological activity. *Phytochemistry* 69, 1084–1110. doi:10.1016/j.phytochem.2007.12.005
- Lawler, L. J. (1984). "Ethnobotany of the Orchidaceae", in *Orchid Biology: reviews and perspectives*. Editor J. Arditti (New York: Cornell University Press), Vol. III, 27–149.
- Leonti, M., Cabras, S., Castellanos, M., Challenger, A., Gertsch, J., and Casu, L. (2013). Bioprospecting: evolutionary implications from a post-olmec pharmacopoeia and the relevance of widespread taxa. *J. Ethnopharmacol.* 147, 92–107. doi:10.1016/j.jep.2013.02.012
- Li, T. S. C., Mazza, G., Cottrell, A. C., and Gao, L. (1996). Ginsenosides in roots and leaves of American Ginseng. *J. Agric. Food Chem.* 44, 717–720.
- Lin, Y. L., Chen, W. P., and Macabalang, A. D. (2005). Dihydrophenanthrenes from *Bletilla formosana*. *Chem. Pharm. Bull.* 53, 1111–1113. doi:10.1248/cpb.53.1111
- Majumder, P. L., Sen, S., and Majumder, S. (2001). Phenanthrene derivatives from the orchid *Coelogyne cristata*. *Phytochemistry* 58, 581–586. doi:10.1016/s0031-9422(01)00287-4
- Miller, M. A., Wayne, P., and Schwartz, T. (2010). "Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees." *2010 Gateway Computing Environments Workshop (GCE)*, (New Orleans, LA: GCE), 1–8.
- Mutsaers, J. A. E. M., Doornbos, L., and Knecht, N. M. (2001). *Bacteriologie voor laboratorium en kliniek* 2. 4th ed. Utrecht: Syntax Media.
- Newman, D. J., Cragg, G. M., and Snader, K. M. (2000). The influence of natural products upon drug discovery. *Nat. Prod. Rep.* 17, 215–234. doi:10.1039/a902202c
- Nixon, K. C. (1999). The parsimony ratchet, a new method for rapid parsimony analysis. *Cladistics* 1, 407–414.
- Orme, D., Freckleton, R., Thomas, G., Thomas, P., Fritz, S., Isaac, N., et al. (2013). CAPER: Comparative analyses of Phylogenetics and evolution in R. *Methods Ecol. Evol.* 3, 145–151. doi:10.1098/rsos.1811182
- Pant, B. (2014). Medicinal orchids and their uses: tissue culture a potential alternative for conservation. *Afr. J. Plant Sci.* 7, 448–467. doi:10.5897/ajps2013.1031
- Pant, B., and Raskoti, B. B. (2013). *Medicinal orchid of Nepal*. Kathmandu: Himalayan Map House Pvt. Ltd.
- Paradis, E., Claude, J., and Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289–290. doi:10.1093/bioinformatics/btg412
- Pedersen, H., Petersen, G., Gravendeel, B., Barkman, T. J., de Boer, H., Sulistyo, B. P., et al. (2020). Phylogenetics of *Dendrochilum* (Orchidaceae): evidence of pronounced morphological homoplasy and predominantly centric endemism. *Taxon* 68, 1173–1188. doi:10.1002/tax.12184
- Peleg, A. Y., and Hooper, D. C. (2010). Hospital-acquired infections due to gram-negative bacteria. *N. Engl. J. Med.* 362, 1804–1813. doi:10.1056/NEJMra0904124
- Pridgeon, A. M., Cribb, P. J., Chase, M. W., and Rasmussen, F. N. (2005). *Genera Orchidacearum, Epidendroideae (Part 1)*, Vol. 4. Oxford, USA: Oxford University Press.
- Priya, K., and Krishnaveni, C. (2005). Antibacterial effect of bulbophyllum neilgherrense wt. (Orchidaceae). An *in vitro* study. *Ancient Sci. Life* 25, 50–52. doi:10.1007/S00044012-0228-2
- Pyakurel, D., and Gurung, K. (2008). Enumeration of orchids and estimation of current stock of traded orchids in Rolpa district : Final report. Rolpa, Nepal: District forest office Rolpa.
- Purkayastha, J. (2016). *Bioprospecting of Indigenous Bioresources of North-East India*. Singapore, Singapore: Springer. doi:10.1007/978-981-10-0620-3
- Qian, C. D., Jiang, F. S., Yu, H. S., Shen, Y., Fu, Y. H., Cheng, D. Q., et al. (2015). Antibacterial biphenanthrenes from the fibrous roots of *Bletilla striata*. *J. Nat. Prod.* 78, 939–943. doi:10.1021/np501012n
- R Core Team (2018). *R: a language and environment for statistical computing*. Vienna, Austria.
- Ramakrishna, A., and Ravishankar, G. A. (2011). Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal. Behav.* 6, 1720–1731. doi:10.4161/psb.6.11.17613
- Robinson, M. M., and Zhang, X. (2011). *Traditional medicines: global situation, issues and Challenges*. Geneva, Switzerland: WHO.
- Saslis-Lagoudakis, C. H., Savolainen, V., Williamson, E. M., Forest, F., Wagstaff, S. J., Baral, S. R., et al. (2012). Phylogenies reveal predictive power of traditional medicine in bioprospecting. *Proc. Natl. Acad. Sci. USA* 109, 15835–15840. doi:10.1073/pnas.1202242109
- Savoia, D. (2012). Plant-derived antimicrobial compounds: alternatives to antibiotics. *Future Microbiol.* 7, 979–990. doi:10.2217/fmb.12.68
- Schuiteman, A., and de Vogel, E. F. (2001). *Orchids of New Guinea*, Vol. I. Illustrated Checklist and Genera CD-ROM. 90-75000-20-0.
- Sikes, D. S., and Lewis, P. O. (2001). Beta software, version 1. PAUPRat:PAUP* implementation of the parsimony ratchet.
- Singh, A., and Duggal, S. (2009). Medicinal orchids: an overview. *Ethnobot. Leaflet* 13, 351–363.
- Singh, S., and Singh, A. K. (2012). Medicinal properties and uses of orchids: a concise review. *Elixir Appl. Bot.* 52, 11627–11634.
- Siqueira, C. F., Cabral, D. L., Peixoto Sobrinho, T. J., De Amorim, E. L., De Melo, J. G., Araújo, T. A., et al. (2012). Levels of tannins and flavonoids in medicinal plants: evaluating bioprospecting strategies. *Evid. Based Complem. Alternat. Med.* 2012, 434782. doi:10.1155/2012/434782
- Smith, W. L., and Wheeler, W. C. (2006). Venom evolution widespread in fishes: a phylogenetic road map for the bioprospecting of piscine venoms. *J. Hered.* 97, 206–217. doi:10.1093/jhered/esj034
- Stamakis, A., Hoover, P., and Rougemont, J. (2008). A rapid bootstrap algorithm for the RAXML web servers. *Syst. Biol.* 57, 758–771. doi:10.1080/10635150802429642
- Subedi, A. (2002). Orchids around Pokhara valley of Nepal. Local Initiatives for Biodiversity, Pokhara, Nepal: . Research and Development.
- Subedi, A., Chaudhary, R. P., van Achterberg, C., Heijerman, T., Lens, F., van Dooren, T. J., et al. (2011). Pollination and protection against herbivory of Nepalese Coelogyninae (Orchidaceae). *Am. J. Bot.* 98, 1095–1103. doi:10.3732/ajb.1000306
- Subedi, A., Kunwar, B., Choi, Y., Dai, Y., van Andel, T., Chaudhary, R. P., et al. (2013). Collection and trade of wild-harvested orchids in Nepal. *J. Ethnobiol. Ethnomed.* 9, 64–10. doi:10.1186/1746-4269-9-64
- Sulistyo, B. P., Boos, R., Coates, J. E., and Gravendeel, B. (2015). *Dendrochilum hampelii* (Coelogyninae, Epidendroideae, Orchidaceae) traded as 'Big Pink' is a new species, not a hybrid: evidence from nrITS, matK and ycf1 sequence data. *PhytoKeys* 56, 83–97. doi:10.3897/phytokeys.56.5432
- Swofford, D. L. (2002). *PAUP. Phylogenetic analysis using Parsimony (and other methods), Version 4.0 Beta 10*. Sunderland: Sinauer Associates.
- Teoh, E. S. (2016). *Medicinal orchids of Asia*. Switzerland: Springer. doi:10.1007/978-3-319-24274-3
- Tóth, B., Hohmann, J., and Vasas, A. (2018). Phenanthrenes: a Promising Group of Plant secondary metabolites. *J. Nat. Prod.* 81, 661–678. doi:10.1021/acs.jnatprod.7b00619
- Vaidya, B., Shrestha, M., and Joshee, N. (2000). "Report on Nepalese orchid species with medicinal Properties," in *Proceeding of Nepal-Japan Joint Symposium-2000*. Editors T. Watanabe, A. Tokano, M. Bista, and H. Saiju (Japan Society for Conservation and Development of Himalaya Medicinal Resources), 146–152.
- Wang, J., Matsuzaki, K., and Kitanaka, S. (2006). Stilbene derivatives from *Pholidota chinensis* and their anti-inflammatory activity. *Chem. Pharm. Bull.* 54, 1216–1218. doi:10.1248/cpb.54.1216
- Washington, J. A. (1996). *Principles of Diagnosis*. Editor S. Baron Galveston (Texas: University of Texas Medical Branch).
- Wati, R. K., van Vugt, R. R., and Gravendeel, B. (2018). A Linnaeus NG interactive key to the species of *Glomera* (Orchidaceae, Coelogyninae) from Southeast Asia. *PhytoKeys* 110, 9–22. doi:10.3897/phytokeys.110.28435
- Webb, C. O., Ackerly, D. D., and Kembel, S. W. (2008). Phylocom: software for the analysis of phylogenetic community structure and trait evolution. *Bioinformatics* 24, 2098–2100. doi:10.1093/bioinformatics/btn358
- Wickham, H. (2011). The split-apply-combine strategy for data analysis. *J. Stat. Softw.* 40, 1–29. doi:10.18637/jss.v040.i01
- Yang, X., Tang, C., Zhao, P., Shu, G., and Mei, Z. (2012). Antimicrobial constituents from the tubers of *Bletilla ochracea*. *Planta Med.* 78, 606–610. doi:10.1055/s-0031-1298264
- Yonzon, R., Kamran, A., and Bhujel, R. B. (2012). "Orchids in ethnobotany", in *International seminar on "Multidisciplinary approaches in angiosperm systematics*, 661–669.
- Yonzon, R., Lama, D., Bhujel, R. B., and Rai, S. (2013). Present availability status, diversity resources and distribution of medicinal orchid species in Darjeeling Himalaya of West Bengal, India. *Int. J. Pharm. Pharmaceut. Sci.* 1, 14–35.

Zhu, F., Qin, C., Tao, L., Liu, X., Shi, Z., Ma, X., et al. (2011). Clustered patterns of species origins of nature-derived drugs and clues for future bioprospecting. *Proc. Natl. Acad. Sci. USA* 108, 12943–12948. doi:10.1073/pnas.1107336108

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Wati, de Graaf, Bogarín, Heijungs, van Vugt, Smets and Gravendeel. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.