

 Open access • Journal Article • DOI:10.1111/JAM.12196

Antimicrobial activity of phenolic compounds identified in wild mushrooms, SAR analysis and docking studies — [Source link](#)

Maria José Alves, Maria José Alves, Isabel C.F.R. Ferreira, Hugo J.C. Froufe ...+3 more authors

Institutions: Instituto Politécnico Nacional, Catholic University of Portugal

Published on: 01 Aug 2013 - Journal of Applied Microbiology (J Appl Microbiol)

Topics: Antimicrobial

Related papers:

- [A Review on Antimicrobial Activity of Mushroom \(Basidiomycetes\) Extracts and Isolated Compounds](#)
- [Antioxidants in Wild Mushrooms](#)
- [p-Coumaric acid kills bacteria through dual damage mechanisms](#)
- [Plant Products as Antimicrobial Agents](#)
- [Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/antimicrobial-activity-of-phenolic-compounds-identified-in-1vwvfmm7ki>

Antimicrobial activity of phenolic compounds identified in wild mushrooms, SAR analysis and docking studies

MARIA JOSÉ ALVES^{1,2,3}, ISABEL C.F.R. FERREIRA^{3,*}, HUGO J.C FROUFE³, RUI M.V. ABREU³, ANABELA MARTINS³, MANUELA PINTADO^{1,*}

^aCBQF-Escola Superior de Biotecnologia - Universidade Católica Portuguesa Porto, Rua Dr. António Bernardino de Almeida, 4200 – 072 Porto, Portugal.

^bCentro Hospitalar de Trás-os-Montes e Alto Douro- Unidade de Chaves, Av. Dr. Francisco Sá Carneiro, 5400-249 Chaves, Portugal.

^cCIMO/ESA, Instituto Politécnico de Bragança, Campus de Santa Apolónia, Apartado 1172, 5301-855 Bragança, Portugal.

*Authors to whom correspondence should be addressed (e-mail: iferreira@ipb.pt, telephone +351-273-303219, fax +351-273-325405; e-mail: mpintado@porto.ucp.pt, telephone +351-22-5580097, fax +351-22-5090351).

Running Head: Antimicrobial activity of phenolic compounds

Abstract

Aim and Methods: Although the antimicrobial activity of extracts from several mushroom species have been reported, studies with the individual compounds present in that extracts are scarce. Herein, the antimicrobial activity of different phenolic compounds identified and quantified in mushroom species from all over the world was evaluated. Furthermore, a structure activity relationship (SAR) analysis and molecular docking studies were performed, in order to provide insights in the mechanism of action of potential antimicrobial drugs for resistant microorganisms.

Results: 2,4-Dihydroxybenzoic and protocatechuic acids were the phenolic compounds with higher activity against the majority of Gram negative and Gram positive bacteria. Furthermore, phenolic compounds inhibited more MRSA than methicillin sensible *Staphylococcus aureus*. MRSA was inhibited by 2,4-dihydroxybenzoic, vanillic, syringic (MICs=0.5 mg/mL) and *p*-coumaric (MIC= 1 mg/mL) acids, while these compounds at the same concentrations had no inhibitory effects against methicillin sensible *Staphylococcus aureus*.

Conclusions: The presence of carboxylic acid (COOH), two hydroxyl (OH) groups in *para* and *ortho* positions of the benzene ring, as also a methoxyl (OCH₃) group in the *meta* position seems to be important for anti-MRSA activity.

Significance and Impact of the Study: Phenolic compounds could be used as antimicrobial agents, namely against some microorganisms resistant to commercial antibiotics.

Keywords: Wild mushrooms; Antimicrobial activity; Clinical isolates; SAR; Docking

Introduction

In recent years, there is an increasing number of reports on phenolic compounds in different mushroom species. Phenolic acids including benzoic and cinnamic acid derivatives have been pointed out as the most common. Among benzoic acid derivatives, *p*-hydroxybenzoic, protocatechuic, gallic, vanillic and syringic acids were identified in different mushroom species (Puttaraju *et al.* 2006; Kim *et al.* 2008; Barros *et al.* 2009; Heleno *et al.* 2011; Reis *et al.* 2011; Vaz *et al.* 2011; Heleno *et al.* 2012) (Table 1). The identification of cinnamic acid and its derivatives such as *p*-coumaric, *o*-coumaric, caffeic, ferulic and chlorogenic acids was also described (Mattila *et al.* 2001; Valentão *et al.* 2005; Puttaraju *et al.* 2006; Barros *et al.* 2008; Kim *et al.* 2008; Heleno *et al.* 2011; Reis *et al.* 2011; Vaz *et al.* 2011; Heleno *et al.* 2012). The presence of some flavonoids such quercetin, rutin and chrysin (Valentão *et al.* 2005; Ribeiro *et al.* 2006; Kim *et al.* 2008; Jayakumar *et al.* 2009; Yaltirak *et al.* 2009) and tannins like ellagic acid (Ribeiro *et al.* 2007) were reported (Table 1).

In vitro and epidemiologic studies suggest that consumption of foods rich in phenolic compounds might significantly decrease the risk of some health problems due to their antioxidant, antimutagenic, anti-inflammatory and antibacterial properties (Surh 2002; Albayrak *et al.* 2010). Nowadays, the evidence that the increasing number of microorganisms resistant to the available antibiotics is an emergent problem and subject for researchers and clinicians from all over the world. In general, it can be observed that the treatment of virus, bacteria, fungi and protozoa with the existent drugs is increasingly difficult. To overlap the disadvantages of the available antimicrobial drugs, other drugs with new mechanisms of action should be developed (Khalafi-Nezhad *et al.* 2005).

Although the antimicrobial activity of extracts from several mushroom species have been reported ([Barros *et al.* 2007](#); [Quereshi *et al.* 2010](#); [Ozen *et al.* 2011](#); [Alves *et al.*, 2012](#)), studies with the individual compounds present in that extracts are scarce, being mainly related to phenolic compounds identified in plant sources ([Kuate *et al.* 2009](#); [Orhan *et al.* 2010](#); [Lou *et al.* 2012](#)).

Therefore, the aim of the present study was to evaluate the antimicrobial activity of most relevant compounds identified and quantified in mushroom species from all over the world. Furthermore, a structure activity relationship (SAR) analysis and molecular docking studies against Penicillin Binding Protein 2a (PBP2a) were performed, in order to give insights in the mechanism of action of potential antimicrobial drugs for resistant microorganisms. Molecular docking is an *in silico* tool that predicts how a ligand (substrate or drug candidate) interacts with a receptor (e.g. proteins involved in several biological processes) and has been successfully applied in several therapeutic programs at the lead discovery stage ([Ghosh *et al.* 2006](#)).

Materials and Methods

Standards and reagents

The culture media Muller Hinton broth (MHB) and Wilkins-Chalgren Broth (WCB) were obtained from Biomerieux (Marcy l' Etoile, France), respectively. The dye *p*-iodonitrotetrazolium chloride (INT) was purchased from sigma–Aldrich (Spruce Street; St. Louis, USA) to be used as microbial growth indicator. Water was treated in a Milli-Q water purification system (TGI Pure Water Systems, USA) before use.

Phenolic compounds

Sixteen phenolic compounds (phenolic acids, flavonoids and tannins) already identified in tens of different wild mushroom species by our research group and by others (**Table 1**), were submitted to antimicrobial activity evaluation against Gram positive and Gram negative bacteria clinical isolates. Compounds were dissolved in water or in water with 1% DMSO (for flavonoids and tannins), at a concentration of 10 mg/mL, and stored at -20 °C for further use (up to 1 week).

Microorganisms and culture media

The microorganisms used were clinical isolates from patients hospitalized in various departments of the Hospital Center of Trás-os-Montes e Alto Douro – Chaves, Portugal. Six Gram positive bacteria (methicillin-susceptible *Staphylococcus aureus* (MSSA) and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from wound exudates, *Staphylococcus epidermidis*, *Enterococcus faecalis* and *Listeria monocytogenes* isolated from blood culture, and *Streptococcus agalactiae* isolated from vaginal swab and five Gram negative bacteria (*Escherichia coli*, *Proteus mirabilis* and *Morganella morganii*, isolated from urine, *Pasteurella multocida* isolated from synozial fluid and *Neisseria gonorrhoeae* isolated from urethral exudate) were used to screen the antimicrobial activity of the selected phenolic compounds. *Escherichia coli* showed resistance to fluoroquinolones (levofloxacin and ciprofloxacin) and ampicillin, being intermedia for amoxicillin/clavulanic acid; *Proteus mirabilis* was resistant to nalidixic acid, levofloxacin, nitrofurantoin, fosfomicin and trimethoprim/sulfasoxazole, and intermediate to gentamicin; *Morganella morganii* showed resistance to ampicillin, amoxicillin/clavulanic acid, cephalothin, cefazolin, cefuroxime, nitrofurantoin, fosfomicin and trimethoprim /sulfasoxazole; MSSA was only resistant to penicillin and

ampicillin, while MRSA was resistant to oxacillin, levofloxacin and ciprofloxacin; *Staphylococcus epidermidis* showed resistance to oxacillin and erythromycin.

All strains were identified using the MicroScan automated methodology - Siemens.

Muller Hinton broth (MHB) and Wilkins-Chalgren Broth (WCB) were used for determination of Minimum Inhibitory Concentration (MIC, lowest concentration of the phenolic compound able to completely inhibit bacterial growth).

Test assays for antimicrobial activity

MIC determinations were performed by the microdilution method and the rapid *p*-iodonitrotetrazolium chloride (INT) colorimetric assay following the methodology suggested by Kuete *et al.* (2011) with some modifications.

Initially, 50 µL of each phenolic compound solution (1mg/mL) filter sterilized was diluted in 450 µL of MHB for all microorganisms except for *Neisseria gonorrhoea* where WCB was used (also with final concentration of 1 mg/mL) and then, 200 µL of this solution was added in each well (96-well microplate). Titrations (eight different final concentrations) were carried out over the wells containing 100 µL of MHB or WCB and afterwards, 10 µL of inoculum (1×10^8 cfu/mL) were added to all the wells.

Two negative (one with MHB or WCB and the other with the phenolic compound) and one positive (with MHB or WCB and the inoculum) controls were performed. The plates were incubated at 37 °C, for 24 h, in an oven (Jouan, Berlin, Germany) or with humidified atmosphere containing 10% CO₂ (NuAire, Plymouth, USA), in the case of *Neisseria gonorrhoeae*.

The MIC of the samples were detected following addition of INT (0.2 mg/mL, 40 µL) and incubation at 37 °C for 30 min. Viable microorganisms reduced the yellow dye to a pink colour. MIC was defined as the lowest phenolic compound concentration that

prevented this change and exhibited complete inhibition of bacterial growth. All the assays were carried out in duplicate.

Compounds and protein structure preparation

ACD/ChemSketch Freeware 12.0 software was used to design 2D structure of the compounds. The software VegaZZ 2.3.1 ([Pedretti *et al.* 2004](#)) was then used to convert all compounds from 2D to 3D structures. AutoDockTools1.5.2 (ADT) ([Sanner 2005](#)) was used to merge nonpolar hydrogens, add Gasteiger charges, and set up rotatable bonds through AutoTors.

The crystal structure of PBP2a (Penicillin Binding Protein 2a) was obtained from the Protein Data Bank (PDB): 1VQQ (PDB entry) ([Lim and Strynadka 2002](#)). The software AutoDockTools was also used to assign polar hydrogens, add gasteiger charges and save the protein structure in PDBQT file format. AutoGrid4 ([Morris *et al.* 2009](#)) was used to create affinity grid maps for all the atoms on the protein and phenolic compounds used.

Molecular docking

AutoDock4 (version 4.2) with the Lamarckian genetic algorithm was used to perform the docking studies. Docking parameters selected for AutoDock4 runs were as follows: 100 docking runs, population size of 200, random starting position and conformation, translation step ranges of 2.0 Å, mutation rate of 0.02, crossover rate of 0.8, local search rate of 0.06, and 2.5 million energy evaluations. Docked conformations were clustered using a tolerance of 2.0 Å RMSD (Root Mean Square Deviation). The molecular docking experiments were performed on a dedicated cluster of 64 Core AMD 2.0 GHz, running on CentOS and using MOLA, a custom designed software for virtual screening

using AutoDock (Abreu et al. 2010). All figures with structure representations were produced using PyMOL (The PyMOL Molecular Graphics System, Version 1.3, Schrödinger, LLC. Available at: (<http://www.pymol.org/>)).

Results

Table 1 presents phenolic compounds that have been identified in different mushroom species from several countries, where it can be observed that different compounds were detected in the same species. Several external factors have been pointed to explain this fact, such as the heterogeneous enzymatic and oxidative decomposition after collection, different stress conditions associated to each sample, and even dissimilar methodologies applied to phenolic compounds extraction (Oke and Aslim 2011; Vaz et al. 2011).

These compounds are well known for their antioxidant properties (Puttaraju et al. 2006; Ribeiro et al. 2007; Kim et al. 2008), but they also revealed antimicrobial activity (Barros et al. 2007; Quereshi et al. 2010; Schwan et al. 2010; Ozen et al. 2011) emerging with potential against multiresistances. Their increasing prevalence is one of the major challenges for the healthcare systems worldwide. Antibiotic resistant infections are associated with a 1.3 to 2-fold increase in mortality compared to antibiotic susceptible infections (Cosgrove and Carmeli 2003). Moreover, antibiotic resistance imposes enormous health expenditure due to the higher treatment costs and longer hospital stays. In addition, the development of new generations of antibiotic drugs is stalling.

In the present study, in the range of tested concentrations (0.78-1000 µg/mL), 2,4-dihydroxybenzoic, protocatechuic, vanillic and *p*-coumaric acids showed antibacterial activity (MIC=1 mg/mL) against *Escherichia coli*, *Pasteurella multocida* and *Neisseria gonorrhoeae* (**Table 2**). It should be highlighted that the *Escherichia coli* isolate used

herein shows resistance to fluoroquinolones (levofloxacin and ciprofloxacin) and ampicillin, being intermedia for amoxicilin/clavulanic acid. Kuete *et al.* (2009) reported a MIC=78 µg/mL for protocatechuic acid isolated from *Ficus ovata* against *Escherichia coli* (β- lactamases positive). The observed difference in MIC values could be related to the use of strains with different susceptibility profiles. *Escherichia coli* resistance to fluoroquinolones and cephalosporins has drastically increased in the last decade (Rogers *et al.* 2011), the mentioned phenolic acids could be an option against this bacteria. Recently, Lou *et al.* (2012) also reported the antimicrobial activity of *p*-coumaric acid (MIC=80 µg/L) against *Escherichia coli*, but also against other Gram negative bacteria such as *Salmonella typhimurium* and *Shigella dysenteriae*; this compound changes the permeability of the cell membrane and has the capacity to bind DNA inhibiting cell function. Other authors (Teke *et al.* 2011) described the antimicrobial activity of vanillic acid against *Escherichia coli* and *Proteus mirabilis*, which is in agreement with the results reported herein. Moreover, the *Proteus mirabilis* strain used in the present study shows resistance to nalidixic acid, ciprofloxacin, nitrofurantoin, fosfomicin and trimethoprim/sulfasoxazole, being intermedia for gentamicin. Nevertheless, it should be highlighted that the strains used herein have different antibiotic resistance profiles, while the ones used in the mentioned study did not revealed relevant resistances; this important feature could be related to the differences observed in MIC values.

Despite the absence of reports regarding the presence of 2,4-dihydroxybenzoic acid in mushrooms and its antimicrobial activity, due to the chemical similarity with other phenolic acids mentioned as antimicrobial compounds, we decided to test it and, as far as we know this is the first report on its activity against Gram negative bacteria.

Gallic acid, ferulic acid and quercetin exhibited activity only against *Pasteurella multocida* and *Neisseria gonorrhoeae*, and the latter was mainly sensible to quercetin

(MIC=0.5 mg/mL; **Table 2**). According to WHO report published in 2001, more than six million cases of gonorrhea (infection caused by *Neisseria gonorrhoeae*) occur in each year, and with increasing levels, mostly in developing countries; furthermore, there is an emergent resistance of this bacteria to the antimicrobial agents used in gonorrhea treatment. Therefore, the mentioned phenolic compounds could be an alternative to be explored for the control of this infection. Studies evaluating the antibacterial activity of mushroom extracts or isolated compounds against *Neisseria gonorrhoeae* are scarce, so it is important to clarify their mechanism of action upon this microorganism as also in other Gram negative cocci.

Although no activity was observed for rutin against the tested Gram negative bacteria (**Table 2**), other authors ([Orhan et al. 2010](#)) reported antimicrobial activity of this compound against different strains of Gram negative bacilli, such as *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Acinetobacter baumannii*.

Once more, 2,4-dihydroxybenzoic and protocatechuic acid were the phenolic compounds with higher activity against the majority of Gram positive bacteria (**Table 3**). Protocatechuic acid showed a MIC of 1 mg/mL for MSSA and MRSA, as also for *Listeria monocytogenes* and *Streptococcus agalactiae*. Other studies reported the antimicrobial activity of this compound against *Staphylococcus aureus* and with lower concentrations (MIC=156 µg/mL; [Kuate et al. 2009](#)). Once more, the strain used herein was resistant to oxacillin and to both fluoroquinolones (ciprofloxacin and levofloxacin), which could be responsible for the higher MIC value observed in comparison to the mentioned study.

Regarding *Staphylococcus*, ferulic and caffeic acids were the only phenolic compounds inhibiting *Staphylococcus aureus*, MRSA and *Staphylococcus epidermidis*.

Nevertheless, other authors reported antimicrobial activity of *p*-coumaric acid, quercetin and rutin against *Staphylococcus aureus* (Kueete *et al.* 2009; Orhan *et al.* 2010; Lou *et al.* 2012). The absence of antimicrobial activity observed in the present study could be related to the different dissolution solvent used, water, and not ethanol:hexane and Tween 80 as used by the mentioned authors. Nevertheless, some of those solvents might have some inherent toxicity and should be carefully used.

Syringic and ellagic acids showed a MIC of 0.5 mg/mL against *Listeria monocytogenes* (Table 3). Cinnamic acid seemed to be the most active upon *Streptococcus agalactiae* (CMI 0.5 mg/mL). Among all the tested phenolic compounds, only 2,4-dihydroxybenzoic acid inhibited *Enterococcus faecalis* (MIC=1 mg/mL); nonetheless, other authors described antimicrobial activity of rutin (MIC=128 mg/mL), protocatechuic acid (MIC=39 µg/mL) and vanillic acid (zone of inhibition 16 mm) (Kueete *et al.* 2009; Orhan *et al.* 2010; Teke *et al.* 2011). Isolates of *Enterococcus faecalis* and *Enterococcus faecium* are the third- to fourth-most prevalent nosocomial pathogen worldwide; acquired resistance, most prominently to penicilin/ampicillin, aminoglycosides (high-level resistance) and glycopeptides are reported in an increasing number of isolates, and the therapeutic spectrum in these cases is limited. Therefore, therapeutic alternatives to treat infections with multi- and vancomycin-resistant enterococci (VRE) are restricted to antibiotics introduced recently into clinical practice such as quinupristin/dalfopristin, linezolid, tigecyclin and daptomycin. However, these drugs are only approved for certain indications and resistance has already been reported (Montero *et al.* 2008; Werner *et al.* 2008), which emphasizes the importance of the discovery of new alternative drugs.

It should be notice that the differences among the results reported by several authors could be related to the use of strains with different resistance profiles, but also to

different methodologies used including different solvents for compound solution preparation or different techniques to determine MICs. In the present study, water was chosen for being the most innocuous solvent; however in the case of flavonoids and tannins water with 1% DMSO was used to assure the total solubility of the compounds. MRSA has been indicated as one of the major causes of nosocomial infections and its increasing prevalence has been observed in the last decade. Furthermore, the treatment of MRSA infections is difficult due to the restrict spectra of efficient antibiotics (Chambers *et al.* 2001). The obtained data in the present study (Table 3) show that phenolic compounds inhibited more MRSA than methicillin sensible *Staphylococcus aureus*. MRSA was inhibited by 2,4-dihydroxybenzoic, vanillic, syringic (MICs=0.5 mg/mL) and *p*-coumaric (MIC= 1 mg/mL) acids, while these compounds at the same concentrations had no inhibitory effects against methicillin sensible *Staphylococcus aureus*. Ferulic acid inhibited both MRSA and methicillin sensible *Staphylococcus aureus*, but in a lower concentration for MRSA (Table 3).

Discussion

Regarding these results, it is interesting to notice that the two *Staphylococcus aureus* tested showed different susceptibility toward the compounds tested, possibly explained by the different resistance mechanisms exhibited by each strain. To understand these differential effects, a structure-activity relationship (SAR) study was carried by analysing the different chemical structure patterns of the evaluated compounds.

Only phenolic acids (benzoic and cinnamic acid derivatives) showed activity, highlighting the importance of the carboxylic group in the molecule structure (proton acceptor). Furthermore, all the compounds with anti-MRSA activity have OH (proton donor) and OCH₃ (proton acceptor) groups in the *para* and *meta* positions of the

benzene ring, respectively (**Table 4**). In the absence of OCH₃ group in the *meta* position (*p*-coumaric acid), the activity decreased. Nevertheless, the absence of the mentioned group in the structure of 2,4-dihydroxybenzoic acid seemed to be overlapped by the OH substitution in *ortho* position of the benzene ring. Only OCH₃ (proton acceptor) or H in position 5 of the benzene ring allowed anti-MRSA activity, since when OH is presented in that position the activity disappears (see the examples of protocatechuic and gallic acids in **Table 4**).

MRSA is resistant to all β -lactam antibiotics and this ability is due to the acquisition of *mecA* gene (Lowy 2003). This gene encodes the PBP2a protein and when it is challenged by β -lactams, MRSA will use the transpeptidase functionality of PBP2a to synthesize the cell wall (Wilke *et al.* 2005).

Since the major difference between MSSA and MRSA is *mecA*, studies of molecular docking were performed using 3D crystal structure of PBP2a (PDB:1QVV) as target to understand the inhibition mechanism of the phenolic compounds with activity against MRSA. The docking results revealed a superimposition of the docking poses for the three benzoic acid derivatives (vanillic, 2,4-dihydroxybenzoic and syringic acids) (**Figure 1**).

The binding pose shows several hydrogen bonds (H-bonds) that validate SAR analysis described above. The carboxylic group is stabilized by H-bonds with the amino (NH₂) group of LYS406 side chain, the hydroxyl (OH) group of SER403 side chain and the carbamide (NH₂CO) group of ASP464 side chain. Furthermore, OH in the *para* position of the benzene ring, which contributes to the anti-MRSA activity of the compounds, establishes a hydrogen bond with Serine (SER461) carbonyl group of the peptidic bond. The OCH₃ group in the *meta* position of the benzene ring (as in vanillic and syringic acids) is stabilized by a hydrogen bond with Glutamate (GLU447) amine

group of the peptidic bond. The OH in *meta* position of the benzene ring (as in 2,4-dihydroxybenzoic acid) is stabilized by a hydrogen bond with Serine (SER462) carbonyl group of the peptidic bond.

Overall, 2,4-dihydroxybenzoic, protocatechuic, vanillic and *p*-coumaric acids were the compounds that showed higher antimicrobial activity against Gram positive and negative bacteria. Cinnamic acid derivatives revealed higher antimicrobial activity against Gram positive and negative coccus.

The presence of carboxylic acid (COOH), two hydroxyl (OH) groups in *para* and *ortho* positions of the benzene ring, as also a methoxyl (OCH₃) group in the *meta* position seems to play an important role in the studied phenolic compounds anti-MRSA activity. The docking studies provided strong evidence that the molecular basis for this activity is probably as PBP2a inhibitors. The mentioned compounds could be a solution for multiresistance problem, but their mechanism of action in different microorganisms should be better understood.

Acknowledgements

The authors are grateful to Fundação para a Ciência e a Tecnologia (FCT, Portugal) and COMPETE/QREN/EU for financial support to this work (research project PTDC/AGR-ALI/110062/2009) and to CIMO (strategic project PEst-OE/AGR/UI0690/2011) and to PEst-OE/EQB/LA0016/2011. They also thank to CHTAD – Hospital Center of Trás-os-Montes e Alto Douro and Siemens for all the support.

References

- Albayrak, S., Aksoy, A., Sagdic, O., Hamzaoglu, E. (2010). Compositions, antioxidant and antimicrobial activities of *Helichrysum* (Asteraceae) species collected from Turkey. *Food Chem* **119**, 114-122.
- Alves, M.J., Ferreira, I.C.F.R., Martins, A., Pintado, M. (2012). Antimicrobial activity of wild mushrooms extracts against clinical isolates resistant to different antibiotics. *J Apl Microbiol* **113**, 466-475.
- Barros, L., Calhella, R.C., Vaz, J.A., Ferreira, I.C.F.R., Baptista, P., Estevinho, L.M. (2007). Antimicrobial activity and bioactive compounds of Portuguese wild edible mushrooms methanolic extracts. *Eur Food Res Technol* **225**, 151-156.
- Barros, L., Dueñas, M., Ferreira, I.C.F.R., Baptista, P., Santos- Buelga, C. (2009). Phenolic acids determination by HPLC-DAD-ESI/MS in sixteen different Portuguese wild mushrooms species. *Food Chem Toxicol* **47**, 1076-1079.
- Chambers, H.F. (2001). The changing epidemiology of *Staphylococcus aureus*? *Emerg Infect Dis* **7**, 178-182.
- Cosgrove, S.E., Carmeli, Y. (2003). The impact of antimicrobial resistance on health and economic outcomes. *Clin Infect Dis* **36**, 1433-1437.
- Ghosh, S., Nie, A.H., An, J., Huang, Z.W. (2006). Structure-based virtual screening of chemical libraries for drug discovery. *Curr Opin Chem Biol* **10**, 194-202.
- Heleno, S.A., Barros, L., Sousa, M.J., Martins, A., Santos-Buelga, C., Ferreira, I.C.F.R. (2011). Targeted metabolites analysis in wild *Boletus* species. *LWT* **44**, 1343-1348.
- Heleno, S.A., Barros, L., Martins, A., Queiroz, M.J.R.P., Santos-Buelga, C., Ferreira, I.C.F.R. (2012). Fruiting body spores and in vitro produced mycelium of *Ganoderma lucidum* from Northeast Portugal: A comparative study of the

antioxidant potential of phenolic and polysaccharidic extracts. Food Res Int **46**, 135-140.

Jayakumar, T., Thomas, P.A. (2009). Geraldine P. *In-vitro* antioxidant activities of an ethanolic extract of the oyster mushroom, *Pleurotus ostreatus*. Innovat Food Sci Emerg Technol **10**, 228-234.

Khalafi-Nezhad, A., Rad, M.N.S., Mohabatkar, H., Asrari, Z., Hemmateenejad, B. (2005). Design, synthesis, antibacterial and QSAR studies of benzimidazole and imidazole chloroaryloxyalkyl derivatives. Bioorg Med Chem **13**, 1931-1938.

Kim, M.Y., Seguin, P., Ahn, J.K., Kim, J.J., Chun, S.C., Kim, E.H., Seo, S.H., Kang, E.Y., Kim, S.L., Park, Y.J., Ro, H.M., Chung, I.M. (2008). Phenolic compound concentration and antioxidant activities of edible and medicinal mushrooms from Korea. J Agric Food Chem **56**, 7265-7270.

Kuete, V., Ango, P.Y., Fotso, G.W., Kapche, G.D., Dzoyem, J.P., Wouking, A.G., Ngadjui, B.T., Abegaz, B.M. (2011). Antimicrobial activities of the methanol extract and compounds from *Artocarpus communis* (Moraceae). BMC Complement Altern Med **25**, 11-42.

Kuete, V., Nana, F., Ngameni, B., Mbaveng, A.T., Keumedjio, F., Ngadjui, B.T. (2009). Antimicrobial activity of the crude extract, fractions and compounds from stem bark of *Ficus ovata* (Moraceae). J Ethnopharmacol **124**, 556-561.

Lim, D., Strynadka, N.C.J. (2002). Structural basis for the beta lactam resistance of PBP2a from methicillin-resistant *Staphylococcus aureus*. Nat Struct Biol **9**, 870-876.

Lou, Z., Wang, H., Rao, S., Sun, J., Ma, C., Li, J. (2012). *p*-Coumaric acid kills bacteria through dual damage mechanisms. Food Control **25**, 550-554.

- Lowy, F.D. (2003). Antimicrobial resistance: the example of *Staphylococcus aureus*. *J Clin Invest* **111**, 1265-1273.
- Mattila, P., Konko, K., Eurola, M., Pihlava, J.M., Astola, J., Vahteristo, L., Hietaniemi, V., Kumpulainen, J., Valtonen, M., Piironen, V. (2001). Contents of vitamins, mineral elements, and some phenolic compounds in cultivated mushrooms. *J Agric Food Chem* **49**, 2343-2348.
- Montero, C.I., Stock, F., Murray, P.R. (2008). Mechanisms of resistance to daptomycin in *Enterococcus faecium*. *Antimicrob Agent Chemother* **52**, 1167-1170.
- Morris, G.M., Huey, R., Lindstrom, W., Sanner, M.F., Belew, R.K., Goodsell, D.S., Olson, A.J. (2009). AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility. *J Comput Chem* **30**, 2785-2791.
- Oke F., Aslim, B. (2011). Protective effect of two edible mushrooms against oxidative cell damage and their phenolic composition. *Food Chem* **128**, 613-619.
- Orhan, D.D., Özçelik, B., Özgen, S., Ergun, F. (2010). Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiol Res* **165**, 496-500.
- Ozen, T., Darcan, C., Aktop, O., Turkekul, I. (2011). Screening of antioxidant, antimicrobial activities and chemical contents of edible mushrooms wildy grown in the Black Sea region of Turkey. *Comb Chem High Throughput Screen* **14**, 72-84.
- Palacios, I., Lozano, M., Moro, C., D'Arrigo, M., Rostagno, M.A., Martínez, J.A., García-Lafuente, A., Guillamón, E., Villares, A. (2011). Antioxidant properties of phenolic compounds occurring in edible mushrooms. *Food Chem* **128**, 674-678.
- Puttaraju, N.G., Venkateshaiah, S.U., Dharmesh, S.M., Urs, S.M., Somasundaram, R. (2006). Antioxidant activity of indigenous edible mushrooms. *J Agric Food Chem* **54**, 9764-9772.

- Quereshi, S., Pandey, A.K., Sandhu, S.S. (2010). Evaluation of antibacterial activity of different *Ganoderma lucidum* extracts. J Sci Res **3**, 9-13.
- Pedretti, A., Villa, L., Vistoli, G. (2004). VEGA – An open platform to develop chemo-bio-informatics applications, using plug-in architecture and script programming. J Comp-Aid Mol Design **18**, 167-173.
- Reis, F.S., Heleno, S.A., Barros, L., Sousa, M.J., Martins, A., Santos-Buelga, C., Ferreira, I.C.F.R. (2011). Toward the antioxidant and chemical characterization of mycorrhizal mushrooms from Northeast Portugal. J Food Sci **76**, 824-830.
- Ribeiro, B., Rangel, J., Valentão, P., Baptista, P., Seabra, R.M., Andrade, P.B. (2006). Contents of carboxylic acids and two phenolics and antioxidant activity of dried Portuguese wild edible mushrooms. J Agric Food Chem **54**, 8530-8537.
- Ribeiro, B., Valentão, P., Baptista, P., Seabra, R.M., Andrade, P.B. (2007). Phenolic compounds, organic acids profiles and antioxidative properties of beefsteak fungus (*Fistulina hepatica*). Food Chem Toxicol **45**, 1805-1813.
- Rogers, B.A., Sidjabat, H.E., Paterson, D.L. (2011). *Escherichia coli* O25b-ST131: a pandemic, multiresistant, community-associated strain. J Antimicrob Chemother **66**, 1-14.
- Sanner, M.F. (2005). A component-based software environment for visualizing large macromolecular assemblies. Structure **13**, 447-462.
- Surh, Y.J. (2002). Anti-tumor promoting potential of selected spice ingredients with antioxidative and anti-inflammatory activities: a short review. Food Chem Toxicol **40**, 1091-1097.
- Teke, G.N., Kuate, J.-R., Kueté, V., Teponno, R.B., Tapondjou, L.A., Tane, P., Giacinti, G., Vilarem, G. (2011). Bio-guided isolation of potential antimicrobial

and antioxidant agents from the stem bark of *Trilepisium madagascariense*. South Afr J Botany **77**, 319-327.

- Valentão, P., Andrade, P.B., Rangel, J., Ribeiro, B., Silva, B.M., Baptista, P., Seabra, R.M. (2005). Effect of the conservation procedure on the contents of phenolic compounds and organic acids in Chanterelle (*Cantharellus cibarius*) mushroom. J Agric Food Chem **53**, 4925-4931.
- Vaz, J.A., Barros, L., Martins, A., Morais, J.S., Vasconcelos, M.H., Ferreira, I.C.F.R. (2011a). Phenolic profile of seventeen Portuguese wild mushrooms. LWT **44**, 343-346.
- Vaz, J.A., Barros, L., Martins, A., Santos-Buelga, C., Vasconcelos, M.H., Ferreira, I.C.F.R. (2011b). Chemical composition of wild edible mushrooms and antioxidant properties of their water soluble polysaccharidic and ethanolic fractions. Food Chem. **126**, 610-616.
- Werner, G., Gfrörer, S., Fleige, C., Witte, W., Klare, I. (2008). Tigecycline-resistant *Enterococcus faecalis* strain isolated from a German ICU patient. J Antimicrob Chemother **61**, 1182-1183.
- WHO. (2001). Global Prevalence and Incidence of Selected Curable Sexually Transmitted Diseases: Overview and Estimates.
- Wilke, M.S., Lovering, A.L., Strynadka, N.C.J. (2005). β -Lactam antibiotic resistance : a current structural perspective. Curr Op Microbiol **8**, 525-533.
- Yaltirak, T., Aslim, B., Ozturk, S., Alli, H. (2009). Antimicrobial and antioxidant activities of *Russula delica* Fr. Food Chem Toxicol **47**, 2052- 2056.

Table 1. Phenolic compounds identified in wild mushrooms and submitted to antimicrobial activity evaluation.

Phenolic compounds	Mushroom species	Country	References
Phenolic acids: benzoic acid derivatives			
<i>p</i> - Hydroxybenzoic acid	<i>Agaricus arvensis</i> , <i>Agaricus bisporus</i> , <i>Agaricus romagnesii</i> , <i>Agaricus silvicola</i> , <i>Amanita caesarea</i> , <i>Amanita muscaria</i> , <i>Amanita pantherina</i> , <i>Amanita rubescens</i> , <i>Armillaria mellea</i> , <i>Auricularia auricula-judae</i> , <i>Boletus aereus</i> , <i>Boletus edulis</i> , <i>Boletus reticulatus</i> , <i>Boletus rhodoxanthus</i> , <i>Boletus satanas</i> , <i>Calocybe gambosa</i> , <i>Cantharellus cibarius</i> , <i>Chroogomphus fulmineus</i> , <i>Citocybe odora</i> , <i>Coprinus comatus</i> , <i>Cortinarius anomalus</i> , <i>Cortinarius collinitus</i> , <i>Cortinarius violaceus</i> , <i>Craterellus cornocopioides</i> , <i>Fistulina hepática</i> , <i>Ganoderma lucidum</i> , <i>Hygrophorus marzuolus</i> , <i>Hygrophorus olivaceo-albus</i> , <i>Ionotus obliquus</i> , <i>Lactarius deliciosus</i> , <i>Lactarius salmonicolor</i> , <i>Lactarius volemus</i> , <i>Lepista nuda</i> , <i>Lentinus edodes</i> , <i>Lycoperdon molle</i> , <i>Phellinus linteus</i> , <i>Pleurotus eryngii</i> , <i>Pleurotus ostreatus</i> , <i>Ramaria botrytis</i> , <i>Russula cyanoxantha</i> , <i>Sarcodon imbricatus</i> , <i>Sparassis crispa</i> , <i>Suillus granulatus</i> , <i>Suillus collinitus</i> , <i>Suillus mediterraneensis</i> , <i>Tricholoma acerbum</i> , <i>Tricholoma equestre</i> , <i>Tricholoma sulphureum</i>	Finland, Korea, Portugal, Spain, Turkea	Mattila <i>et al.</i> 2000; Ribeiro <i>et al.</i> 2006, 2007; Kim <i>et al.</i> 2008; Barros <i>et al.</i> 2009; Heleno <i>et al.</i> 2011; 2012; Oke <i>et al.</i> 2011; Palacios <i>et al.</i> 2011; Reis <i>et al.</i> 2011; Vaz <i>et al.</i> 2011a, 2011b
Protocatechuic acid	<i>Agaricus bisporus</i> , <i>Agaricus blazei</i> , <i>Amanita caesarea</i> , <i>Amanita pantherina</i> , <i>Auricularia polytricha</i> , <i>Boletus edulis</i> , <i>Boletus rhodoxanthus</i> , <i>Boletus satanas</i> , <i>Cantharellus cibarius</i> , <i>Cantherallus clavatus</i> , <i>Calocybe gambosa</i> , <i>Chroogomphus fulmineus</i> , <i>cortinarius anomalus</i> , <i>Craterellus cornocopioides</i> , <i>Fistulina hepática</i> , <i>Flammulina velutipes</i> , <i>Ganoderma lucidum</i> , <i>Helvella crispa</i> , <i>Hygrophorus agathosmus</i> , <i>Hygrophorus marzuolus</i> , <i>Hydnum repandum</i> , <i>Ionotus obliquus</i> , <i>Lactarius deliciosus</i> , <i>Lactarius sangifluus</i> , <i>Lentinus edodes</i> , <i>Lentinus squarrulosus</i> , <i>Lentinus sajor caju</i> , <i>Lepista nuda</i> , <i>Macrolepiota procera</i> , <i>Morchella anguiticeps</i> , <i>Morchella conica</i> , <i>Mycena haematopus</i> , <i>Pleurotus djamor</i> , <i>Pleurotus eryngii</i> , <i>Phellinus linteus</i> , <i>Pleurotus ostreatus</i> , <i>Pleurotus sajor-caju</i> , <i>Ramaria botrytis</i> , <i>Russula brevepis</i> , <i>Sparassis crispa</i> , <i>Suillus collinitus</i> , <i>Suillus mediterraneensis</i> , <i>Termitomyces heimii</i> , <i>Termitomyces microcarpus</i> , <i>Termitomyces mummiformis</i> , <i>Termitomyces shimperi</i> , <i>Termitomyces tylerance</i>	Finland , India, Korea, Portugal, Spain	Puttaraju <i>et al.</i> 2006; Kim <i>et al.</i> 2008; Barros <i>et al.</i> 2009; Heleno <i>et al.</i> 2011; Oke <i>et al.</i> 2011; Palacios <i>et al.</i> 2011; Reis <i>et al.</i> 2011; Vaz <i>et al.</i> 2011a
Gallic acid	<i>Auricularia auricula-judae</i> , <i>Agaricus bisporus</i> , <i>Agaricus blazei</i> , <i>Auricularia polytricha</i> , <i>Boletus edulis</i> , <i>Calocybe gambosa</i> , <i>Cantharellus cibarius</i> , <i>Cantherallus clavatus</i> ,	India, Korea, Spain Turkey	Puttaraju <i>et al.</i> 2006; Kim <i>et al.</i> 2008;

	<i>Craterellus cornocopioides, Flammulina velutipes, Ganoderma lucidum, Geastrum arinarius, Helvella crispa, Hygrophorus marzuolus, Hydnum repandum, Ionotus obliquus, Lactarius deliciosus, Lactarius sangifluus, Lentinus edodes, Lentinus sajor caju, Lentinus squarrulosus, Macrolepiota procera, Morchella anguiticeps, Morchella conica, Pleurotus djamor, Pleurotus eryngii, Pleurotus ostreatus, Phellinus linteus, Pleurotus sajor-caju, Russula brevipes, Russula delica, Sparassis crispa, Termitomyces heimii, Termitomyces microcarpus, Termitomyces mummiformis, Termitomyces shimperi, Termitomyces tylerance</i>		Yaltirak <i>et al.</i> 2009; Oke <i>et al.</i> 2011; Palacios <i>et al.</i> 2011;
Vanillic acid	<i>Auricularia auricula-judae, Auricularia polytricha, Cantherallus clavatus, Helvella crispa, Hydnum repandum, Lactarius sangifluus, Lentinus squarrulosus, Lentinus sajor caju, Lycoperdon molle, Macrolepiota procera, Morchella conica, Pleurotus sajorcaju, Pleurotus djamor, Pleurotus eryngii, Russula brevipes, Termitomyces heimii, Termitomyces microcarpus, Termitomyces shimperi, Tricholoma acerbum</i>	India, Portugal, Turkey	Puttaraju <i>et al.</i> 2006; Barros <i>et al.</i> 2009; Oke <i>et al.</i> 2011
Syringic acid	<i>Agaricus blazei, Cantherallus clavatus, Auricularia auricula-judae, Hydnum repandum, Lactarius sangifluus, Lentinus sajor caju, Macrolepiota procera, Morchella conica, Morchella anguiticeps, Pleurotus eryngii, Pleurotus djamor, Russula brevipes, Sparassis crispa, Termitomyces mummiformis, Termitomyces tylerance, Termitomyces microcarpus</i>	India, Korea, Turkey	Puttaraju <i>et al.</i> 2006; Kim <i>et al.</i> 2008; Oke <i>et al.</i> 2011
Cinnamic acid and derivatives			
Cinnamic acid	<i>Agaricus arvensis, Agaricus bisporus, Agaricus blazei Agaricus silvicola, Agaricus romagnesii, Amanita caesarea, Amanita muscaria, Amanita pantherina, Armillaria mellea, Boletus aereus, Boletus edulis, Boletus purpureus, Boletus reticulatus, Boletus rhodoxanthus, Boletus satanas, Calocybe gambosa, Cantharellus cibarius, Cantherallus clavatus, Chroogomphus fulmineus, Citocybe odora, Coprinus comatus, cortinarius anomalus, Cortinarius collinitus, Cortinarius violaceus, Fistulina hepática, Ganoderma lucidum, Hygrophorus agathosmus, Hydnum repandum, Hygrophoropsis aurantiaca, Hygrophorus olivaceo-albus, Lactarius aurantiacus, Lactarius quietus, Lactarius salmonicolor, Lactarius sangifluus, Lentinus squarrulosus, Lentinus edodes, Lactarius volemus, Lycoperdon perlatum, Pleurotus eryngii, Macrolepiota procera, Mycena haematopus, Pleurotus sajor-caju, Pleurotus djamor, Sparassis crispa, Russula caerulea, Russula sardonia, Suillus collinitus, Suillus luteus, Suillus mediterraneensis, Termitomyces heimii, Termitomyces mummiformis, Termitomyces shimperi, Tricholoma atrosquamosum, Tricholoma sulphureum, Tricholoma ustale</i>	Finland, India, Korea, Portugal, Turkea	Mattila <i>et al.</i> 2000; Valentão <i>et al.</i> 2005; Puttaraju <i>et al.</i> 2006; Kim <i>et al.</i> 2008; Barros <i>et al.</i> 2009; Heleno <i>et al.</i> 2011, 2012; Oke <i>et al.</i> 2011; Reis <i>et al.</i> 2011; Vaz <i>et al.</i> 2011a, 2011b
<i>p</i> -Coumaric acid	<i>Agaricus arvensis, Agaricus bisporus, Agaricus silvicola, Amanita muscaria, Amanita</i>	India, Korea,	Puttaraju <i>et al.</i> , 2006;

	<i>pantherina, Boletus aereus, Boletus edulis, Calocybe gambosa, Cantharellus cibarius, Chroogomphus fulmineus, Citocybe odora, Coprinus comatus, Cortinarius collinitus, Fistulina hepática, Ganoderma lucidum, Geastrum arinarius, Hygrophorus agathosmus, Hygrophorus marzuolus, Lactarius sangifluus, Lentinus sajor caju, Lepista nuda, Macrolepiota procera, Pleurotus djamor, Pleurotus ostreatus Sparassis crispa, Termitomyces heimii, Tricholoma atosquamosum</i>	Portugal, Spain	Ribeiro <i>et al.</i> 2007; Kim <i>et al.</i> 2008; Barros <i>et al.</i> 2009; Heleno <i>et al.</i> 2011, 2012; Palacios <i>et al.</i> 2011; Reis <i>et al.</i> 2011; Vaz <i>et al.</i> 2011a, 2011b
<i>o</i> -Coumaric acid	<i>Ionotus obliquus</i>	Korea	Kim <i>et al.</i> 2008
Caffeic acid	<i>Auricularia auricula-judae, Agaricus bisporus, Boletus edulis, Calocybe gambosa, Cantharellus cibarius, Cantherallus clavatus, Fistulina hepática, Flammulina velutipes, Hygrophorus marzuolus,, Lactarius sangifluus, Lactarius deliciosus Lentinus sajor caju, Lentinus squarulosus, Morchella anguiticeps, Morchella conica, Macrolepiota procera, Phellinus linteus, Pleurotus djamor, Pleurotus eryngii, Russula brevipes, Russula delica, Sparassis crispa, Termitomyces heimii, Termitomyces microcarpus, Termitomyces shimperi Termitomyces tylerance</i>	India, Korea, Portugal, Spain , Turkey	Valentão <i>et al.</i> 2005; Puttaraju <i>et al.</i> 2006; Ribeiro <i>et al.</i> 2007; Kim <i>et al.</i> 2008; Yaltirak <i>et al.</i> 2009; Oke <i>et al.</i> 2011; Palacios <i>et al.</i> 2011
Ferulic acid	<i>Agaricus bisporus, Cantherallus clavatus, Calocybe gambosa, Cantharellus cibarius, Craterellus cornocopioides , Flammulina velutipes, Ionotus obliquus, Lactarius deliciosus, Lactarius sangifluus, Lentinus squarulosus, Morchella conica, Macrolepiota procera, Pleurotus djamor, Pleurotus ostreatus Pleurotus sajor-caju, Sparassis crispa, Termitomyces heimii, Termitomyces microcarpus, Termitomyces shimperi</i>	India, Korea, Spain	Puttaraju <i>et al.</i> 2006; Kim <i>et al.</i> 2008; Palacios <i>et al.</i> 2011
5- <i>O</i> -cafeoylquinic acid.	<i>Agaricus bisporus, Boletus edulis, Calocybe gambosa, Cantharellus cibarius, Flammulina velutipes, Lactarius deliciosus, Pleurotus ostreatus, Phellinus linteus</i>	Korea, Portugal, Spain	Valentão <i>et al.</i> 2005; Kim <i>et al.</i> 2008; Palacios <i>et al.</i> 2011
Flavonoids			
Quercetin	<i>Agaricus blazei, Flammulina velutipes, Ganoderma lucidum, Ionotus obliquus, Sparassis crispa , Suillus luteus, Suillus granulatus</i>	Korea, Portugal	Ribeiro <i>et al.</i> 2006; Kim <i>et al.</i> 2008
Rutin	<i>Cantharellus cibarius, Pleurotus ostreatus, Russula delica</i>	India, Turkey	Valentão <i>et al.</i> 2005; Jayakumar <i>et al.</i> 2009; Yaltirak <i>et al.</i> 2009
Chrysin	<i>Pleurotus ostreatus</i>	India	Jayakumar <i>et al.</i> 2009
Tannins			
Ellagic acid	<i>Fistulina hepatica</i>	Portugal	Ribeiro <i>et al.</i> 2007

Table 2. MIC values (mg/mL) of wild mushroom phenolic compounds against clinical isolates of Gram negative bacteria.

Phenolic compounds	<i>Escherichia coli</i>	<i>Proteus mirabilis</i>	<i>Morganella morganii</i>	<i>Pasteurella multocida</i>	<i>Neisseria gonorrhoeae</i>
Benzoic acid derivatives					
<i>p</i> -Hydroxybenzoic acid	> 1	> 1	> 1	> 1	> 1
2,4-Dihydroxybenzoic acid	1	> 1	> 1	1	1
Protocatechuic acid	1	> 1	> 1	1	1
Gallic acid	> 1	> 1	> 1	1	1
Vanillic acid	1	1	> 1	1	1
Syringic acid	> 1	> 1	> 1	1	> 1
Cinnamic acid derivatives					
Cinnamic acid	>1	> 1	> 1	> 1	1
<i>p</i> -Coumaric acid	1	> 1	> 1	1	1
<i>o</i> -Coumaric acid	> 1	> 1	> 1	> 1	> 1
Caffeic acid	> 1	> 1	> 1	1	> 1
Ferulic acid	> 1	> 1	> 1	1	1
Chlorogenic acid	> 1	> 1	> 1	> 1	>1
Flavonoids					
Quercetin	> 1	> 1	> 1	1	0,5
Rutin	> 1	> 1	> 1	> 1	> 1
Chrysin	> 1	> 1	> 1	> 1	> 1
Tannins					
Ellagic acid	> 1	> 1	> 1	1	> 1
Reference compounds					
Imipenem	≤ 1	2	4	nt	nt
Ceftriaxon	nt	nt	nt	≤ 1	≤ 1



nt-not tested.

Table 3. MIC values (mg/mL) of the wild mushroom phenolic compounds against clinical isolates of Gram positive bacteria.

Phenolic compounds	MSSA	MRSA	<i>Staphylococcus epidermidis</i>	<i>Enterococcus faecalis</i>	<i>Listeria monocytogenes</i>	<i>Streptococcus agalactiae</i>
Benzoic acid derivatives						
<i>p</i> -Hydroxybenzoic acid	> 1	> 1	> 1	> 1	> 1	> 1
2,4-Dihydroxybenzoic acid	> 1	0.5	> 1	1	> 1	1
Protocatechuic acid	1	1	> 1	> 1	1	1
Gallic acid	> 1	> 1	> 1	> 1	> 1	> 1
Vanillic acid	> 1	0.5	> 1	> 1	1	1
Syringic acid	> 1	0.5	> 1	> 1	0.5	> 1
Cinnamic acid derivatives						
Cinnamic acid	> 1	> 1	> 1	> 1	> 1	0.5
<i>p</i> -Coumaric acid	> 1	1	> 1	> 1	> 1	> 1
<i>o</i> -Coumaric acid	> 1	> 1	> 1	> 1	> 1	1
Caffeic acid	1	1	1	> 1	> 1	> 1
Ferulic acid	1	0.5	1	> 1	> 1	1
Chlorogenic acid	> 1	> 1	> 1	> 1	1	> 1
Flavonoids						
Quercetin	> 1	> 1	> 1	> 1	1	> 1
Rutin	> 1	> 1	> 1	> 1	1	> 1
Chrysin	> 1	> 1	> 1	> 1	> 1	> 1
Tannins						
Ellagic acid	> 1	> 1	> 1	> 1	0.5	> 1
Reference compounds						
Gentamicin	≤ 1	4	≤ 1	nt	nt	nt
Penicillin	nt	nt	nt	2	0.25	≤ 0.03

MSSA (Methicillin-susceptible *Staphylococcus aureus*); MRSA (Methicillin-resistant *Staphylococcus aureus*).

Table 4. Phenolic acids identified in mushrooms submitted to structure activity relationship (SAR) analysis.

		<i>Substitutions</i>				
Benzoic acid derivatives	X	R¹	R²	R³	R⁴	
2,4-Dihydroxybenzoic acid	COOH	OH	H	OH	H	
<i>p</i> -Hydroxybenzoic acid	COOH	H	H	OH	H	
Protocatechuic acid	COOH	H	H	OH	OH	
Gallic acid	COOH	H	OH	OH	OH	
Vanillic acid	COOH	H	OCH ₃	OH	H	
Syringic acid	COOH	H	OCH ₃	OH	OCH ₃	
		<i>Substitutions</i>				
Cinnamic acid derivatives	X	R¹	R²	R³	R⁴	
Cinnamic acid	CHCHCOOH	H	H	H	H	
<i>p</i> -Coumaric acid	CHCHCOOH	H	H	OH	H	
<i>o</i> -Coumaric acid	CHCHCOOH	OH	H	H	H	
Caffeic acid	CHCHCOOH	H	OH	OH	H	
Ferulic acid	CHCHCOOH	H	CH ₃ O	OH	H	




Figure 1. 2,4-Dihydroxybenzoic acid (orange), syringic acid (yellow) and vanillic acid (cyan) docking poses (lines) in PBP2a (carton blue). Hydrogen bonds are present in dash.