

Original Research Article

Antimicrobial and Antioxidant Activities of the Essential Oils of Some Aromatic Medicinal Plants (*Pulicaria inuloides*-Asteraceae and *Ocimum forskolei*-Lamiaceae)

Nabil Qaid M Al-Hajj^{1,2}, Hong Xing Wang^{1*}, Chaoyang Ma¹, Zaixiang Lou¹, Mohanad Bashari¹ and Riyadh Thabit¹

¹State Key Laboratory of Food Science and Technology, School of Food Science and Technology, Jiangnan University, Wuxi 214122, PR China, ²Marine Science and Biological Research Authority, Aden, PO Box 1231, Aden, Yemen

*For correspondence: **Email:** whx1964@126.com; **Tel:** +8651085917795; **Fax:** +8651085876799

Received: 24 March 2014

Revised accepted: 11 July 2014

Abstract

Purpose: To determine the antimicrobial and antioxidant activities of *Pulicaria inuloides* and *Ocimum forskolei* essential oils.

Methods: Steam distillation of the aerial parts of *P. inuloides* and *O. forskolei* was performed using a Clevenger apparatus. Essential oils were analyzed by gas chromatography–mass spectrometry. Total phenolic content and antioxidant activities were determined by 2,2-diphenyl-1,1-picrylhydrazyl (DPPH) and β -carotene bleaching assays. Disc diffusion and microtiter broth microdilution assays were employed to determine antimicrobial activity.

Results: The chemical compounds in *P. inuloides* essential oil include 2-cyclohexen-1-one, 2-methyl-5-(1-methyl) (55.1 %) and benzene, methyl- (20.6 %). The major components identified in *O. forskolei* essential oil included bicyclo [3.1.1] hept-2-ene,2, (22.4 %) and naphthalene 1,2,3,4,4a,5,6, (19.3 %). *P. inuloides* showed a higher total phenol content than *O. forskolei* (144 ± 5.32 vs. 54.6 ± 30 mg GAE/g extract), higher antioxidant activity (92.92 ± 0.10 % vs. 26.76 ± 0.11 % scavenging activity; IC₅₀, 4.5 ± 0.05 vs. 73.03 ± 0.05) and β -carotene bleaching (90.77 ± 0.21 % vs. 41.03 ± 6.35 % inhibition). *P. inuloides* essential oil inhibited all tested microorganisms except *Salmonella typhimurium* and *Shigella dysenteriae* with a minimum inhibitory concentration (MIC) of 3.0 μ g/mL against *Escherichia coli*. *O. forskolei* essential oil inhibited only *Candida albicans*.

Conclusion: *P. inuloides* essential oil possesses significant antioxidant and antimicrobial activities.

Keywords: Essential oil; Phenolic content; Antioxidant; Antimicrobial activity, *Pulicaria inuloides*, *Ocimum forskolei*, *Salmonella typhimurium*, *Shigella dysenteriae*

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

The genus *Pulicaria*, which belongs to the family Asteraceae (tribe Inuleae), consists of more than 77 species found throughout the world. Members of this genus contain various bioactive compounds such as monoterpenes, flavonoids, acetylenes, isocomene, and sesquiterpene lactones [1]. The leaves of *Pulicaria inuloides* are

used to flavor foods and to make an herbal tea [2]. Biological actions reported for *Pulicaria* species include the antibacterial and antispasmodic activities of *P. undulate*, *P. odora*, and *P. dysenterica* [3]. In addition, members of this genus have been traditionally used to repel insects, to reduce influenza and common cold symptoms, and to treat back pain, intestinal disorders, and inflammation [4].

The genus *Ocimum*, which belongs to the Lamiaceae family, is the most well-studied genus of all the aromatic herbs and contains up to 160 species [5]. *Ocimum*, also called basil, grows to a height of about one meter and has an obtusely quadrangular stem and opposite leaves, which are used in folk medicine as a tonic and vermifuge [6]. The plant oils have been used to alleviate mental fatigue, colds, spasms, rhinitis, wasp stings, and snakebites [6]. The medicinal and aromatic properties of basil are derived from the essential oil, which accumulates primarily in the leaves and flowers. The fresh and dried herb is used as an aromatic spice, and its bioactive components harbor antimicrobial, antimutagenic, and fungistatic activities [7].

Because the chemical composition of the essential oils of *P. inuloides* and *O. forskolei* have not yet been reported, we used gas chromatography-mass spectrometry (GC-MS) to determine their total phenolic content, as well as their antioxidant, antibacterial, and antifungal activities.

EXPERIMENTAL

Plant collection and identification

The aerial parts of *P. inuloides* (flowers) and *O. forskolei* (leaves) were collected in August 2013 during the flowering stage in the Sana'a and Dammar areas of Yemen. The plant specimens were air-dried and taxonomically identified by Prof. Abdellah Amine (College of Agriculture, Sana'a University). A voucher specimen (SANB N. 196, DAMO N. 179) of the plant material was deposited at the Department of Biology in the College of Agriculture at Sana'a University.

Extraction of essential oil

The fresh aerial parts of *P. inuloides* and *O. forskolei* (3 kg) were ground in a blender, and essential oils were obtained by hydrodistillation at 100 °C using a Clevenger-type apparatus for 4 h with 3 L distilled water. The extracted oils were dried over sodium sulfate (Na_2SO_4) and stored at 4 °C until use.

Analysis of essential oil

The essential oil components were identified by GC-MS [8] using a Varian 1200L system (electronic impact mode, ionization potential 70 eV, ion source temperature 200 °C, mass range 35–500 Da) fitted with a relatively nonpolar capillary column (DB-5, 30 m × 0.25 mm, 0.25 μm film thickness). The injection port and

interface were held at 220 °C and 260 °C, respectively. The column temperature was increased from 50 °C to 220 °C (15 °C/min) and then the temperature was maintained at 220 °C for 25 min. Helium was the carrier gas. Most of the compounds were identified using two different analytical methods: (a) KI, Kováts Indexes in reference to n-alkanes (C8- C32) [9]; and (b) mass spectra (authentic chemicals and Wiley spectral library collection). Identification was considered tentative when it was based on mass spectral data only.

Determination of total phenolic content

Total phenolic content was estimated as milligrams of gallic acid equivalent (mg GAE) per gram plant extract, as previously described [10]. In brief, a 100 μL aliquot of dissolved extract was transferred to a volumetric flask containing 46 mL distilled H_2O , and 1 mL Folin-Ciocalteu reagent was added. After 3 min, 3 mL Na_2CO_3 (2 %) was added. The sample was incubated at 25 °C for 2 h, after which absorbance was measured at 760 nm. Gallic acid (0.2–1 mg/mL; Sigma) was used as the standard for the calibration curve, and total phenolic content was expressed as mg GAE/g extract ($Y = 0.0792x + 0.00302$; $R^2 = 0.9739$) [10].

Determination of radical scavenging activity

The antioxidant activities of the essential oils of *P. inuloides* and *O. forskolei* were determined by the 2, 2-diphenyl-1,1-picrylhydrazyl (DPPH) free radical scavenging assay, as previously described [11]. Each concentration of essential oil was mixed individually into 1 mL methanol solution containing 0.1 mM DPPH radicals. The reaction mixture was shaken thoroughly, incubated in the dark for 30 min at room temperature, and measured at an absorbance of 517 nm (2100, Unico, Shanghai, China). The free radical scavenging activity was calculated as follows using equation 1:

$$\text{DPPH (\%)} = \{(A_n - A_m)/A_m\}100 \dots \dots \dots (1)$$

where A_n is absorbance of the control (without essential oil), and A_m is absorbance of the sample.

β-carotene bleaching assay (BCB)

Antioxidant activities of the essential oils were determined as previously described [11], with some modifications. The β-carotene (0.1 mg) was added to a boiling flask together with linoleic acid (20 mg) and Tween 40 dissolved in chloroform. After evaporating the chloroform

under vacuum at 50 °C using a rotary evaporator, 50 mL oxygenated distilled water was added, and the mixture emulsified for 1 min. Thereafter, 5 mg of each essential oil was added separately to 4.8 mL of the emulsion. Absorbance at 470 nm was measured using a spectrophotometer before (t = 0 h) and after a 2-h incubation at 50 °C (t = 2 h). All measurements were performed in duplicate, and antioxidant activity, assessed as percent inhibition of β -carotene bleaching, was calculated using equation 2:

$$\% \text{ inhibition} = \{AA(2h) - AC(2h) / (AC(0h) - AC(2h))\} 100 \dots (2)$$

where AA(2h) is absorbance of the sample at t = 2 h, AC(2h) is absorbance of the control at t = 2 h, and AC(0h) is absorbance of the control at t = 0 h, and AC(2h) is absorbance of the control at t = 2 h.

Assessment of antimicrobial activity

Microorganisms

Staphylococcus aureus 6538, *Streptococcus pneumoniae* ATCC 25922, *Bacillus subtilis* ATCC 6633, *Escherichia coli* ATCC 25922, *Shigella dysenteriae* 51302, *Salmonella typhimurium* 50013 and *Candida albicans* ATCC 10231 were purchased from the China General Microbiological Culture Collection Center (Beijing, China).

Disc diffusion assay

Antimicrobial activity of the essential oils was determined by using the disc diffusion assay [12] with some modifications. Tryptic soy agar was inoculated with the microorganism (10^4 colony-forming units/mL). A 6-mm paper filter disc impregnated with 20 μ L essential oil diluted in dimethyl sulfoxide was placed on the agar, and the oil was allowed to diffuse into the medium for 30 min at room temperature. The plates were then incubated at 37 °C for 24 h (bacteria) or at 32 °C for 72 h (yeast). The zone of inhibition was recorded as the mean \pm standard deviation (SD) of triplicate experiments. Ampicillin (10 μ g) and gentamicin (10 μ g) were used as reference antibiotics for bacteria, and nystatin (100 μ g) was used as the reference antifungal agent for *C. albicans*.

Determination of minimal inhibitory concentration (MIC)

The MIC of the essential oil was determined using the microtiter broth microdilution assay described by Amsterdam [13]. The essential oils

were diluted to 50 μ L/mL and subjected to a serial dilution in a microtiter plate containing tryptic soy broth (for bacteria) or Sabouraud dextrose broth (for yeast). The bacterial and yeast strains were suspended in the liquid culture medium at a final concentration of 10^4 colony-forming units/mL. After incubation at 37 °C for 24 h (bacteria) or at 32 °C for 72 h (yeast), optical density was measured at 520 nm using a spectrophotometer. MIC was defined as the lowest concentration of the essential oil at which the microorganisms did not exhibit visible growth.

Statistical analysis

Experiments were conducted at least in triplicate. Groups were compared by analysis of variance, and differences between mean values were evaluated by Fisher LSD test; $p < 0.05$ was considered significant. Statistical analyses were carried out using SPSS version 19.0 (SPSS, Chicago, IL, USA).

RESULTS

Chemical composition of the essential oils

The compositions of the essential oils obtained by hydrodistillation of the flowers and leaves of *P. inuloides* and *O. forskolei* were determined by GC-MS. Ten components accounted for 87.468 % of the *P. inuloides* essential oil, with 2-cyclohexen-1-one, 2-methyl-5-(1-methyl (55.1 %), benzene, methyl- (20.6 %), and Z-Citral (2.915 %) being the major constituents (Table 1). Ten components were identified in the essential oil of *O. Forskolei*, with Bicyclo [3.1.1] hept-2-ene, 2, (22.4 %), naphthalene, 1,2,3,4,4a,5,6, (19.3 %), and phytol (6.7 %) being the major constituents (Table 2).

Total phenolic content and antioxidant activity

The total phenolic content of *P. inuloides* essential oil (144 ± 5.32 mg GAE/100 g extract) was considerably higher than that of *O. forskolei* essential oil (54.6 ± 3.0 mg GAE/100 g extract) (Table 3). In addition, the antioxidant activity of *P. inuloides* essential oil (90.77 ± 0.21 % inhibition) was stronger than that of *O. forskolei* essential oil (41.03 ± 6.35 % inhibition), as assessed by the BCB assay. The half maximal inhibitory concentration (IC_{50}) was graphically obtained to determine concentration needed to inhibit 50 % of free DPPH radicals. The IC_{50} values of *P. inuloides* and *O. forskolei* were 4.95 ± 0.05 and 73.54 ± 0.05 μ g/mL, respectively.

Table 1: Main components of the essential oil of *Pulicaria inuloides*

RT (min)	Area (%)	Compound	Content (%)
5.785	3.68	Benzene, methyl- (CAS)	20.6
18.966	9.83	2-Cyclohexen-1-one, 2-methyl-5-(1-methyl)	55.1
19.081	1.95	Zingiberene	1.094
19.182	5.20	Z-Citral	2.915
19.355	2.09	alpha-Farnesene	1.169
19.684	3.09	Cyclohexene, 3-(1,5-dimethyl-4-hexenyl)-	1.734
19.961	2.74	7-Oxabicyclo[4.1.0]heptane, 1-methyl- (C	1.538
20.609	3.15	Geranyl propionate	1.511
22.097	1.76	(-)-Caryophyllene oxide	0.985
26.679	1.47	Pentacosane	0.822
			Total = 87.468

RT = retention time

Table 2: Main components of the essential oil of *Ocimum forskolei*

RT (min)	Area (%)	Compound	Content (%)
17.532	2.83	Bicyclo[3.1.1] hept-2-ene, 2,	22.369
17.554	1.12	Bicyclo[3.1.1] hept-2-ene, 2,	8.877
17.694	4.442	trans-Caryophyllene	3.496
18.498	3.74	alpha-Humulene	2.956
19.189	2.45	Naphthalene, 1,2,3,4,4a,5,6,	19.327
19.28	7.86	Azulene, 1,2,3,5,6,7,8,8a-oc	6.212
19.341	3.83	Naphthalene, 1,2,3,4,4a,5,6,	3.03
19.367	5.04	alpha-Selinene	3.983
19.701	7.37	Naphthalene, 1,2,3,4,4a,5,6,	5.827
28.098	8.48	Phytol	6.704
			Total = 82.781

RT = retention time

Table 3: Essential oil antioxidant activity and total phenolic content of *Pulicaria inuloides* and *Ocimum forskolei*

Botanical name	Total phenolic content (mg GAE/g DW)	Inhibition of DPPH (%)	IC ₅₀ (µg/ml)	β-carotene bleaching (% inhibition)
<i>P. inuloides</i>	144±5.32 ^a	92.92±0.10 ^a	4.95±0.05 ^a	90.77±0.21 ^a
<i>O. forskolei</i>	54.6±3.0 ^b	26.76±0.11 ^b	73.54±0.05 ^b	41.03±6.35 ^b

Different letters indicate significant differences between the two essential oils ($p < 0.05$). DW - dry weight. Results are expressed as mean ± SD

Antimicrobial activity

The *in vitro* bacteriostatic activity of *P. inuloides* essential oil was greater than that of *O. forskolei* essential oil, as assessed by the disc diffusion assay (Table 4) and by the microtiter broth microdilution assay (Table 5).

DISCUSSION

To the best of our knowledge, the chemical compositions of *P. inuloides* and *O. forskolei* essential oils have not previously been reported. However, essential oils of plants belonging to the same genera (*P. dysenterica* and *O. basilicum*) show qualitative and quantitative differences, which can be attributed to different growth conditions, genetic factors, geographical variations, and analytical procedures [3].

Medicinal plants have been extensively studied for their antioxidant and free radical scavenging activity in the last few decades. Among these natural antioxidants, phenolic compounds have been shown to quench oxygen-derived free radicals by donating a hydrogen atom or an electron to the free radical [11]. The total phenolic content of the essential oil of *O. forskolei* (54.6 ± 30 mg GAE/g) determined in this study is in agreement with that previously reported for *O. basilicum* (55 ± 122 mg GAE/g) [13]. Phenolic compounds may contribute directly to antioxidant activity.

Compared with *O. forskolei* essential oils, *P. inuloides* essential oils strongly inhibited β-carotene bleaching and had a higher total phenolic content in our study, suggesting greater

Table 4: Antimicrobial activity of *Pulicaria inuloides* and *Ocimum forskolei* essential oils

Test microorganism	<i>Pulicaria inuloides</i> (zone of inhibition, mm)	<i>Ocimum forskolei</i> (zone of inhibition, mm)	Standard antibiotic ^b (zone of inhibition, mm)
Gram-positive bacteria			Ampicillin
<i>Staphylococcus aureus</i>	16.6 ± 1.16	ND	25 ± 0.9
<i>Streptococcus pneumoniae</i>	18.2 ± 2.24	ND	25.2 ± 2.18
<i>Bacillus subtilis</i>	18.0 ± 0.4	ND	20.0 ± 1.1
Gram-negative bacteria			Gentamicin
<i>Shigella dysenteriae</i>	ND	ND	17.7 ± 1.22
<i>Salmonella typhimurium</i>	ND	ND	26 ± 1.1
<i>Escherichia coli</i>	14.7 ± 1.37	ND	25 ± 0.9
Yeast			Nystatin
<i>Candida albicans</i>	26 ± 2.5	10.1 ± 1.85	26.4 ± 1.20

^bStandard antibiotics used as positive control; ND = Not detectable, essential oil has no antimicrobial activity against this microorganism

Table 5: Minimal inhibitory concentration (MIC) of the essential oils of *Pulicaria inuloides* and *Ocimum forskolei* against bacteria and yeast

Microorganism	MIC (µg/mL)	
	<i>Pulicaria inuloides</i>	<i>Ocimum forskolei</i>
<i>Staphylococcus aureus</i>	6.25	ND
<i>Streptococcus pneumoniae</i>	3.7	ND
<i>Bacillus subtilis</i>	3.0	ND
<i>Shigella dysenteriae</i>	ND	ND
<i>Salmonella typhimurium</i>	ND	ND
<i>Escherichia coli</i>	3.0	ND
<i>Candida albicans</i>	3.12	35.3

ND = Not detectable, i.e., essential oil has no antimicrobial activity against this microorganism

antioxidant capacity [11]. In addition, *P. inuloides* essential oils exhibited higher radical scavenging activities than *O. forskolei* (IC₅₀, 4.95 ± 0.05 µg/mL vs. 73.54 ± 0.05 µg/mL). These results are consistent with a previous report on the radical scavenging activities of *Pulicaria* species [14]; however, there are few studies reporting β-carotene assay results for essential oils [14]. Differences in values may be due to differences in geographical areas, genus, reproductive stage, climate, season of harvest, and extraction methods [15].

Similar to the essential oils described in this report, the essential oils of *Thymus* species contain the aromatic monoterpenes caryophyllene oxide and thymol, and their biological activities are often attributed to these compounds [2]. In this study, *P. inuloides* essential oils demonstrated higher antibacterial activities against all bacteria tested except *Salmonella typhimurium* and *Shigella dysenteriae*. Furthermore, the Gram-positive bacteria *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Bacillus subtilis* were more

sensitive to this essential oil than the Gram-negative bacteria *Escherichia coli*. Similarly, a previous study reported that the essential oils of *Pulicaria astephanocarpa* showed high antimicrobial activity against *Candida albicans* and all tested bacteria except *Pseudomonas aeruginosa* [13]. Our results suggest that *P. inuloides* essential oils may be useful in the treatment of infectious diseases caused by *S. aureus*, *B. subtilis*, *S. pneumoniae*, *E. coli*, and *C. albicans*.

The essential oils of *O. forskolei* demonstrated no antibacterial activity against all tested bacteria (Table 4), which is consistent with a previous report [16]. That study reported that extracts derived from another *Ocimum* species, *O. gratissimum*, showed no activity against 11 tested bacterial strains, including *S. aureus* (four strains), *E. coli* (two strains), *Pseudomonas aeruginosa* (one strain), *Proteus* spp. (three strains), and *Shigella* (one strain). Similarly, the essential oils of *O. basilicum* were ineffective against gram-positive and gram-negative bacteria tested in another study [17].

In the present study, we found that *P. inuloides* essential oils demonstrated activity against *C. albicans*. This activity may be related to its high level of oxygenated monoterpenes, especially its major constituent, 2-cyclohexen-1-one, 2-methyl-5-(1-methyl) (55.1 %). In addition, the presence of two phenolic isomers of thymyl acetate (0.143 and 0.354 % of the total content) may synergistically contribute to the antifungal activity of *Pulicaria* essential oils. Phenolic compounds in essential oils (e.g., carvacrol, thymol, and eugenol) are thought to be primarily responsible for their biological properties [18]. The antimicrobial activity of these phenolic compounds can be attributed to the presence of

an aromatic nucleus and a phenolic OH group, which can form hydrogen bonds with disulfide groups at the active sites of target fungal enzymes, resulting in their deactivation [18].

MIC values indicate that *P. inuloides* essential oils have a greater inhibitory action against Gram-positive bacteria than Gram-negative bacteria. A previous study showed that the antimicrobial effects of spices and herbs against *C. albicans*, *E. coli*, and other Gram-negative bacteria were due to their complex chemical composition, which included compounds such as thymol, carvacrol, methyl eugenol, linalool, α -pinene, 1, 8-cineole, and camphor [19]. This discrepancy in antibacterial potential may be caused by variations in chemical composition, which may be influenced by the distillation and extraction techniques as well as geographical origin.

In our study, *O. forskolei* essential oils were completely inactive against all tested microorganisms other than *C. albicans*. This result may be due to evaporation of the oil during the boiling process. The antifungal property of the oil was likely due to the eugenol [18]. Nanasombat and Lohasupthawee [19] described the antibacterial action of the essential oils derived from another *Ocimum* species, *O. basilicum*, against clinically determined multidrug-resistant isolates of *Staphylococcus*, *Enterococcus*, and *Pseudomonas*, reporting MIC values of 0.0030 % to 0.0007 % (v/v). In another study, *O. gratissimum* essential oils demonstrated significant antimicrobial action, whereas essential oils of *O. basilicum* were found to have less activity, and *O. sanctum* oils were completely inactive against microorganisms [19]. Our findings were consistent with the results of these studies analyzing *Ocimum* species. After reviewing the literature, we found no studies evaluating the antimicrobial activity of essential oils and/or plant extracts of *P. inuloides* or *O. forskolei*. Therefore, we believe our study is the first to report on this topic, thus making a significant contribution to the scientific field.

CONCLUSION

The essential oils of *P. inuloides* exerted strong antimicrobial actions against gram-positive bacteria and *C. albicans*, whereas *O. forskolei* essential oils inhibited the growth of *C. albicans* only. The strong antioxidant activity of *P. inuloides* essential oil suggests its potential as a food additive and candidate for clinical

development as a new broad-spectrum bioactive compound.

ACKNOWLEDGEMENT

This work was supported by the National Science and Technology Program of China (no. 2014BAD33B05). The authors express their thanks to the students of the Nutrition Laboratory of School of Food Science and Technology at Jiangnan University, China for their technical assistance.

REFERENCES

1. Hanbali FEL, Akssira M, Ezoubeiri A, Gadhi CEA, Mellouki F, Benherra f. Chemical composition and antibacterial activity of essential oil of *Pulicaria odora* L. *J Ethnopharmacol.* 2005; (99): 399-401.
2. Ross SA, El SKA, El SMA, Hamann MT, Abdel HOB, Ahmed AMM. Phytochemical analysis of *Geigeriaalata* and *Francoeuria crispa* essential oils. *Planta Med* 1997; 63: 479–482.
3. Elkamali HH, Ahmed AH, Mohammed AS, Yahia AAM, Eltayebi H, Ali AA. Antibacterial properties of essential oil from *Nigella citrates* leaves and *Pulicaria Undulata* Aerial Parts. *Fitoterapia* 1998; 69: 77-78.
4. Stavri M, Mathew KT, Gordon A, Shnyder SD, Falconer RA, Gibbons S. Guaianolide sesquiterpenes from *Pulicaria acrispa* (Forssk.) Oliv. *Phytochemistry* 2008; 69: 1915-1918.
5. Sajjadi SE. Analysis of the essential oils of two cultivated basil (*Ocimum basilicum* L.) from Iran. *Daru J Pharm Sci.* 2006;14 (3) 128-130.
6. Baytop T. Treatment with plants in Turkey. *Istanbul Univ.*1984; Publ 3255.
7. Dambolena JS, Zunino MP, López AG, Rubinstein HR, Zygadlo JA, Mwangi JW, Kariuki ST. Essential oils composition of *Ocimum basilicum* L. and *Ocimum gratissimum* L. from Kenya and their inhibitory effects on growth and fumonis in production by *Fusarium verticillioides* *Innovative food science and emerging Technologies* 2010;11: 410-414.
8. Kumar KA, Patel J, Choudhary R K. Chemical composition and antimicrobial activity of the essential oil of *Desmostachya bipinnata* linn. *Int J Phytomed* 2010; 2(4) 436-439.
9. Ralchenko, YU, Kramida AE, Reader J. NIST atomic spectra database (version 4.0). National Institute of Standards and Technology, Gaithersburg, MD, 2010.
10. Avanmardi J, Stushnoff C, Locke E, Vivanco JM. Antioxidant activity and total phenolic content of Iranian *Ocimum* accessions. *Food Chem* 2003; 83(4): 547-550.
11. Pratt, DE. Natural antioxidants of soybeans and other oil seeds. *Autoxidation in food and biological systems.* Springer US 1980; 283-293.

12. Zhu SYY, Yang HD, Yu Y, Ying Y, Zou GL. Chemical composition and antimicrobial activity of the essential oils of *Chrysanthemum indicum*. *J Ethnopharmacol.* 2005; 96: 151–158.
13. Amsterdam D. Susceptibility testing of antimicrobials in liquid media. *Antibiot Lab Med* 1996; 4: 52-111.
14. Alzoreky N, Kazuhiko N. Antioxidant activity of some edible Yemeni plants evaluated by ferryl myoglobin BTS± Assay. *Food Sci Technol Res* 2001; 7: 141–144.
15. Wu X, Gu L, Holden J, Haytowitz BD, Gebhardt, ES, Beecher G. Development of a database for total antioxidant capacity in foods: a preliminary study. *J Agric Food Chem.* 2004; 17: 407-422.
16. Chah KF, Eze CA, Emuelosi CE, Esimone, CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. *J Ethnopharmacol.* 2006; 104(1): 164-167.
17. Algabr MN, Mekkiou R, Ameddah S, Menad A, Boumaza O, Seghiri R, Benayache F. Antioxidant activities from the aerial parts of *Pulicaria jaubertii*. *American Eurasian Network for Scientific Information* 2010; 4(1): 63-70.
18. Ultee A, Bennik MHJ, Moezelaar R. The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and Environmental Microbiology* 2002; 68: 1561-1568.
19. Nanasombat S, Lohasupthawee P. Antibacterial activity of crude ethanolic extracts and essential oils of spices against *Salmonellae* and other enterobacteria. *KMITL Sci Tech* 2005; 5: 527-538.
20. Burt S. Essential oils: their antibacterial properties and potential applications in foods. A review. *Int J Food Microbiol* 2004; 94: 223-253.