

Antimicrobial and physico-chemical effects of essential oils on fermented milk during preservation

Tovidé Sènou Noël¹, Adeoti Kifouli^{1*}, Yèhouéno Boniface², Dahouénon-Ahoussi Edwige², Baba-Moussa Farid¹, Toukourou Fatiou¹

¹Laboratoire de Microbiologie et des Technologies Alimentaires, Faculté des Sciences et Techniques/Université d'Abomey-Calavi, Cotonou, Benin

²Laboratoire d'Étude et de Recherche en Chimie Appliquée, EPAC/ UAC.

*: corresponding author ; zoulade@yahoo.fr

Original submitted in on 3rd February 2016. Published online at www.m.elewa.org on 31st March 2016
<http://dx.doi.org/10.4314/jab.v99i1.12>

ABSTRACT

Objectives: Milk and milk products are known to be the good media for development of many microorganisms. Some essential oils are known to have antimicrobial activities against bacteria, mould and fungi. With the aim of contributing to the preservation of the fermented dairy products, the present study explored the use of essential oils for biopreservation in food pathogen control. The objectives were to study the effect of two essential oils extracted from local plants *Xylopiya aethiopica* (Dunal) A.Rich. and *Pimenta racemosa* (Mill.) J.W. Moore on the physico-chemical, microbiological and organoleptic characteristics of the fermented milk.

Methodology and Results: At the first level, essential oils were extracted from the leaves of plants and the minimum inhibitory concentration (MIC) of each one was detected using the serial dilution method and Muller Hinton Broth medium. The references strains used were *Staphylococcus aureus* ATCC 25923 and *Escherichia coli* ATCC 25922. Based on the MICs determined, essential oils were added to fermented milk and determination of its physicochemical characteristic, evolution of the microbial flora (mesophilic flora, lactic flora, coliforms, *Staphylococcus aureus*) were done during two weeks of preservation. The results showed that *Xylopiya aethiopica* had a lowest minimum inhibitory concentration. After fifteen days of preservation, a decrease of mesophilic flora and lactic bacteria was observed and a disappearance was noticed for coliforms, thermotolerant coliforms and *Staphylococcus aureus* during the same time. These results prove the efficiency of those essential oils in fermented milk preservation. Moreover, microorganisms' inhibition was more pronounced with *Xylopiya aethiopica* essential oil than the *Pimenta racemosa* one.

Conclusion and applications of findings: The study results showed a possible way of using essential oils as an alternative to chemical additives used for preservation of food products. As there are natural products, there could take an important part in the food safety policies.

Key words: Essential oils, biopreservation, fermented milk

INTRODUCTION

In Benin, agriculture contributes to 37% on gross domestic product (GDP) among which livestock stands to 6% of this part (Alkoiret *et al.*, 2011). The milk produced from this livestock is quantitatively low

and varies from 1L to 2.5 L per day according to the breeds (Dossou *et al.*, 2006). Indeed, milk is a high nutritious food, which contains lactose, protein, minerals, vitamins, fats and casein (Mahaut *et al.*, 2000; Vierling, 2003). In most of developing countries, local populations in rural areas consume milk fresh. In urban areas it is consumed both fresh and as other milk-derived products in which the most known is yogurt. In Benin, milk mostly produced by pastoral community, plays an important role in their daily food nutrition and contributes up to 50% of household (Ogodja *et al.*, 1991) income through milk selling or by-products. Besides the low production of milk in Benin, the major problem is related to its preservation. To avoid the problem of preservation, milk is transformed into traditional cheese so called "waragashi" in local language. The milk pH close to neutral and water content around 90% to 92% makes it quickly perishable and susceptible to microorganisms attacks such as bacteria, moulds and yeasts. During milk storage, chemical reactions occur due to the presence of microorganisms, which induce deterioration of nutritional and organoleptic qualities and reducing the duration of preservation. Spoilage and contamination may occur in the milk chain because of poor hygiene, long periods of transportation and lack of appropriate storage facilities. Despite its nutritional value, milk also serves as a good medium for the growth of many microorganisms. To avoid the short-term conservation of milk and its by-products, it is important to inhibit or slow down these mechanisms

MATERIAL AND METHODS

Collection of plant material and extraction of essential oils: The plant material is constituted by the fresh leaves of *Xylopiya aethiopicica* and of *Pimenta racemosa* harvested respectively in Djassin Tokpa (Porto-Novo) and in Malawi (Adjarra) in the department of Ouémé. These Fresh leaves were then used for essential oils extraction. The essential oils were extracted from fresh leaves by the method of hydro-distillation in a distiller type Clevenger. Obtained essential oils were dried over anhydrous sodium sulphate and preserved at 4° C until their use.

Determination of Minimal Inhibitory Concentration: The method of microdilution using microplates of 96 wells

which occurred during milk deterioration. In the fermented products, such as yogurt, maintain vitality of lactic flora is very important. Therefore, different methods and use of natural or chemical antioxidant and preservatives are applied to reach this purpose (Abdenouri *et al.*, 2008) in order to extend the duration of preservation. Also, the limitation imposed by the food industry and the regulatory authorities about the use of certain synthetic food additives led to a renewed interest in the search for alternatives, as natural antimicrobial compounds, in particular those of vegetable origin (Hammer *et al.*, 1999). Essential oils appeared for many years to be an alternative in foods preservation process. Essential oil as well as compound by-products possess important activities of which the most studied is antimicrobial activity. Indeed, the antioxidant, bactericidal and fungicidal activities of essential oil extracted from several plants were studied by several researchers (Rasooli & Mirmostafa, 2002; Tepe *et al.*, 2005; Giordani *et al.*, 2005 ; Sacchetti *et al.*, 2005 ; Valero & Frances , 2006). The study aims to explore the use of essential oils in biopreservation of food pathogen control. The objectives are to study the effect of two essential oils extracted from local plants *Xylopiya aethiopicica* and *Pimenta racemosa* on the physico-chemical, microbiological and organoleptic characteristics of the fermented milk. *Pimenta racemosa* and *Xylopiya aethiopicica* are plants that have antimicrobial properties. Both essential oils were reported to have antimicrobial activities on *Staphylococcus aureus* and *Escherichia coli* strains.

according to Bajpai *et al.* (2008a) and reported by Yèhouénou *et al.* (2010ab) was used. Two (2) ml of Mueller Hinton Broth (MHB) added by phenol red at the concentration of 0.02 g/l were used to dilute 40 µl of essential oil. Two (2) drops of Tween 80 were added to facilitate the mixture. The mixture of tested essential oil and Muller-Hinton Broth (MHB) medium was used as a negative control and the positive control was carried out with a mixture of tested microorganism. After making serial dilutions well by well, the microplate was covered and incubated at 37°C +/- 1°C for twenty four (24) hours. **Collection of biological material:** The fresh milk was bought early the morning, just after milking in stock farms

of the city of Abomey-Calavi. Samples were carried cold and stored in the refrigerator before the manipulations.

Figure 1 show the diagram used for production of fermented milk in which essential oil was added.




Figure 1: Production of fermented milk containing essential oil

Physico-chemical analysis: Physico-chemical analysis concerned pH determination and peroxide index. For pH determination, the pH of fresh milk, fermented milk and fermented milk added with some essential oil was raised (found) every day during fifteen (15) days of preservation. Peroxide index was measured according to the standard method NT. 118.22 (1996). This value was estimated every three (03) days during the fifteen days of preservation.

Microbiological analysis: Milk samples were analyzed for their microbiological quality. Microbial flora of fresh milk, fermented milk without oil and fermented milk added

with each type of essential oil were traced during fifteen (15) days storage at ambient temperature. Finally, the four types of samples were used for microbiological analysis. Fresh milk was used as a control. Enumeration of total mesophile flora using Plate Count Agar (PCA) with plates incubation at 30°C had been done for 72 hours. Coliforms were determined using violet red bile lactose agar (VRBL) after 24 hours ± 2 at 37°C and 44°C. The enumeration of total coliform was determined according to the NF standard method V08-050-199. The Staphylococci are determined according to the standard (ISO 15213-2003) by using the Baird-Parker (BP)

medium after incubation at 37°C for 24 hours. Enumeration of the lactic flora was made on the Man-Rogosa-Sharpe agar (M.R.S) medium with incubation at 30°C during 72 hours. Preparation of decimal dilution of each sample has been done using the maximum recovery diluents.

Data analysis: Data obtained were statistically analyzed using Statistical Package for Social Scientists (SPSS)

RESULTS AND DISCUSSION

Minimal inhibitory concentration (MIC) of the two essential oils: The MICs of *Xylopi*a *aethi*opica essential oil (EO) for the two microorganisms are 0.23 mg/ml and 0.46 mg/ml, while those for *Pimenta racemosa* are 1.10 mg/ml and 0.55 mg/ml respectively for *Staphylococcus aureus* and *Escherichia coli* are (Table 1). MIC values

version 17.0. Data were expressed as mean ± standard deviations of three replicate determinations (Pillai & Nair, 2013). The data obtained for each sample were evaluated for significant differences in their means with analysis of variance (ANOVA). Critical difference at p = 0.05 was estimated.

obtained for *Xylopi*a *aethi*opica near those obtained by Yèhouénou *et al.* (2010a) for the same microorganisms that were respectively 0.27 mg/ml and 0.54 mg/ml. the study noted that the tested strains were more sensitive to *Xylopi*a *aethi*opica essential oil than to *Pimenta racemosa*.

Table 1: Determination of minimal inhibitory concentration of tested oils

Tested strains	MIC (mg/ml)	
	<i>Pimenta racemosa</i>	<i>Xylopi</i> a <i>aethi</i> opica
<i>Staphylococcus aureus</i> ATCC 25923	1.10	0.23
<i>Escherichia coli</i> ATCC 25922	0.55	0.46

Effect of essential oils on milk flora during storage

Total mesophilic flora: The aerobic mesophilic flora of fresh milk determined before fermentation was 2,5.10⁵ CFU (Table 2). This value was higher than the standard recommend for fresh milk, which is 5.10⁴. During fermentation after 5, 10 and 15 days of storage at 4°C, the value determined were respectively 10⁴, 3.10³ and 2.10³ CFU/ml. This flora did not exceed the critical threshold of 5.10⁴ CFU / ml during the fifteen days of preservation. During the period of storage, mesophile flora of fermented milk added with essential oils also

decreased significantly (Table 3). The total number of flora estimated were lower than those obtained with fresh milk and fresh milk fermented without oil. Analysis of variance showed a significant difference between fermented milk added with essential oils and which fermented without essential oils. Indeed, fermented milk added with essential oil of *Pimenta racemosa* and *Xylopi*a *aethi*opica had respectively 5.10² CFU/ml and 3,6.10² CFU/ml after fifteen days of preservation. This result showed the low level of microbial proliferation in which added with *Xylopi*a *aethi*opica oil than the other.

Table 2: Microbial flora of fresh milk

	Fresh milk	Standard
Mesophile flora	2,5.10 ⁵	5.10 ⁴
Lactic flora	7.10 ³	-
Coliforms	2,8.10 ²	10 ²
Thermotolerant coliforms	1,6.10 ³	1
<i>Staphylococcus aureus</i>	1,9.10 ³	10 ²

Coliforms: As total aerobic mesophile flora was determined, the coliforms had been enumerated for fresh milk fermented, fermented milk added with essential oils. Table 3 showed the results obtained for each sample during the same period of preservation. The level of coliform decreased with the time of preservation in the

three types of sample. Moreover, this decrease is more important in the fermented milk added with essential oils especially with *Pimenta racemosa* compared to the other. Indeed, enumerated coliform was below the critical threshold, which is the 10² CFU/ml. This result showed that essential oils of these species contributed to the

reduction of coliforms. In the same line, variation of the rate of the thermo-tolerant coliforms (TTC) in fermented milk preserved with or without essential oil was also determined. The number of unit forming colonies of the TTC decreased, in a fast way during the fifteen days of preservation samples. However, the number of TTC was lower than that required by the standard, which is 1 CFU / ml.

Lactic flora: The lactic flora (FL) of the milk fermented without essential oil after 5, 10 and 15 days of preservation in 4°C is respectively 10^7 , 4.10^6 and 10^6

CFU / ml. During the period of storage, the FL knew a significant decrease for the two (2) lots of fermented milk preserved with essential oils. The fermented milk preserved with *Pimenta racemosa* oil presented for 5, 10 and 15 days respectively a rate of 28.10^5 , 8.10^5 and 14.10^4 CFU / ml and a respective rate of 12.10^5 , 44.10^4 and 9.10^4 CFU / ml for that preserved with *Xylopiya aethiopic*. This flora exceeded the critical threshold of 10^2 CFU / ml during the fifteen days of preservation for three lots (Table 3).

Table 3: Enumeration of specific flora of fermented milk with and without essential oils

	Fermented milk without oils CFU/ml	Fermented milk + <i>Pimenta racemosa</i> CFU/ml	Fermented milk + <i>Xylopiya aethiopic</i> CFU/ml	Standard CFU/ml References
Mesophile flora				
5 days	10^4	5.10^3	10^3	5.10^4
10 days	3.10^3	10^3	9.10^2	
15 days	2.10^3	5.10^2	$3,6 10^2$	
Lactic flora				
5 days	10^7	28.10^5	12.10^5	
10 days	4.10^6	8.10^5	44.10^4	
15 days	10^6	14.10^4	9.10^4	
Coliforms				
5 days	12	< 1	5	10^2
10 days	6	< 1	< 1	
15 days	5	< 1	< 1	
Thermotolerant coliforms				
5 days	< 1	< 1	< 1	1
10 days	< 1	< 1	< 1	
15 days	< 1	< 1	< 1	
<i>Staphylococcus aureus</i>				
5 days	10	7	< 1	
10 days	9	2	< 1	10^2
15 days	9	< 1	< 1	

Enumeration of *Staphylococcus sp.* Flora: The variations of the rate of staphylococci in fermented milk preserved with or without essential oils are enumerated. The number of colonies (CFU/ml) of staphylococci decreases in a fast way in the samples during the fifteen days of preservation. However, the number of staphylococci is lower than that required by the standard, which is 10^2 CFU/ml for the three lots.

Influence of essential oils on Physico-chemical characteristics of milk

Evolution of pH during storage: The values of pH had been determined during the same period of preservation for all samples. pH varied from 4.73 to 4.96, 4.03 to 4.16 and 4 to 4.2 respectively for milk fermented without additive, fermented milk preserved with *Pimenta racemosa* and fermented milk preserved with *Xylopiya aethiopic* (Figure 2). The lowest values of pH were observed for the milk preserved with essential oils although all pH values were acids. Nevertheless, analysis of variance showed no difference between values obtained for the completely fermented milk.




Figure 2: Evolution of pH in fermented milk during preservation

Evolution of peroxide index: The peroxide index showed an increasing during the first ten (10) days of preservation for all fermented milk apart from addition or not of essential oils. After this, a decrease was observed during the last days resulting in the production on secondary metabolites derived from peroxide decomposition (Figure 3). The consequence is the production of smell rancid and bad taste. Using ANOVA test and application of Duncan' test, a significant difference was observed between fermented milk and

which added with essential oils. On the other hand, there was a significant difference between fermented milk preserved with essential oils. *Pimenta racemosa*' oil has antioxidant activity and contributed to reduce the degree of oxidation of the fat through inhibition of free radical formation (Eymard, 2003). Similar results had been reported for *Xylopiya aethiopic* with a lower effect than those observed with *Pimenta racemosa* (Asekun & Adeniyi, 2004; Noudogbessi et al., 2011).




Figure 3: Evolution of peroxide index of fermented milk fat content during preservation

DISCUSSION

The low pH observed for all fermented milk could be explained on the one hand by fermentation due by lactic bacteria and on the second hand by the action of essential oils. Indeed, essential oils had an acid property, which is in relation with the peroxide index. These results were proved by the pH value obtained with fermented milk added by essential oils were low pH were obtained. Many studies showed the biological activities of essential oils (Deans & Ritchie, 1987 ; Bajpai *et al.*, 2008a,b;) and their multiple applications inside foods as antimicrobials (Lambert *et al.*, 2001; Smith *et al.*, 2001; Kékè *et al.*, 2009; Yèhouéno *et al.*, 2010b). It was observed a decrease of microbial flora such as mesophile germs, coliforms, lactic bacteria and *Staphylococci* flora species in all fermented milk added with essential oils when compared with the control. The highest value was found in fermented milk without oil, while the lowest was detected in fermented milk obtained with essential oils. Results from the Analysis of Variance (ANOVA) suggested that there was a significant difference ($P < 0.05$) in bacterial loads between the two groups of samples. The use of essential oil of *Pimenta racemosa* and *Xylopi aethiopica* in the preservation of the fermented milk allowed us to notice that the efficiency of these oils on the bacterial flora is reduced when it is mixed with milk. The reduction of the microbial activity of oils on the bacteria of the fermented milk could be explained by the reduction of the contact area with the bacteria, which is doubtless function of the rate of miscibility of oil with milk. However, we noticed a real contribution of these oils to the reduction of the bacterial flora of the milk when we compare samples tests with the control without oil. In the fermented milk preserved without oil, we also noticed a decrease of the bacterial flora during the time of preservation except *Staphylococcus*' flora that remained intact. The persistence of this microbial flora could be explained by an insufficiency of acidity of milk resulted from fermentation and essential oil action to inhibit the

CONCLUSION

Results of this study indicated that there is a great possibility to use essential oils as biopreservatives in fermented milk. Indeed, both of essential oils tested showed inhibitory activity against microorganisms. It occurs that, essential oils could be used to control microbial contaminants of foods and could be able to

growth of the *Staphylococcus* which have the minimal pH of growth around 4.2. Besides, for fermented milk preserved with essential oil, the efficiency of microbial flora decrease resulted from combination of acidity and essential oils. Indeed, for the milk added by essential oil of *Pimenta Racemosa*, it needed at least ten days to observe the disappearance of *Staphylococcus* when only five days were required for *Xylopi aethiopica*. These results showed a significant contribution of essential oils in inhibition of microbial flora. Indeed, according to Guylène *et al.* (1998), oil extracted from *Pimenta racemosa* were bacteriostatic and fungistatic. The same results had been reported for *Xylopi aethiopica* (Karioti *et al.*, 2004). Moreover, antimicrobial activity of *Xylopi aethiopica* is more pronounced than that of *Pimenta racemosa* when compared the sample preserved with essential oils. This result could be explained by the fact that essential oil of *Xylopi aethiopica* contains more bacteriostatic substances than *Pimenta racemosa*' oil. Microbial activity of this oil was proved by several authors. Okigbo & Nmeko (2005) showed the antifungal and antimicrobial power of essential oil extracted from *Xylopi aethiopica* leaves. And its bactericidal activity was demonstrated by many authors (Alitonou, 2006; Noudogbessi *et al.*, 2008; Yèhouéno *et al.*, 2010a). Indeed, with their great potential to control pathogen in food materials, essential oils had been recognized as alternative in foods preservation by several authors (Burt, 2004; Dahouenon-Ahoussi *et al.*, 2010). According to Kouninki *et al.* (2007), action of *Xylopi aethiopica* could be explained by its chemical composition. On the other hand, Samaddar *et al.* (2015) reported that the use of plant extracts and essential oils in consumer goods is expected to increase in the future because volatile oils can be considered as a natural alternative to synthetic food preservatives and could be used to enhance food safety and shelf life.

become useful tools for applications in foods preservation systems. The challenges for future are to determine the conditions, which avoid the loss of lactic bacteria; to preserve organoleptic characteristic of fermented milk after incorporation of essential oil; and preserve the effects of essential oils on food borne pathogens.

REFERENCES

- Abdenouri N, Idliman A, Kouhila M, 2008. Étude hygroscopique du lait en poudre. Revue des énergies renouvelables, Alger. 35-44.
- Alitonou G, 2006. Huiles essentielles extraites de plantes aromatiques acclimatées au Bénin : étude chimique, évaluation biologique et applications potentielles. Thèse de Doctorat en cotutelle, Université d'Abomey-Calavi et Université Montpellier II. 283 pp
- Alkoiret IT, Yari HM, Gbangboche AB, Lokossou R, 2011. Reproductive performance and milk production of girolando cows in the ranch of kpinnou, south-west of Benin Republic. Journal of Animal and Veterinary Advances 10 (19): 2588-2592.
- Asekun OT. and Adeniyi BA, 2004. Antimicrobial and cytotoxic activities of the fruit essential oil of *Xylopiya aethiopicica* from Nigeria. Fitoterapia 75 (3-4): 368–370.
- Bajpai KV, Dung NT, Kwon JO, Kang SC, 2008a. Analysis and the potential application of essential oil and leaf extracts of *Silene Americana* L. to control food spoilage and food-borne pathogens, Journal of Food Technology 227: 1613-1620.
- Bajpai VK, Rahman A, Kang SC, 2008b. Chemical composition and inhibitory parameters of essential oil and extracts of *Nandina domestica*. Thumb to control food-borne pathogenic and spoilage bacteria, International Journal of Food Microbiology 125: 117-122.
- Burt S, 2004. Essential oils: their antibacterial properties and potential applications in foods. A review : International Journal of Food Microbiology 94: 223-253.
- Dahouénon-Ahoussi E, Sessou P, Wotto DV, Yèhouénon B, Kinsoudé E, Kpatinvoh B, Soumanou M, Souhounhloúé D, 2010. Mise au point d'une technologie de production d'une boisson locale Africa drink a base d'huiles essentielles de menthes verte et poivrée. Bulletin d'information de la Société Ouest Africaine de Chimie 7 : 39-53.
- Deans SG. and Ritchie G, 1987. Antibacterial properties of plant essential oils. International Journal of Food Microbiology 5: 165-180.
- Dossou J, Hounzangbe-Adote S, Soule H, Chabi BI, 2006. Production and processing of fresh milk into Fulani cheese in Benin. Good Practice Guide. GRET-CAD/FSA. 33 pp.
- Eymard S, 2003. Mise en évidence et suivi de l'oxydation des lipides au cours de la conservation et la transformation du Chinchard : choix des procédés. Thèse de doctorat Biochimie. École polytechnique de l'université de Nantes France.
- Giordani R, Regli P, Kaloustian J, Mikail C, Abou L, Portugal H, 2005. Antifungal effect of various essential oils against *Candida albicans*. Potential of antifungal action of amphotericin B by essential oil from *Thymus vulgaris*. In P. R.18 (12): 990-995.
- Guylène SA, Jacqueline A, Paul B, Joëlle L, 1998. Antibacterial and antifungal activities of the essential oils of *Pimenta racemosa* var. *racemosa* P. Miller (J.W. Moore) (*Myrtaceae*). Journal of Essential Oil Research 10: 161-164.
- Hammer KA, Carson CF, Ridley CV, 1999. Antimicrobial activity of essential oils and other plants extract. Journal of applied microbiology 86: 985-990.
- Karioti A, Hadjipavlou-Litina D, Mensah MLK, Theophilus C, Skaltsa FH, 2004. Composition and Antioxidant Activity of the Essential Oils of *Xylopiya aethiopicica* (Dun) A. Rich. (*Annonaceae*) Leaves, Stem Bark, Root Bark, and Fresh and Dried Fruits, Growing in Ghana. J. Agric. Food Chem. 52 (26): 8094–8098
- Kèkè M, Yèhouénon B, de Souza C, Sohounhloúé D, 2009. Evaluation of hygiene and nutritional quality of peulh cheese treated by *Sorghum vulgaris* (L) and *Pimenta racemosa* (Miller) extracts. Scientific Study & Research – Chemistry & Chemical Engineering, Biotechnology, Food Industry 10 (1): 29-46;
- Kouninki H, Haubruge E, Noudjou FE, Lognay G, Malaisse F, Ngassoum MB, Goudou A, Mapongmetsem PM, Ngamo LS, Hance T, 2007. Toxicity of some terpenoids of essential oils of *Xylopiya aethiopicica* from Cameroon against *Sitophilus zeamais* Motsch. Journal of Applied Entomology 131: 269-274.
- Lambert RJW, Skandamis PN, Coote PJ, Nychas JEG, 2001. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. Journal of Applied Microbiology 91: 453-462.
- Mahaut M, Jeantet R, Brulé G, Schuck P, 2000. Les produits laitiers industriels. Tec. et Doc. Lavoisier, Paris. pp 52-55.

- Noudogbessi JP, Dansou K, Sohounhloué DCK, 2008. Composition Chimique et Propriétés Physico-chimiques des Huiles Essentielles de *Pimenta racemosa* (Miller) et de *Chromolaena odorata* (L. Robinson) Acclimatées au Bénin. Journal de la Société Ouest-Africaine de Chimie : 026 : 11 - 19.
- Noudogbessi JP, Natta AK, Avlessi F, Sohounhloué DCK, Figueredo G, Chalchat JC, 2011. Chemical composition of the essential oils extracted from two *annonaceae* required in Beninese pharmacopeia. Australian Journal of Basic and Applied Science 5 (2) : 34-40.
- Ogodja JO, Hounsou-Vè G, Dehoux JP, 1991. Part I : Rôle et activité de la femme Peulh dans son ménage dans le sud Borgou au Bénin. Part II : Commercialisation du lait et des produits laitiers dans le sud Borgou en République du Bénin. Projet de Développement Pastoral Intégré dans le Borgou. Parakou Bénin. 20 pp.
- Okigbo RN. and Nmeka IA, 2005. Control of yam tuber rot with leaf extracts of *Xylopi aethiopica* and *Zingiber officinale*. African Journal of Biotechnology 4 (8): 804-807.
- Pillai LS. and Nair BR, 2013. Proximate composition, Mineral elements and Antinutritional factors in *Cleome viscosa* L. and *Cleome burmanni* W. & A. (*Cleomaceae*). Int. J. Pharm. Pharm. Sci. 5(1): 384-387.
- Samaddar M, Ram C, Sen M, 2015. Assessment of storage stability of essential oil enriched flavoured milk. Indian J. Dairy Sci 68(4): 357-363.
- Rasooli I. and Mirmostafa SA, 2002. Antibacterial properties of *Thymus pubescens* and *Thymus serpyllum* essential oils. Fitoterapia 73(3): 244-250.
- Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R, 2005. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. Food Chemistry 91 (4): 621-632.
- Smith-Palmer A, Stewart J, Fyfe L, 2001. The potential application of plant essential oils as natural food preservatives in soft cheese. Food Microbiology 18: 463-470.
- Tepe B, Sokmen M, Akpulat HA, Daferera D, Polissiou M, Sokmen A, 2005. Antioxydative activity of the essential oils of *thymus sipyleus* and *thymus sipyleus* subsp. *Sipyleus* var. *rosulans*. Journal of Food Engineering 66 (4): 447-454.
- Valero M. and Frances E, 2006. Synergistic bactericidal effect of carvacrol, cinnamaldehyde or thymol and refrigeration to inhibit *Bacillus cereus* in carrot broth. Food microbiology 23 (1) : 68-73.
- Vierling E, 2003. Science des aliments. Aliments et boissons, Filières et produits. Doin éditeurs. Bordeaux cedex. p 25- 36.
- Yèhouénou B, Noudogbessi JP, Sessou P, Avlessi F, Sohounhoué D, 2010a. Étude chimique et activités antimicrobiennes d'extraits volatils des feuilles et fruits de *Xylopi aethiopica* (Dunal) A. Rich. contre les pathogènes des denrées alimentaires. Journal de la Société Ouest-Africaine de Chimie 029 : 19-27.
- Yèhouénou B, Wotto V, Bankolé H, Sessou P, Noudogbessi JP, Sohounhloué D, 2010b. Chemical study and antimicrobial activities of volatile extracts from fresh leaves of *Crassocephalum rubens* (juss & jack) s. moore against food-borne pathogens. Scientific Study & Research : Chemistry & Chemical Engineering, Biotechnology, Food Industry Vol 11 (3): 343-351.