

01 May 2020

Antimicrobial Resistance Needs to be Combated at Primary Levels of Prevention by Nurses

Sarah Oerther

Daniel B. Oerther

Missouri University of Science and Technology, oertherd@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/civarc_enveng_facwork



Part of the [Civil and Environmental Engineering Commons](#)

Recommended Citation

S. Oerther and D. B. Oerther, "Antimicrobial Resistance Needs to be Combated at Primary Levels of Prevention by Nurses," *Nursing Open*, vol. 7, no. 3, pp. 678-679, Wiley-Blackwell Publishing Ltd, May 2020. The definitive version is available at <https://doi.org/10.1002/nop2.445>

This Editorial is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Civil, Architectural and Environmental Engineering Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.



Antimicrobial resistance needs to be combated at primary levels of prevention by nurses

Antimicrobial resistance (AMR) is a threat to public health (O'Neil, 2014; World Health Organization, 2018). AMR occurs when antimicrobial therapies become ineffective at killing infections caused by bacteria, fungi, parasites and viruses, and this leads to drug-resistant infections (World Health Organization, 2018). Drug-resistant infections are impacting all populations (Centers for Disease Control, 2018; Logan & Bonomo, 2016; World Health Organization, 2018; Zerr et al., 2014). Inappropriate use of antimicrobial therapies is believed to be a significant contributing factor to the emergence and spread of AMR (O'Neil, 2014; World Health Organization, 2018).

Drug-resistance infections contribute to longer hospital stays, higher medical costs and increased incidence of morbidity (World Health Organization, 2018; Zetts, Stoesz, Smith, & Hyun, 2018). Though there is advocacy for the prevention of AMR, interventions have enjoyed limited success because of primary care providers' fears and patient's perceived demand for pills to treat all illnesses (Martínez-González et al., 2017; Wood et al., 2012). For instance, primary care providers want to meet the patients' need to feel satisfied with their care (Martínez-González et al., 2017; Wood et al., 2012; Zetts et al., 2018). Primary care providers also may fear that patients may become ill without the use of antimicrobial therapies (Martínez-González et al., 2017; Wood et al., 2012; Zetts et al., 2018). AMR needs to be combated with primary levels of prevention by nurses.

1 | BACKGROUND

In the United States (US), 23,000 deaths and two million illnesses were caused by antibiotic-resistant bacterial infections (Centers for Disease Control, 2018). If AMR continues to rise, it is estimated that by 2050 up to 10 million people will die annually from drug-resistant infections (O'Neil, 2014). The indirect economic losses attributed to AMR could result in a global loss of 60–100 trillion USD between now and 2050 (O'Neil, 2014).

One of the main risk factors for AMR in the United States is inappropriate prescription of antimicrobial therapies by primary care providers (Zetts et al., 2018). Researchers have found that approximately 154 million primary care visits in the United States result in an antibiotic prescription and approximately 30 per cent, or 47 million, of these prescriptions are not necessary (PEW, 2016).

Researchers have shown that patients' demand for antimicrobial therapies varies depending on cultural background and geographic location (Larson et al., 2009; Morgan & Hart, 2009; Zetts et al., 2018). Researchers have found Hispanic cultures request antibiotics more frequently (Larson et al., 2009). This could be because antibiotics are available over-the-counter in some Latin American countries (Larson et al., 2009). Geographically, southern states in the United States average 920 prescriptions per 1,000 people compared to western states in the United States where residents receive an average of 632 antibiotic prescriptions per 1,000 people (Zetts et al., 2018). However, people in extremely rural communities in the United States tend to wait longer before seeing a provider for treatment for upper respiratory symptoms (Morgan & Hart, 2009). Researchers have found that people in rural communities are more accepting of a provider's recommendations to treat an infection with supportive care rather than a prescription for antibiotics (Morgan & Hart, 2009).

2 | PRIMARY PREVENTION

Interventions at the primary level are needed long term to prevent the spread of AMR (Centers for Disease Control, 2018; Logan & Bonomo, 2016; World Health Organization, 2018; Zerr et al., 2014). Nursing interventions at the primary level need to focus on overcoming existing barriers to care, including gaps in knowledge about best practices, patients' expectations for antibiotics, the need to see patients in a short amount of time and primary care providers' concerns regarding patient satisfaction when they do not prescribe an antimicrobial therapy (Sanchez, Fleming-Dutra, Roberts, & Hicks, 2016).

Appropriate nursing interventions at the primary level are also needed to overcome cultural and geographic barriers (Hicks et al., 2015). For example, people living in rural communities are less likely to listen to AMR prevention campaigns (Hicks et al., 2015). Researchers believe this barrier exists because prevention campaigns do not incorporate rural cultural considerations (Hicks et al., 2015). Patients living in rural areas need educational programmes designed to address rural social norms (Hicks et al., 2015).

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Nursing Open* published by John Wiley & Sons Ltd.

3 | CONCLUSION

Researchers have demonstrated that the primary level of prevention of AMR has a clear impact and is cost-effective (Ball, Dains, Flynn, Solomon, & Stewart, 2019; Hersh, Jackson, & Hicks, 2013). Implementing primary prevention needs further development by nurses to decrease the number of people receiving unnecessary antimicrobial therapies from primary care providers (Ball et al., 2019; Hersh et al., 2013). Nurses also need to create appropriate health education programmes about AMR to improve patients' knowledge and practices towards safe use antimicrobial therapies, especially in rural areas and among Latino populations in the United States (Larson et al., 2009; Zetts et al., 2018).

ACKNOWLEDGEMENTS

None.

AUTHOR CONTRIBUTIONS

All authors contributed equally to this article.

Sarah Oerther¹ 

Daniel B. Oerther² 

¹School of Nursing, Saint Louis University, Saint Louis, MO, USA

²King's College London, London, UK

Correspondence

Sarah Oerther, School of Nursing, Saint Louis University,
Saint Louis, MO, USA.

Email: sarah.oerther@gmail.com

ORCID

Sarah Oerther  <https://orcid.org/0000-0002-9990-6739>

Daniel B. Oerther  <https://orcid.org/0000-0002-6724-3205>

REFERENCES

- Ball, J. W., Dains, J. E., Flynn, J. A., Solomon, B. S., & Stewart, R. W. (2019). *Seidel's Guide to Physical Examination* (9th ed). St. Louis: Mosby-Year Book.
- Centers for Disease Control (2018). *Antibiotic / Antimicrobial Resistance*. Retrieved from <https://www.cdc.gov/drugresistance/index.html>
- Hersh, A. L., Jackson, M. A., & Hicks, L. A. (2013). Principles of judicious antibiotic prescribing for upper respiratory tract infections in pediatrics. *Pediatrics*, 132(6), 1146–1154. <https://doi.org/10.1542/peds.2013-3260>
- Hicks, L. A., Bartoces, M. G., Roberts, R. M., Suda, K. J., Hunkler, R. J., Taylor, T. H., & Schrag, S. J. (2015). US outpatient antibiotic prescribing variation according to geography, patient population, and provider specialty in 2011. *Clinical Infectious Diseases*, 60(9), 1308–1316. <https://doi.org/10.1093/cid/civ076>
- Larson, E., Ferng, Y.-H., Wong, J., Alvarez-Cid, M., Barrett, A., Gonzalez, M. J., ... Morse, S. S. (2009). Knowledge and Misconceptions Regarding Upper Respiratory Infections and Influenza Among Urban Hispanic Households: Need for Targeted Messaging. *Journal of Immigrant and Minority Health*, 11(2), 71–82. <https://doi.org/10.1007/s10903-008-9154-2>
- Logan, L. K., & Bonomo, R. A. (2016). Metallo- β -Lactamase (MBL)-producing Enterobacteriaceae in United States children. *Open Forum Infectious Disease*, 3(2), ofw090. <https://doi.org/10.1093/ofid/ofw090>
- Martínez-González, N. A., Coenen, S., Plate, A., Colliers, A., Rosemann, T., Senn, O., & Neuner-Jehle, S. (2017). The impact of interventions to improve the quality of prescribing and use of antibiotics in primary care patients with respiratory tract infections: A systematic review protocol. *British Medical Journal Open*, 7(6), <https://doi.org/10.1136/bmjopen-2017-016253>
- Morgan, K., & Hart, A. M. (2009). Families in rural settings: Values regarding acute respiratory infections. *Families, Systems, & Health*, 27(1), 85–97. <https://doi.org/10.1037/a0014754>
- O'Neil, J. (2014). *Antimicrobial Resistance: Tackling A Crisis for the Health*. Retrieved from https://amr-review.org/sites/default/files/AMRReviewPaper-Tacklingacrisisforthehealthandwealthofnations_1.pdf
- PEW (2016). *Antibiotic Use in Outpatient Settings*. Retrieved from <https://www.pewtrusts.org/en/research-and-analysis/reports/2016/05/antibiotic-use-in-outpatient-settings>
- Sanchez, G. V., Fleming-Dutra, K. E., Roberts, R. M., & Hicks, L. A. (2016). *Core Elements of Outpatient Antibiotic Stewardship*. Retrieved from https://www.cdc.gov/antibiotic-use/community/pdfs/16_268900-A_CoreElementsOutpatient_508.pdf
- Wood, F., Phillips, C., Brookes-Howell, L., Hood, K., Verheij, T., Coenen, S., ... Butler, C. C. (2012). Primary care clinicians perceptions of antibiotic resistance: A multi-country qualitative interview study. *Journal of Antimicrobial Chemotherapy*, 68(1), 237–243. <https://doi.org/10.1093/jac/dks338>
- World Health Organization (2018). *Antimicrobial Resistance*. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance>
- Zerr, D. M., Qin, X., Oron, A. P., Adler, A. L., Wolter, D. J., Berry, J. E., ... Weissman, S. J. (2014). Pediatric infection and intestinal carriage due to extended-spectrum-cephalosporin-resistant enterobacteriaceae. *Antimicrobial Agents and Chemotherapy*, 58(7), 3997–4004. <https://doi.org/10.1128/aac.02558-14>
- Zetts, R. M., Stoesz, A., Smith, B. A., & Hyun, D. Y. (2018). Outpatient antibiotic use and the need for increased antibiotic stewardship efforts. Retrieved from <https://pediatrics.aappublications.org/content/141/6/e20174124>. <https://doi.org/10.1542/peds.2017-4124>