

Research Article

Antimicrobial Resistance of *Salmonella enterica* Serovars Typhi and Paratyphi Isolates from a General Hospital in Karawaci, Tangerang, Indonesia: A Five-Year Review

Nata Pratama Hardjo Lugito¹ and Cucunawangsih²

¹Department of Internal Medicine, Faculty of Medicine, Pelita Harapan University, Tangerang, Banten, Indonesia

²Department of Microbiology, Faculty of Medicine, Pelita Harapan University, Tangerang, Banten, Indonesia

Correspondence should be addressed to Nata Pratama Hardjo Lugito; nata.lugito@uph.edu

Received 12 January 2017; Revised 26 March 2017; Accepted 27 March 2017; Published 6 April 2017

Academic Editor: Giuseppe Comi

Copyright © 2017 Nata Pratama Hardjo Lugito and Cucunawangsih. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Typhoid and paratyphoid fever known as enteric fever pose important global public health problem, with 21.6 million cases and approximately 250,000 deaths annually. It is a prevalent disease in Indonesia, but data on the antimicrobial resistance pattern is limited. This study aim was to provide data on the antimicrobial resistance pattern of *S. Typhi* and *S. Paratyphi* bloodstream isolates in a general hospital in Karawaci, Tangerang, Banten, Indonesia, during the period of January 2011 to December 2015. Susceptibility against antimicrobials was detected according to the guidelines of the Clinical and Laboratory Standards Institute (CLSI). Out of a total of 168 isolates 55.4% were *S. Typhi* and 44.6% *S. Paratyphi* A. Most of the isolates, 92.9%, were from children aged 6–18 years and adult population. There was low resistance of *S. Typhi* against ampicillin, trimethoprim-sulfamethoxazole, ceftriaxone, ciprofloxacin, and levofloxacin, similar to previous studies in Indonesia. In the 2011–2015 period, resistance rates against most antimicrobials and MDR rate of *S. Typhi* and *S. Paratyphi* were low, emphasizing that there is a distinct epidemiological dynamic of the enteric fever in Indonesia.

1. Introduction

Enteric fever, both typhoid and paratyphoid fever are important global public health problems, with 21.6 million cases and approximately 250,000 deaths annually [1, 2]. Typhoid and paratyphoid fever are caused by *Salmonella enterica* serovars Typhi (*Salmonella* ser. Typhi) and Paratyphi, respectively. It is estimated that more than 90% of typhoid fever cases were reported in South and Southeast Asian countries [1].

In many parts of the world, the changing modes of presentation and the development of multidrug resistance have made enteric fever increasingly difficult to diagnose and treat. Multidrug resistant (MDR) *S. Typhi* is defined as *S. Typhi* strains which are resistant to all the three first-line recommended drugs for treatment, *that is*, chloramphenicol, ampicillin, and trimethoprim-sulfamethoxazole [3]. There had been reports on resistance of *Salmonella* species against

antimicrobial used, beginning with the report of chloramphenicol resistance in 1972 to the report of multidrug resistant strains [4, 5]. The first MDR strains were discovered in Southeast Asia in the late 1980s and have since spread throughout the region [6]. The use of first-line antimicrobial for the treatment of enteric fever such as ampicillin, chloramphenicol, and trimethoprim-sulfamethoxazole had been recommended to be replaced with quinolone or 3rd generation cephalosporins [7]. In Indonesia, MDR *S. Typhi* was reported in Surabaya, in 2009 [8]. A multicentric study conducted across five Asian countries endemic for typhoid including Indonesia reported that the prevalence of MDR *S. Typhi* ranged from 7% to 65% [9].

Enteric fever is a prevalent disease in Indonesia, but data on the antimicrobial resistance pattern is limited. The aim of this study was to provide data on the antimicrobial resistance pattern of *S. Typhi* and *S. Paratyphi* isolates from patients

admitted to a general hospital in Karawaci, Tangerang, Banten, Indonesia during the period of 2011–2015.

2. Material and Method

2.1. Study Setting. This retrospective descriptive study was conducted in Siloam Hospital in Karawaci, Tangerang, Banten, Indonesia. The hospital is a private teaching hospital with 274 beds, affiliated to Faculty of Medicine, Pelita Harapan University. The samples in this study were all bloodstream isolates of *S. Typhi* and *S. Paratyphi* collected at the Microbiology Laboratory of the hospital.

2.2. Specimen, Culture, and Identification. Venous blood samples were collected from patients admitted to hospital from January 2011 to December 2015. Blood was inoculated into enriched soybean-casein digest broth with resins in BACTEC aerobic plus/F (Becton-Dickinson, New Jersey, USA) bottles. For patients with body weight less than 12.8 kg of weight, BACTEC Peds Plus/F bottles were used. When there was bacterial growth indicated by the BACTEC machine, blood culture bottles were subcultured onto a MacConkey agar plate.

2.3. Susceptibility Testing. Susceptibility of *Salmonella* isolates against antimicrobials was detected using agar dilution method according to the guidelines of the Clinical and Laboratory Standards Institute (CLSI). Minimal inhibitory concentrations (MICs) for each antibiotic were determined by VITEK 2 compact (bioMérieux, Marcy-l'Etoile, France). Susceptibility interpretations were based on CLSI M100-S23 clinical breakpoints [10]. The antibiotics used included carbapenems (imipenem and meropenem), penicillins (ampicillin and amoxicillin clavulanate), cephalosporins (ceftriaxone and ceftazidime), fluoroquinolones (ciprofloxacin and levofloxacin), trimethoprim-sulfamethoxazole, and aminoglycosides (amikacin and gentamicin). Results were included in the analysis only when the corresponding QC isolates tested were within the acceptable range according to CLSI guidelines. Chloramphenicol susceptibility for *S. Typhi* and *S. Paratyphi* was not tested according to the policy of the Indonesian Microbiology Association.

2.4. Statistics. The data were analyzed statistically using Statistical Package for Social Sciences (SPSS, version 24.0.0.0). Categorical data were presented as frequency (percentage), while numerical data as mean \pm SD or median (range).

2.5. Ethical Considerations. The study protocol was approved by the Ethical Committee of Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

3. Results

3.1. Distribution of Isolates. A total of 168 *Salmonella enterica* serovar Typhi and Paratyphi isolates were collected in the study period, including 93 (55.4%) *S. Typhi* and 75 (44.6%) *S. Paratyphi A*. There were male preponderance over female (100 (59.5%) versus 68 (40.5%)). The median age was 19

TABLE 1: Demographic characteristics of *Salmonella* isolates in Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

	<i>n</i>	%
Serovars		
<i>Salmonella enterica</i> serovar Typhi	93	55.4
<i>Salmonella enterica</i> serovar Paratyphi A	75	44.6
Sex		
Male	68	40.5
Female	100	59.5
Age group		
≤5 years	8	4.8
6–18 years	68	40.5
19–59 years	88	52.4
≥60 years	4	2.4

years (range 1 to 80 years). Most of the isolates, 156 (92.9%), were from children aged 6–18 years and adult population, 8 (4.8%) were from children below 5 years of age, and 4 (2.4%) were from adult above 60 years of age. There was no mortality. Demographic characteristics of *Salmonella* isolates were shown in Table 1.

3.2. Antimicrobial Susceptibility. *Salmonella Typhi* in this study showed low resistance against most antimicrobials such as ampicillin, trimethoprim-sulfamethoxazole, ceftriaxone, ciprofloxacin, and levofloxacin (5.4%, 8.6%, 0.0%, 1.1%, and 3.2%, resp.). There were also low resistance against the carbapenems (meropenem and imipenem). A decrease of resistance rates against ampicillin, amoxicillin clavulanate, cefotaxime, trimethoprim-sulfamethoxazole, and ampicillin and trimethoprim-sulfamethoxazole was observed between 2011 to 2015. Other antimicrobials showed stable resistance rates. The resistance pattern of *S. Typhi* is shown in Table 2. The minimum inhibitory concentration (MIC) to antimicrobials of *Salmonella Typhi* isolates is shown in Table 3.

Salmonella Paratyphi A in this study showed low resistance against most antimicrobials such as ampicillin, trimethoprim-sulfamethoxazole, ceftriaxone, cefotaxime, ciprofloxacin, and levofloxacin (6.7%, 8.0%, 1.3%, 2.7%, 0.0%, and 1.3%, resp.). Similarly, there was no resistance against the carbapenems. A decrease of resistance rates against trimethoprim-sulfamethoxazole was observed. Other antimicrobials showed stable resistance rates. The resistance pattern of *S. Paratyphi A* is shown in Table 4. The minimum inhibitory concentration (MIC) to antimicrobials of *Salmonella Paratyphi A* isolates is shown in Table 5.

4. Discussion

The *Salmonella* serovars found in this study were 55.4% *S. Typhi* and 44.6% *S. Paratyphi A*, which was in line with a study in Indonesia that showed the predominance of *S. Typhi* over *S. Paratyphi* [11]. Studies in India and China stated that the reason for increasing prevalence of *S. Paratyphi A* over *S. Typhi* in recent years was the increased use of Vi polysaccharide vaccine [12]. Even though, in this study,

TABLE 2: Antimicrobial resistance pattern of *Salmonella* Typhiisolates in Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

Antimicrobial	2011 (n = 14)		2012 (n = 16)		2013 (n = 17)		2014 (n = 27)		2015 (n = 19)		Total (n = 93)	
	n	%	n	%	n	%	n	%	n	%	n	%
AMP	2	14.3	2	12.5	1	5.9	0	0.0	0	0.0	5	5.4
AMC	1	7.2	1	6.3	1	5.9	0	0.0	0	0.0	3	3.2
CRO	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
CT	2	14.3	1	6.3	0	0.0	0	0.0	0	0.0	3	3.2
MER	0	0.0	0	0.0	0	0.0	1	3.7	0	0.0	1	1.1
IPM	0	0.0	0	0.0	0	0.0	1	3.7	0	0.0	1	1.1
TMP-SMX	5	35.7	1	6.3	1	5.9	1	3.7	0	0.0	8	8.6
AMP + TMP-SMX	2	14.3	1	6.3	1	5.9	0	0.0	0	0.0	4	4.3
CIP	0	0.0	1	6.3	0	0.0	0	0.0	0	0.0	1	1.1
LEVX	0	0.0	1	6.3	1	5.9	1	3.7	0	0.0	3	3.2

AMP, ampicillin; AMC, amoxicillin clavulanate; CRO, ceftriaxone; CT, cefotaxime; MER, meropenem; IPM, imipenem; TMP-SMX, trimethoprim-sulfamethoxazole; AKN, amikacin; GEN, gentamicin; CIP, ciprofloxacin; LEVX, levofloxacin.

TABLE 3: Minimum inhibitory concentrations to antimicrobials of *Salmonella* Typhiisolates in Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

Antimicrobial	MIC ($\mu\text{g}/\text{mL}$)	2011 (n = 14)		2012 (n = 16)		2013 (n = 17)		2014 (n = 27)		2015 (n = 19)		Total (n = 93)	
		n	%	n	%	n	%	n	%	n	%	n	%
AMP	≤ 2	11	78,6	10	62,5	11	64,7	20	74,1	16	84,2	68	73,1
	4	1	7,1	4	25,0	3	17,6	4	14,8	2	10,5	14	15,1
	8	0	0,0	0	0,0	2	11,8	3	11,1	1	5,3	6	6,5
	≥ 32	2	14,3	2	12,5	1	5,9	0	0,0	0	0,0	5	5,4
AMC	≤ 2	13	92,9	15	93,8	16	94,1	27	100,0	19	100,0	90	96,8
	4	1	7,1	1	6,3	1	5,9	0	0,0	0	0,0	3	3,2
CRO	≤ 1	14	100,0	16	100,0	17	100,0	27	100,0	19	100,0	93	100,0
	> 1	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
CT	≤ 1	12	85,7	15	93,8	17	100,0	27	100,0	19	100,0	90	96,8
	> 1	2	14,3	1	6,3	0	0,0	0	0,0	0	0,0	3	3,2
MER	≤ 0.25	14	100,0	16	100,0	17	100,0	26	96,3	19	100,0	92	98,9
	> 0.25	0	0,0	0	0,0	0	0,0	1	3,7	0	0,0	1	1,1
IPM	≤ 1	14	100,0	15	93,8	17	100,0	26	96,3	19	100,0	91	97,8
	2	0	0,0	1	6,3	0	0,0	0	0,0	0	0,0	1	1,1
	> 2	0	0,0	0	0,0	0	0,0	1	3,7	0	0,0	1	1,1
TMP-SMX	≤ 20	9	64,3	15	93,8	16	94,1	26	96,3	19	100,0	85	91,4
	≥ 320	5	35,7	1	6,3	1	5,9	1	3,7	0	0,0	8	8,6
CIP	≤ 0.25	12	85,7	15	93,8	12	70,6	21	77,8	19	100,0	79	84,9
	0,5	0	0,0	0	0,0	1	5,9	0	0,0	0	0,0	1	1,1
	1	2	14,3	0	0,0	4	23,5	6	22,2	0	0,0	12	12,9
	4	0	0,0	1	6,3	0	0,0	0	0,0	0	0,0	1	1,1
LEVX	≤ 0.12	14	100,0	14	87,5	13	76,5	19	70,4	19	100,0	79	84,9
	0,25	0	0,0	0	0,0	1	5,9	0	0,0	0	0,0	1	1,1
	0,5	0	0,0	1	6,3	2	11,8	1	3,7	0	0,0	4	4,3
	1	0	0,0	0	0,0	0	0,0	6	22,2	0	0,0	6	6,5
	2	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
	≥ 8	0	0,0	1	6,3	1	5,9	1	3,7	0	0,0	3	3,2

MIC, minimum inhibitory concentrations; AMP, ampicillin; AMC, amoxicillin clavulanate; CRO, ceftriaxone; CT, cefotaxime; MER, meropenem; IPM, imipenem; TMP-SMX, trimethoprim-sulfamethoxazole; AKN, amikacin; GEN, gentamicin; CIP, ciprofloxacin; LEVX, levofloxacin.

TABLE 4: Antimicrobial resistance pattern of *Salmonella* Paratyphi A isolates in Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

Antimicrobial	2011		2012		2013		2014		2015		Total	
	(n = 14)		(n = 22)		(n = 17)		(n = 18)		(n = 4)		(n = 75)	
	n	%	n	%	n	%	n	%	n	%	n	%
AMP	1	7.1	2	9.1	1	5.9	1	5.6	0	0.0	5	6.7
AMC	1	7.1	0	0.0	1	5.9	1	5.6	0	0.0	3	4.0
CRO	0	0.0	1	4.5	0	0.0	0	0.0	0	0.0	1	1.3
CT	1	7.1	1	4.5	0	0.0	0	0.0	0	0.0	2	2.7
MER	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
IPM	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
TMP-SMX	5	35.7	0	0.0	0	0.0	1	5.6	0	0.0	6	8.0
AMO + TMP-SMX	1	7.1	0	0.0	0	0.0	0	0.0	0	0.0	1	1.3
CIP	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
LEVX	0	0.0	0	0.0	0	0.0	1	5.6	0	0.0	1	1.3

AMP, ampicillin; AMC, amoxicillin clavulanate; CRO, ceftriaxone; CT, cefotaxime; MER, meropenem; IPM, imipenem; TMP-SMX, trimethoprim-sulfamethoxazole; AKN, amikacin; GEN, gentamicin; CIP, ciprofloxacin; LEVX, levofloxacin.

TABLE 5: Minimum inhibitory concentrations to antimicrobials of *Salmonella* Paratyphi A isolates in Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia.

Antimicrobial	MIC ($\mu\text{g/mL}$)	2011		2012		2013		2014		2015		Total	
		(n = 14)		(n = 22)		(n = 17)		(n = 18)		(n = 4)		(n = 75)	
		n	%	n	%	n	%	n	%	n	%	n	%
AMP	≤ 2	13	92,9	18	81,8	13	76,5	13	72,2	2	50,0	59	78,7
	4	0	0,0	2	9,1	1	5,9	1	5,6	1	25,0	5	6,7
	8	0	0,0	0	0,0	2	11,8	3	16,7	1	25,0	6	8,0
	≥ 32	1	7,1	2	9,1	1	5,9	1	5,6	0	0,0	5	6,7
AMC	≤ 2	13	92,9	22	100,0	16	94,1	17	94,4	4	100,0	72	96,0
	4	1	7,1	0	0,0	1	5,9	1	5,6	0	0,0	3	4,0
CRO	≤ 1	14	100,0	21	95,5	17	100,0	18	100,0	4	100,0	74	98,7
	> 1	0	0,0	1	4,5	0	0,0	0	0,0	0	0,0	1	1,3
CT	≤ 1	13	92,9	21	95,5	17	100,0	18	100,0	4	100,0	73	97,3
	> 1	1	7,1	1	4,5	0	0,0	0	0,0	0	0,0	2	2,7
MER	≤ 0.25	14	100,0	22	100,0	17	100,0	18	100,0	4	100,0	75	100,0
	> 0.25	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
IPM	≤ 1	14	100,0	21	95,5	17	100,0	18	100,0	4	100,0	74	98,7
	2	0	0,0	1	4,5	0	0,0	0	0,0	0	0,0	1	1,3
	> 2	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
TMP-SMX	≤ 20	9	64,3	22	100,0	17	100,0	17	94,4	4	100,0	69	92,0
	≥ 320	5	35,7	0	0,0	0	0,0	1	5,6	0	0,0	6	8,0
CIP	≤ 0.25	13	92,9	21	95,5	14	82,4	13	72,2	3	75,0	64	85,3
	0,5	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
	1	1	7,1	1	4,5	3	17,6	5	27,8	1	25,0	11	14,7
	4	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
LEVX	≤ 0.12	14	100,0	21	95,5	15	88,2	10	55,6	2	50,0	62	82,7
	0,25	0	0,0	0	0,0	0	0,0	1	5,6	0	0,0	1	1,3
	0,5	0	0,0	1	4,5	2	11,8	2	11,1	2	50,0	7	9,3
	1	0	0,0	0	0,0	0	0,0	4	22,2	0	0,0	4	5,3
	2	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0	0	0,0
	≥ 8	0	0,0	0	0,0	0	0,0	1	5,6	0	0,0	1	1,3

MIC, minimum inhibitory concentrations; AMP, ampicillin; AMC, amoxicillin clavulanate; CRO, ceftriaxone; CT, cefotaxime; MER, meropenem; IPM, imipenem; TMP-SMX, trimethoprim-sulfamethoxazole; AKN, amikacin; GEN, gentamicin; CIP, ciprofloxacin; LEVX, levofloxacin.

there were no data on the patients' history of vaccination, the data of this study showed that salmonellosis affected mostly young and middle-aged population as 92.9% of these populations have more contact with water and food that were contaminated with *Salmonella* [13]. The increased public standard of living resulted the reduction in the frequency of *Salmonella* isolation in a study in China, due to the reduced consumption of contaminated water and food [14].

The antimicrobial resistance rates of *S. Typhi* differ among countries in the world. A study in Pakistan showed that resistance rates of *S. Typhi* and Paratyphi were 88.2% for fluoroquinolone, 66.1% for ampicillin, and 66.5% for trimethoprim-sulfamethoxazole [15]. Another study in China showed that resistance rates of *S. Typhi* and Paratyphi were 13.5% and 5.9% for ciprofloxacin, 5.4% and 0.8% for levofloxacin, 5.4% and 1.4% for sulfamethoxazole, 10.8% and 2.0% for ampicillin, 5.4% and 0.0% for ceftriaxone, and 0.0% for meropenem and imipenem [14]. A study in Nepal showed 1.8% resistance of *S. Typhi* against ampicillin and 3.9% of *S. Paratyphi* against ciprofloxacin, while there was no resistance against trimethoprim-sulfamethoxazole [16]. A study in Asian countries showed that resistance rates in Bangladesh were 39.5% for ciprofloxacin, 68.4% for ampicillin, 57.9% for trimethoprim, and 68.4% for sulfamethoxazole, in Vietnam 0.0% for ciprofloxacin, and 80.4% for ampicillin, trimethoprim, and sulfamethoxazole, while in Indonesia they were 0.0% for ciprofloxacin, 1.8% for ampicillin and trimethoprim, and 3.6% for sulfamethoxazole [17]. A community-based study from 2001 to 2003 in Indonesia showed low resistance rates of *S. Typhi* and *S. Paratyphi*, where only 2.5% isolates of *S. Typhi* were resistant against ampicillin, while there was no resistance against trimethoprim-sulfamethoxazole, ceftriaxone, or ciprofloxacin [11]. A hospital-based study from 2006 to 2010 in Indonesia also found similar results; *S. Typhi* showed no resistance against trimethoprim-sulfamethoxazole, ciprofloxacin, and meropenem, 0.9% against ceftriaxone, 1.6% against cefotaxime, and 1.9% against ampicillin [18]. This study also found similar results to studies in Indonesia; the low resistance of *S. Typhi* and *S. Paratyphi* isolates from 2011 to 2015 against most antimicrobials such as ampicillin, trimethoprim-sulfamethoxazole, ceftriaxone, ciprofloxacin, and levofloxacin (5.4%, 8.6%, 0.0%, 1.1%, 3.2% and 6.7%, 8.0%, 1.3%, 2.7%, 0.0%, and 1.3%, resp.). In this study, the frequency of MDR *S. Typhi* would be no more than 4.3%. The resistance rate of *S. Typhi* isolates against ampicillin and trimethoprim-sulfamethoxazole was only 4.3%, so even if all the isolates were resistant to chloramphenicol, the MDR rate would not be higher than that. A study stated that there were differences in the resistance rates between *S. Paratyphi A* or *S. Typhi*; therefore they have to be considered as different diseases [19]. This study, on the contrary, indicated no significant different resistance rates between *S. Typhi* and *S. Paratyphi A*.

The two antimicrobial resistance mechanisms in *S. Typhi* were the plasmid-mediated mechanism and the chromosomal-mediated mechanism [20]. Plasmids that transfer the antimicrobial resistance in *S. Typhi* are the plasmids of incompatibility group (Inc) HII. The first IncHII plasmid

was isolated in Mexico City in 1972 and rendered resistance to chloramphenicol, tetracycline, streptomycin, and sulphonamides. Afterwards, MDR *S. Typhi* associated with IncHII plasmids were found worldwide. The chromosomal-mediated drug resistance against fluoroquinolones was the result of selective selection on the bacterial population by the usage of the antimicrobial. The single point mutation in the quinolone resistance determining region (QRDR) of the topoisomerase gene *gyrA* that encodes DNA gyrase was the resistance gene against fluoroquinolones. Other resistance genes of *S. Typhi* were *cat* for chloramphenicol, *tetA* for tetracycline, *bla* for ampicillin, *strA-strB* for streptomycin, and *dfrA* and *sulI* for trimethoprim and sulfonamide [20]. A study found very closely related plasmids that have a common backbone carrying identical resistance insertions and were present in *S. Typhi* and *S. Paratyphi A*. This study determined the DNA sequence of an IncHII plasmid, pAKU_1, encoding MDR in a *S. Paratyphi A*. It strongly suggests that there had been transfer of a plasmid between *S. Typhi* and *S. Paratyphi A* [21].

The explanation of different resistance pattern in the world was that there are different strains of *S. Typhi* and *S. Paratyphi* in different regions which also have different genes that contributed to resistance against antimicrobials [17]. A study in 2008 on haplotypes and pathotypes of *S. Typhi* in Jakarta, Indonesia, found 9 serovar strains of haplotypes which were H1, H8, H42, H45, H50, H52, H59, H84, and H85. Haplotypes H59 and H8 were dominating the isolates, 53% and 24%, respectively. Haplotypes of H42, H50, H59, and H85 were also found in previous studies, whereas H85 had been isolated in 1987 and 2003. Haplotype H59 which is associated with *j* and *z66* flagellum expression seemed to be the specific phenotype in Indonesia, as it has been consistently isolated over a 30-year period. A study on the DNA profiles of *S. Typhi* in Indonesia showed clear differences according to regions but all the *S. Typhi* isolates showed similar phenotypes which were susceptible to chloramphenicol, ampicillin, and trimethoprim-sulfamethoxazole [22]. The MDR rate in this study was 4.3%, which confirmed the result of previous studies that MDR rate in Indonesia is low. Only one strain of H58 that was resistant to fluoroquinolone was found from a French traveler returning from Indonesia in 2003. Mutations in DNA gyrase were not found in the study, which indicates that there was no recent clonal expansion of H58 in Indonesia. Studies found that MDR *S. Typhi* strains belong to haplotype H58, which spread widely over the Indochina Peninsula, the Indian subcontinent, and Africa but is not prevalent in Indonesia [23–28]. It could be concluded that there is a distinct epidemiological dynamic of the enteric fever in Indonesia. As stated above, there was evidence of transfer of a plasmid responsible for multidrug resistance between *S. Typhi* and *S. Paratyphi A* [21]. If the haplotype H58 *S. Typhi* is not found in Indonesia, the IncHII plasmid that could transfer multidrug resistance to *S. Paratyphi* would also be absent. This would explain the nonsignificant difference of resistance rates between *S. Typhi* and *S. Paratyphi A* in this study.

According to the 2006 Indonesian Typhoid Fever Disease Control Guidelines, chloramphenicol, ampicillin, or

trimethoprim-sulfamethoxazole is used as first-line antimicrobials, and if there is resistance to the first-line antimicrobials then the second line antimicrobials; ceftriaxone, cefixime, or quinolones is indicated [29]. In Indonesia, a study on hospitalized adult typhoid patients found that ceftriaxone was the antimicrobial most commonly used followed by levofloxacin, ciprofloxacin, meropenem, metronidazole, and ampicillin-sulbactam [30]. The frequent use of second line antimicrobials for enteric fever in Indonesia could also be the explanation for the low MDR and low resistance rates for first-line antimicrobials in this study, where a study in Nepal found that the discontinuation of chloramphenicol and trimethoprim-sulfamethoxazole for long periods might be the cause of reemergence of sensitive *S. Typhi* and *S. Paratyphi* [16].

There were limitations of this study that should be mentioned. The samples of this study were blood collected from admitted patients in a hospital; thus the antimicrobial resistance pattern was restricted to the patients of the hospital. The resistance pattern must be interpreted cautiously, as isolates might not represent the entire population of Banten province or Indonesia. Another issue was the unavailability of data on chloramphenicol susceptibility because of the policy of the Indonesian Microbiology Association not to test chloramphenicol for *S. Typhi* and *S. Paratyphi* anymore. Other antimicrobial not tested was nalidixic acid. In vitro nalidixic acid is more appropriate to test for in vivo fluoroquinolone resistance.

5. Conclusions

The study discovered that in the 2011–2015 period, the resistance rates against most antimicrobials and even MDR rate of *S. Typhi* and *S. Paratyphi* were low, in accordance with previous studies in different regions of Indonesia in the 2001–2003, 2006–2010, and 2008 periods. It emphasizes that there is a distinct epidemiological dynamic of the enteric fever in Indonesia. To prevent the occurrence of resistance and MDR *S. Typhi*, it is imperative to maintain continuous monitoring of antimicrobial resistance and follow a rational prescription of antimicrobials based on the local antimicrobial pattern.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.




Acknowledgments

The authors wish to acknowledge the assistance of Dewi Purnamasari M.D., Resident Medical Officer at Siloam Hospital, Karawaci, Tangerang, Banten, Indonesia, who collected the epidemiology, clinical, and microbiological data from medical records and Microbiology laboratory.

References

- [1] J. A. Crump, S. P. Luby, and E. D. Mintz, "The global burden of typhoid fever," *Bulletin of the World Health Organization*, vol. 82, no. 5, pp. 346–353, 2004.
- [2] A. Kothari, A. Pruthi, and T. D. Chugh, "The burden of enteric fever," *Journal of infection in developing countries*, vol. 2, no. 4, pp. 253–259, 2008.
- [3] S. Kumar, M. Rizvi, and N. Berry, "Rising prevalence of enteric fever due to multidrug-resistant *Salmonella*: an epidemiological study," *Journal of Medical Microbiology*, vol. 57, no. 10, pp. 1247–1250, 2008.
- [4] C. K. J. Paniker and K. N. Vimala, "Transferable chloramphenicol resistance in salmonella typhi," *Nature*, vol. 239, no. 5367, pp. 109–110, 1972.
- [5] M.-L. Ackers, N. D. Puhr, R. V. Tauxe, and E. D. Mintz, "Laboratory-based surveillance of *Salmonella* serotype typhi infections in the United States: antimicrobial resistance on the rise," *Journal of the American Medical Association*, vol. 283, no. 20, pp. 2668–2673, 2000.
- [6] S. H. Mirza, N. J. Beeching, and C. A. Hart, "Multi-drug resistant typhoid: a global problem," *Journal of Medical Microbiology*, vol. 44, no. 5, pp. 317–319, 1996.
- [7] S. Le Hello, A. Bekhit, S. A. Granier et al., "The global establishment of a highly-fluoroquinolone resistant *Salmonella enterica* serotype Kentucky ST198 strain," *Frontiers in Microbiology*, vol. 4, article 395, 2013.
- [8] D. Yanagi, G. C. D. Vries, D. Rahardjo et al., "Emergence of fluoroquinolone-resistant strains of *Salmonella enterica* in Surabaya, Indonesia," *Diagnostic Microbiology and Infectious Disease*, vol. 64, no. 4, pp. 422–426, 2009.
- [9] R. L. Ochiai, C. J. Acosta, M. C. Danovaro-Holliday et al., "A study of typhoid fever in five Asian countries: disease burden and implications for controls," *Bulletin of the World Health Organization*, vol. 86, no. 4, pp. 260–268, 2008.
- [10] Clinical and Laboratory Standards Institute, *Performance Standards for Antimicrobial Susceptibility Testing*, M100-S23, Clinical & Laboratory Standards Institute: CLSI, Wayne, Pa, USA, 2013.
- [11] N. H. Punjabi, M. D. Agtini, R. L. Ochiai et al., "Enteric fever burden in North Jakarta, Indonesia: a prospective, community-based study," *Journal of Infection in Developing Countries*, vol. 7, no. 11, pp. 781–787, 2013.
- [12] S. Dutta, S. Das, U. Mitra et al., "Antimicrobial resistance, virulence profiles and molecular subtypes of *Salmonella enterica* serovars Typhi and paratyphi A blood isolates from Kolkata, India during 2009–2013," *PLoS ONE*, vol. 9, no. 8, Article ID e101347, 2014.
- [13] B. M. Taban, S. A. Aytac, N. Akkoc, and M. Akcelik, "Characterization of antibiotic resistance in salmonella enterica isolates determined from ready-to-eat (RTE) salad vegetables," *Brazilian Journal of Microbiology*, vol. 44, no. 2, pp. 385–391, 2013.
- [14] J. Yaxian, Z. Hui, N. Hua et al., "Antimicrobial resistance surveillance of *Salmonella* isolates from the First People's Hospital of Yunnan Province, China," *Journal of Infection in Developing Countries*, vol. 9, no. 4, pp. 333–337, 2015.
- [15] F. N. Qamar, A. Azmatullah, A. M. Kazi, E. Khan, and A. K. M. Zaidi, "A three-year review of antimicrobial resistance of *Salmonella enterica* serovars Typhi and Paratyphi A in Pakistan," *Journal of Infection in Developing Countries*, vol. 8, no. 8, pp. 981–986, 2014.
- [16] H. J. Chand, K. R. Rijal, B. Neupane, V. K. Sharma, and B. Jha, "Re-emergence of susceptibility to conventional first line drugs in *Salmonella* isolates from enteric fever patients in Nepal," *Journal of Infection in Developing Countries*, vol. 8, no. 11, pp. 1483–1487, 2014.
- [17] C.-S. Chiou, T.-L. Lauderdale, D. C. Phung et al., "Antimicrobial resistance in *Salmonella enterica* serovar Typhi isolates from

- Bangladesh, Indonesia, Taiwan, and Vietnam,” *Antimicrobial Agents and Chemotherapy*, vol. 58, no. 11, pp. 6501–6507, 2014.
- [18] A. Alam, “Pola resistensi *Salmonella enterica* Serotipe typhi, departemen Ilmu Kesehatan anak RSHS, Tahun 2006–2010,” *Sari Pediatri*, vol. 12, no. 5, pp. 296–301, 2011.
- [19] J. Wain, L. T. D. Nga, C. Kidgell et al., “Molecular analysis of incHII antimicrobial resistance plasmids from *Salmonella* serovar Typhi strains associated with typhoid fever,” *Antimicrobial Agents and Chemotherapy*, vol. 47, no. 9, pp. 2732–2739, 2003.
- [20] S. A. Zaki and S. Karande, “Multidrug-resistant typhoid fever: a review,” *Journal of Infection in Developing Countries*, vol. 5, no. 5, pp. 324–337, 2011.
- [21] K. E. Holt, N. R. Thomson, J. Wain et al., “Multidrug-resistant *Salmonella enterica* serovar Paratyphi A harbors IncHII plasmids similar to those found in serovar Typhi,” *Journal of Bacteriology*, vol. 189, no. 11, pp. 4257–4264, 2007.
- [22] L. H. Moehario, “The molecular epidemiology of *Salmonella* Typhi across Indonesia reveals bacterial migration,” *Journal of Infection in Developing Countries*, vol. 3, no. 8, pp. 579–584, 2009.
- [23] K. E. Holt, S. Baker, S. Dongol et al., “High-throughput bacterial SNP typing identifies distinct clusters of *Salmonella* Typhi causing typhoid in Nepalese children,” *BMC Infectious Diseases*, vol. 10, article 144, 2010.
- [24] P. Roumagnac, F.-X. Weill, C. Dolecek et al., “Evolutionary history of *Salmonella* Typhi,” *Science*, vol. 314, no. 5803, pp. 1301–1304, 2006.
- [25] S. Kariuki, G. Revathi, J. Kiiru et al., “Typhoid in Kenya is associated with a dominant multidrug-resistant *Salmonella enterica* serovar typhi haplotype that is also widespread in Southeast Asia,” *Journal of Clinical Microbiology*, vol. 48, no. 6, pp. 2171–2176, 2010.
- [26] K. E. Holt, J. Parkhill, C. J. Mazzoni et al., “High-throughput sequencing provides insights into genome variation and evolution in *Salmonella* Typhi,” *Nature Genetics*, vol. 40, no. 8, pp. 987–993, 2008.
- [27] K. E. Holt, C. Dolecek, T. T. Chau et al., “Temporal fluctuation of multidrug resistant *Salmonella* Typhi haplotypes in the mekong river delta region of Vietnam,” *PLoS Neglected Tropical Diseases*, vol. 5, no. 1, article e929, 2011.
- [28] S. Baker, K. Holt, E. Van De Vosse et al., “High-throughput genotyping of *Salmonella enterica* serovar Typhi allowing geographical assignment of haplotypes and pathotypes within an urban district of Jakarta, Indonesia,” *Journal of Clinical Microbiology*, vol. 46, no. 5, pp. 1741–1746, 2008.
- [29] Minister of Health Republic of Indonesia, “Guidelines for control of typhoid fever,” Decree Number 364/MENKES/SK/V/2006, 2006.
- [30] A. B. Anggraini, C. Opitasari, and S. Qamp, “The use of antibiotics in hospitalized adult typhoid patients in an Indonesian hospital,” *Health Science Journal of Indonesia*, vol. 1, pp. 40–43, 2014.



Hindawi

Submit your manuscripts at
<https://www.hindawi.com>

