

Antioxidant and hypoglycemic activities of extract and fractions of Rambutan seeds (*Nephelium lappaceum* L.)

Slyvia Soeng^{1†}, Endang Evacuasiyany¹, Wahyu Widowati^{1†}, Nurul Fauziah²

¹Faculty of Medicine, Maranatha Christian University, Bandung, West Java, Indonesia

²Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia

ARTICLE INFO

Article history:

Received: January 2015

Accepted: June 2015

Available online:

August 2015

Keywords:

α -glucosidase

Antidiabetic

Antioxidant

DPPH

Nephelium lappaceum L.

SOD

†Corresponding author:

Jl. Prof. Drg. Suria Sumantri
no 65 Bandung 40164,
Indonesia.

wahyu_w60@yahoo.com;

s.soeng@yahoo.com

ABSTRACT

Objective : This research was done to measure the antioxidants and hypoglycemic activities of NLS extract and fractions.

Methods: The ethanol extract of rambutan seeds were prepared by maseration method and the fractions (n-hexane, ethyl acetate, buthanol and water) by separation of extract based on the polarity. The antioxidant activity was determined by using superoxide dismutase value (SOD), 1,1-diphenyl-2-picryl-hydrazyl (DPPH) radical scavenging activity. The hypoglycemic activity was determined by using inhibition of α -glucosidase test. The DPPH scavenging and hypoglycemic activities were analized by median of Inhibitory Concentration (IC-50).

Result : The highest SOD activity showed that ethyl acetate and aqueous fraction of NLS were 3.3771 μ g/ml and 3.0374 μ g/ml. Meanwhile DPPH assay showed that both NLS extract and fractions had low DPPH scavenging activity. Hypoglycemic activity showed that extract of NLS had highest activity as α -glucosidase inhibitor.

Conclusion : NLS extract and fractions have high SOD antioxidant value but low DPPH scavenging activity and can be used as potential hypoglycemic agent.

1. Introduction

Diabetes Mellitus (DM) is common disease associated with markedly increase mortality rate. The number of DM has reached to 285 million, approximately 6.4% of the world's adult population, more than 3.8 million people die yearly [1,2,3]. Diabetes can be characterized by hyperglycemia resulting from defects in insulin action and insulin secretion or both. The most common type of DM is DM II, which accounts for 85 to 95% of all cases and constitutes a major and growing public health problem [4]. DM II is generally managed through intensive therapy that consists of lifestyle and sequential addition of oral antihyperglycemic agents (OHAs). One therapeutic approach to decrease postprandial hyperglycemia in diabetic state is by retarding absorption of glucose through inhibition of

carbohydrate hydrolyzing enzymes, like α -glucosidase and α -amylase in the digestive tract [5]. α -glucosidase inhibitors delay absorption of complex carbohydrates and thus inhibit postprandial glucose peaks thereby leading to decreased postprandial insulin levels [6].

Oxidative stress in hyperglycemia-related diabetic patient causing excessive amount of free radical which may damage antioxidant defense [7].

These drugs can induce hypoglycemia, lose their efficacy, have prominent side effects and trigger diabetic complications. Plants have been suggested as source of potentially antidiabetic drugs [8].

Nephelium lappaceum L. known as rambutan is one of variety tropical fruit which commonly consumed in south-east Asia. Rambutan has antioxidant activity and high phenolic content [9].

Therefore, we conducted our research to evaluate antioxidant and hypoglycemic activity of rambutan's seed (NLS) extract and fractions.

2. Materials and Methods

2.1 Extract and fraction preparation

Extraction was done based on maceration method [10,11,12] and fractions were done as modified partition [10,13]. NLS was collected from Kesamben-Blitar plantation, East Java, Indonesia. Four hundred gram of dried and milled NLS were soaked in destilated ethanol 70% then were evaporated and resulted 11.25 % of crude extract (45 g). Fifty gram of NLS ethanolic extract was partitioned with n-hexan and water (1:1), yielded hexane fraction 23.37 g (46,74 %), The residue was partitioned with ethyl acetate and water (1:1) yielded ethyl acetate fraction 2.3 g (7.71%), the residue was partioned with buthanol and water (1:1) yielded buthanol fraction 2.85 (9.56 %), the residue was water fraction 1.3 g (2.6 %).

2.2 DPPH scavenger test

The DPPH scavenger test was done by introducing 50 μ l extract and fractions of NLS in methanol with final concentrations (0.19 μ g/ml; 0.391; 0.781; 1.563; 3.125; 6.25; 12.5; 25; 50; to 100 μ g/ml) in 96 well microplate and were added 200 μ l DPPH 0.077 mmol in DMSO. The mixture was shaken vigorously and incubated at room temperature and dark room for 30 min, and then measured at 517 nm absorbance using a microplate reader (Multi Go Skan). Negative controls used DPPH 250 μ l, blank used 250 μ l methanol [11,12,13]. The DPPH scavenger activity (%):

$$\text{scavenging \%} = \frac{A_c - A_s}{A_c} \times 100$$

Aa: sample absorbance

Ac: negative control absorbance (without sample)

2.3 Superoxide Dismutase (SOD) test

The SOD test was done by a SOD assay kit (Cayman) comprised of assay buffer, sample buffer, radical detector, SOD standard, and xanthine oxidase. SOD standards were prepared by introducing 200 μ l diluted radical detector and 10 μ l SOD standard (7-level standard) per well [12]. Samples were dissolved in DMSO in

concentrations 500, 125, 31.25 μ g/ml. The sample well contained 200 μ l diluted radical detector and 10 μ l sample. Samples and standards wells were added 20 μ l diluted xanthine oxidase. The mixtures were shaken carefully for few seconds, incubated for 20 minutes at room temperature, SOD activity was measured on a microplate reader at 440-460 nm [12]. The SOD value was calculated using the equation from the linear regression of standard curve substituting linear lineared rate (LR) for each sample. One unit is defined as the amount of enzyme to yield 50% dismutation of the superoxide radical [12]. Furthermore, standard curves were constructed based on the value of the LR and SOD value be calculated:

The calculation was done by calculating the value of SOD linearized rate/LR (LR Std A= Abs Std A/Abs Std A: LR Std B= Abs Std A/Abs Std B)

2.3 α -glucosidase inhibitor test

The α -glucosidase inhibitor activity was

$$\text{SOD (U/ml)} = \left[\frac{\text{Sample LR-y intercept}}{\text{Slope}} \right] \times 0,23\text{ml}/0,01\text{ml} \times \text{Sample dilution}$$

tested by the modified method of [14,15]. Samples were diluted in DMSO 10% with various concentrations (500 μ g/ml; 125; 31.25; 7.81; 1.9 μ g/ml). Amount of 5 μ L of sample, 25 μ l of 200 mM p-nitrophenyl- α -glucopyranoside, 45 μ l phosphate buffer saline (pH 7), 25 μ l of *Saccharomyces* sp. yeast α - glucosidase were introduced in the microplate and incubated at 37^o C for 5 min. The reaction were stopped by adding 100 μ L of 200 mM Na₂CO₃ and then was measured at 400 nm using a microplate reader (Multi Go Skan). Controls without inhibitors were checked, as a reference. The α -glucosidase inhibitory activity could be calculated as follows:

$$\text{scavenging \%} = \frac{A_c - A_s}{A_c} \times 100$$

Ac, as represent the absorbance at 400 nm of the control, sample respectively

3. Result

3.1 DPPH scavenger activity

The DPPH free radical scavenger activity of extract and fractions are representative of antioxidant activity. The IC₅₀ is the concentration of antioxidant needed to scavenge 50% of the DPPH free radical [12]. Based on the IC₅₀ (Table

1.) and Figure 1 showed that NLS extract had the lowest DPPH scavenger activity with IC₅₀ 341.20 µg/ml, ethyl acetate fraction had highest activity with IC₅₀ 104.03 µg/ml.

Based on the IC₅₀ values of DPPH scavenger activity (Table 1.) showed that extract and fractions of NLS had low antioxidant activity. The IC₅₀ value was used to determine an antioxidant

Table 1. DPPH scavenger activity (IC₅₀) of extract and fractions of NLS. The DPPH scavenger activity test were measured triplicate for each sample. Linear equations, coefficient of regression (R²), and IC₅₀ were calculated.

Samples	Linear equation	R ²	IC50 (µM)
Extract	Y=0.1415X+1.7203	0.9285	341.20
Hexane fraction	Y=0.2842X+4.3993	0.9002	160.45
Ethyl acetate fraction	Y=0.4519X+2.9909	0.9887	104.03
Buthanol fraction	Y=0.155X+3.9125	0.8904	297.34
Water fraction	Y=0.2188X+5.6823	0.9023	202.55




Figure 1. DPPH scavenger activity of extract and fractions of NLS diluted in methanol to achieve the final concentrations 100; 50; 25; 12.5; 6.25; 3.125; 1.563; 0.781; 0.391; 0.19 µg/ml.

activity by DPPH test, which the smallest the IC₅₀ value is highest antioxidant activity [13].

3.2 The SOD activity

Superoxide anion (O₂^{•-}) is one of the most important radical formed in aerobic cells due to leakage of the electron transport chain. Although less reactive but radical O₂^{•-} is initiation radical oxidation, it is a precursor to form hydroxyl radical (OH[•]) are highly reactive through Fenton reaction and Haber-Weiss [16]. Superoxide

dismutase (SOD) as an antioxidant activity would be increased by changing the superoxide anion (O₂^{•-}) into hydrogen peroxide (H₂O₂) and oxygen (O₂) [17]. The SOD activity of extract and fractions of NLS in trapping O₂^{•-} can be seen in Table 2, Figure 2.

Based on the results (Table 2.) showed that lower concentrations of sample reduced SOD value. The highest SOD value at a concentration of 500 µg/ml was buthanol fraction (3.377 U/ml), while the lowest was hexane and ethyl acetate fractions (1.334-1.472 U/ml). The highest SOD

Table 2. Mean and Tukey HSD post hoc test of SOD activity of extract and fractions (U/ml) was measured in triplicate for sample. (Linear equation, coefficient of regression (R²) of SOD standard and SOD activity of extract and fractions were calculated)

Samples	Concentrations (µg/ml)		
	500	125	31.25
Extract	1.780±0.118 b	1.296±0.036 c	0.807±0.032 bc
Hexane fraction	1.334±0.091 a	1.038±0.014 b	0.726±0.033 b
Ethyl acetate	1.472±0.164 a	0.759±0.005 a	0.577±0.052 a
Buthanol fraction	3.377±0.175 d	1.874±0.077 d	0.876±0.026 c
Water fraction	3.037±0.095 c	2.314±0.024 e	1.587±0.049 d

Data are presented as mean ± standard deviation. Different letters in the same column (among samples) are significant at P < 0.05 (Tukey's HSD post hoc test)

value at 125 µg/ml was water fraction 2.314 U/ml and the lowest was ethyl acetate fraction 0.759 U/ml. The highest SOD value at 31.25 µg/ml was water fraction 1.587 U/ml), the lowest was ethyl acetate (0.577 U/ml). Overall in the three concentrations the highest SOD value were buthanol and water fractions of NLS. Table 2 showed that the higher concentration was used the higher sample could reduce SOD value.

3.3 The α -glucosidase inhibitor activity

Alpha-glucosidase is a key enzyme in carbohydrate digestion, it catalyzes the hydrolysis of 1,4- α -glucosidic bonds within carbohydrates with release α -glucose and trigger the gaining blood glucose levels after meal. Alpha-glucosidase inhibitors can delay the intestinal carbohydrate

Table 3. α -glucosidase inhibitor (IC_{50}) of extract and fractions of NLS. The α -glucosidase inhibitor activity test were measured triplicate for each sample. Linear equations, coefficient of regression (R^2), and IC_{50} were calculated.

Samples	Linear equation	R^2	IC_{50} (μ M)
Extract	$Y=1.8448X+31.705$	0.9507	9.92
Hexane fraction	$Y=2.6678X+6.7865$	0.8728	16.20
Ethyl acetate fraction	$Y=2.2174X+26.866$	0.8003	10.43
Buthanol fraction	$Y=2.7092X+15.662$	0.8587	12.67
Water fraction	$Y=2.5254X+14.2$	0.8324	14.18




Figure 2. α -glucosidase inhibitor activity of extract and fractions of NLS diluted in DMSO to achieve the final concentrations 12.5; 3.125; 0.781; 0.195; 0.049 µg/ml

absorption and slow the gaining blood glucose levels [3]. The α -glucosidase inhibitors of extract and fractions can be seen in Table 3, Figure 2.

Based on (Table 3) showed that extract and fractions of NLS had high α -glucosidase inhibitor activity. The highest α -glucosidase inhibitor activity was extract with IC_{50} 9.92 µg/ml.

4. Discussion

Extract and fractions of NLS had low DPPH free radical scavenger activity (Table 1., Fig. 1), but had high SOD value (Table. 2). Based on Table 1 showed that NLS seed extract had lowest radical scavenging activity. This data was consistent with previous research that rambutan seed contained low of total phenolic compound,

rambutan peel extract contained 542.2 mg catechin/g and rambutan seed extract contained 58.5 mg catechin/g. The rambutan peel extract exhibited higher antioxidant activity than seed extracts [18], rambutan peel possessed high DPPH free radical scavenging activity (IC_{50} of 8.87 µg/ml) [19], high polyphenols content of rambutan peel contributes towards high free radical scavenging activity [20]. The result of DPPH assay in our study showed that ethyl acetate fraction which had the highest free radical scavenging activity with IC_{50} 104.3 µg/ml, in previous study the highest DPPH scavenger activity was methanolic fraction with IC_{50} 4.94 µg/ml [21]. While the reason for the different activity could be due to different solvent for extraction and fractionation [22]. Previous study

exhibited that ethanolic extract of white saffron had lower DPPH scavenger activity compared to water extract of white saffron [22, 23]. Different solvent resulted different compound and bioactivity. Water extract of Forsythia korean flowers exhibited a higher phenolic content than ethanolic extract [24].

Previous studies reported that rambutan seed possessing a relatively high amount of fat with values between 14-41 g/100 g [25], the rambutan seed possesses a relatively high amount of fat between 17-39%. The fat content of rambutan seed were saturated fatty acid (SFA) 50.7% and monosaturated fatty acid (MUFA) 48.1%, the main fatty acid were oleic acid 40.3%, arachidic acid 34.5%, stearic acid 7.1% [26]. Previous data that using HPLC of rambutan seed contained high triacylglycerol with AOO (Arachidoyl-Dioleoylglycerol) 49.84%, ASO (Arachidoyl-Stearoyl-Oleoylglycerol) 15.058%, AOP (Arachidoyl-Oleoyl-Palmitoglycerol) 12.822% [27]. The natural antioxidant in lipid-containing product and lipid-based product such as oil, fat, margarine, butter in rambutan seed are considered insufficient antioxidant activities [26], fermentation and roasting process in rambutan seed fat can improve the antioxidant activity and total phenolics compound of rambutan seed fat [26,28].

Extract of NLS showed the highest α -glucosidase inhibitor activity (Table 3., Figure 2.). This data was validated with previous research that high dose of rambutan seed infusion (3.12 g/kg bw) had significant effect in reducing the blood glucose and improve pancreatic beta cells of diabetic mice [29]. Extract and fractions of NLS were more active as α -glucosidase inhibitor activity compared to drug acarbose with $IC_{50}=3500 \mu\text{g/ml}$ [30], more active than glucobay with $IC_{50} 24.44 \mu\text{g/ml}$ [15]. α -glucosidase is an enzyme for carbohydrate digestion and absorption and has been used as therapeutic target for its modulating action to reduce postprandial hyperglycemia. Hyperglycemia is a risk factor for the development of oxidative stress-related diabetes mellitus [28, 29]. In this present study showed that NLS extract and fractions have potency to be hypoglycemic agent due to it α -glucosidase inhibitor activity.

5. Conclusion

This study has demonstrated the antioxidant and α -glucosidase inhibitory activities of NLS extract and fractions. The potential antioxidant

activities are ethyl acetate fraction for highest DPPH radical scavenger activity and water fraction, buthanol fraction for highest SOD value. NLS extract and fractions have beneficial effect to be hypoglycemic agent by inhibiting α -glucosidase activities.

Acknowledgment

We gratefully acknowledge the financial support of the Directorate General of Higher Education, National Ministry of Republic Indonesia for research grant of Hibah Bersaing 2012, 2014

Conflict of Interest

The authors report no conflicts of interest

References

1. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pr* 2010; 87: 4-14.
2. Van Dieren S, Beulens JW, van der Schouw JT, Grobbee DE, Neal B. The global burden of diabetes and its complications: an emerging pandemic. *Eur J Cardiovasc Prev Rehabil* 2010;17 Suppl 1:S3-8.
3. Wu C, Shen J, He P, Chen Y, Li L, Zhang L, Li Y, Fu Y, Dai R, meng W, Deng Y. The α -glucosidase inhibiting isoflavones isolated from *Belamcanda chinensis* leaf extract. *Rec Nat Prod* 2012; 6(2):110-120
4. Cheplick S, Kwon YI, Bhowmik P, Shetty K. Phenolic linked variation in strawberry cultivars for potential dietary management of hyperglycemia and related complications of hypertension. *Bioresour Technol* 2010; 101: 404-413.
5. Manaharan T, Ling LT, Appleton D, Cheng HM, Masilamani T, Palanisamy U. Antioxidant and antihyperglycemic potential of *Peltophorum pterocarpum* plant parts. *Food Chem* 2011; 129:1355-1361.
6. Van de Laar FA, Lucassen PLBJ, Akkermans RP, Van de Lisdonk EH, Rutten GEHM, Van Weel C. Alpha-glucosidase inhibitors for type 2 diabetes mellitus: a systematic review. *Chin J Evid-based Med* 2006; 6(5):335-351
7. Johansen JS, Harris AK, Rychly DJ, Ergul A: Oxidative stress and the use of antioxidants in diabetes: Linking basic science to clinical practice. *Cardiovasc Diabetol* 2005, 4(5):1-11
8. Mukherjee PK, Maiti K, Mukherjee K, Houghton, P.J. Leads from Indian medicinal plants with hypoglycemic potentials. *J Ethnopharmacol* 2006;106:1-28.
9. Thitilertdecha, N.; Teerawutgulrag, A.; Rakariyatham, N. Antioxidant and antibacterial activities of *Nepheium lappaceum* L. extracts. *LWT Food Sci Technol* 2008; 41: 2029-2035
10. Widowati W, Ratnawati H, Rusdi UD, Winarno W, Kasim F. The antiplatelet aggregation effect of extract and ethyl acetate fraction of velvet bean seed (*Mucuna pruriens* L.) in dyslipidemic rat. *Agritech* 2011; 31(1):52-59
11. Widowati W, Wijaya L, Wargasetia TL, Yellianty Y. Antioxidant, anticancer, and apoptosis-inducing effects of Piper extracts in HeLa cells. *J Exp Integrat Med* 2013; 3(3):225-230
12. Widowati W, Herlina T, Ratnawati H, Mozef T. Antioxidant activities and platelet aggregation inhibitor of black tea (*Camellia sinensis* L.) extract and fractions. *Med Plants* 2011;3(1):21-26.

13. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin J Sci Technol* 2004; 26(2): 211-219.
14. Kim YM, Wang MH, Rhee HI. A novel α -glucosidase inhibitor from pine bark. *Car Res* 2004; 339:715-717
15. Widowati W, Ratnawati H, Retnaningsih CH, Lindayani L, Rusdi DU, Winarno W. Free radical scavenging and α -glucosidase inhibitor activity of *Mucuna pruriens* L. *JFI* 2011; 5(3):117 -124.
16. Halliwell B, Gutteridge JMC. *Free radicals in biology and medicine*. New York: Oxford University Press: 1999.
17. Sairam RK, Srivastava GC, Agarwal S, Meena RC. Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes. *Biologia Plantarum* 2005; 49(1): 85-91
18. Thitilertdecha N, Teerawutgulrag A, Rakariyatham N. Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. *Food Sci Tech* 2008; 41: 2029-2035
19. Samuagam L, Sia CM, Akowuah GA, Okechukwu PN, Yim HS. The effect of extraction conditions on total phenolic content and free radical scavenging capacity of selected tropical fruits' peel. *Health Environ J* 2013; 4(2):80-102.
20. Lim YY, Murtijaya J. Antioxidant properties of *Phyllanthus amarus* extracts as affected by different drying methods. *LWT-Food Sci Tech*. 2007; 40: 1664-1669.
21. Thitilertdecha, N., Teerawutgulrag, A., Rakariyatham, N. Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. *Swiss Soc Food Sci Technol*. 2008
22. Widowati W, Sardjono CT, Wijaya L, Laksmiawati DR, Darsono L. Free radicals scavenging activities of spices and curcumin. *Proceedings 2nd Int Symposium on Temulawak*. ISBN 978-979-25-1209-0; 2011:178-181
23. Pujimulyani D, Wazyka A, Anggrahini S, Santoso U. Antioxidative properties of white saffron extract (*Curcuma mangga* Val) in the β -carotene bleaching and DPPH-radical scavenging methods. *Indonesian Food Nut Prog* 2004; 11(2):35-40.
24. Yang XN, Kang SC. In vitro antioxidant activity of the water and ethanol extracts of *Forsythia koreana* flowers. *Nat Prod Res* 2012; 26(4):375-379.
25. Sirisompong W, Jirapakkul W, Klinkesorn U. Response surface optimization and characteristics of rambutan (*Nephelium lappaceum* L.) kernel fat by hexane extraction. *LWT - Food Sci Tech* 2011; 44:1946-1951
26. Issara U, Zzaman W, Yang TA. Rambutan seed fat as a potential source of cocoa butter substitute in confectionary product. *Int Food Res Tech* 2014; 21(1): 25-31.
27. Harahap SN, Ramli N, Vafaei N, Said M. Physicochemical and nutritional composition of rambutan (*Nephelium lappaceum* L) seed and seed oil. *Pakistan J Nut* 2011; 1073-1077.
28. Febrianto NA, Abdullah WNW, Yang TA. Effect of fermentation time and roasting process on the antioxidant properties of rambutan (*Nephelium lappaceum*) seed fat. *Archives Des Sci* 2012; 65: 694-702.
29. Rahayu L, Zakir L, Keban SA. The Effect of rambutan seed (*Nephelium lappaceum* L.) infusion on blood glucose and pancreas histology of mice induced with alloxan. *JFI* 2013;11(1):28-35.
30. Palanisamy U, Manaharan T, Teng LL, Radhakrishnan AKC, Subramaniam T, Masilamani T. Rambutan rind in the management of hyperglycemia. *Food Res Int* 2011; 44 (2011): 2278-2282.
31. Tiwari AK, Rao JM: Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects. *Curr Sci* 2002; 83:30-38.
32. Kim J-S, Kwon C-S, Son K-H: Inhibition of α glucosidase and amylase by luteolin, a flavonoid. *Biosci Biotechnol Biochem* 2000; 64:2458-2461.