



## Antioxidant, Anticancer and Antibacterial Activity of *Withania somnifera* Aqueous Root Extract

D. A. Barnes<sup>1</sup>, R. Barlow<sup>1</sup>, P. Singh Nigam<sup>1</sup> and R. Owusu-Apenten<sup>1\*</sup>

<sup>1</sup>School of Biomedical Sciences, University of Ulster, Coleraine, Cromore Road, Coleraine BT52 1SA, UK.

### Authors' contributions

This work was carried out in collaboration between all authors. Authors ROA and PSN jointly designed the study and wrote the protocol. Authors DAB and RB performed all experiments and data analysis. Author DAB wrote the 1<sup>st</sup> draft. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/JABB/2016/22523

#### Editor(s):

(1) Joana Chiang, Department of Medical Laboratory Science and Biotechnology, China Medical University, Taiwan.

#### Reviewers:

(1) Abdullahi Hassan Kawo, Bayero University, Kano, Nigeria.

(2) Rodrigo Crespo Mosca, Sao Paulo University, Brazil.

Complete Peer review History: <http://sciencedomain.org/review-history/12257>

Short Research Article

Received 7<sup>th</sup> October 2015  
Accepted 29<sup>th</sup> October 2015  
Published 10<sup>th</sup> November 2015

### ABSTRACT

**Aims:** To evaluate total antioxidant capacity, anticancer activity and antibacterial effects *Withania somnifera* aqueous-root extracts.

**Study Design:** *In vitro* study.

**Place of Study:** School of Biomedical Sciences, Ulster University, UK.

**Methodology:** Total antioxidant capacity (TAC) of whole powder and freeze dried *W. somnifera* aqueous-root extracts was determined using FRAP, DPPH, Folin and ABTS assays. Anticancer activity was accessed using MDA-MB-231 breast cells and Sulforhodamine B staining for cell viability. Antibacterial activity was by disk diffusion assay with penicillin, amoxicillin and streptomycin as positive controls.

**Results:** The TAC for *W. somnifera* extract was 86, 47, 195, or 443 gallic acid equivalents per 100g dry basis (mgGAE/ 100 g) using FRAP, DPPH, Folin or ABTS assays, respectively. Corresponding TAC values for freeze dried *W. somnifera* aqueous-root extract were, 418, 553, 1898 or, 1770 (mgGAE/100 g). *W. somnifera* aqueous-root extract inhibited MDA-MB-231 cell proliferation in a dose-dependent manner with IC<sub>50</sub> = 0.19 mg/ml (21 µM GAE). Nil antibacterial effects were detected for freeze dried *W. somnifera* extract (0-1 mg/ml) across six species of bacteria tested.

\*Corresponding author: E-mail: [r.owusu-apenten@ulster.ac.uk](mailto:r.owusu-apenten@ulster.ac.uk);

**Conclusion:** *Withania somnifera* root water extract showed significant antioxidant and anticancer activity for MDA-MB-231 breast cancer cells but no antibacterial activity under the conditions of this study.

**Keywords:** *Withania somnifera*; antioxidant capacity; anticancer; antibacterial.

## 1. INTRODUCTION

*Withania somnifera* (Ashwaganda, Indian ginseng or Winter cherry) is a member of the Solanaceae family found naturally in Africa, Asia (temperate/ tropical) and Europe [1]. Health uses of *W. somnifera* can be traced from 5000BCE to Hindu Ayurveda medical practices. Recent chemical analysis showed diverse components, including alkaloids, steroidal lactones, and flavonoids; [2-5]. The in-vitro antioxidant activity of *W. somnifera* roots extracts were determined as part of routine investigations [6-9] with both extracts and isolated components reported to inhibit breast cancer cells [10-14]. The antimicrobial activity of *W. somnifera* extracts received increasing interest over the recent past as shown by representative studies [7,15-19].

Currently, a few studies examined the anticancer effect of root extracts *W. somnifera* [20-22] and comparatively less research dealt with aqueous extracts [23]. To our knowledge, no publication has compared the antioxidant, anticancer and antimicrobial activity for *W. somnifera* within a single study. To fill the perceived research gap, the aims of this study were to evaluate *W. somnifera* aqueous root extracts, for antioxidant, anticancer and antibacterial effects *in-vitro*.

## 2. MATERIALS AND METHODS

### 2.1 Materials and Extract Preparation

USDA approved organic *W. somnifera* root powder was purchased from a licensed herbal dealer. Chemicals and reagents including 2, 2-diphenyl-1-picrylhydrazyl, Folin Denis reagent and 2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) were obtained from Sigma-Aldrich. Nutrient broth and nutrient agar were both purchased from Oxoid. Aqueous root extract was prepared by adding 1 g of whole *W. somnifera* root powder to 19 ml of water, and stirring for 1 hour, centrifuging (13000 rpm for 20 minutes), freezing at -80°C overnight and freeze drying.

### 2.2 In vitro Assay for Total Antioxidant Capacity

Assays for total antioxidant capacity (TAC) were adapted to microplate analysis as described previously [24]. The ferric Reducing Ability of Plasma (FRAP) assay was adapted from [24] and samples were read at 593nm in a microplate reader. The Folin assay for total phenol was performed as described in [25] and read at 760nm using a microplate reader. The DPPH (2, 2-diphenyl-1-picrylhydrazyl) assay was adapted from [26] and samples were read at 515nm using a microplate reader. The ABTS (2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) assay was adapted from [27] and the final solution mixtures measured at 734nm using a microplate reader. All assays were calibrated with 0-3 mM gallic acid as reference antioxidant.

### 2.3 Anticancer Assay

MDA-MB-231 breast cancer cells were grown in DMEM with 10% FBS, 1% PenStrep and 1% non-essential amino acids. Trypsinized cells were plated (10,000 per well) and allowed attach overnight. Freeze dried *W. somnifera* (5 mg/ml) in complete culture medium was filter sterilized and 1:5 serial dilutions were added to cells for 72 hrs. Cells viability was determined using sulforhodamine B staining [28].

### 2.4 Antibacterial Methods

Three Gram positive (*B. cereus*, *S. aureus*, *S. epidermis*), and Gram negative (*Salmonella typhi*, *E. coli* and *Pseudomonas aeruginosa*) with verified, identity and phenotypes were used. Antimicrobial activity was evaluated by disk diffusion assay as described by [17]. Control antibiotic disks contained either streptomycin, amoxicillin or, penicillin; two blank discs were placed, at equal distances on the agar plate. Freeze dried plant extract (1 mg/ml) was added to discs one drop at until the disk was saturated.

### 2.5 Statistical Analysis

Statistical analysis was carried out using Microsoft excel and the IBM SPSS v22 statistic

package for Microsoft Windows for ANOVA analysis. Significance level was at ( $P=0.05$ ) across all tests.

### 3. RESULTS AND DISCUSSION

#### 3.1 Total antioxidant Capacity Assay

The total antioxidant capacity (TAC) for *W. somnifera* root aqueous-extract and the freeze dried extract (Table 1) reflected the method of analysis (ABTS > Folin > FRAP ≥ DPPH) and values were significantly different for all assays ( $p=0.05$ ). The total phenols values (95 & 1898 mg GAE/100 g) for whole water extract and freeze dried extract, compare with 60 -1107 mg GAE/100 g reported previously [6,7]. The total phenols value for the *W. somnifera* root extracted with 80% hot methanol was 600 mg GAE/100 g [9]. The FRAP values (86 - 418 mg GAE/100 g) compares well with 167-200 mg GAE/100 g (converted from 500-600 mg trolox equivalent/ 100 g) reported for *W. somnifera* methanolic extract [9]. ABTS analysis for *W. somnifera* root water-extract produced a TAC of 218 mgGAE/100 g (from 938 mg trolox equivalents/ 100 g) [7] compared with 443 and

1770 mg GAE/100 g from this study. In summary, the TAC for *W. somnifera* is influenced by the extraction conditions (solvent used, temperature and time of extraction), plant maturity and botanical part of the plant selected. *W. somnifera* leaves had 3-4 times higher phenols and TAC compared with roots for the fully mature plant [9] whilst other investigators found no systematic difference [8,17]. The present paper provides a unique comparison of TAC for the *W. somnifera* root extracts using four different assays, which is important because different antioxidant methods employ different mechanisms [29].

#### 3.2 Anticancer Activity towards MDA-MB-231 Breast Cancer Cells


Breast cancer is the most common cancer among women globally with 1.67 million new cases diagnosed in 2012 and accounting for 25% of all cancers [30]. *W. somnifera* leaf extracts and isolated components were reported to inhibit breast cancer cells [10-14] but few studies examined the anticancer effect of root extracts [20-22] and studies using aqueous-extracts are limited [23]. In the present study (Fig. 1), freeze

**Table 1. Antioxidant activity for whole powder (WP) and freeze dried *Withania somnifera* aqueous extracts across four *in-vitro* antioxidant assays**

Assay*	LDR (µM)	$\epsilon$ ( $M\text{ cm}^{-1}$ )	$R^2$	TAC ‡(WP)	TAC‡(FD)
FRAP	0-25	195014	0.997	86±11	418 ±12
DPPH	0-25	92886	0.989	47±8	553.1±31
Folin Denis	0-75	25836	0.981	195±11	1898±57
ABTS	0-20	50642	0.988	443±28	1770±57

Results from 3-replicates,  $n \geq 48$  data points, WP = whole powder, FD = freeze dried extract

\*LDR =Range of concentrations or which calibration graph is straight,  $\epsilon$  ( $M\text{ cm}^{-1}$ ) = molar absorptivity,  $R^2$  = Correlation coefficient. ‡Total antioxidant capacity (TAC) expressed as gallic acid equivalents (GAE) per 100g dry weight. Within each column all TAC are significantly different at ( $P=0.05$ )




**Fig. 1. Effect of different doses of aqueous extracts from *W. somnifera* root powder (0.004, 0.02, 0.1, 0.5, 2.5 mg/ml) on MDA-MB-231 cell viability with 72hr treatment**  
 Bars shows means (+/- SEM) for 3 independent experiments ( $n= 18$ ) per treatment, Different letters on each column show significant differences ( $P=0.05$ )

dried *W. somnifera* aqueous root extracts were found to be cytotoxic to MDA MB 231 breast cancer cells with 50% inhibitory concentration (IC<sub>50</sub>) at 0.19 mg/ml for 72 hr. treatment. Cell viability was just over 10% at the highest concentration used. By comparison, a previous report found the IC<sub>50</sub> was 0.48 mg/ml and 0.032 mg/ml for aqueous and ethanoic extracts from the roots of *W. somnifera* tested with MDA-MB-231 cell line though there were detailed differences in experimental method [23]. There is general agreement that both leaf and root extracts from *W. somnifera* possess anticancer activity [31]. Interestingly a previous study showed that leaf and root extract had similar anticancer activity though more exhaustive studies are needed [14].

Whether the anticancer and TAC activity of *W. somnifera* are related is uncertain To address this issue, we estimated the IC<sub>50</sub> value for freeze dried *W. somnifera* extract in terms of total phenols content of samples ( IC<sub>50</sub> (mol/l) = 0.19 (gGAE/l) \* 18.89 (gGAE/g sample)/ 170.1 (g/mol)) which is 21 μM GAE. Interestingly, the IC<sub>50</sub> for pure gallic acid for MDA-MB-231 cells was 47 μM with 24 hr treatment [32].

### 3.3 Antibacterial Activities

Previous investigations found, that *W. somnifera* extracts show antimicrobial activity towards both Gram negative and Gram positive strains but, that the degree of effectiveness varied with plant components (leaf > root extracts), extraction solvent (methanol, chloroform > water > petroleum ether) and dose of bioactive agent applied [7,15-19]. The present study (Fig. 2) show that *W. somnifera* aqueous-root extract had no antimicrobial effects at the dose tested (1mg/ml; 1mg addition/disc). Of particular note, previous investigations using disc diffusion testing required a concentration range of 20-100 mg/ml of extract with 4-5 mg solid per disc or well [16,17]. A methanolic *W. somnifera* root extract (5mg/ml and 200 μl or 1 mg addition per disc) showed antimicrobial activity but lower concentrations were not tested [8]. Aqueous and chloroform extracts of *W. somnifera* fruit extracts (10 mg/ml; 100 μl or 1 mg addition per disc) showed antimicrobial results [33]. Finally, recent investigations using 100 mg/ml extract (10 mg addition/ disc) found aqueous, ethanol and acetone extracts of *W. somnifera* had broadly similar antimicrobial activity [17,34].



**Fig. 2. Zones of inhibition for disc diffusion assay of *Withania somnifera* extract using Gram +ve (*B. cereus*, *S. aureus*, *S. epidermis*) and Gram -ve (*Salmonella typhi*, *E. coli* and *Pseudomonas aeruginosa* 17) bacteria**

Extract 1 mg/ml (1 mg/ disk), 3 control antibiotics and PBS as an additional control. Error bars are standard deviation. 6 mm is size of antibiotic disk. Number of replicates n=2

#### 4. CONCLUSION

*Withania somnifera* aqueous root extract was demonstrated to have potent antioxidant activity. Freeze drying the *W. somnifera* root extract further increased the TAC per dry weight basis by 5-11 fold and is expected to also preserve antioxidant activity. Aqueous-root extracts *W. somnifera* showed anticancer activity (IC<sub>50</sub>~0.19 mg/ml) but no antibacterial ability was detectable at such doses. This preliminary study is the first to compare, antioxidant, anticancer and antimicrobial activity of *W. somnifera* extracts. The results, supported by emerging literature, suggest that the effective dose for *W. somnifera* anticancer activity is 1-order of magnitude lower compared to concentrations for antimicrobial activity according to the strain of bacterial tested in this study. Further research is required to improve current understanding of the anticancer and antimicrobial properties *Withania*.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. USDA ARS National Genetic Resources Program. Germplasm Resources Information Network - (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland. Taxon: *Withania somnifera* (L.) Dunal; 2015. Available:<http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?102407> (Cited 2015 May).
2. Bano A, Sharma N, Dhaliwal HS, Sharma VA. Systematic and comprehensive review on *Withania somnifera* (L.) Dunal-An Indian ginseng. Br J Pharm Res. 2015; 7(2):63-75.
3. Dar NJ A, Hamid A, Ahmad M. Pharmacologic overview of *Withania somnifera*, the Indian Ginseng. Cell Mol Life Sci. 2015;1-16.
4. Vanden Berghe W, Sabbe L, Kaileh M, Haegeman G, Heyninck K. Molecular insight in the multifunctional activities of Withaferin A. Biochem Pharmacol. 2012; 84(10):282-91.
5. Mishra LC, Singh BB, Dagenais S. Scientific basis for the therapeutic use of *Withania somnifera* (Ashwagandha): A review. Altern Med Rev. 2000;5(4): 334-346.
6. Sumathi S, Padma PR. Antioxidant status of different parts of *Withania somnifera*. Plant Archives, 2008;8(1):69-72.
7. Mehrotra V, Mehrotra S, Kirar V, Shyam R, Misra K, Srivastava AK, Nandi SP. Antioxidant and antimicrobial activities of aqueous extract of *Withania somnifera* against methicillin-resistant *Staphylococcus aureus*. J. Microbiol. Biotechnol. Res. 2011;1:40-45.
8. Alam N, Hossain M, Mottalib MA, Sulaiman SA, Gan SH, Khalil MI. Methanolic extracts of *Withania somnifera* leaves, fruits and roots possess antioxidant properties and antibacterial activities. BMC Complement Altern Med. 2012;12:175.
9. Fernando IDNS, Abeysinghe DC, Dharmadasa RM. Determination of phenolic contents and antioxidant capacity of different parts of *Withania somnifera* (L.) Dunal. From three different growth stages. Ind Crops Prod. 2013;50:537-539.
10. Jayaprakasam B, Zhang YJ, Seeram NP, Nair MG. Growth inhibition of human tumor cell lines by withanolides from *Withania somnifera* leaves. Life Sci. 2003;74(1): 125-132.
11. Kaur K, Rani G, Widodo N, Nagpal A, Taira K, Kaul SC, Wadhwa R. Evaluation of the anti-proliferative and anti-oxidative activities of leaf extract from *in vivo* and *in vitro* raised Ashwagandha. Food Chem Toxicol. 2004;42(12):2015-20.
12. Subbaraju GV, Vanisree M, Rao CV, Sivaramakrishna C, Sridhar P, Jayaprakasam B, Nair MG. Ashwagandhanolide, a bioactive dimeric thiowithanolide isolated from the roots of *Withania somnifera*. J Nat Prod. 2006; 69(12):1790-2.
13. Stan SD, Zeng Z, Singh SV, Withaferin A. A constituent of Indian Ayurvedic medicinal plant *Withania somnifera*, causes G2 phase and mitotic arrest in human breast cancer cells. Nutr Cancer. 2008; 60(Suppl 1):51-60.
14. Yadav B, Bajaj A, Saxena M, Saxena AK. *In vitro* anticancer activity of the root, stem and leaves of *Withania somnifera* against various human cancer cell lines. Ind J Pharm Sci 2010;72(5):659-U321.
15. Jaffer HJ, Jawad A-LM, Saber HS, Al-Naib A. Evaluation of antimicrobial activity of

- Withania-somnifera* extracts. Fitoterapia. 1988(6):497-500.
16. Arora S, Dhillon S, Rani G, Nagpal A. The in vitro antibacterial/synergistic activities of *Withania somnifera* extracts. Fitoterapia. 2004;75(3):385-388.
  17. Owais M, Sharad KS, Shehbaz A, Saleemuddin M. Antibacterial efficacy of *Withania somnifera* (Ashwagandha) an indigenous medicinal plant against experimental murine salmonellosis. Phytomedicine. 2005;12(3):229-35.
  18. Datta S, Kumar Pal NK, Nandy AK. Inhibition of the emergence of multi drug resistant *Staphylococcus aureus* by *Withania somnifera* root extracts. Asian Pac J Trop Med. 2011;4(11):917-20.
  19. Sundaram S, Dwivedi P, Purwar S. *In vitro* evaluation of antibacterial activities of crude extracts of *Withania somnifera* (Ashwagandha) to bacterial pathogens. Asian J Biotech. 2011;3(2):194-199.
  20. Khazal K, Samuel T, Hill DL, Ge X, Grubbs C. Effect of an extract of *Withania somnifera* on human breast cancer Cells. Faseb J. 2012;26:852.12
  21. Khazal KF, Samuel T, Hill DL, Grubbs CJ. Effect of an extract of *Withania somnifera* root on estrogen receptor-positive mammary carcinomas. Anticancer Res. 2013;33(4):1519-23.
  22. Khan MA, Ahmad R, Trivedi A, Srivastava AN. Determination of combined effect of pH and *Withania somnifera* on human breast cancer cell line MDA-MB-231. Austin J Cancer Clin Res. 2015;2(6):1050.
  23. Maliyakkal N, Udupa N, Pai KS, Rangarajan A. Cytotoxic and apoptotic activities of extracts of *Withania somnifera* and *Tinospora cordifolia* in human breast cancer cells. IJARNP. 2013;6(4):1-10.
  24. Bolanos de la Torre AA, Henderson T Nigam PS, Owusu-Apenten RK. A universally calibrated microplate Ferric Reducing Antioxidant Power (FRAP) assay for foods and applications to Manuka honey. Food Chem. 2015;174:119-23.
  25. Ainsworth EA, Gillespie KM. Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. Nat Protoc. 2007; 2(4):875-7.
  26. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. LWT - Food Sci Technol. 1995;28(1):25-30.
  27. Walker RB, Everette JD. Comparative reaction rates of various antioxidants with ABTS radical cation. J Agric Food Chem. 2009;57(4):1156-61.
  28. Vichai V, Kirtikara K. Sulforhodamine B colorimetric assay for cytotoxicity screening. Nature Protocol. 2006;1(3): 1112-6.
  29. Thaipong K, Boonprakob U, Crosby K, Cisneros-Zevallos L, Byrne DH. Comparison of ABTS, DPPH, FRAP and ORAC assays for estimating antioxidant activity from guava fruit extracts. J Food Comp Anal. 2006;19(6-7):669-675.
  30. Cancer Research UK. Breast cancer statistics; 2015. Available:<http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/breast-cancer#heading-Zero> (Cited 2015 Aug)
  31. Rai M, Jogee PS, Agarkar G, Santos CA. Anticancer activities of *Withania somnifera*: Current research, formulations and future perspectives. Pharm Biol. 2015;1-9.
  32. Devi YP, Uma A, Narasu ML. Anticancer activity of gallic acid on cancer cell lines, HCT15 and MDA MB 231. IMPACT: IJRANSS. 2014;2(5):269-272.
  33. Singariya P, Mourya KK, Kumar P. Antimicrobial activity of the crude extracts of *Withania somnifera* and *Cenchrus setigerus* *In-vitro*. Pharmacogn J. 2012; 4(27):60-65.
  34. Al-Ani N, Hadi SA, Nazar R. Antimicrobial activities of *Withania Somnifera* crude extract. Sci. Agri. 2013;4(3):74-76.

© 2016 Barnes et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
 The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/12257>