





Research Article

Antioxidant Effect of a Polyphenol-Rich Murtilla (*Ugni molinae* Turcz.) Extract and Its Effect on the Regulation of Metabolism in Refrigerated Boar Sperm

Ignacio Jofré ¹, Magdalena Cuevas ², Leticia Signori de Castro,³
João Diego de Agostini Losano ⁴, Mariana Andrade Torres,⁵ Marysol Alvear,⁶
Erick Scheuermann,⁷ André Furugen Cesar Andrade,⁵ Marcilio Nichi ⁴,
Mayra Elena Ortiz Assumpção ³, and Fernando Romero ¹

¹Laboratory of Neurosciences and Biological Peptides (BIOREN-CEGIN-CEBIOR-UFRO) Medicine Faculty, Universidad de La Frontera, Francisco Salazar, 01145 Temuco, Chile

²Department of Physiology, University of Concepción, PO Box 160-C, Concepción, Chile

³Laboratory of Spermatozoa Biology, Department of Animal Reproduction, School of Veterinary Medicine and Animal Science, University of Sao Paulo, Avenida Prof. Dr. Orlando Marques de Paiva 87, Cidade Universitária, 05508-270 Sao Paulo, SP, Brazil

⁴Laboratório de Andrologia, Departamento de Reprodução Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, Brazil

⁵Swine Research Center, School of Veterinary Medicine and Animal Science, University of São Paulo, Pirassununga, São Paulo, Brazil

⁶Department of Chemical Sciences and Natural Resources, Faculty of Engineering and Sciences, Universidad de La Frontera, Av. Francisco Salazar, 01145 Temuco, Chile

⁷Department of Chemical Engineering, Universidad de La Frontera, Francisco Salazar, 01145 Temuco, Chile

Correspondence should be addressed to Fernando Romero; fernando.romero@ufrontera.cl

Received 30 September 2018; Revised 22 March 2019; Accepted 28 April 2019; Published 3 June 2019

Academic Editor: Joël R. Drevet

Copyright © 2019 Ignacio Jofré et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The production of reactive oxygen species (ROS) in boar spermatozoa increases in refrigeration; this can have an impact on sperm quality and fertilization capacity. We evaluated the effect of polyphenol-rich aqueous extract of murtilla (*Ugni molinae* Turcz) on boar sperm stored at 17°C in order to reduce oxidative stress and improve sperm quality in the long term. Five experiments were performed: first, characterization of the polyphenol content from five genotypes of murtilla; second, determination of the genotype with the best antioxidant effect (MT-Ex); third, the antioxidant capacity on O₂⁻ and lipid peroxidation; fourth, the influence of MT-Ex on motility, calcium movement, cAMP, and metabolic parameters; and fifth, analysis of long-term refrigeration. The average phenolic content was 344 ppm; gallic acid, catechin, quercetin, myricetin, and kaempferol were detected. All extracts evaluated presented a concentration-dependent antioxidant effect. MT-Ex reduces intracellular O₂⁻/peroxides but low lipid peroxidation. MT-Ex in nonstimulated ROS conditions reduces sperm motility, mitochondrial membrane potential, cAMP, and ATP, but the succinate dehydrogenase activity remained normal; also, we observed a reduction in calcium movement in *in vitro* sperm capacitation. The long-term analyses showed that MT-Ex improved sperm motility decay and reduced membrane damage and ROS at 168 h. Based on this study, we propose MT-Ex as a supplement in semen extenders.

1. Introduction

The world pork market grows larger every year. Pork is considered a good and cheap alternative source of animal

protein and a rich source of B complex vitamins. In the swine reproduction industry, artificial insemination (AI) is the best system for increasing the number of animals and decreasing the interval between generations, a combination of factors

which improves breeding programs. One of the most used systems in the swine industry is the refrigeration of semen dosage and it is recommended to maintain sperm viability during transportation/storage, and its efficiency is determined by many factors including the medium, the semen extender used, and the refrigeration time and temperature (usually 17°C) [1]. Nevertheless, this process leads to a decrease in fertility rates due to reduced sperm motility and viability [2] and increased production of reactive oxygen species (ROS) [3–5]. The purpose of an extender is to maintain the stability of the spermatozoa during the refrigeration period, reducing the number of dead spermatozoa as a result of cold stress and osmotic variations [6]. In this sense, the study of oxidative stress in boar semen allows evaluating the effects of antioxidants in the long term unlike other biological models.

Boar spermatozoa have been described as very sensitive to thermal shock because the membrane has a high content of polyunsaturated fatty acids [7]. In addition, low temperatures (9–15°C) generate a decrease in membrane phospholipid phases and damage the membrane proteins, affecting sperm viability [8] and increasing their sensitivity to lipid peroxidation. ROS have physiological functions at low concentrations, such as sperm activation [9], hyperactivation [10], and acrosome reaction [11]; at high levels, however, they may activate capacitation and reduce motility [12], generating oxidation and DNA fragmentation [13].

Antioxidants are compounds that control the effects of ROS and protect the cells against massive oxidation. Nonenzymatic antioxidants reduce ROS and indirectly protect the membrane from lipid peroxidation [14, 15]; these antioxidants, which may be found in extracellular milieu [15], can neutralize the hydroxyl radical and superoxides and prevent sperm agglutination [16]. Studies have demonstrated that supplementation of the semen extender with antioxidants (enzymatic and nonenzymatic), such as α -tocopherol, butylated hydroxytoluene, superoxide dismutase, catalase, cysteine, and glutathione, generates an improvement in boar sperm quality [17–21]. Molecules found in berries containing high concentrations of phenols and anthocyanin have been described as potential reducing agents against oxidative stress under *in vitro* conditions. Murtilla or murta (*Ugni molinae* Turcz.), a plant of the Myrtaceae family, is endemic to Chile and Argentina; its fruit and leaves are rich in phenolic components [22–26]. Some reports indicate that these flavonoids have a broad antioxidant activity, intervening in the signalling in cancer, cardiovascular, neuroprotection, and anti-inflammatory pathways [27]. The aims of the present work were to compare the antioxidant effect of the extracts from five different murtilla genotypes; to select the genotype with the best antioxidant capacity; to evaluate the influence of murtilla extract (MT-ex) on sperm parameters such as ATP/cAMP content, SDH activity, and ROS production; and to explore its long-term effect at 196 h, this mechanism being still unknown.

2. Materials and Methods

2.1. Bioethics Procedures and Ethics Statement. This study was carried out at the Center of Biotechnology in Reproduction,

Faculty of Medicine, and the Scientific and Technological Bioresource Nucleus of La Frontera University, Chile (CEBIOR-BIOREN-CEGIN-UFRO). The study and the experimental methods were approved by the Scientific Ethics Committee (CEC) of La Frontera University and performed under the guidelines and regulations of this entity (Protocol 102/13, Acta N80/2013).

2.2. Reagents. Folin-Ciocalteu, Na_2CO_3 , FeSO_4 , gallic acid, formic acid, acetonitrile, H_2O_2 , $[\text{Fe}(\text{CN})_6]^{3-}$, and NaOH were obtained from Merck. Luminol (3-aminophthalhydrazide, 5-amino-2,3-dihydro-1,4-phthalazinedione) and ATP were obtained from Sigma-Aldrich. Phosphate-buffered saline (PBS), Hank's buffered saline solution, and all fluorescent dyes were obtained from Thermo Fisher.

2.3. Plant Material and Extraction Procedure. Fresh fruits of five genotypes of murtilla (*Ugni molinae* Turcz.) (G14-4, G19-1, G22-1, G23-1, and G27-1) were obtained from the Agricultural Research Institute, Vilcún, Chile (INIA Carillanca). The fruits were harvested in April 2012 at the INIA Carillanca experimental station located close to Puerto Saavedra, Araucanía Region, Chile (38°45'S, 73°21' W). Six grams of fruit of each genotype was ground in a mortar and transferred to a bottle containing 20 mL⁻¹ of prewarmed (30°C) distilled water. The mixture was shaken in an incubator (GFL 3032, Germany) at 170 rpm, 30°C, for 20 min and vacuum-filtered (Whatman N° 1 filter paper). The aqueous extract was stored under refrigeration, protected from light and oxygen, until each analysis.

2.4. Experiment 1: Chemical Characterization

2.4.1. Determination of Total Polyphenol Content. The total polyphenol content was determined using the Folin-Ciocalteu method as described by Alfaro et al. [28]. Aqueous murtilla fruit extract (40 μL) was mixed with distilled water (3.16 mL) in a test tube, and then 200 μL of Folin-Ciocalteu reagent was added. After 5 min at 20°C, 600 μL of 20% Na_2CO_3 was added to the reaction mixture, which was maintained at 20°C for 120 min in the dark. The absorbance was measured at 765 nm using a spectrophotometer (Spectronic Genesys 5, Sweden), and the results were expressed as μg of gallic acid equivalent per mL ($\mu\text{gGAE mL}^{-1}$) of aqueous extract.

2.4.2. Phenolic Compound Concentration by HPLC Analysis. Phenolic compounds were identified from the five murtilla genotypes using the Merck Hitachi High-Performance Liquid Chromatography system (LaChrom, Tokyo, Japan) coupled to an L-7100 pump and L-4250 UV-VIS detector. A 5 μm C18 RP Inertsil ODS-3 column (GL Sciences Inc., Tokyo, Japan) was used with a 250 mm \times 4.60 mm i.d., maintained at 25°C. The original extract was concentrated to dryness using a rotary evaporator (Büchi R-210, Germany) at 140 rpm, 30°C, and then resuspended in 5 mL⁻¹ of methanol:formic acid (99:1, v/v). The sample extract was filtered through a 0.45 μm filter, and 20 μL was injected for polyphenol analysis. The identification of compounds was confirmed both by comparison of their retention time

with pure standards and by coinjection. A linear gradient solvent system consisting of 1% formic acid (A) and acetonitrile (B) was used at a flow rate of 1 mL⁻¹ min as follows: 0-2 min, 100% A; 2-15 min, 80% A/20% B; 15-20 min, 70% A/30% B; 20-30 min, 40% A/60% B; and 30-35 min, 100% A. Phenolic compounds were detected at 280 nm [28].

2.5. Experiment 2: Antioxidant Capacity of the Extracts. The second experiment was carried out in two assays: (1) antioxidant screening to determine which of the 5 murtilla genotype extracts presented the best antioxidant capacity in refrigerated sperm samples stored in the MR-A semen extender and (2) determining the influence of time and concentration of antioxidant response under refrigeration (17°C), to reveal the maximum and minimum time limits to induce the antioxidant effect using a mathematical model to explain the antioxidant behaviour of the murtilla genotype extract selected in the first assay (MT-Ex).

2.5.1. Semen Processing. Sperm samples from ten boars (5 collections each) aged 15 to 28 months from Sociedad Agrícola y Ganadera Pehuén Ltda. (Victoria, Chile) were used in experiments 2, 3, and 4. The sperm samples were collected weekly from each boar using the gloved-hand technique. Semen samples were purified, and the sperm-rich fraction was placed in an MR-A (Minitube®) semen extender prewarmed to 37°C; the concentration was adjusted to 25 × 10⁶ cells/mL and refrigerated at 17°C. In attempt to normalize the possible individual variations, we performed seminal analysis of fresh ejaculate from the 10 boars, which included total motility (>70%), concentration, and morphology (<20% abnormal spermatozoa), and according to the parameters established by the Brazilian College of Animal Reproduction, all of them were approved by these criteria.

2.5.2. Oxidative Stimulation in Boar Sperm. The ideal oxidative stimulation condition was performed with H₂O₂ as a prooxidant. The stimulation was performed with 122.7 μM, during 6 h of incubation according to a standard curve of stimulation (data not shown). All stimulated samples were compared with a control without stress stimulation (without H₂O₂).

2.5.3. Antioxidant Screening. For antioxidant screening, we used the luminol assay [29] modified which evaluates oxidation in the whole sample, where in the presence of luminol and iron the sample generates a luminescence directly proportional to the oxidation status. Samples with 25 × 10⁶ cells (1 mL of final volume) were incubated with 122.7 μM of H₂O₂ for 6 h to stimulate the oxidant condition; then, the samples were washed by centrifugation (300g, 10 minutes) to remove the H₂O₂ in excess and diluted in 1 mL of MR-A. Concentrations of the five different murtilla genotypes were then added (0.0001 to 100 μgGAE mL⁻¹), and the samples were incubated for 6 h at 17°C. At the end of the incubation period, each reaction (150 μL) was centrifuged at 700g for 10 min, the supernatant was discarded, and the pellet was suspended in 300 μL of Luminol (600 μM in PBS 1x) and incubated for 15 min at 37°C in a Luminometer microplate reader (Luminoskan,

Thermo Scientific). Then, 25 μL of the developed solution was added using an automated dispenser (0.1 M of [Fe(CN)₆]³⁻ in 0.1 M of NaOH, pH 8) and immediately measured with an integration of 1000 milliseconds (ms). The data were compared with a control treated with H₂O₂ and without antioxidant treatment. Transformation of the X-axis to log₁₀ was performed followed by a nonlinear regression (dose response) to determine the EC50 (effective concentration to reduce stress by 50%) of the antioxidant effect of each extract evaluated. The extract which presented the lowest dispersion of data was selected as the best antioxidant; subsequent experiments were executed with the murtilla extract selected, G14-4, hereafter “MT-Ex”.

2.5.4. Time and Concentration Effect of MT-Ex. The method was performed using 25 × 10⁶ cells in MR-A semen extender. Each sample was incubated with 122.7 μM of H₂O₂ for 6 h at 17°C to stimulate ROS production, then centrifuged at 700g for 10 min. The resulting pellets were resuspended in the MR-A semen extender supplemented with 0.0001–100 μgGAE mL⁻¹ of MT-Ex and incubated for 30 min to 6 h at 17°C. The antioxidant effect was evaluated following the Luminol protocol described above.

2.6. Experiment 3: Effect of MT-ex on Oxidative Stress and Sperm Motility. To detect the superoxide anion and lipid peroxidation, we used 25 × 10⁶ cells in the MR-A semen extender, divided into 4 groups: “control”—sperm samples incubated in the MR-A semen extender, “oxidant”—sperm samples treated with an oxidant agent, “MT-Ex”—sperm samples treated with 0.0315 μgGAE mL⁻¹ as antioxidant treatment (concentration obtained in experiment 2), and “MT-Ex + oxidant”—sperm samples coincubated with oxidant agent and 0.315 μgGAE mL⁻¹ as antioxidant treatment. The oxidant agent used was 122.7 μM of H₂O₂ to stimulate superoxide anion/hydrogen peroxide production and 100 μM of FeSO₄ to stimulate lipid peroxidation. All groups were incubated for 30 minutes at 17°C.

2.6.1. Intracellular Superoxide-Anion Reduction. The capacity of the selected extract to reduce superoxide anion and hydrogen peroxide was measured using dihydroethidium (DHE) as a fluorescent indicator of ROS in combination with SYTOX Green as a membrane damage indicator [30]. At the end of incubation, all groups were washed using MR-A with pre-warming and centrifuged for 10 minutes at 900g. Staining was performed with 2.2 μM of DHE and 0.04 μM of SYTOX Green for 20 minutes at 37°C in the MR-A semen extender. At the end of incubation, the samples were suspended in 300 μL of prewarmed PBS (1x) and immediately analysed in a flow cytometer (FACSCanto II, BD) using the spectra 488/583 (DHE) and 488/523 (SYTOX Green) analysing 10,000 events per sample. The data was analysed in Flowing Software 2.0, gating the SYTOX Green negative events (viable cells); the population was then assessed with high DHE fluorescence (viable cells with high ROS production).

2.6.2. Lipid Peroxidation. Membrane lipid peroxidation was determined using the BODIPY C-11 fluorescent sensor, which exhibits a red basal fluorescence in the nonoxidized

state, while oxidation shows a green fluorescence. The method was carried out according to Aitken et al. with some modifications [31]. Thirty minutes before oxidant and antioxidant treatment, the sperm samples were incubated with 100 nmol of BODIPY C-11 in MR-A for 30 minutes at 37°C. The samples were washed twice with prewarmed MR-A and centrifuged at 700g for 10 minutes to remove unbound BODIPY C-11 residues. The same samples were divided into the 4 treatment groups described above. At the end of treatment incubation, each sample was washed with PBS 1x and immediately analysed in a microplate reader (Synergy HT, BioTek) to detect the mean fluorescence using the wavelength 488/590 for the nonoxidized fraction and 488/530 for the oxidized fraction according to Drummen et al. [32].

2.7. Experiment 4: Effect of Murtilla Extract on Sperm Metabolism. The same groups described in experiment 3 were used in this experiment. We characterised sperm motility, succinate dehydrogenase activity, mitochondrial membrane potential, and ATP content.

2.7.1. Sperm Motility. Sperm motility evaluation was performed in two steps: first, an assay to determine the concentration effect of MT-Ex on sperm motility. The MR-A semen extender was supplemented with 0 to 10 µgGAE mL⁻¹, then 25 × 10⁶ cells/ml were added to the solution and incubated for 30 minutes at 17°C. The second step was carried out with the same groups described in experiment 3. In both cases, the temperature was increased to 37°C five minutes before the end of incubation to recover sperm motility. Two microliters of sample was loaded in a slide chamber (Leja), and the motile, static, and progressive populations were analysed (%) in Automatic Sperm Analyzer System according to the following parameters: 45 frames at a frame rate of 60 frames/s; minimum contrast = 46; minimum cell size = 7 pixels; motility > 45 µm/s; progressive motility > 45 µm/s; and straightness > 45%.

2.7.2. Mitochondrial Membrane Potential (MMP). The mitochondrial membrane potential was evaluated by a JC-1 probe [33] determining the sperm population with basal mitochondrial membrane potential (MMP) in oxidant and antioxidant conditions. At the end of the treatment, 2 × 10⁵ cells were incubated with 76.6 µg/mL of JC-1 and 0.5 µM propidium iodide for 5 minutes at 37°C. At the end of incubation, the cells were suspended in 300 µL of prewarmed PBS (1x) and immediately analysed in a flow cytometer (FACSCanto II BD) analysing 10,000 events per sample.

2.7.3. Succinate Dehydrogenase Activity and MTT Reduction Assay. The succinate dehydrogenase activity assay was based on reduction of tetrazolium salts to insoluble formazan in the presence of succinate by the mitochondrial enzyme succinate dehydrogenase, which allowed the metabolic rate related to aerobic respiration to be determined [34]. Each sample of intact cells was incubated with MTT (3.5 µg/mL) for 1 hour at 37°C; at the end of incubation, 100 µL of DMSO was added to lysate cells, the formazan crystals were solubilized, and the reaction was stopped. The

colorimetric reaction was measured in a microplate reader using an absorbance of 514 nm (Synergy HT, BioTek). The data was compared with the nontreated group (100% normal enzymatic activity).

2.7.4. ATP Content. The influence of MT-Ex on ATP content was measured using a luminescence-dependent indicator (CellTiter, Promega) [35]. At the end of incubation, 100 µL of each sperm sample (5 × 10⁵ cells) was incubated with 100 µL of the ATP probe mixture following the manufacturer's recommendations. The solution was incubated for 10 minutes in a shaker (180 rpm), and the luminescence was immediately measured in a luminometer (Luminoskan, Thermo Scientific) with integration of 1000 ms. The luminescence measured was compared with the standard curve for ATP (0.0001 to 1000 nmol), based on a linear regression, and the data were interpreted as nmol of ATP in 1 × 10⁶ cells.

2.7.5. cAMP and Calcium Movement. cAMP was determined using the commercial kit cAMP-Screen Direct (Invitrogen) [36]. At the end of incubation, the sperm samples were centrifuged at 700g for 10 minutes and washed with HBSS 1x (Ca²⁺- and Mg²⁺-free). cAMP was measured following the manufacturer's instruction for suspended cells. The data were compared with a standard curve for cAMP (0.0006 to 6000 pmol), analysed by sigmoidal equation, and interpreted as pmol of cAMP in 1 × 10⁶ cells.

The transient calcium was measured to determine the effect of MT-Ex on the capacitation state of sperm samples. The measurement method used was the green fluorescent probe FLUO-4 AM (Invitrogen) to detect the calcium signal [37]. First, buffers were prepared to stimulate sperm capacitation following Supplementary Table S1. Before stimulation, the sperm samples (25 × 10⁶ cells) were incubated with 1 µM of FLUO-4 AM (without pluronic acid F-127) for 45 minutes at 37°C in the MR-A semen extender. The samples were divided into 4 groups and washed twice (700g, 10 minutes) to remove the unbound stain. Each group was diluted in each prepared buffer and maintained at 17°C until analysis. Then 15 × 10⁶ cells were placed in a microplate (with optical glass bottom), and the fluorescence was recorded for 50 min at 60-second intervals using a microplate reader (BioTek Synergy HT) in the 488/523 spectrum.

2.7.6. TUNEL Assay. The In Situ Cell Death Detection kit (Roche, Mannheim, Germany), which is based on the terminal deoxynucleotidyl transferase- (TdT-) mediated dUTP nick-end labeling (TUNEL) technique, was used to analyse the sperm DNA fragmentation. At the end of the incubation, 5 × 10⁶ cells were recuperated of each treatment and incubated with 2 mmol⁻¹ of dithiothreitol (1 mL final volume) for 45 min at room temperature in order to relax the chromatin and allow the enzyme better access to the DNA [38]. Subsequently, the cells were washed twice with PBS, fixed in 2% paraformaldehyde for 15 min at 4°C, and permeabilized with 0.1% Triton X-100 for 10 min at room temperature. Then, the sperm were incubated with the TUNEL reaction solution and incubated for 1 h at 37°C. Finally, the cells were washed twice, resuspended in DPBS, and analyzed

by flow cytometry. The results were expressed as the percentage of FITC-positive cells.

2.8. Experiment 5: Long-Term Refrigeration. In this experiment, long-term analyses were performed to evaluate sperm motility, viability, mitochondrial membrane potential, and ROS and O_2^- production. The samples were divided into two groups: control—semen diluted in the MR-A semen extender; MT-Ex—semen diluted in the MR-A semen extender and supplemented with $0.315 \mu\text{gGAE mL}^{-1}$ of MT-Ex. Both groups were refrigerated at 17°C for 168 h and evaluated at 24 h intervals.

2.8.1. Semen Processing. Five mature boars ($n = 15$; 5 animals \times 3 collections), aged 15 to 28 months, were used in this study; they were obtained from the Swine Research Center, School of Veterinary Medicine and Animal Science, University of São Paulo, Pirassununga, São Paulo, Brazil. The sperm samples were collected weekly from each boar using the gloved-hand technique. Semen samples were purified, and the sperm-rich fraction was placed in the MR-A extender supplemented with $0.315 \mu\text{gGAE mL}^{-1}$ prewarmed to 37°C and then refrigerated at 17°C (25×10^6 cells/ml). In attempt to normalize the possible individual variations, we performed seminal analysis of fresh ejaculate from the 10 boars, which included total motility ($>70\%$), concentration, and morphology ($<20\%$ abnormal spermatozoa), and according to the parameters established by the Brazilian College of Animal Reproduction, all of them were approved by these criteria.

2.8.2. Sperm Motility. Motility parameters were evaluated using the Computer Assisted Sperm Analysis (CASA) system (IVOS, v. 12.2, Hamilton Thorne Research, Beverly, MA). The parameters used were 45 frames at a frame rate of 60 frames/s; minimum contrast = 46; minimum cell size = 7 pixels; motility $> 45 \mu\text{m/s}$; progressive motility $> 45 \mu\text{m/s}$; and straightness $> 45\%$. In brief, each slide was heated to 37°C ; $7 \mu\text{L}$ of the sample was placed in the slide and covered with a coverslip. A minimum of six fields were selected for analysis. The data obtained to observe differences between control and MT-Ex were the total motile population.

2.8.3. Sperm Characteristics. Aliquots of 2×10^5 cells were obtained every 24 h to evaluate mitochondrial membrane potential (JC-1), and production of superoxide anion/hydrogen peroxides (DHE) following the procedures described above. One extra protocol was performed in this experiment, to determine ROS production: determination of total ROS by CellROX Green/PI [39]. Aliquots of 2×10^5 cells were stained with $5 \mu\text{M}$ of CellROX Green for 30 minutes at 37°C ; in the last 10 minutes, PI was added to a final concentration of $6 \mu\text{M}$. All fluorescent reactions were performed in a Guava easyCyte flow cytometer (Merck Millipore) analysing 10,000 events per sample using a 488 laser and fluorescence channels according to the manufacturer's recommendations.

2.9. Statistical Analysis. For experiment 1, the data were analysed using one-way analysis of variance (ANOVA). The values obtained at $P < 0.05$ were considered significant. The

differences between means were determined using Tukey's multiple comparison tests. The results were expressed as the means of the measurements and their corresponding standard deviations. For experiment 2, the concentration-response analysis was performed by polynomial regression to determine EC50. The EC50 obtained for each genotype was subjected to analysis of variance (ANOVA), and the significance of the difference between means was determined by Tukey's test ($P < 0.05$). The Time and Concentration analysis was evaluated by ANOVA to determine the interaction between variables. Once the absence of the interaction between variables was verified, a prediction analysis of the surface plot and equation that describes the mathematical model was carried out using the software Curve Expert (Microsoft 2.2). The equation was solved by log transformation of the concentration parameter and full cubic polynomial regression. The EC50 analysis for each time was performed by polynomial regression using GraphPad Prism. For experiment 3, evaluation of O_2^- /peroxide production and lipid peroxidation was subjected to analysis of variance (ANOVA) and the significance of the difference between means was determined by Tukey's test ($P < 0.05$). For experiment 4, sperm motility data were analysed by log transformation of the concentration variable and sigmoidal nonlinear regression. The mitochondrial membrane potential, SDH activity, ATP, and cAMP content were subjected to analysis of variance (ANOVA), and the significance of the difference between means was determined by Tukey's test ($P < 0.05$). Calcium analysis was performed by linear regression comparing the linear trends. For experiment 5, long-term analysis was performed by calculation of means and S.D., and statistical analysis of the results was performed using GraphPad Prism 5.0. Analysis of repeated measures of variance was done to compare semen motility, mitochondrial membrane potential, viability, ROS production, and mitochondrial O_2^- between the control and MT-Ex groups and over time, followed by a Bonferroni significant difference test (to locate differences). Values were considered significant when $P < 0.05$.

3. Results

3.1. Experiment 1: Chemical Characterization. The total polyphenol content in MT-Ex ranged from 85.5 to $406.5 \mu\text{gGAE mL}^{-1}$; the highest value was obtained from genotype 19-1 (Table 1). Phenolic compounds identified by HPLC analysis were gallic acid, catechin, quercetin-3- β -D-glucoside, myricetin, quercetin, and kaempferol; their concentrations for the five genotypes are shown in Table 1. The flavonol catechin was the most abundant constituent in our samples with 2.69, 0.521, 10.01, 2.28, and $1.81 \mu\text{g/mL}$ in the G14-4, G19-1, G22-1, G23-1, and G27-1 genotypes, respectively. The concentrations of gallic acid, quercetin-3- β -D-glucoside, myricetin, quercetin, and kaempferol in MT-Ex ranged from 0.009 to $3.21 \mu\text{g/mL}$.

3.2. Experiment 2: Extract Antioxidant Capacity

3.2.1. Antioxidant Screening. We evaluated the antioxidant potential of fruit extract from five *Ugni molinae* genotypes,

TABLE 1: Total polyphenol content (TPC, $\mu\text{gGAE mL}^{-1}$) in murtilla (*Ugni molinae* Turcz.) fruit aqueous extract obtained from genotypes INIA 14-4, 19-1, 22-1, 23-1, and 27-1. Phenolic compound concentration ($\mu\text{g/mL}$) in murtilla (*Ugni molinae* Turcz.) fruit aqueous extract obtained from each genotype determined by HPLC analysis. Values are mean \pm standard deviation.

	Genotype				
	14-4	19-1	22-1	23-1	27-1
TPC ($\mu\text{gGAE mL}^{-1}$)	85.5 \pm 0.7	406.5 \pm 0.2	314.5 \pm 0.4	285.2 \pm 0.3	264 \pm 0.3
Gallic acid ($\mu\text{g mL}^{-1}$)	0.059 \pm 0.02	0.013 \pm 0.01	n.d.	n.d.	n.d.
Catechin ($\mu\text{g mL}^{-1}$)	2.696 \pm 0.7	0.521 \pm 0.7	10.015 \pm 0.9	2.287 \pm 1.1	1.812 \pm 0.9
Quercetin-3- β -D-glucoside ($\mu\text{g mL}^{-1}$)	0.141 \pm 0.1	3.211 \pm 0.3	0.074 \pm 0.01	0.131 \pm 0.05	0.050 \pm 0.02
Myricetin ($\mu\text{g mL}^{-1}$)	0.115 \pm 0.001	0.026 \pm 0.009	0.083 \pm 0.004	0.119 \pm 0.01	0.026 \pm 0.000
Quercetin ($\mu\text{g mL}^{-1}$)	0.009 \pm 0.001	0.471 \pm 0.2	0.007 \pm 0.002	n.d.	0.010 \pm 0.001
Kaempferol ($\mu\text{g mL}^{-1}$)	0.014 \pm 0.008	0.022 \pm 0.006	0.035 \pm 0.008	0.010 \pm 0.001	0.025 \pm 0.003

n.d.: not determined.




FIGURE 1: Antioxidant activity of MT-Ex. (a) Extracellular antioxidant effect of five genotypes of MT-Ex (log of 0.0001 to 100 $\mu\text{gGAE mL}^{-1}$) in sperm cells pretreated with 122.7 μM of H_2O_2 as oxidant agent. The EC50 for each genotype indicates the effective concentration to reduce oxidative stress by 50%. (b) Time and concentration evaluation of antioxidant activity of murtilla genotype 14-4 8 (MT-Ex) in sperm samples pretreated with H_2O_2 as oxidant treatment.

using H_2O_2 as prooxidant agent to reveal the minimum and maximum concentrations of antioxidant effect. In all genotypes evaluated, the results showed a concentration-dependent effect when the antioxidant concentration increased (Figure 1(a)). We observed similar tendencies in the sigmoidal curves and EC50 calculated for each genotype: 16.6 $\mu\text{gGAE mL}^{-1}$ (G14-4), 28.9 $\mu\text{gGAE mL}^{-1}$ (G19-1), 15.3 $\mu\text{gGAE mL}^{-1}$ (G22-1), 23.2 $\mu\text{gGAE mL}^{-1}$ (G23-1), and 24.8 $\mu\text{gGAE mL}^{-1}$ (G27-1). The average of the five genotypes was 21.76 $\mu\text{gGAE/mL}^{-1}$. Genotype G14-4 was selected as the ideal extract in the characterization and antioxidant analysis because of the low dispersion of the data in the curve (R^2 : 0.995). The results below were performed using only G14-4, hereafter "MT-Ex".

3.2.2. Time and Concentration Effects of MT-Ex. Evaluation of the surface plot revealed that the effect was concentration-dependent (Figure 1(b)) and partially time-dependent. This profile is explained in a mathematical model according to the calculated equation (1) where x_1 corresponds to time (in hours) and x_2 to the log of the concentration in $\mu\text{gGAE mL}^{-1}$.

$$y = 22.09 - 16.79x_1 + 3.91x_2 + 3.43x_1^2 + 0.2x_2^2 + 0.72x_1^3 + 0.05x_2^3. \quad (1)$$

The EC50 calculated for each hour showed an effect indirectly proportional to the time of antioxidant treatment:

the EC50 values at 30 minutes and 1, 2, 4, and 6 h were 0.315, 0.055, 0.030, 0.003, and 0.006 $\mu\text{gGAE mL}^{-1}$, respectively (Table 2). We determined that the optimum concentration and time, with low variation of effective concentration for the increase in exposure time, was 0.315 $\mu\text{gGAE mL}^{-1}$ for 30 minutes. This combination was used in subsequent analyses.

3.3. Experiment 3: Effect of Murtilla Extract on Sperm Oxidative Stress Parameters

3.3.1. Intracellular Superoxide-Anion Reduction. The EC50 which produced an antioxidant effect (0.315 $\mu\text{gGAE mL}^{-1}$ for 30 min) on O_2^- production was evaluated by the DHE probe. As expected, the oxidant group presented the highest percentage of population with high O_2^- production ($93.6\% \pm 4.1$). This was followed by the MT-Ex + oxidant group ($37.67\% \pm 5.6$) and the control ($30.3\% \pm 3.2$) with no statistical difference between them. The lowest percentage (statistically significant difference) of viable cells with high superoxide production ($19.3\% \pm 4.9$) was found in the antioxidant group (MT-Ex) (Figure 2(a)).

3.3.2. Lipid Peroxidation. Lipid peroxidation was evaluated by the BODIPY C-11 probe; we observed a different response profile between treatment groups when compared to superoxide-anion reduction. In this case, the oxidant and MT-Ex + oxidant groups presented the highest fluorescence intensity (4938 ± 455.2 and 3796 ± 147.2), while the control and MT-Ex groups presented lower fluorescence intensities (2921 ± 64.6 and 2879 ± 44.4 , respectively) (Figure 2(b)).

3.4. Experiment 4: Effect of MT-Ex on Sperm Metabolism

3.4.1. Sperm Motility. The effect of MT-Ex on sperm motility was evaluated in two steps. First, we examined the effect of increasing concentrations of MT-Ex on sperm motility during refrigerated incubation: we observed a concentration-dependent decrease in total motility and a consequent increase in static cells, without any significant change in the progressive population (Figure 3).

In the second trial, we determined the influence of MT-ex on sperm motility in oxidant and antioxidant conditions. A reduction in sperm motility was observed in all groups treated. As expected, the oxidant group resulted in a reduction in the motile sperm population from $89.6\% \pm 3.1$ (control) to $56.3\% \pm 2.6$ (oxidant); however, treatments with MT-Ex on cells without oxidative stimulation resulted in a reduction in sperm motility ($65.3\% \pm 1.5$), while the addition of MT-Ex to oxidized sperm did not improve the motility parameter; on the contrary, a substantial reduction was observed with respect to the oxidant group ($46.3\% \pm 1.9$) (Table 3).

3.5. Mitochondrial Membrane Potential (MMP). The mitochondrial membrane potential was evaluated with the JC-1 probe as a measure of the influence of MT-Ex on spermatozoa in oxidant and antioxidant conditions. We observed that the cell count with basal MMP in the control group was the highest with 88.6%. The addition of H_2O_2 as an oxidative

TABLE 2: Predicted EC50: predicted EC50 for each time analysed according to sigmoidal dose-response analysis.

Time (hrs)	EC50 MT-Ex G.14-4 ($\mu\text{gGAE mL}^{-1}$)	SD
0.5	0.315	0.094
1	0.055	0.019
2	0.030	0.014
4	0.003	0.009
6	0.006	0.009

The data shown represent the effective concentration to reduce the oxidative stress by 50% at each hour evaluated.

stimulator reduced this parameter, independently of the presence or absence of MT-Ex, with similar results between groups (34.1% and 32.7%, respectively). However, the addition of MT-Ex in nonoxidized sperm presented a higher percentage of basal MMP (65.7%) when compared to groups with H_2O_2 , but still lower than in the control group (Figure 4(a)).

3.6. Succinate Dehydrogenase and MTT Reduction Assay. The evaluation of succinate dehydrogenase activity showed that the presence of MT-Ex resulted in no significant differences compared to control (96.3%). However, for the oxidant group, the percentage of cells was reduced to 41.5%, while in the MT-Ex + oxidant group, the MMP did not present substantial improvements (Figure 4(b)).

3.7. ATP Content. The effect of MT-Ex on ATP content was evaluated under different incubation conditions. We observed a reduction in ATP content in all treatments under oxidant and antioxidant conditions. The ATP content in the group treated with MT-Ex without oxidant stimulation was reduced from 2.29 ± 0.61 nmol (control group) to 1.06 ± 0.23 nmol (MT-Ex group). On the other hand, the ATP contents in the groups treated with H_2O_2 (oxidant) and with MT-Ex and H_2O_2 (MT-Ex + oxidant) were reduced by 0.20 ± 0.03 nmol and 0.23 ± 0.04 nmol, respectively (Figure 4(c)).

4. cAMP and Calcium Movement

We evaluated the effect of MT-Ex and H_2O_2 on the cAMP content of sperm samples under refrigeration and subsequently the influence of MT-Ex on calcium movement. The results suggest a reduction in the cAMP content in all the treated groups: in oxidant conditions, the cAMP concentration was reduced, reaching 0.19 pmol ± 0.03 , while in the MT-Ex group the concentration was 0.95 pmol ± 0.21 . However, treatment with MT-Ex did not improve the loss of cAMP (Figure 5(a)) in oxidized cells.

In order to corroborate the effect of cAMP as a physiological indicator of important processes in capacitation and acrosomal reaction, we evaluated the intracellular calcium movement of sperm samples maintained in different buffers. The results suggest that the fluorescent signal of FLUO-4 AM increased in sperm samples in the capacitation buffer, indicating an increase in intracellular calcium, whereas in the




FIGURE 2: Antioxidant effect of MT-Ex. (a) Antioxidant capacity of MT-Ex on peroxides and superoxide anion production. (b) Effect of MT-Ex on lipid peroxidation. In both analyses: “control”—sperm samples incubated in MR-A semen extender; “oxidant”—sperm samples treated with oxidant agent; “MT-Ex”—sperm samples treated with $0.0315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment; and “MT-Ex + oxidant”—sperm samples coincubated with oxidant agent and $0.315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment. The oxidant agents were H_2O_2 and FeSO_4 to produce O_2^- /perox and lipid peroxidation, respectively. Differences observed were $*P < 0.05$ and $**P < 0.01$ with respect to the “oxidant” group.




FIGURE 3: Effect of MT-Ex on sperm motility. Analysis of motile, progressive, and static population treated with MT-Ex for 30 min. A sigmoidal representation of the effect of different concentrations of MT-Ex on sperm motility. Samples treated with MT-Ex were stored at 17°C with MT-Ex and sperm motility was triggered by increasing the temperature to 37°C . Differences observed were $*P < 0.05$ and $**P < 0.01$ with respect to the “0” group.

basal condition without calcium stimulation, a small increase in fluorescence was observed as expected. Surprisingly, the addition of $10 \mu\text{gGAE mL}^{-1}$ of MT-Ex to the capacitation buffer reduced the fluorescence of the calcium indicator significantly as compared to the sperm samples without addition of MT-Ex. The sperm samples maintained in TALP-HEPES and supplemented with MT-Ex presented a similar behaviour. For all cases, it was observed that as time increased, there was an increase in the fluorescent signal; however, the greatest change in fluorescence over time was observed in the basal condition (Figure 5(b)).

4.1. DNA Fragmentation. We evaluated the influence of MT-Ex on DNA fragmentation under basal and oxidant

conditions. Only $3.24 \pm 0.58\%$ of the population showed DNA fragmentation in the basal group, while in oxidant conditions this parameter increased at $7.02 \pm 1.21\%$ ($P < 0.001$). The MT-Ex supplementation in nonoxidizing conditions did not show significant differences with respect to basal conditions, while the antioxidant treatment decreased the count of sperm cells with fragmented DNA with respect to the oxidant group ($P < 0.005$) (Figure 6).

4.2. Experiment 5: Long-Term Effect. Long-term analysis was carried out to determine the antioxidant capacity and verify the stabilizing properties of MT-Ex over time. In the study, it was observed that the addition of MT-Ex to the MR-A semen extender increased the number of motile cells at 168 h ($17.5 \pm 8.3\%$) as compared to the control ($1.3 \pm 1.9\%$). The reduction rates of sperm motility in the two groups were control 12.17% per day and MT-Ex 9.08% per day (Figure 7(a)).

The analysis of sperm viability was performed by a propidium iodide probe, which depends on membrane stability. We observed an increase in cell survival in samples with long exposure times, with significant differences observed between groups at 144 to 168 h. At the last time evaluated, the MT-Ex group presented an undamaged cell count of $17.1\% \pm 3.6$, versus $5.2\% \pm 1.2$ of cells with intact membrane in the control group. The reduction in viability over time was control -15.05% per day and MT-Ex -11.45% per day (Figure 7(b)).

The MMP analysis showed a higher angular coefficient of fall in the control group than in the MT-Ex group (-13.2% and -3.22% per day, respectively), indicating that populations with basal MMP are rapidly reduced over time in the absence of MT-Ex. Thus, the addition of MT-Ex to the MR-A semen extender delayed the loss of membrane potential in the last hours of the analysis ($16.3\% \pm 3.5$), in contrast to the control which presented a substantial decrease in populations with basal potential ($2.4\% \pm 1.4$) (Figure 7(c)).

Oxidative stress production was lower in the samples treated with MT-Ex than in the control. It was observed that the endogenous production of ROS in viable cells after 168 h reached $79.1\% \pm 5.3$ in the presence of MT-Ex while the value

TABLE 3: Effect of sperm motility on sperm samples with and without oxidant stimulation. The groups were “control”—sperm samples incubated in the MR-A semen extender, “oxidant”—sperm samples treated with oxidant agent, “MT-Ex”—sperm samples treated with $0.0315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment, and “MT-Ex + oxidant”—sperm samples coincubated with oxidant agent and $0.315 \mu\text{gGAE mL}^{-1}$ antioxidant treatment. Significant differences are represented for each parameter evaluated independently in treatment groups. Different letters correspond to significant differences between treatments but independently of group motility parameters.

Group	Control	Oxidant	MT-Ex	MT-Ex + oxidant
Static (%)	11.2 ± 1.5^a	42.4 ± 1.3^b	32.9 ± 1.2^c	51.3 ± 1.2^d
Motile (%)	89.6 ± 3.1^a	56.3 ± 2.6^b	65.3 ± 1.5^c	46.3 ± 1.9^d
Progressive (%)	25.6 ± 2.7^a	5.3 ± 1.6^b	21.1 ± 0.9^a	2.4 ± 2.1^b




FIGURE 4: Analysis of metabolic parameters in sperm samples treated with MT-Ex. (a) Effect of MT-Ex on mitochondrial membrane potential. The data shown represent the cell count with normal potential measured by the JC-1 probe. (b) Succinate dehydrogenase activity evaluated by MTT assay. (c) Determination of ATP content. The groups were “control”—sperm samples incubated in the MR-A semen extender, “oxidant”—sperm samples treated with oxidant agent, “MT-Ex”—sperm samples treated with $0.0315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment, and “MT-Ex + oxidant”—sperm samples coincubated with oxidant agent and $0.315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment. Significant differences are represented between control and groups. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$.




FIGURE 5: Analysis of cAMP and calcium movement of sperm samples treated with MT-Ex. (a) cAMP content of sperm samples in oxidant and antioxidant conditions. (b) Evaluation of calcium release of sperm samples in different incubation conditions. Basal: TALP-HEPES; Cap.Buffer: capacitating buffer ($\text{CaCl}_2 + \text{HCO}_3$); MT: TALP-HEPES + $10 \mu\text{gGAE mL}^{-1}$ of MT-Ex; MT+Cap.Buffer: capacitating buffer + $10 \mu\text{gGAE mL}^{-1}$ of murtilla. Significant differences are represented between basal and groups. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. $N = 5$.

in the control was 89.8 ± 2.6 (Figure 7(d)). ROS production increased over time in both groups, in the control group by 5.56% per day and in the MT-Ex group by 2.1% per day.

Antioxidant treatment with MT-Ex reduced the normal increase in ROS production in the longest refrigeration time (168 h). On the other hand, the evaluation of mitochondrial




FIGURE 6: Analysis of DNA fragmentation in sperm samples treated with MT-Ex. The data shown represent the cell count with FITC + fluorescence. The groups were “control”—sperm samples incubated in the MR-A semen extender, “oxidant”—sperm samples treated with oxidant agent, “MT-Ex”—sperm samples treated with $0.0315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment, and “MT-Ex + oxidant”—sperm samples coincubated with oxidant agent and $0.315 \mu\text{gGAE mL}^{-1}$ as antioxidant treatment. Significant differences are represented between control and groups. * $P < 0.005$, ** $P < 0.01$, and *** $P < 0.001$.

O_2^- production showed a reduction in this parameter in both groups, with no significant difference when the storage time was increased (Figure 7(e)).

5. Discussion

The ability of antioxidant agents to control the harmful effects of reactive oxygen species on spermatozoa has been extensively investigated, to further their use as the main regulators of these oxidant molecules. However, the main focus of the present study was on sperm quality rather than the influence of antioxidant agents on metabolic functionality and sperm physiology [40–43]. Here, for the first time we evaluated the antioxidant and metabolic effects of murtilla as an antioxidant supplement in boar MR-A semen extender in an attempt to improve sperm refrigeration based on the behaviour of sperm metabolism.

In previous studies, various genotypes evaluated presented differences in antioxidant activity measured by the DPPH test [22, 23]. These variations in the composition and concentration of phenolic compounds normally involve distinct biological effects due to synergic actions of certain compounds. The total polyphenol contents determined for murtilla fruit aqueous extract from five genotypes (G14-4, G19-1, G22-1, G23-1, and G27-1) in the present study were lower than in the murtilla fruit aqueous extract from three other locations in Chile with different climatic conditions [44]. The phenolic compounds identified (Table 2) have commonly been found in murtilla fruit and leaves [22–26, 45]. The flavonol catechin, which is the most abundant constituent, has been considered a valuable chemical marker for antioxidant activity in naturally occurring agents [44].

Boonsorn et al. reported that the addition of catechin in an extender for boar spermatozoa could improve sperm motility, viability, and acrosome integrity and reduce malondialdehyde levels during incubation periods. Therefore, catechin may be used as an antioxidant to reduce sperm abnormalities and improve sperm quality in boar semen [46]. On the other hand, results obtained by Tvrda et al. showed that quercetin, another natural flavonol, is able to prevent the decline in spermatozoa viability, functional activity, and antioxidant capacity [47].

Some of these compounds have been reported in fruit extracts from the berries of other endemic Chilean plants (*Berberis microphylla*, *Luma chequen*, *Luma apiculata*, and *Amomyrtus meli*), with antioxidant activity shown by DPPH radical bleaching and FRAP test [48]. In a previous study, we demonstrated the high performance of MT-Ex as an antioxidant in endothelial cells and as a cardioprotective agent which generates an antioxidant effect at concentrations in the range of picograms to micrograms of phenols [49].

Our studies of antioxidant capacity showed that although the domesticated murtilla samples were of different genotypes, all five presented antioxidant potential at similar concentrations, demonstrated by their EC50 values; similarities were found in their gradients and kinetics, but not in the concentrations of phenols detected by HPLC. Measurement of oxidative stress by the Luminol technique was sensitive to differences between the genotypes evaluated; however, the dispersion of the data represented by their biological replicates did not represent a substantial difference between the samples. The use of this technique to evaluate the pharmacokinetics of time and concentration determined through a mathematical model allows the behaviour of antioxidant activity to be assessed and understood exclusively under conditions of refrigeration (17°C) in the MR-A semen extender.

The superoxide anion is known to be a highly reactive molecule, unstable over time. It is one of the main stimulators of oxidative damage [50] and in conjunction with H_2O_2 is one of the initiators of lipid peroxidation in sperm [51]. In our study, we evaluated the antioxidant effect of cells previously treated with H_2O_2 as an oxidizing agent, which were subsequently washed to study the intracellular antioxidant potential of MT-Ex. We observed that the addition of MT-Ex to cells under oxidative stimulation by H_2O_2 significantly reduced the production of O_2^- . However, we observed that the addition of MT-Ex at EC50 concentrations reduced the population with high ROS production by more than 50%, while MT-Ex supplementation in cells without oxidative stimulation did not result in a reduction in the basal oxidation count. Thus, ROS reduction was intrinsically limited to the cells after induced stress, while the oxidative levels of cells with normal endogenous ROS production were not affected, suggesting that MT-Ex reduces the impact of H_2O_2 , when the cells already initiated a process of substantial ROS increase, indicating then that the scavenging effect of MT-Ex should be limited only under conditions where the sample presents a high endogenous production of ROS. However, due to the large conformation of the MT-Ex components, it is not possible to suggest a specific target different to H_2O_2 and O_2^- in ROS reduction.




FIGURE 7: Long-term analysis. Effect of MT-Ex on sperm parameters, from 24 to 168 h of incubation at 17°C. In all analyses, white circles represent the control group—sperm samples stored in the MR-A semen extender, and black circles represent the MT-Ex group—sperm samples stored in the MR-A semen extender supplemented with MT-Ex. (a) Motile sperm population, (b) viable sperm count, (c) sperm population with basal mitochondrial membrane potential (MMP), (d) viable sperm cells with high ROS production, and (e) viable cells with high O₂⁻/peroxide production. Significant differences are represented between control and MT-Ex. **P* < 0.05, ***P* < 0.01, and ****P* < 0.001.

We did not observe a substantial reduction in lipid peroxidation by MT-Ex in sperm samples stimulated by FeSO₄ (Fenton's pathway) or in samples in the basal condition. Studies in rat liver microsomes have shown that the application of high concentrations of flavonoids (75–100 μM) in media supplemented with Fe³⁺ and EDTA favour the production of OH and therefore the triggering of lipid peroxidation; the concentrations of flavonoids applied in our assays were too low to stimulate endogenous production of ROS and induce peroxidation, as postulated by Laughton et al.'s study [52].

On the other hand, evaluation of the effect of MT-Ex on metabolic parameters showed a clear influence of the extract on the normal metabolism of spermatozoa under refrigeration conditions. An effect on MMP was observed, with a substantial reduction in the sperm count with basal MMP in samples without oxidant stimulation, while in spermatozoa under oxidative conditions, treatment with MT-Ex to reduce the oxidizing effect did not prove effective in stabilizing MMP.

Furthermore, we did not observe any regulatory or modulating effects of MT-Ex on succinate dehydrogenase by the MTT assay in either oxidant or antioxidant conditions. This

enzyme (complex II of electron transport chain) is involved in the transformation of succinate to fumarate to generate an influx of electrons during the oxidative phosphorylation process in aerobic respiration [53]. We did not observe that any substantial change in enzyme capacity was mediated by the extract, at least in reduction of tetrazolium salt in formazan crystal, suggesting an independence of action on succinate dehydrogenase.

In the functional context, a study in isolated mitochondria reported important effects of the flavonoids quercetin, taxifolin, catechin, and galangin, which act as inhibitors of the respiratory chain of mitochondria or cause uncoupling [54]. Other authors report an important effect of ATP synthesis in bovine sperm, demonstrating the influence of quercetin, kaempferol, catechin, and epicatechin gallate in inhibiting ATPse/synthase. Quercetin generates an improvement in motility without modifying the ATP content and cellular respiration; however, motility and ATP production decrease after less than 2 hours [55]. We likewise found a substantial reduction of ATP in oxidant and antioxidant conditions. On this basis, we evaluated ATP production as a direct measure of metabolism functionality, especially as it is one of the adenosines involved in sperm motility.

Depression of this adenosine is classically described as the effect of oxidative stress; if ROS are not quickly reduced by antioxidant enzymes, they lead to a decrease in the ATP content due to inefficient phosphorylation [56–58]. We observed that the ATP content was substantially decreased in the presence of H_2O_2 at concentrations of $122.7 \mu M$; however, the application of MT-Ex to sperm samples without ROS stimulation by H_2O_2 generated a significant reduction of ATP (>1 nm of ATP).

Some polyphenols, such as epigallocatechin gallate, epicatechin, quercetin, and kaempferol, have been associated with inhibition of ATP synthase complex (T0), occluding proton movement in subunits T0 and T1 [59]. Other studies have demonstrated the influence of flavonoids on complex I and cytochrome C; modulation of epicatechin, quercetin, and kaempferol is observed in these complexes, retarding H_2O_2 production [60].

It is not possible to generate an efficient balance when an oxidative process is triggered, and this could be directly related to sperm motility. Some studies suggest that the addition of catechin ($50 \mu M$) could improve sperm motility in bovine spermatozoa [46]. In our study, we verified the effect of MT-Ex in boar sperm motility, suggesting that MT-Ex may have induced a reduction of the motile population. Interestingly, one of the chemical compounds of our extract contained catechin. Based on this, we suggest that there may be a synergistic/antagonistic or additive effect of the different components of the extract that is not necessarily present when the compounds are applied in isolation. The effect of the extract on sperm motility may well stimulate different pathways, blocking or activating the molecular machinery involved in sperm motility. In fact, we observed an important modulating effect in calcium movement, with a reduction in the intensity of the FLUO-4 AM indicator; this suggests inhibition/blocking of capacitating activation and reduction of the cAMP count.

Furthermore, it has been reported that polyphenols increase cell viability, decrease intracellular Ca^{2+} levels and ROS formation, and improve mitochondrial membrane potential in cells [61]; however, the mechanisms involved in this process are not completely elucidated, especially in the control of intracellular free calcium according to the time and concentrations of antioxidant treatment [62]. On the other hand, new reports have shown that quercetin (quercetin-3-O-glucoside) is able to interfere with different targets of cAMP signalling [63]. Our results suggest an influence of MT-Ex on calcium movement which may block hypermotility, cAMP-mediated capacitation, and the acrosome reaction, allowing the useful life of the boar sperm sample to be extended. In the same way, these parameters are involved as an important physiological indicator of fertility capacity, as well as the stability of the DNA [64]; in this sense, our results indicate a partial reduction in DNA fragmentation induced by oxidative stimulation and subsequent reduction of this status with MT-Ex, but without reaching the basal levels considered normal for the group without oxidative stimulation, indicating a differential index of protection, probably by the low concentration of MT-Ex reaching the DNA to reduce the oxidization of DNA and subsequent

fragmentation, in comparison with the intra/extracellular milieu, supporting the idea of partial pharmacologic efficacy delimited by the subcellular localization.

On the other hand, our results under long-term refrigeration showed beneficial effects on sperm viability and motility, with improved results in both parameters after long exposure times (168 hrs). This effect has been observed in long-term refrigeration using cysteine as an antioxidant, where motility and cell viability are substantially improved, while adverse effects were observed with tocopherol, which reduces mitochondrial potential, motility, and viability [65, 66]. Our initial results showed a reducing effect of mitochondrial parameters, reflected in the long incubation analyses; however, in contrast to the control, application of the extract led to a smaller loss of potential over time and reduced the production of oxidizing agents, suggesting a positive effect from the formulation of a seminal dose until the end of the analysis. In the incubation period, however, the endogenous production of total O_2^- was not improved by the addition of the antioxidant, suggesting other oxidant species involved in the oxidization of sperm cell in the long term. Recent studies have shown that the glycolytic pathway is able to support ATP production and maintain the sperm in conditions to generate motility after the decoupling of oxidative phosphorylation [35, 67], so this regulation would allow greater efficiency in the refrigeration of boar semen.

6. Conclusion

The five genotypes of murtilla tested contain different phenolic compounds in different concentrations but present the same pattern of antioxidant activity. The antioxidant activity of MT-Ex controls the harmful prooxidant effect of H_2O_2 and reduces the impact of the oxidizing agent on DNA fragmentation; however, it also generates an important decrease in sperm metabolism, especially ATP production. This could directly affect the fertilization potential due to the reduction in sperm motility and the calcium pathway to activate capacitation and subsequent acrosome reaction. Nevertheless, we consider that the metabolic decrease and subsequent modulation of sperm capacitation are an advantage in extending the useful life of boar sperm samples refrigerated at $17^\circ C$.

Data Availability

All data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare no conflicts of interest.

Acknowledgments

The authors are grateful to La Frontera University and University of Sao Paulo, Faculty of Veterinary Medicine, Fondo de Fomento al Desarrollo Científico y Tecnológico-Chile (FONDEF AF-01-1007) and FIA-Fundación para la Investigación Agraria (PYT0035-2013). The PhD fellowship

was supported by CONICYT 21151406 and UFRO, Scientific and Technological Bioresource Nucleus (BIOREN-UFRO).

Supplementary Materials

Supplementary 1: composition of sperm TALP-HEPES and modification of the composition for each treatment. Basal TALP-HEPES maintains the basal condition of sperm samples without generating any physiological modification to induce capacitation. CAP.BUFFER includes 2.0 mM of CaCl_2 for activating capacitation by the calcium pathway. MT corresponds to TALP-HEPES and $10 \mu\text{gGAE mL}^{-1}$ of MT-Ex. MT + CAP.BUFFER contains the same characteristic of capacitating buffer but includes $10 \mu\text{gGAE mL}^{-1}$ of extract to evaluate the calcium response in sperm samples. (*Supplementary Materials*)

References

- [1] A. Lopez Rodriguez, A. Van Soom, I. Arsenakis, and D. Maes, "Boar management and semen handling factors affect the quality of boar extended semen," *Porcine Health Management*, vol. 3, no. 1, p. 15, 2017.
- [2] E. T. Donnelly, N. McClure, and S. E. M. Lewis, "Cryopreservation of human semen and prepared sperm: Effects on motility parameters and DNA integrity," *Fertility and Sterility*, vol. 76, no. 5, pp. 892–900, 2001.
- [3] N. Am-in, R. N. Kirkwood, M. Techakumphu, and W. Tantasuparuk, "Effect of storage for 24 h at 18°C on sperm quality and a comparison of two assays for sperm membrane lipid peroxidation," *Canadian Journal of Animal Science*, vol. 90, no. 3, pp. 389–392, 2010.
- [4] L. Fraser, J. Strzeżek, K. Wasilewska, and C. S. Pareek, "Sperm DNA damage in relation to lipid peroxidation following freezing-thawing of boar semen," *South African Journal of Animal Science*, vol. 47, no. 2, pp. 213–218, 2017.
- [5] A. Kumaresan, G. Kadirvel, K. M. Bujarbaruah et al., "Preservation of boar semen at 18°C induces lipid peroxidation and apoptosis like changes in spermatozoa," *Animal Reproduction Science*, vol. 110, no. 1-2, pp. 162–171, 2009.
- [6] A. K. Blasse, H. Oldenhof, M. Ekhlesi-Hundrieser, W. F. Wolkers, H. Sieme, and H. Bollwein, "Osmotic tolerance and intracellular ion concentrations of bovine sperm are affected by cryopreservation," *Theriogenology*, vol. 78, no. 6, pp. 1312–1320, 2012.
- [7] A. Maldjian, F. Pizzi, T. Gliozzi, S. Cerolini, P. Penny, and R. Noble, "Changes in sperm quality and lipid composition during cryopreservation of boar semen," *Theriogenology*, vol. 63, no. 2, pp. 411–421, 2005.
- [8] A. M. Simpson, M. A. Swan, and I. G. White, "Susceptibility of epididymal boar sperm to cold shock and protective action of phosphatidylcholine," *Gamete Research*, vol. 17, no. 4, pp. 355–373, 1987.
- [9] P. Leclerc, E. De Lamirande, and C. Gagnon, "Regulation of protein-tyrosine phosphorylation and human sperm capacitation by reactive oxygen derivatives," *Free Radical Biology and Medicine*, vol. 22, no. 4, pp. 643–656, 1997.
- [10] E. De Lamirande and C. Gagnon, "A positive role for the superoxide anion in triggering hyperactivation and capacitation of human spermatozoa," *International Journal of Andrology*, vol. 16, no. 1, pp. 21–25, 1993.
- [11] R. J. Aitken, M. Paterson, H. Fisher, D. W. Buckingham, and M. Vanduin, "Redox regulation of tyrosine phosphorylation in human spermatozoa and its role in the control of human sperm function," *Journal of Cell Science*, vol. 108, no. 5, pp. 2017–2025, 1995.
- [12] H. D. Guthrie and G. R. Welch, "Determination of intracellular reactive oxygen species and high mitochondrial membrane potential in Percoll-treated viable boar sperm using fluorescence-activated flow cytometry," *Journal of Animal Science*, vol. 84, no. 8, pp. 2089–2100, 2006.
- [13] S. Lopes, A. Jurisicova, J. G. Sun, and R. F. Casper, "Reactive oxygen species: potential cause for DNA fragmentation in human spermatozoa," *Human Reproduction*, vol. 13, no. 4, pp. 896–900, 1998.
- [14] R. C. Werthmann, M. J. Lohse, and M. Bunemann, "Temporally resolved cAMP monitoring in endothelial cells uncovers a thrombin-induced [cAMP] elevation mediated via the Ca^{2+} -dependent production of prostacyclin," *The Journal of Physiology*, vol. 589, no. 1, pp. 181–193, 2011.
- [15] R. A. Saleh and A. Agarwal, "Oxidative stress and male infertility: from research bench to clinical practice," *Journal of Andrology*, vol. 23, no. 6, pp. 737–752, 2002.
- [16] P. Das, A. R. Choudhari, A. Dhawan, and R. Singh, "Role of ascorbic acid in human seminal plasma against the oxidative damage to the sperms," *Indian Journal of Clinical Biochemistry*, vol. 24, no. 3, pp. 312–315, 2009.
- [17] E. Brzezińska-Ślebodzińska, A. B. Ślebodziński, B. Pietras, and G. Wiczorek, "Antioxidant effect of vitamin E and glutathione on lipid peroxidation in boar semen plasma," *Biological Trace Element Research*, vol. 47, no. 1-3, pp. 69–74, 1995.
- [18] S. Cerolini, A. Maldjian, P. Surai, and R. Noble, "Viability, susceptibility to peroxidation and fatty acid composition of boar semen during liquid storage," *Animal Reproduction Science*, vol. 58, no. 1-2, pp. 99–111, 2000.
- [19] F. J. Pena, A. Johannisson, M. Wallgren, and H. Rodriguez Martinez, "Antioxidant supplementation in vitro improves boar sperm motility and mitochondrial membrane potential after cryopreservation of different fractions of the ejaculate," *Animal Reproduction Science*, vol. 78, no. 1-2, pp. 85–98, 2003.
- [20] J. Roca, M. A. Gil, M. Hernandez, I. Parrilla, J. M. Vazquez, and E. A. Martinez, "Survival and fertility of boar spermatozoa after freeze-thawing in extender supplemented with butylated hydroxytoluene," *Journal of Andrology*, vol. 25, no. 3, pp. 397–405, 2004.
- [21] M. M. Satorre, E. Breininger, M. T. Beconi, and N. B. Beorlegui, " α -Tocopherol modifies tyrosine phosphorylation and capacitation-like state of cryopreserved porcine sperm," *Theriogenology*, vol. 68, no. 7, pp. 958–965, 2007.
- [22] S. Alfaro, A. Mutis, R. Palma, A. Quiroz, I. Seguel, and E. Scheuermann, "Influence of genotype and harvest year on polyphenol content and antioxidant activity in murtilla (*Ugni molinae Turcz*) fruit," *Journal of Soil Science and Plant Nutrition*, vol. 13, no. 1, pp. 67–78, 2013.
- [23] T. R. Augusto, E. S. S. Salinas, S. M. Alencar, M. A. B. R. D'arce, A. C. de Camargo, and T. M. F. de Souza Vieira, "Phenolic compounds and antioxidant activity of hydroalcoholic extracts of wild and cultivated murtilla (*Ugni molinae Turcz.*)," *Food Science and Technology*, vol. 34, no. 4, pp. 667–679, 2014.
- [24] M. Junqueira-Gonçalves, L. Yáñez, C. Morales, M. Navarro, R. A. Contreras, and G. Zúñiga, "Isolation and characterization of phenolic compounds and anthocyanins from murta

- (Ugni molinae Turcz.) fruits. Assessment of antioxidant and antibacterial activity,” *Molecules*, vol. 20, no. 4, pp. 5698–5713, 2015.
- [25] E. Scheuermann, M. Ihl, L. Beraud et al., “Effects of packaging and preservation treatments on the shelf life of murtila fruit (Ugni molinae Turcz) in cold storage,” *Packaging Technology and Science*, vol. 27, no. 3, pp. 241–248, 2014.
- [26] C. Shene, A. K. Reyes, M. Villarroel, J. Sineiro, M. Pinelo, and M. Rubilar, “Plant location and extraction procedure strongly alter the antimicrobial activity of murta extracts,” *European Food Research and Technology*, vol. 228, no. 3, pp. 467–475, 2009.
- [27] P. M. Aron and J. A. Kennedy, “Flavan-3-ols: nature, occurrence and biological activity,” *Molecular Nutrition & Food Research*, vol. 52, no. 1, pp. 79–104, 2008.
- [28] S. Alfaro, A. Mutis, A. Quiroz, I. Seguel, and E. Scheuermann, “Effects of drying techniques on murtila fruit polyphenols and antioxidant activity,” *Journal of Food Research*, vol. 3, no. 5, pp. 73–82, 2014.
- [29] A. Agarwal and A. Majzoub, “Laboratory tests for oxidative stress,” *Indian Journal of Urology*, vol. 33, no. 3, pp. 199–206, 2017.
- [30] R. Mahfouz, N. Aziz, R. Sharma, M. Bykova, E. Sabanegh, and A. Agarwal, “Assessment of intracellular human sperm reactive oxygen species after hydrogen peroxide exposure using four different probes,” *Fertility and Sterility*, vol. 90, pp. S320–S321, 2008.
- [31] R. J. Aitken, J. K. Wingate, G. N. De Iuliis, and E. A. McLaughlin, “Analysis of lipid peroxidation in human spermatozoa using BODIPY C11,” *Molecular Human Reproduction*, vol. 13, no. 4, pp. 203–211, 2007.
- [32] G. P. C. Drummen, L. C. M. van Liebergen, J. A. F. Op den Kamp, and J. A. Post, “C11-BODIPY 581/591, an oxidation-sensitive fluorescent lipid peroxidation probe: (micro)spectroscopic characterization and validation of methodology,” *Free Radical Biology and Medicine*, vol. 33, no. 4, pp. 473–490, 2002.
- [33] T. R. dos Santos Hamilton, C. M. Mendes, L. S. de Castro et al., “Evaluation of lasting effects of heat stress on sperm profile and oxidative status of ram semen and epididymal sperm,” *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 1687657, 12 pages, 2016.
- [34] R. Hosamani, G. Krishna, and Muralidhara, “Standardized Bacopa monnieri extract ameliorates acute paraquat-induced oxidative stress, and neurotoxicity in prepubertal mice brain,” *Nutritional Neuroscience*, vol. 19, no. 10, pp. 434–446, 2016.
- [35] J. D. A. Losano, J. F. Padín, I. Méndez-López et al., “The stimulated glycolytic pathway is able to maintain ATP levels and kinetic patterns of bovine epididymal sperm subjected to mitochondrial uncoupling,” *Oxidative Medicine and Cellular Longevity*, vol. 2017, Article ID 1682393, 8 pages, 2017.
- [36] Y. He, Q. Zou, H. Chen, S. Weng, T. Luo, and X. Zeng, “Lead inhibits human sperm functions by reducing the levels of intracellular calcium, camp, and tyrosine phosphorylation,” *The Tohoku Journal of Experimental Medicine*, vol. 238, no. 4, pp. 295–303, 2016.
- [37] K. R. Gee, K. A. Brown, W.-N. U. Chen, J. Bishop-Stewart, D. Gray, and I. Johnson, “Chemical and physiological characterization of fluo-4 Ca²⁺-indicator dyes,” *Cell Calcium*, vol. 27, no. 2, pp. 97–106, 2000.
- [38] L. A. Mitchell, G. N. De Iuliis, and R. J. Aitken, “The TUNEL assay consistently underestimates DNA damage in human spermatozoa and is influenced by DNA compaction and cell vitality: development of an improved methodology,” *International Journal of Andrology*, vol. 34, no. 1, pp. 2–13, 2011.
- [39] L. S. de Castro, P. M. de Assis, A. F. P. Siqueira et al., “Sperm oxidative stress is detrimental to embryo development: a dose-dependent study model and a new and more sensitive oxidative status evaluation,” *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 8213071, 12 pages, 2016.
- [40] A. Basioura, C. M. Boscos, I. Parrilla, G. Tsousis, and I. A. Tsakmakidis, “Effect of astaxanthin on the quality of boar sperm stored at 17°C, incubated at 37°C or under in vitro conditions,” *Reproduction in Domestic Animals*, vol. 53, no. 2, pp. 463–471, 2018.
- [41] M. J. Maya-Soriano, E. Taberner, M. Sabes-Alsina, M. Piles, and M. Lopez-Bejar, “Absence of beneficial effects on rabbit sperm cell cryopreservation by several antioxidant agents,” *Zygote*, vol. 23, no. 1, pp. 1–10, 2015.
- [42] L. Vichas, I. A. Tsakmakidis, D. Vafiadis, G. Tsousis, E. Malama, and C. M. Boscos, “The effect of antioxidant agents’ addition and freezing method on quality parameters of frozen thawed ram semen,” *Cell and Tissue Banking*, vol. 19, no. 1, pp. 113–121, 2018.
- [43] M. Spinaci, V. Muccilli, D. Bucci et al., “Biological effects of polyphenol-rich extract and fractions from an oenological oak-derived tannin on in vitro swine sperm capacitation and fertilizing ability,” *Theriogenology*, vol. 108, pp. 284–290, 2018.
- [44] A. Rashidinejad, E. J. Birch, and D. W. Everett, “Antioxidant activity and recovery of green tea catechins in full-fat cheese following gastrointestinal simulated digestion,” *Journal of Food Composition and Analysis*, vol. 48, pp. 13–24, 2016.
- [45] M. Rubilar, M. Pinelo, M. Ihl, E. Scheuermann, J. Sineiro, and M. J. Nunez, “Murta leaves (Ugni molinae Turcz) as a source of antioxidant polyphenols,” *Journal of Agricultural and Food Chemistry*, vol. 54, no. 1, pp. 59–64, 2006.
- [46] T. Boonsorn, W. Kongbuntad, N. A. Nakkong, and W. Aengwanich, “Effects of catechin addition to extender on sperm quality and lipid peroxidation in boar semen,” *American-Eurasian Journal of Agricultural and Environmental Science*, vol. 7, no. 3, pp. 283–288, 2010.
- [47] E. Tvrdá, E. Tusimova, A. Kovacik, D. Paal, L. Libova, and N. Lukac, “Protective effects of quercetin on selected oxidative biomarkers in bovine spermatozoa subjected to ferrous ascorbate,” *Reproduction in Domestic Animals*, vol. 51, no. 4, pp. 524–537, 2016.
- [48] A. Brito, C. Areche, B. Sepúlveda, E. Kennelly, and M. Simirgiotis, “Anthocyanin characterization, total phenolic quantification and antioxidant features of some Chilean edible berry extracts,” *Molecules*, vol. 19, no. 8, pp. 10936–10955, 2014.
- [49] I. Jofré, C. Pezoa, M. Cuevas et al., “Antioxidant and vasodilator activity of Ugni molinae Turcz. (Murtila) and its modulatory mechanism in hypotensive response,” *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 6513416, 11 pages, 2016.
- [50] D. B. Zorov, M. Juhaszova, and S. J. Sollott, “Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release,” *Physiological Reviews*, vol. 94, no. 3, pp. 909–950, 2014.
- [51] J. F. Griveau, E. Dumont, P. Renard, J. P. Callegari, and D. Le Lannou, “Reactive oxygen species, lipid peroxidation and enzymatic defence systems in human spermatozoa,” *Journal of Reproduction and Fertility*, vol. 103, no. 1, pp. 17–26, 1995.

- [52] M. J. Laughton, B. Halliwell, P. J. Evans, J. Robin, and S. Hoult, "Antioxidant and pro-oxidant actions of the plant phenolics quercetin, gossypol and myricetin: effects on lipid peroxidation, hydroxyl radical generation and bleomycin-dependent damage to DNA," *Biochemical Pharmacology*, vol. 38, no. 17, pp. 2859–2865, 1989.
- [53] J. Hroudová and Z. Fišar, "Control mechanisms in mitochondrial oxidative phosphorylation," *Neural Regeneration Research*, vol. 8, no. 4, pp. 363–375, 2013.
- [54] D. J. Dorta, A. A. Pigoso, F. E. Mingatto et al., "The interaction of flavonoids with mitochondria: Effects on energetic processes," *Chemico-Biological Interactions*, vol. 152, no. 2-3, pp. 67–78, 2005.
- [55] L. Nass-Arden and H. Breitbart, "Modulation of mammalian sperm motility by quercetin," *Molecular Reproduction and Development*, vol. 25, no. 4, pp. 369–373, 1990.
- [56] A. Agarwal, G. Virk, C. Ong, and S. S. du Plessis, "Effect of oxidative stress on male reproduction," *The World Journal of Men's Health*, vol. 32, no. 1, pp. 1–17, 2014.
- [57] A. K. Bansal and G. S. Bilaspuri, "Impacts of oxidative stress and antioxidants on semen functions," *Veterinary Medicine International*, vol. 2011, Article ID 686137, 7 pages, 2011.
- [58] A. Ferramosca, S. P. Provenzano, D. D. Montagna, L. Coppola, and V. Zara, "Oxidative Stress Negatively Affects Human Sperm Mitochondrial Respiration," *Urology*, vol. 82, no. 1, pp. 78–83, 2013.
- [59] J. Zheng and V. D. Ramirez, "Inhibition of mitochondrial proton F0F1-ATPase/ATP synthase by polyphenolic phytochemicals," *British Journal of Pharmacology*, vol. 130, no. 5, pp. 1115–1123, 2000.
- [60] R. Lagoa, I. Graziani, C. Lopez-Sanchez, V. Garcia-Martinez, and C. Gutierrez-Merino, "Complex I and cytochrome C are molecular targets of flavonoids that inhibit hydrogen peroxide production by mitochondria," *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, vol. 1807, no. 12, pp. 1562–1572, 2011.
- [61] L. Chen, X. Yang, H. Jiao, and B. Zhao, "Tea catechins protect against lead-induced ROS formation, mitochondrial dysfunction, and calcium dysregulation in PC12 cells," *Chemical Research in Toxicology*, vol. 16, no. 9, pp. 1155–1161, 2003.
- [62] Z. Liu, L. P. Zhang, H. J. Ma, C. Wang, M. Li, and Q. S. Wang, "Resveratrol reduces intracellular free calcium concentration in rat ventricular myocytes," *Acta Physiologica Sinica-Chinese Edition*, vol. 57, no. 5, pp. 599–604, 2005.
- [63] B. Pavan, A. Capuzzo, and G. Forlani, "Quercetin and quercetin-3-o-glucoside interact with different components of the camp signaling cascade in human retinal pigment epithelial cells," *Life Sciences*, vol. 121, pp. 166–173, 2015.
- [64] K. Takeda, K. Uchiyama, M. Kinukawa, T. Tagami, M. Kaneda, and S. Watanabe, "Evaluation of sperm DNA damage in bulls by TUNEL assay as a parameter of semen quality," *The Journal of Reproduction and Development*, vol. 61, no. 3, pp. 185–190, 2015.
- [65] F. Chatiza, P. W. Mokwena, T. L. Nedambale, and C. Pilane, "Effect of antioxidants (taurine, cysteine, tocopherol) on liquid preserved Kolbroek boar semen characteristics," *African Journal of Biotechnology*, vol. 17, no. 4, pp. 65–72, 2018.
- [66] T. Vongpralub, P. Thananurak, C. Sittikasamkit et al., "Comparison of Effects of Different Antioxidants Supplemented to Long-Term Extender on Boar Semen Quality Following Storage at 17 Degrees C," *The Thai Journal of Veterinary Medicine*, vol. 46, no. 1, pp. 119–126, 2016.
- [67] J. D. A. Losano, D. S. R. Angrimani, A. Dalmazzo et al., "Effect of mitochondrial uncoupling and glycolysis inhibition on ram sperm functionality," *Reproduction in Domestic Animals*, vol. 52, no. 2, pp. 289–297, 2017.



Hindawi

Submit your manuscripts at
www.hindawi.com

