

Antioxidants for elevated sperm DNA fragmentation: a mini review

Ahmad Majzoub¹, Ashok Agarwal², Sandro C. Esteves³

¹Department of Urology, Hamad Medical Corporation, Doha, Qatar; ²American Center for Reproductive Medicine, Cleveland Clinic, Cleveland, OH, USA; ³ANDROFERT, Andrology and Human Reproduction Clinic, Referral Center for Male Reproduction, Campinas, SP, Brazil

Correspondence to: Ahmad Majzoub. Associate Consultant, Department of Urology, Hamad Medical Corporation, Doha, Po Box 3050, Qatar. Email: dr.amajzoub@gmail.com.

Response to: Cho CL. Commentary: sperm DNA fragmentation testing in action. *Transl Androl Urol* 2017;6:S647-8.

Submitted Jun 27, 2017. Accepted for publication Jun 28, 2017.

doi: 10.21037/tau.2017.07.09

View this article at: <http://dx.doi.org/10.21037/tau.2017.07.09>

We would like to thank Dr. Cho for his insightful commentary on our article discussing the utility of sperm DNA fragmentation (SDF) testing for fertility evaluation (1). The author elegantly summarized the principle concepts of the article and provided his expert views of the topic. Dr. Cho rightly commented that “One of the major huddles of clinical application of a test lies in the lack of treatment modalities. And the development of treatment strategies, e.g., the use of antioxidants and sperm selection methods, must go hand in hand with advancement of sperm DNA testing which was also included in the guideline.” We certainly agree with his view on this point, especially since SDF testing has been criticized based on the unavailability of effective treatment modalities. In reply, we would like to highlight the existing available evidence on antioxidant use in patients with high SDF.

Recently published research has increased our understanding of how SDF affects male fertility (1). Sperm DNA is bound to protamine, allowing it to exist in a compact state. This arrangement helps protect the DNA against the damaging processes that can occur during sperm transport through the male and female reproductive tracts (2). Several factors have been shown to cause SDF; errors in chromatin packaging and remodeling that can occur during spermatogenesis (3), excessive ROS production, decreased seminal antioxidants (4), and other apoptotic events that take place during epididymal transit.

Other causes of SDF include environmental exposure to pollutants and chemo-radiation, conditions such as varicocele and infection and inflammations of the reproductive tract, lifestyle factors such as drug and cigarette use, and patient factors such as advanced age

(5,6). Both *in vitro* and *in vivo* studies have confirmed that elevated SDF may negatively influence fertility by affecting fertilization, early embryo development, implantation, and pregnancy (7-10).

Antioxidants are readily available compounds that can be consumed through diet or as an oral supplement. They are easily accessible and the most common form of therapy prescribed for the treatment of male subfertility (11,12). Although a number of studies have documented an improvement in semen parameters with antioxidant supplementation, properly powered randomized placebo-controlled trials are still needed before establishing their clear role in human reproduction (13,14). A Cochrane review of 48 randomized controlled clinical trials that included 4,179 subfertile men reported considerable variability in the antioxidant effect on semen parameters. However, the review authors reported a statistically significant improvement in the live birth rate (OR, 4.21; 95% CI, 2.08 to 8.51; $P < 0.0001$) and clinical pregnancy rate (OR, 3.43; 95% CI, 1.92 to 6.11; $P < 0.0001$) (15).

In order to study the effectiveness of antioxidant therapy on SDF measures, we conducted a literature search using PubMed as a search engine and the keywords “sperm DNA fragmentation” AND “Antioxidants”. Publications were reviewed selecting only the ones examining SDF measures before and after antioxidant therapy (16-27) (Table 1).

In general, while most reports were small sized and used short treatment protocols, they conveyed a beneficial effect for antioxidants on measures of SDF or semen parameter results (Table 1). Greco *et al.* (21) randomized 64 patients with unexplained infertility and elevated levels of SDF into either a treatment group (1 g vitamin C + 1 g vitamin E

Table 1 Review of studies examining the effect of antioxidant therapy on sperm DNA fragmentation

Study	Supplements (daily intake)	Study design	n	Duration	SDF testing method	Result
Tunc <i>et al.</i> (16)	Vitamin C (100 mg), E (400 IU), Se (26 µg), Zn (25 mg), folic acid (0.5 mg) and garlic (1 mg)	Prospective observational study	50	3 months	TUNEL, CMA3 assay	Significant decrease in SDF [-4% (3.1–3.5%) [†] , P<0.01] and protamine packaging [+ 4.6% (4.4–5.8%) [†] , P<0.01]. No significant improvement in semen parameters
Ménézo <i>et al.</i> (17)	b-carotene (18 mg), vitamins C (400 mg), E (400 mg), Zn (500 µmol) and Se (1 µmol)	Prospective observational study	58	3 months	SCSA	Significant decrease in SDF (-19.1%, P<0.01)
Kodama <i>et al.</i> (18)	Vitamins C (200 mg), E (200 mg), glutathione (400 mg)	Prospective observational study for the treatment group	14	2 months	8-hydroxy-2'-deoxyguanosine	Significant improvement in sperm concentration (+7±1 million/mL, P<0.05). Significant decrease in oxidative DNA damage (-0.4%±0.1%, P<0.05)
Ornu <i>et al.</i> (19)	Group 1: Zn (400 mg); group 2: Zn (400 mg) + vitamin E (20 mg); group 3: Zn (400 mg) + vitamins E (20 mg) + C (10 mg)	Randomized placebo-controlled study	45 (treatment—group 1: n=11; group 2: n=12; group 3: n=14); 8 placebo group	3 months	SCSA	Improvement in SDF measures with treatment*. Significant improvement in sperm motility with the treatment group (group 1: +25%±12%; group 2: +24%±6%; group 3: +25%±9%, P<0.01)
Greco <i>et al.</i> (20)	Vitamin C (1,000 mg) + vitamin E (1,000 mg)	Prospective observational study	29 (patients with high SDF and prior failed ICSI)	2 months	TUNEL	Significant decrease SDF levels (-15.8%±3.6%, P<0.001). Insignificant effect on semen parameters. Significant improvement in ICSI clinical pregnancy (48.2% versus 6.9%, P<0.05) and implantation (19.6% versus 2.2%, P<0.01) rates compared with the pretreatment ICSI outcomes
Greco <i>et al.</i> (21)	Vitamin C (1,000 mg) + vitamin E (1,000 mg)	Randomized placebo-controlled study	64 men with unexplained infertility and high SDF levels	2 months	TUNEL	Significant decrease in SDF levels (-13%±0.5%, P<0.01). Insignificant effect on semen parameters
Martínez-Soto <i>et al.</i> (22)	Docosahexaenoic acid (DHA) (1,500 mg)	Randomized, double blind, placebo-controlled, parallel-group study	57 (32 treatment group and 25 placebo group)	10 weeks	TUNEL	Significant decrease in SDF levels (-17.2%±2.8%, P<0.001) in treatment group vs. (+11.2%±1.9%, P>0.05) in the placebo group. Insignificant effect on semen parameters
Fraga <i>et al.</i> (23)	Vitamin C (250 mg)	Prospective observational study	10	15 weeks	8-hydroxy-2'-deoxyguanosine	Vitamin C depletion/repletion was associated with inverse relationship with seminal vitamin C levels and oxo8dG measures (P<0.01)

Table 1 (continued)

Table 1 (continued)

Study	Supplements (daily intake)	Study design	n	Duration	SDF testing method	Result
Abad et al. (24)	L-carnitine (1,500 mg); vitamin C (60 mg); coenzyme Q10 (20 mg); vitamin E (10 mg); zinc (10 mg); vitamin (200 µg) B9 selenium (50 µg); vitamin B12 (1 µg)	Prospective observational study	20 infertile patients diagnosed with asthenoteratozoospermia	3 months	SCD measurement done following various periods of sperm storage (0, 2, 6, 8 and 24 h) at 37 °C	Significant decrease in SDF levels (at 0 h: -8.4%±6.7%, P<0.01; at 2 h: -8.07%±5.03%, P<0.01; at 6 h: -8.58%±0.81%, P<0.01; at 8 h: -9.03%±2.79%, P<0.01; at 24 h: -20.95%±8.59%, P<0.01). Significant improvement in DDS (-1.6%±0.9%, P=0.04). Significant improvement in sperm concentration (+0.1±6.47 million/mL, P=0.04), motility (+6.5%±2.8%, P=0.02) and morphology (+1.5%±2.4%, P=0.04)
Vani et al. (25)	Vitamin C (1,000 mg) 5 consecutive days in a week	Prospective comparative study	240 (120 men exposed to lead, and 120 healthy human subjects)	3 months	Comet	Significant decrease in alkaline-labile sites and mean tail length of the comet when compared to the control group (P<0.01). Significant improvement in all semen parameters (P<0.01)
Gual-Frau et al. (26)	L-carnitine (1,500 mg); vitamin C (60 mg); coenzyme Q10 (20 mg); vitamin E (10 mg); zinc (10 mg); vitamin B9 (200 µg), selenium (50 µg), vitamin B12 (1 µg)	Prospective observational study	20 infertile men with grade 1 varicocele	3 months	SCD	Significant decrease in SDF levels (-22.1%, P=0.02). 31.3% fewer highly degraded sperm cells (P=0.07), significant increase in sperm concentration (P=0.04)
Piomboni et al. (27)	Beta-glucan (20 mg), fermented papaya (50 mg), lactoferrin (97 mg), vitamin C (30 mg), and vitamin E (5 mg)	Prospective observational study	36 men with leukocytospermia; 15 controls	3 months	Acridine orange (AO) staining	No significant decrease in SDF. Significant increase in sperm morphology (+12.8%±1.3%, P<0.01) and total progressive motility (+15.8%±1.1%, P<0.01). Significant reduction in leukocyte number (-1.3±0.7 million/mL, P<0.05)

†, interquartile range; *, no values were provided. SDF, sperm DNA fragmentation; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling; CMA3, chromomycin A3; SCSA, sperm chromatin structure assay; DDS, DNA degraded sperm.

daily for 2 months) or a placebo group. Although semen parameters between both groups were not statistically significantly different, the percentages of SDF were lower ($P<0.001$) in the treatment group.

In another study, combination antioxidant therapy with zinc and selenium resulted in a statistically significant decrease in SDF by 19% ($P<0.0004$) (17). Kodama *et al.* (18) investigated the effect of vitamins C (200 mg) and E (200 mg) and glutathione (400 mg) on measures of sperm DNA oxidation, using the 8-hydroxy-2'-deoxyguanosine assay, in 14 infertile men. After a treatment period of 2 months the authors revealed a significant decrease in oxidative DNA damage (-0.4 ± 0.1 , $P<0.05$) together with a significant improvement in sperm concentration ($+7\pm 1$ million/mL, $P<0.05$). Abad *et al.* (24) examined the effects of oral antioxidant therapy on the dynamics of SDF, measured with sperm chromatin dispersion (SCD), following various periods of sperm storage (0, 2, 6, 8 and 24 h) at 37 °C. A combination antioxidant regimen was used on 20 infertile men with asthenoteratozoospermia for a period of 3 months. In addition to finding a significant improvement in sperm concentration ($P=0.04$), progressive motility ($P=0.002$) and normal morphology ($P=0.04$), the authors also reported significant reductions in SDF measures at each experimental point ($P<0.05$). Moreover, a significant reduction in the portion of sperm with highly degraded DNA (DDS) damage was detected after antioxidant therapy ($P=0.04$).

Antioxidant therapy also seems to improve the outcome of intracytoplasmic sperm injection (ICSI) in patients with elevated SDF levels. Greco *et al.* (20) treated 29 patients with high SDF and a prior history of failed ICSI with vitamins C (1,000 mg) and E (1,000 mg) for a period of 3 months. While no significant effect was documented on conventional semen parameters, SDF levels significantly declined ($-15.8\%\pm 3.6\%$, $P<0.001$). In addition, ICSI clinical pregnancy rates (48.2% versus 6.9%, $P<0.05$) and implantation rates (19.6% versus 2.2%, $P<0.01$) were significantly higher than pretreatment ICSI outcomes.

The utility of antioxidant therapy in patients with high SDF secondary to varicocele has been investigated as well. In one study, 20 infertile men with low grade varicocele and high SDF levels, measured with sperm chromatin structure assay (SCSA), were treated with a combination antioxidant regimen containing L-carnitine, vitamin C, coenzyme Q10, vitamin E, zinc, vitamin B9, selenium and vitamin B12 for a period of 3 months (26). The authors reported a significant decrease in SDF levels (-22.1% , $P=0.02$) and a significant

increase in sperm concentration ($P=0.04$). After assessing the amount of highly DDS cells, the authors revealed that 31.3% fewer DDS cells were observed after antioxidant therapy ($P=0.07$).

While certainly further studies of larger sample size and superior design are still required, this brief review reveals that antioxidant therapy appears to be a reasonable option that can be used either solely or as part of a treatment strategy individualized to each patient's clinical presentation.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Agarwal A, Majzoub A, Esteves SC, et al. Clinical utility of sperm DNA fragmentation testing: practice recommendations based on clinical scenarios. *Transl Androl Urol* 2016;5:935-50.
2. Erenpreiss J, Spano M, Erenpreisa J, et al. Sperm chromatin structure and male fertility: biological and clinical aspects. *Asian J Androl* 2006;8:11-29.
3. Shamsi MB, Kumar R, Dada R. Evaluation of nuclear DNA damage in human spermatozoa in men opting for assisted reproduction. *Indian J Med Res* 2008;127:115-23.
4. Sharma RK, Said T, Agarwal A. Sperm DNA damage and its clinical relevance in assessing reproductive outcome. *Asian J Androl* 2004;6:139-48.
5. Sharma R, Biedenharn KR, Fedor JM, et al. Lifestyle factors and reproductive health: taking control of your fertility. *Reprod Biol Endocrinol* 2013;11:66.
6. Majzoub A, Esteves SC, Gosalvez J, et al. Specialized sperm function tests in varicocele and the future of andrology laboratory. *Asian J Androl* 2016;18:205-12.
7. Saleh RA, Agarwal A, Sharma RK, et al. Evaluation of nuclear DNA damage in spermatozoa from infertile men with varicocele. *Fertil Steril* 2003;80:1431-6.
8. Saleh RA, Agarwal A, Nada EA, et al. Negative effects of increased sperm DNA damage in relation to seminal oxidative stress in men with idiopathic and male factor infertility. *Fertil Steril* 2003;79 Suppl 3:1597-605.
9. Lewis SE, John Aitken R, Conner SJ, et al. The impact of

- sperm DNA damage in assisted conception and beyond: recent advances in diagnosis and treatment. *Reprod Biomed Online* 2013;27:325-37.
10. Agarwal A, Cho CL, Esteves SC. Should we evaluate and treat sperm DNA fragmentation? *Curr Opin Obstet Gynecol* 2016;28:164-71.
 11. Agarwal A, Majzoub A. Role of antioxidants in male infertility. *BJUI Knowledge* 2016;e-learning module:1-9.
 12. Agarwal A, Majzoub A. Role of Antioxidants in Assisted Reproductive Techniques. *World J Mens Health* 2017;35:77-93
 13. Gharagozloo P, Aitken RJ. The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy. *Hum Reprod* 2011;26:1628-40.
 14. Zini A, Al-Hathal N. Antioxidant therapy in male infertility: fact or fiction? *Asian J Androl* 2011;13:374-81.
 15. Showell MG, Mackenzie-Proctor R, Brown J, et al. Antioxidants for male subfertility. *Cochrane Database Syst Rev* 2014;(12):CD007411.
 16. Tunc O, Thompson J, Tremellen K. Improvement in sperm DNA quality using an oral antioxidant therapy. *Reprod Biomed Online* 2009;18:761-8.
 17. Ménéz YJ, Hazout A, Panteix G, et al. Antioxidants to reduce sperm DNA fragmentation: an unexpected adverse effect. *Reprod Biomed Online* 2007;14:418-21.
 18. Kodama H, Yamaguchi R, Fukuda J, et al. Increased oxidative deoxyribonucleic acid damage in the spermatozoa of infertile male patients. *Fertil Steril* 1997;68:519-24.
 19. Omu AE, Al-Azemi MK, Kehinde EO, et al. Indications of the mechanisms involved in improved sperm parameters by zinc therapy. *Med Princ Pract* 2008;17:108-16.
 20. Greco E, Romano S, Iacobelli M, et al. ICSI in cases of sperm DNA damage: beneficial effect of oral antioxidant treatment. *Hum Reprod* 2005;20:2590-4.
 21. Greco E, Iacobelli M, Rienzi L, et al. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl* 2005;26:349-53.
 22. Martínez-Soto JC, Domingo JC, Cordobilla B, et al. Dietary supplementation with docosahexaenoic acid (DHA) improves seminal antioxidant status and decreases sperm DNA fragmentation. *Syst Biol Reprod Med* 2016;62:387-95.
 23. Fraga CG, Motchnik PA, Shigenaga MK, et al. Ascorbic acid protects against endogenous oxidative DNA damage in human sperm. *Proc Natl Acad Sci U S A* 1991;88:11003-6.
 24. Abad C, Amengual MJ, Gosalvez J, et al. Effects of oral antioxidant treatment upon the dynamics of human sperm DNA fragmentation and subpopulations of sperm with highly degraded DNA. *Andrologia* 2013;45:211-6.
 25. Vani K, Kurakula M, Syed R, et al. Clinical relevance of vitamin C among lead-exposed infertile men. *Genet Test Mol Biomarkers* 2012;16:1001-6.
 26. Gual-Frau J, Abad C, Amengual MJ, et al. Oral antioxidant treatment partly improves integrity of human sperm DNA in infertile grade I varicocele patients. *Hum Fertil (Camb)* 2015;18:225-9.
 27. Piomboni P, Gambera L, Serafini F, et al. Sperm quality improvement after natural anti-oxidant treatment of asthenoteratospermic men with leukocytospermia. *Asian J Androl* 2008;10:201-6.

Cite this article as: Majzoub A, Agarwal A, Esteves SC. Antioxidants for elevated sperm DNA fragmentation: a mini review. *Transl Androl Urol* 2017;6(Suppl 4):S649-S653. doi: 10.21037/tau.2017.07.09