

The Journal of Phytopharmacology

(Pharmacognosy and phytomedicine Research)

Review Article

ISSN 2320-480X

JPHYTO 2020; 9(4): 281-290

July- August

Received: 07-05-2020

Accepted: 22-06-2020

©2020, All rights reserved

doi: 10.31254/phyto.2020.9412

Lakshmi Narayanan Venu

Cholayil Private Limited, Research & Development Centre, 31-A/24, 4th Cross Main Road, SIDCO Industrial Estate (North), Ambattur, Chennai – 600098, Tamil Nadu, India

Anoop Austin

Cholayil Private Limited, Research & Development Centre, 31-A/24, 4th Cross Main Road, SIDCO Industrial Estate (North), Ambattur, Chennai – 600098, Tamil Nadu, India

Correspondence:

Anoop Austin

Cholayil Private Limited, Research & Development Centre, 31-A/24, 4th Cross Main Road, SIDCO Industrial Estate (North), Ambattur, Chennai – 600098, Tamil Nadu, India

Email: [austin\[at\]cholayil.com](mailto:austin[at]cholayil.com)

Antiviral efficacy of medicinal plants against respiratory viruses: Respiratory Syncytial Virus (RSV) and Coronavirus (CoV) / COVID 19

Lakshmi Narayanan Venu, Anoop Austin*

ABSTRACT

Respiratory syncytial virus (RSV) and Coronavirus disease (CoV) / Covid 19 are droplet infections caused by a group of RNA viruses infecting respiratory system and communicates through sneezing or physical contact. Due to the pandemic effect of Covid 19, it is essential to identify active components, which could be an active compound, which would be beneficial to the community such as respiratory syncytial virus (RSV) and coronavirus. 40 medicinal plants were screened basis the antiviral property pertaining to RNA virus and were dealt in detail to identify the plant which could be a source for developing a novel treatment for Respiratory Syncytial Virus and Covid 19. This review attempts to address the importance of herbs for the treatment of respiratory virus because rapid spread of the infection and preventing the use of allotropic medicine.

Keywords: Antiviral Medicinal plants, Covid 19, Corona virus, Respiratory syncytial virus, Viral inhibiting plants.

INTRODUCTION

Acute respiratory disease (ARD) are one among the most acute morbidities and mortalities in large proportion mortalities worldwide, among which acute viral respiratory tract infection accounts to 80 % [1] approximately. Among those, Influenza Virus, Respiratory Syncytial Virus (RSV), Coronavirus, Adenovirus, and Rhinovirus are some of the major viral pathogens [2]. These are highly pathogenic in nature and they emerging and reemerging coronaviruses in nature which cause epidemics or pandemics, such as the Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV), Middle East Respiratory Syndrome Coronavirus (MERS-CoV), etc, and most recently, a novel coronavirus 2019 have posed great threat to global public health [3]. Among those, Respiratory syncytial virus (RSV) is a single-stranded RNA virus which belongs to the family Paramyxoviridae [4]. In temperate climates, the disease peaks in the winter [5]. The structure of RSV is that of a bilipid-layer-envelope surrounding a ribonucleoprotein core, with several membrane proteins, one of which functions in attachment to host cells, and one of which functions in fusion to host cells. There is only one serotype of RSV, but it is classified into two strains, "A" and "B" with differences consisting of variation in the structure, most especially the attachment protein [6]. Respiratory syncytial virus (RSV) will affect persons of all ages and is one of the major causes of serious Lower Respiratory Tract Infections in younger group of children. Children below the age group of one year are especially likely to develop lower respiratory involvement, with up to 40% of primary infections resulting in bronchiolitis [7]. Among those, coronavirus primarily targets the Human respiratory system, which is one of the major pathogens for respiratory system. Coronaviruses is from the family Orthocoronavirinae [8]. This group of viruses can cause diseases in mammals and birds. Four coronaviruses which are endemic to humans are HCoV-229E, HCoV-OC43, HCoV-NL63, HCoV-HKU1 and they are mainly associated with mild respiratory illnesses and Severe Acute Respiratory Syndrome coronavirus (SARS-CoV), Middle East Respiratory Syndrome Coronavirus (MERS-CoV) and COVID-19 ((SARS-CoV-2) are present as emerging infections causing severe respiratory syndrome [9]. The genomic sequence of the COVID-19 demonstrates about 80% identity towards SARS-CoV and 50% to MERS-CoV [10-11] and interestingly it has been observed that both SARS-CoV and MERS-CoV originate in bats [12].

Outbreaks of SARS-CoV and MERS-CoV are characterized and considered to be a great threat towards the public health. In November, 2002, the SARS-CoV emerged in China causing global anxiety as the outbreak rapidly spread, and by July, 2003, had resulted in over 8000 cases in 26 countries [13].

In 2012, a novel human coronavirus, which was termed as MERS-CoV, emerged in the Middle East to cause fatal human infections in three continents. This has a high fatality rate similar to SARS-CoV and its contagious ability to spread from person to person. Both viruses also have close relationships with bat coronaviruses [14-16]. During late December 2019, a cluster of patients were admitted to hospitals in the city of Wuhan Province, with an initial diagnosis of pneumonia of unknown etiology. Initially they were epidemiologically linked to a seafood and wet animal wholesale market in Wuhan, China [17-19]. On April 15, 2020, a total of 2,000,066 cases have been confirmed COVID-19 in 210 countries and territories around the world. The case fatality rate was calculated to be 6.3 % (126754 / 2,000,066) [19]. Despite this verity of the global health threat and economic burden posed by respiratory virus infection, there is no effective Covid 19 specific antiviral agents, which can manage or treat the virus. Though Ribavirin is and FDA approved drug of choice for RSV, it has its own limitations owing to several factors, including toxicity and difficulty of administration [20]. Chloroquine is effective in preventing the spread of corona virus in cell culture [21].

The conventional treatment for these disease are not yet identified and worldwide many rationale treatments are advocated along with documented side effects in patients [22]. With this rationale a suitable drug is warranted and plants with already established antiviral properties from Medicinal Plants could also be established to identify new molecules in the management of such Pandemic situation [23]. Many of those Medicinal plants are widely used in various traditional

systems of medicine proving its benefit in successful management in various viral ailments [24]. Among those some are localized like Tibetan medicine and others, like Ayurveda, Siddha, Unani (AYUSH) medicines are also gaining popularity worldwide and are being increasingly used in various parts of the world [25].

Antiviral activity of herbal medicine against selected viruses

Many medicinal plants are being evaluated internationally, and are being advocated for strong antiviral activities. Among those some are being treat animals and people who suffer from viral infection [26]. Study demonstrates that during 1952 the search of antiviral agents development were started in Europe by the Boots drug company at Nottingham, England. They were the pioneres in examining the actions of 288 plants against influenza A virus in embryonated eggs. Among that they found that 12 of them suppressed virus multiplications [27]. During the past two to three decades, there are enumerable broad-based screening programmes initiated in different parts of the globe to evaluate the antiviral activity of medicinal plants both *in vitro* and *in vivo* assays [28].

Basis this the medicinal plants were screened basis their antiviral potential, basis which can inhibit the RNA virus, which could be of beneficial in developing a drug in the management of Covid 19 / RSV. The suggestive plants which can be further subjected for detailed evaluation 40 medicinal plants were found to be effective basis the antiviral properties and are tabulated in Table 1.

Table 1: Medicinal Plants demonstrating active antiviral properties against various viruses

S. No.	Botanical Name	Family Name	Antiviral benefit	Phytochemical composition	References
1	<i>Acacia modesta</i> Wall	Mimosoideae	HCV	4-hydroxy benzoic acid, palmitone, dihydroflavonols, catechins, chalcones, anthocyanidins, Quercetin, kaempferol, terpenoids, essential oils, Lupeol, Betulin, amyrin, pentacyclic triterpenes, βs itosterol, Neurolathrogen and α-amino-β-oxalylaminopropionic	[29, 30]
2	<i>Allium sativum</i> L.	Alliaceae	H1N1, Influenza B, HSV 1, HSV 2	Ajoene, allyl alcohol, Diallyl disulphide, Diallyl disulfide (DADS), diallyl sulfide (DAS), Alliin Organo sulfur compounds like Allicin, Diallyl trisulfide, Ajoene	[31-33]
3	<i>Allium porrum</i> L.	Alliaceae	Adenovirus	Dipropyl disulfide and Dipropyl trisulfide quercetin, Zalcitabine, Allicin, Ribavirin	[34]
4	<i>Amaryllis belladonna</i> L.	Amaryllidaceae	RSV	Phenanthridine Lycorine Tazetine	[35]
5	<i>Andrographis paniculata</i> (Burm.f.) Nees	Acanthaceae	HIV	Andrographaloid, Neoandrographolide, 1,5-dimethyl-1,5-cyclooctadiene and 2-hydroxyethyl benzoate	[36, 37]
6	<i>Artemisia annua</i> L.	Asteraceae	CMV, HSV-1, HSV-2, EBV, HBV, HIV-1, HIV-2, HCV, Influenza virus	Artemisinin, Artesunate, Scopoletin	[38]
7	<i>Blumea laciniata</i> (Wall. ex Roxb.) DC.	Asteraceae	HSV-1 & RSV	Borneol, β-caryophyllene, Germacrene D, sabinene	[39, 40]
8	<i>Blumea laciniata</i> (Wall. ex Roxb.) DC.	Asteraceae	RSV	Protocatechuic acid, chrysoeriol, apigenin, 4-hydroxy-3,5-dimethoxybenzoic acid, scopolet.	[40, 41]
9	<i>Boerhavia diffusa</i> L.	Nyctaginaceae	RNA Virus	Punarnavine, 2-glucopyranose-4-hydroxy-5-[P-hydroxyphenyl]- propionyl diphenyl methane, steroids, triterpenoids, alkaloids and flavonoids.	[42,43]
10	<i>Bryonia alba</i> L.	Cucurbitaceae	Human CoV, H1N1,	Bryonicine, Saponarin, Vitexin, Isovitexin, lutanarin, Isoorientin, Glycosides 22-deoxocucurbitosides A and B, Arvenin IV	[36, 44]

11	<i>Camellia sinensis</i> (L.) Kuntze	Theaceae	Bovine CoV, HIV	Epigallocatechin gallate, Epigallocatechin, Epicatechin gallate, Epicatechin, Catechin	[45]
12	<i>Citrus reticulata</i> Blanco	Rutaceae	SARS-CoV-2, Dengue Virus serotypes 1-4, HBV, HCV	Tangeretin, Nobiletin, Hesperetin, Tangeretin, Naringenin, Nobiletin	[46]
13	<i>Curcuma zedoaria</i> (Christm.) Roscoe	Zingiberaceae	H1N1, Influenza B virus	Germacrone, Curcumin	[47-49]
14	<i>Echinacea purpurea</i> (L.) Moench	Asteraceae	H1N1, HSV-1, Influenza	Caffeic acid Cichoric acid alkylamides	[50, 51]
15	<i>Elephantopus scaber</i> L.	Asteraceae	RSV	3,4-dihydroxy benzaldehyde, p-coumaric acid, Vanillic acid, Syringic acid, Isovanillic acid, p-hydroxybenzoic acid, Ferulic acid, 3-methoxy-4-hydroxyl cinnamic aldehyde, Tricin 2-hydroxybenzolate acid	[40, 52]
16	<i>Eupatorium perfoliatum</i> L.	Asteraceae	H1N1, Influenza virus	Caffeic acid derivatives, Flavonoids, Sesquiterpene Lactones, Tannins.	[36]
17	<i>Galanthus nivalis</i> L.	Amaryllidaceae	SARS-CoV	Ethyl-a-D-glucopyranoside, Ethyl-a-D-ribose, Ethyl linoleate. <i>O N⁴</i> -hydroxycytidine, <i>N</i> -(4-fluorophenylsulphonyl)-L-valyl-L-leucinal.	[53]
18	<i>Glycyrrhiza glabra</i> L.	Fabaceae	Kaposi's sarcoma-associated virus, HAV, HBV, CoV, Influenza virus, HIV-1, SARS	Glycyrrhizic acid, Glycyrrhizin, Glycyrrhetic acid, Chalcone	[54, 55]
19	<i>Houttuynia cordata</i> Thunb.	Saururaceae	SARS-CoV	Houttuynoside A (1) and Houttuynamide A, Anthocyanins,	[56]
20	<i>Isatis indigotica</i> L.	Brassicaceae	H1N1, SARS-CoV, Foot-and-mouth disease, Rabies,	Epigoitrin, Indirubin	[57-61]
21	<i>Justicia adhatoda</i> L.	Acanthaceae	Influenza	Alkaloids, lignans, flavonoids, and terpenoids (Iridoids, Diterpenoids, and Triterpenoids)	[36, 62]
22	<i>Laggera pterodonta</i> (DC.) Benth.	Asteraceae	H1N1, H9N2	Pterodontic acid	[63]
23	<i>Lindera aggregata</i> (Sims) Kosterm.	Lauraceae	SARS-CoV	Benzylisoquinoline alkaloids	[64]
24	<i>Lycoris radiata</i> (L'Her.) Herb.	Amaryllidaceae	SARS-CoV	Lycorine	[64]
25	<i>Mussaenda pubescens</i> Dryand	Rubiaceae	RSV	Triterpene esters, 3-palmitoyllupeol, 3-benzoylepi-betulin and β -sitosterol	[40, 65]
26	<i>Narcissus tazetta</i> L.	Amaryllidaceae	RSV, Influenza A (H1N1, H3N2, H5N1) and Influenza B virus, Japanese encephalitis, Dengue, Yellow Fever, HIV Type 1	Pseudolycorine, Galanthamine, 1-hydroxygalanthine. Isoquinolone derivatives	[66]
27	<i>Ocimum tenuiflorum</i> L.	Lamiaceae	Orthomyxovirus, Paramyxovirus	Eugenol, Methyl eugenol, Cirsilineol, Circimaritin, Isothymusin, Apigenin, Rosameric acid, Terpenoid compounds	[36, 67]
28	<i>Phyllanthus amarus</i> Schum. & Thonn.	Euphorbiaceae	HBV	Lignins alkaloids, Hydrolysable tannins, Flavonoids, Ellagic acid, polyphenols.	[68, 69]
29	<i>Piper longum</i> L.	Piperaceae	HBV	Pentenoyl Piperidine, Piperolactam A, Pirrolidine, Turmerone, Piperine Demethoxycurcumin Bisdemethoxycurcumin, Glycosides longumosides A and B,	[70]
30	<i>Piper nigrum</i> L.	Piperaceae	Human Para influenza virus	Polyphenols, Piperidine, Piperine	[70, 71-74]
31	<i>Pyrrhosia lingua</i> (Thunb.) Farw.	Polypodiaceae	SARS-CoV	Chlorogenic acid, Flavonoids	[75, 76]
32	<i>Rosmarinus officinalis</i> L.	Lamiaceae	hRSV, H1N1, Influenza B virus	Rosmarinic acid, Carnosol, Carnosic	[77-80]
33	<i>Schefflera heptaphylla</i> (L.)	Araliaceae	RSV, H1N1	Two highly active pure triterpenoids	[81]
34	<i>Schefflera heptaphylla</i> (L.)	Araliaceae	H1N1, CVB3, HSV-1	Dicaffeoylquinic acids	[82]

35	<i>Scrophularia scorodonia</i> L.	Scrophulariaceae	HSV-1, VSV, PV-1	Iridoids, Phenylpropanoids, Scordioside, Phenolic acids, Flavonoids, Saponins.	[75, 83]
36	<i>Scutellaria indica</i> L.	Lamiaceae	RSV	Scutellarin, Luteolin, Naringenin, Wogonoside, Apigenin, Hispidulin, Wogonin, Chrysin	[40]
37	<i>Selaginella sinensis</i> (Desvaux) Spring	Selaginellaceae	RSV	Alkaloid, Phenolic compounds, Terpenoid, Amentoflavone	[84]
38	<i>Solanum nigrum</i> L.	Solanaceae	HCV	Alkaloids, Saponins, polyphenols, tannins, terpenoids, Solasolidine	[30, 85]
39	<i>Withania somnifera</i> (L.) Dunal	Solanaceae	H1N1	Alkaloids, Withasomnine, Visamine, Amino acids, Withaferin A	[86-88]
40	<i>Zingiber officinale</i> Roscoe	Zingiberaceae	HCV	6-gingerol, 6- paradols and 6-shogaol	[36, 85]

CMV – Cytomegaloviruses, CoV – Corona virus, CVB3 - Cocksackie B3, EBV - Epstein-Barr virus, HAV - Hepatitis A virus, HBV - Hepatitis B virus, HCV - Hepatitis C virus, HSV 1 - Herpes simplex virus 1., HSV 2 - Herpes simplex virus 2, HIV - Human Immuno Deficiency Virus, HIV-1 - Human Immuno Deficiency Virus 1, HIV-2 - Human Immuno Deficiency Virus 2, hRSV – Human Orthopneumovirus, H1N1 -Influenza A virus, H9N2 - Avian influenza A virus, PV-1 - Poliovirus Type 1, RSV - Respiratory Syncytial Virus, SARS - Severe Acute Respiratory Syndrome, SARS-CoV - Severe Acute Respiratory Syndrome- Corona virus, VSV - Vesicular stomatitis virus,

From the table it is evident that the medicinal plant belongs to 22 families, which are found to be effective in antiviral property ingredients basis various studies carried out by various authors. Methanolic extract of *A. modesta* demonstrated a dose dependent inhibition against arachidonic acid induced platelet aggregation at a dose of 2.5 mg/ml provided IC₅₀ was 0.80 mg/ml, which is beneficial in viral infections causing platelet aggregation. Further compounds like long chain alcohols [octacosanol, nonaicosanol and hentriacontanol], hydrocarbons [hentriacontane and octacosane], 4-hydroxy benzoic acid, palmitone, dihydroflavonols, catechins, chalones, anthocyanidins, Quercetin, kaempherol, mono-, sesqui-, di-, tri-, tetra-terpenoids, essential oils, phytosterols, Lupeol, Betulin, camyrin, pentacyclic triterpenes, β sitosterol, Neurolathrogen and α -amino- β -oxalylaminopropionic acid are being reported for its therapeutic benefit. RNA expression using acetone extract exhibiting antiviral activity against HCV at a concentration of 100 μ g, when evaluated via real time RT-PCR [89].

Allium sativum has a strong inhibitory effect on virus multiplication, due to the flavonoid composition, which is also observed to block the formation of protein and genetic materials in the virus, which minimizes influenza A and B viral infections and also effective against cytomegalovirus, rhinovirus, HIV, herpes simplex virus 1, herpes simplex virus 2, viral pneumonia, and rotavirus. It is also observed to significantly minimize the occurrence of the common cold virus. Compounds like Ajoene, allyl alcohol, and diallyl disulfide are found effective against HIV infected cells. Organosulfur compounds like allicin, diallyl trisulfide, and ajoene are the main chemicals which impart antiviral property. Diallyl disulfide (DADS), diallyl sulfide (DAS), and alliin considerably reduced inflammation during dengue virus infection. Quercetin is well-known to act against the entry of the virus in the host cell. For example, Hemagglutinin and neuraminidase are envelope glycoproteins responsible for entry of the Influenza virus. This glycoprotein helps in attachment and membrane fusion of the virus to the host cell. The process of membrane fusion further facilitates the release of the viral ribonucleic proteins into the cytosol. Ribonucleic protein is then transported into the nucleus, which is the site for genome replication. In an interesting study, quercetin was observed to interact with Haemagglutinin protein, which resulted in the inhibition of virus entry into the cell. Similarly, quercetin reduced Enterovirus infection by blocking viral attachment stage of viral infection. It is also an inhibitory effect on quercetin on viral replication. Further, quercetin derivative have been proved to inhibit the translation of polio-virus RNA. The process of formation of multiple copies of polio-virus using the minus-RNA strand was blocked by quercetin. This was attributed to a reduction in viral RNA polymerase, an enzyme essential to initiate the formation of the viral genome. Similarly, quercetin was observed to inhibit the translation process of the hepatitis C virus. SARS-CoV protease, which is required for multiplication of the SARS virus, was inhibited in the presence of quercetin. It is also observed that quercetin

derivatives can increase zinc uptake, which can inhibit RNA Polymerase. This will be beneficial in Covid 19 treatment since absorption of zinc will help to destroy the viral protein. Quercetin also has the potential to disrupt the activation of RNA polymerase by reducing the processing of polyprotein by Rhinovirus proteases. Therefore, the application of quercetin is efficient in enhancing the immune response in the host cells. It has been observed that during the replication process, RNA viruses, including Rhinovirus release protease which attack host cells by cleaving eukaryotic initiation factor (eIF)-4GI and eIF4GII. This ultimately blocks cap-dependent host cell protein synthesis and therefore facilitates viral multiplication. Quercetin has the ability to reduce the cleavage of eIF4GII and minimize the formation of viral capsid protein. This affects the replication of RhinoVirus replication. Similarly, allicin is the chemical present, which also can act against Virus. Allicin can pass through the phospholipid membrane of the cell and can further contribute in inhibiting viral multiplication. It can modulate the immune system in response to viral infection. It has been proved that it blocks the release of pro-inflammatory cytokines like Interleukine 6 (IL-6) and Tumor Necrosis Factor- α (TNF- α). Apart from inhibiting cytokines, allicin has a property to alter transcription of the nuclear factor kappa B (NF- κ B) and DNA binding activity. It has a high amount of selenium and sulphur, which impart antioxidant effect by reacting with intracellular thiol compounds [90].

Chemical compounds like quercetin, zalcitabine, allicin, and ribavirin are found in *A. porrum*, which are effective against adenovirus. By MTT assay and PRD (Plaque reduction) methods exhibited the highest level of antiviral activity [91]. *A. belladonna* has been identified with Twenty-six different Amaryllidaceae alkaloids, among those 1-*O*-acetylcaranine, 3-*O*-acetylhamayne, buphanamine were found to be effective in *in vitro* screening against parasitic protozoa *Plasmodium falciparum*) using methanol extract, which emphasis its benefit in Covid 19 with the report of beneficial of Chloroquine [92].

Andrographaloide from *A. paniculata* are active against HIV, selected pathogenic bacterial organism and possess immune-regulating activities. Alcoholic (90 %) extract demonstrated antiviral activity cell line A549 against Simian Retro Virus (SRV) by RT-PCR analysis, and was nontoxic and comparable with standard drug, Lamivudine, which could benefit in the management of Covid. Immuno-stimulant activity was described by its ability which induces lymphocytes cell proliferation using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Interestingly at a lower concentration of 1 μ g/mL the extract stimulated 38 % of lymphocyte cell proliferation, which is quiet promising for further evaluation on Covid 19 [93].

Artemisinin derived from *A. annua*, which can be metabolized in the liver by microsomal cytochrome P 450 monooxygenase enzymes,

CYP2B6 and CYP3A4 is active against malaria and other viral diseases. It provides a good *in vitro* and *in vivo* evidence as antiviral against DNA viruses like cytomegaloviruses, human herpesvirus 6, herpes simplex viruses 1 and 2, Epstein-Barr virus and Hepatitis B virus. On the contrary, it was not potent against Polyomaviruses and papilloma viruses., RNA viruses such as HIV 1 and 2, hepatitis C virus, influenza virus etc. Further artesunate exerted synergistic inhibition of antiviral drugs clinically also exert cross-resistance to ganciclovir-resistant HCMV. Specific terpenoids and lignoids with activity against SARS [94]. Alcoholic extract was found to be effective against 2005 SARS-CoV. Like SARS-related coronavirus strain, SARS-CoV-2 is also member of the same subgenus and the similarities between these two virus, conclude that artemisinin derivatives could be beneficial drug for further evaluation [95].

Human RSV was inhibited by aqueous extract derived from the following plants namely, *B. laciniata*, *E. scaber*, *L. pterodonta*, *M. pubescens*, *S. octophylla* and *S. utellaria indica* [96]. Root extracts of *B. diffusa* was very potential against potato virus X. Further a purified glycoprotein (70 - 80% protein and 8-13% carbohydrates; Mol. Wt 16 - 20 kDa) from *B. diffusa* exhibited strong antimicrobial activity against RNA (ribonucleic acid) bacteriophages and also active against tobacco mosaic virus [97].

B. alba demonstrated a very good inhibitory effects against human coronavirus 229E and influenza A virus, which was mainly through interference in early stage of viral replication, such as absorption and penetration and attenuating aberrant pro-inflammatory cytokine production. These two viruses are cultured in human cells, human fetal lung fibroblasts MRC-5 and A549 cells, respectively [98].

The beneficial effects of *C. sinensis* are mainly attributed to the presence of a type of polyphenols known as catechins and formed by several isomers including (-) - epigallocatechin gallate (EGCG), (-) - epigallocatechin, (-) - epicatechin gallate, (-) - epicatechin, and (+) - catechin. The catechins have a wide range of antiviral activity against a variety of viruses that act by interfering with its replication cycle. Polyphenolic catechins from *C. sinensis* demonstrated potential inhibitors of HIV reverse transcriptase (RT). EGCG attenuates neuronal damage mediated by HIV infection in the presence of IFN-gamma both *in vitro* and *in vivo* and mildly reduce activated microgliosis and enhance neuron survival [99].

C. reticulata contains several methoxy flavonoids, such as hesperetin, tangeretin, naringenin, and nobiletin which perform low binding energy which are comparable with the reference ligands, lopinavir and nafamostat to the three essential receptors. Hesperetin is used in SARS-CoV-2 which binds with ACE2 receptor, Naringenin is beneficial against Dengue virus (DENV) serotypes 1-4, which Inhibits virus infection and replication Tangeretin in Hepatitis C virus (HCV) reduces HCV secretion in infected cells *in vitro* and *in vivo*. Nobiletin in Hepatitis B virus (HBV) Inhibits HBsAg production and HBV replication *in vitro* and *in vivo*. With this benefit this can be tried to combat beta coronavirus, included SARS-CoV-2 [100].

A major component of essential oil, Germacrone, extracted from the rhizome of *C. zedoaria*, inhibited influenza viral multiplication and was active against H1N1 and H3N2, Influenza A and B viruses, in a dose-dependent manner. On evaluation of the viral protein expression and RNA synthesis were decreased both in MDCK and A549 cell lines treated with germacrone. Further it also exhibited an inhibitory role on viral replication cycle and interestingly it also had an effective protection of mice from lethal infection and viral load / titres of the lung were markedly reduced. Germacrone and Oseltamivir in combination exhibited an additive effect on the inhibition of influenza viral infection, both *in vitro* and *in vivo*. Further Curcumin and Curcuminol can also inhibit NF- κ B and TGF- β 1/Smads signaling pathways [101].

Methanol and aqueous extracts derived from *E. purpurea* partially protected Influenza A virus and Herpes simplex virus type 1, Caffeic

acid, a constituent prevented replication of vesicular stomatitis virus, a membrane containing RNA virus in a mouse cell line. Hexane root extract was active against herpes simplex virus, and was identified due to Cichoric acid present in it. The aqueous fraction containing caffeic acid derivatives and alkylamides showed impressive activity against HSV and influenza. Ethanol extract was as active against HSV [102].

β sesquiphellandrene, is one of the important sesquiterpene compounds showing anti-rhinoviral activity which is found in *E. scaber* Further phenolic acids and flavonoids are also isolated from various fractions. Out of which 3,4-dihydroxy benzaldehyde, p-coumaric acid, vanillic acid, syringic acid, isovanillic acid, p-hydroxybenzoic acid, ferulic acid, 3-methoxy-4-hydroxyl cinnamic aldehyde, triclin, syringic acid, E-3-(3-ethoxy-4-hydroxyphenyl) acrylic acid, 2-hydroxybenzolate acid, are phenolic compounds, which were purified by ethanol fraction. Towards the activity against RSV a new dicaffeoyl derivative, 1 α , 2 β -O-dicaffeoylcyclopentan-3 β -ol showed *in vitro* inhibition which was quite promising Similarly methyl esters of four dicaffeoylquinic acids 4, 5 -di-O-caffeoylquinic acid, 3, 4, di-O-caffeoylquinic acid also possessed strong activity against RSV activity with a marked IC₅₀, which was lower than that of ribavirin [103], which could be subjected for further evaluation.

Hydroalcoholic extracts from the aerial parts of *E. perfoliatum* inhibited the growth of H1N1 and Influenza virus with an inhibitory concentration (IC₅₀) of 7 μ g/mL and 14 μ g/mL, and a selectivity index (SI) of 52 and 26, respectively, and the antiviral activity might be due to polyphenolic compounds which protect cells from IAV infection by inhibiting viral attachment to the host cells. Hence it appears to be a promising compound for the pandemic evaluation for covid 19 [104].

Compounds from *G. nivalis* namely *O*^N-hydroxycytidine, calpain inhibitors (*N*-(4-fluorophenylsulphonyl)-L-valyl-L-leucinal), aglycon derivatives and mannose-specific plant lectins inhibit SARS-CoV replication with an EC₅₀ of 1 μ M and selectivity index of ≥ 100 . The effect was comparable with drugs like Vancomycin, Eremomycin and Teicoplanin [105].

Glycyrrhizic acid from *Glycyrrhiza glabra* demonstrated inhibition on the replication of SARS-CoV *in vivo* and also increased interferon- γ production. Kaposi's sarcoma-associated viral elimination via apoptosis and anti-viral effects are also being described against DNA and RNA viruses, like Hepatitis A virus, Hepatitis B virus, Coronavirus, Influenza virus, HIV-1 etc. [106].

H. cordata is a promising source for many flavonoid molecules including anthocyanins, flavones and flavanones. Aqueous extract increases the proliferation of mouse splenic lymphocytes significantly and dose-dependently and increased the proportion of CD4(+) and CD8(+) T cells. It also possessed significant inhibitory effects on SARS-CoV 3C-like protease (3CL (pro)) and RNA-dependent RNA polymerase (RdRp), which are very beneficial on the drug development [107].

Epigoitrin, a natural alkaloid from *I. indigotica*, protects against influenza infection and also relieves stress. Further interestingly it also elevated mitochondria antiviral signaling (MAVS) protein expression and subsequently increased the production of IFN- β and interferon inducible transmembrane 3 (IFITM3), thereby helping to fight viral infections, which could assist in reducing the susceptibility to influenza virus via mitochondrial antiviral signaling [108].

In vitro antiviral effect of *J. adhatoda* methanolic and aqueous extracts demonstrated inhibition of Influenza virus by Hemagglutination assay with 100% at the concentration of 10 and 5mg/ml respectively with an inhibition of 33% and 16.67% in the simultaneous assay. These results suggest the strong basis against influenza virus, which inhibits viral replication, attachment, and can be further evaluated as viral prophylaxis [109].

In vitro antiviral activity of pterodontic acid, derived from *L. pterodonta*, inhibits Influenza A viral replication by blocking nuclear export of viral RNP complexes, and attenuating the inflammatory response by inhibiting and activation by NF- κ B pathway. In addition, in H1N1 it significantly attenuates the expression of pro-inflammatory molecules like IL-6, MIP-1 β , MCP-1, and IP-10 and also downregulate expression of cytokines and chemokines which are induced by the Avian Influenza A virus (H9N2) [110].

L. aggregata exhibited a significant inhibition against SARS-CoV strain with an EC₅₀ of 2.4 ± 0.2 and CC₅₀ values range from 886.6 ± 35.0 μ g/ml. [111]. Similarly, Lycorine isolated from *Lycoris radiata* L. initially used for certain other indication, showed strong anti-SARS-CoV activity.

Pseudolycorine, Galanthamine, and 11-hydroxygalanthine are isolated from *N. tazetta*. Isoquinolone derivatives were active against RNA containing flavivirous namely Japanese encephalitis, Dengue and Yellow Fever. It is also effective against Lentivirus namely Human Immuno Deficiency virus Type 1 [113] (Bjarne Gabrielsen, *et. al.*, 1992).

In vitro cytotoxicity with aqueous extract of *O. tenuiflorum* revealed moderate inhibitory activity Orthomyxovirus, whereas alcohol extract demonstrated moderate inhibitory activity against Paramyxovirus, Methanol and chloroform extracts showed significant inhibition of orthomyxovirus. Further crude extract and terpenoid compounds showed significant virucidal activity in comparison with virus control and maintained its effect for longer period of time of up to 72 h post-incubation, suggesting antiviral properties to control viral infections [114].

P. amarus exerts antiviral effect by interacting with transcription factors C/EBP α and β and also HBV enhancer I thereby inhibiting the HBV polymerase activity and mRNA transcription. *In vitro* study of *P. amarus* at a dosage of 1 mg/ml⁻¹ concentration was inhibiting the secretion of HBsAg. This disruption against HBV polymerase activity, mRNA transcription, and replication supports its role as an antiviral agent. Extracts were found active against both HBV and WHV surface antigens namely HBsAg and WHsAg which were inhibited by their corresponding antibodies. In addition it also inhibited, WHV DNA-polymerase activity *in vitro*, in a dose-dependent manner. This activity could be attenuated due to inhibition of Immune complexes and virus replication, which are normally restricted to Liver parenchymal cells. Flavonoid compound ellagic acid present in *P. amarus* inhibits immunotolerance against hepatitis B virus e-antigen [115]. Therapeutically HBV infections can be halved by *P. amarus* towards the spreading of virus and improving the immune complexes which could restore to normal liver in its cellular functions.

Alcohol extract of *P. longum* possessed *in vitro*. Activity against HBV. Six new compounds, two new glycosides longumosides A and B and two new amide alkaloids erythro-1-[1-oxo-9(3,4-methylenedioxyphenyl)-8,9-dihydroxy-2E-nonenyl]-piperidine, threo-1-[1-oxo-9(3,4-methylenedioxyphenyl)-8,9-dihydroxy-2E-nonenyl]-piperidine, as well as two compounds 3 β ,4 α -dihydroxy-2-piperidinone, 5,6-dihydro-2(1H)-pyridinone, were also able to demonstrate antiviral activity *in vitro*. Piperine was able to demonstrate remarkable inhibitor against HBV and also against the secretion of HBsAg (Hepatitis B virus surface antigen) and HBeAg (Hepatitis B virus e antigen). Interestingly their Selectivity Index (SI) values were 15.7 and 16.8, respectively, which is also well established as a drug to improve bioavailability [116].

Anti-viral activity of *P. nigrum* in chloroform extract showed higher activity than methanolic extract against Human Para influenza virus on HeLa cell line and was due to the presence of Piperidine [117]. It also contains Piperine and the need to incorporate this compound into formulations and adjunctive therapy is to enhance the bioavailability of various (chemo) therapeutic drugs [118]. *P. lingua* exhibited significant inhibition effects on virus-induced CPE against SARS-CoV and a dose

dependent antiviral activity was determined in serial dilutions of compounds [119].

Marked antiviral activity by Amantadine was revealed from the methanol extract of *R. officinalis* which inhibited the replication of enterovirus and was effective than Amantadine, a common antiviral agent with higher therapeutic index and greater safety [96]. Further various compounds were also tested on the chemical compounds from *R. officinalis*, out of which carnosic acid displayed a potent anti-hRSV activity and was illustrating effect against both A- and B-type viruses. It suppressed the replication of hRSV and suppressed viral gene expression without inducing type-I interferon production or affecting cell viability, suggesting that it can directly affect viral factors, in a concentration-dependent manner. Carnosic acid after injection of 8 hours also in a time course analysis demonstrated an effective blockage on hRSV gene expression suggesting that it can inhibit the replication of hRSV directly [120].

Interestingly *S. heptaphylla* exhibited potent antiviral activity against RSV. Fractionation and isolation from *S. heptaphylla* identified two triterpenoids, namely 3 α -hydroxylup-20(29)-ene-23,28-dioic acid and 3-*epi*-betulinic acid 3-*O*-sulfate. These two triterpenoids demonstrated a wide range of antiviral activity against RSV by cytopathic effect (CPE) reduction method with an IC₅₀ value of 6.25 μ g/mL. Further for Influenza A (H1N1) virus, Coxsackie B3 (Cox B3) virus and Herpes simplex virus type 1 (HSV-1) respectively the IC₅₀ values were 25 to 31.3, 12.5 to 20 and 18.8 and 25 μ g/mL, respectively. Newer compounds like three caffeoylquinic acid derivatives, namely 3-*O*-caffeoylquinic acid, 3,4-di-*O*-caffeoylquinic acid and 3,5-di-*O*-caffeoylquinic acid also were effective against RSV with of 2.33 μ M and 1.16 μ M respectively by plaque reduction assay. Dicafeoylquinic acids against HEp-2 cells exhibited minimal cytotoxicity with a median cytotoxic concentration (CC₅₀) higher than 1000 μ M and maximal non-cytotoxic concentration (MNCC) of the two dicafeoylquinic acids were about 96.7 μ M, suggesting its non-cytotoxicity against RSV. But it was not having antiviral activity against influenza A (Flu A), Coxsackie B3 (Cox B3), and Herpes simplex type one (HSV-1) viruses. Moreover, they could not inhibit RSV attachment to host cells, thereby can not protect HEp-2 cells from RSV infection, which further reinforce that these compounds exerted their anti-RSV effects by inhibiting virus cell fusion in early stage, and at the end stage by inhibiting cell to cell fusion of RSV replication cycle [121].

S. scorodonia was effective in *in vitro* against certain strains of virus namely, Herpes simplex type I (HSV-1), Vesicular stomatitis virus (VSV) and Poliovirus type 1, where Scorodioside showed 47.8 % inhibition at 500 μ /ml [122].

Amentoflavone and three other flavonoids isolated from the ethanol extract of *S. sinensis*. illustrated activity against RSV, with an IC₅₀ of 5.5 μ /ml. It is also interesting that the contents of amentoflavone in nine species of *Selaginella* were it was high in *S. sinensis* and the content was 1.13% [123].

Methanol and chloroform extracts of *S. nigrum* (SN) exhibited a marked inhibition of Hepatitis C Virus (HCV) against HCV NS3 protease by transfecting HCV NS3 protease plasmid into liver cells. Among those two extracts, chloroform extract in a dose dependent manner was able to decrease the expression or function of HCV NS3 protease and on the contrary GAPDH remained constant, which can be advocated with interferon as a better option to treat chronic HCV [124].

Withaferin A an active constituent of *W. somnifera* can attenuate the neuraminidase of H1N1 influenza. Further alcoholic extract is potential and increase the total WBC count and increase the bone marrow cellularity, which is very vital in the pandemic of viral infection to improve the immunity [125].

Z. officinale extract *in vitro* in HepG2 cell line (Hepatocellular carcinoma) was effective against HCV. Viral replications were

inhibited which were detected by amplification of viral RNA segments and further inhibited viral replication inside the HCV-infected HepG2 cells with an inhibitory dose of 100µg/ml. *Z. officinale*, rhizome are reported to contain pungent vanillyl ketones. These including [6]-gingerol and [6]-paradol, compounds which are being credited to have marked therapeutic and preventive health benefits, including anti-cancer and antiviral properties [126].

World Health Organization (WHO), has clearly stated on usage of traditional medicine as a primary health care need as first step medication states that 80% of people from India, 85% from Burma and 90% from Bangladesh are found to use herbal based products as their primary stage of health care management. Basis this many novel compounds are being identified from those plants and are also subjected to various antiviral studies, worldwide. Understanding this scenario, in this review, we are evaluating various approaches currently available or under development for antiviral medicine which can be recommended for the management / control and treatment for RSV and Covid 19. Nowadays, the risk of epidemic and pandemic spread of virus is panicking and is spreading among continents and countries, very quickly. Though there are viral drugs available, due to the pandemic situation of Covid 19, no confirmed drug is available and with the existing antiviral and symptomatic treatment is advocated. Due to mutagenic activity of viral ailments, they are found to be difficult to control and there are still relatively few drugs for treatment of viral diseases [127]. Ribavirin is one such drug approved for the treatment of Respiratory Syncytial Virus which are beneficial for children against bronchiolitis [128]. As on the current prevailing scenario, there is no specific antiviral therapy available for covid 19. Since with the emergence of SARS-CoV which has triggered a largest thrust of antiviral research for corona, on this basis a detailed review against antivirals specific to SARS-CoV could provide a lead to identify novel therapy for the humans [129]. Though currently many antiviral agents which were effective against SARS, like Ribavirin, Lopinavir-Ritonavir, and Oseltamivir, were not showing any progressive proof against Covid 19 [130] provided the general strategies which are advocated include bed rest and supportive treatment, including antiviral therapy [131], antibiotics, immunomodulating therapy and nutritional supplements like zinc and Vitamin C [132, 133]. Another study demonstrated that Remdesivir and Chloroquine are highly effective in the control of corona infection *in vitro*, but cinically there are many challenges [21, 134]. Chloroquine is effective in preventing the spread of corona virus in cell culture. Favorable inhibition of virus spread was observed when the cells were either treated with chloroquine prior to or after corona infection. Synthetic drugs which are being prescribed for viral infections are seldom proving unsatisfactory results and limited narrow spectrum of activity, toxicity and resist antiviral strains [135, 136]. The medicinal plants dealt with are well established towards their antiviral activity and in specific to RNA virus, which can be further screened and could be a possible solution to treat Covid 19 and also can be coprescribed along with the existing which required further evaluation and also, could reduce the dosage by its bioavailability, which is an well-established fact.

Medicinal plants were screened and evaluated basis their antiviral property with special emphasis to Respiratory syncytial virus and RNA virus in order to identify the probable medicinal plants, which could be a right candidate in controlling the Covid 19. Basis this, the study was carried out and interestingly, *A. belladonna*, *B. laciniata*, *E. scaber*, *S. indica*, *S. sinensis* and *S. heptaphylla*, *M. pubescens* were effective against Respiratory syncytial virus. *A. paniculata*, *A. annua* and *Glycyrrhiza glabra* were effective against Human Immuno Deficiency Virus. *A. modesta*, *A. annua*, *C. reticulata*, *S. nigrum* and *Z. officinale* were found to be effective in treating Hepatic C. Influenza was found to be inhibited by *Allium sativum*, *A. annua*, *B. alba*, *C. sinensis*, *C. zeodaria*, *E. purpurea*, *E. perfoliatum*, *Lindigotica*, *Piper nigrum*, *S. heptaphylla*, *Rosmarinus officinalis*, *Schefflera heptaphylla*, *Glycyrrhiza glabra*, *Justicia adathoda*, *Laggera pterodonta* *Withania somnifera*. Sesquiterpene Lactones from *E. perfoliatum* is effective against H1N1 virus. Similarly, *A. porrum* against Adenovirus and

Purnavine from *B. diffusa* and Scorodioside, from *S. scorodonia* are effective against RNA virus. It is also interesting to note that lutoanin, Isoorientin, from *B. alba*, Epicatechin, Catechin from *C. sinensis* and Glycyrrhizin and Glycyrrhizic acid from *G. glabra* are found to be effective in Human Coronavirus, which we can also revisit in finding a novel candidate to fight the pandemic. Nobiletin, from *C. reticulata*, Houttuynoside A and Houttuynamide from *H. cordata*, Epigoitrin, Indirubin from *I. indigotica*, Benzylisoquinoline alkaloids from *L. aggregata*, Lycorine from *L. radiata*, Glucopyranoside from *G. nivalis* Glycyrrhizin from *Glycyrrhiza glabra* and Chlorogenic acid from *P. lingua* are also proven to be effective against SARS-CoV, which is further supporting to evaluate and identify new molecules in the management of viral disease. These observations clearly elucidate that the viral property of those plants could be suggestive in selecting a novel compound and further evaluating, which can through in-depth knowledge, which could help in identifying the right candidate against the viral pandemic.

CONCLUSION

Communicable diseases are very common in spreading the pandemic of Covid 19 and can be a concern to all age groups worldwide. Ever since ages, most of the viral diseases are being considered as intractable to selective antiviral drug, the reason behind is that it has replicative cycle of the virus which are being assumed to be closely interwoven with normal cell metabolism, further it attempts to suppress virus reproduction which could severely harm or doomed to kill the uninfected cell as well. The current review substantiates the usage of herbal drugs and compounds which are mainly having antiviral property with special activity against RNA virus and plants in general are considered to be with lesser side effects, relatively low cost and additional important advantage of their easy availability The current review demonstrates medicinal plants, and compounds like Lycorine, Sabinene, Apigenin, Naringenin, Andrographolide, Artesunate, Scopoletin, Glycyrrhizin, Carnosic acid and similar active compounds which were identified from the selected list of plants screened in this study, which could help in developing a novel drug in the management of Respiratory syncytial virus and Covid 19, which is the need of the hour.

REFERENCES

1. Mahony JB, Petrich A, Smieja M. Molecular diagnosis of respiratory virus infections. *Crit Rev Clin Lab Sci*. 2011; 48:217-249.
2. Azar MM, Landry ML. Detection of influenza A and B viruses and respiratory syncytial virus by use of Clinical Laboratory Improvement Amendments of 1988 (CLIA)-waived point-of-care assays: a paradigm shift to molecular tests. *J Clin Microbiol*. 2018; 56:0-18.
3. Hu B, Ge X, Wang LF, Shi Z. Bat origin of human coronaviruses. *Virology*. 2015; 12:221.
4. Falsey AR, Walsh EE. Respiratory syncytial virus infection in adults. *Clin Microbiol Rev* 2000; 13:371-84.
5. Centers for Disease Control and Prevention (CDC). Brief report: respiratory syncytial virus activity--United States, July 2006-November 2007. *MMWR Morb Mortal Wkly Rep* 2007; 56:1263-1265.
6. Krause CI. The ABCs of RSV. *Nurse Pract*. 2018; 43(9):20-26.
7. Omer SB, Bednarczyk R, Kazi M, Guterman LB, Aziz F, Allen KE, *et al*. Assessment and Validation of Syndromic Case Definitions for Respiratory Syncytial Virus Testing in a Low Resource Population. *Pediatr. Infect. Dis. J*. 2019; 38(3):57-59.
8. De Groot RJ, Baker SC, Baric R, Enjuanes L, Gorbalenya AE, Holmes KV, Perlman S, *et al*. Middle East Respiratory Syndrome Coronavirus (MERS-CoV): Announcement of the Coronavirus Study Group. *J. Virol*. 2013; 87(14):7790-7792.
9. Chan PK, Chan MC. Tracing the SARS - coronavirus. *J Thorac Dis*. 2013; 5(S2):118-121.
10. Ren LL, Wang YM, Wu ZQ, Xiang ZC, Guo L, Xu T. Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chinese Med J*. 2020.
11. R. Lu, X. Zhao, J. Li, P. Niu, B. Yang, H. Wu. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *J. Lancet*. 2020; 395(10224):565-574.

12. J. Cui, F. Li, Z.L. Shi. Origin and evolution of pathogenic coronaviruses. *Nat. Rev. Microbiol.* 2019; 17:181-192.
13. Peiris JS, Lai ST, Poon LL. Coronavirus as a possible cause of severe acute respiratory syndrome. *Lancet.* 2003; 361:1319-1325.
14. Zaki AM, Van Boheemen S, Bestebroer TM. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med.* 2012; 367:1814-1820.
15. Chan JF, Li KS, To KK. Is the discovery of the novel human betacoronavirus 2c EMC/2012 (HCoV-EMC) the beginning of another SARS-like pandemic. *J Infect.* 2012; 65:477-489.
16. Chan JF, Lau SK, Woo PC. The emerging novel Middle East Respiratory Syndrome Coronavirus: the “knowns” and “unknowns”. *J Formos Med Assoc.* 2013; 112(7):372-381.
17. Bogoch, A. Watts, A. Thomas-Bachli, C. Huber, M.U.G. Kraemer, K. Khan, Pneumonia of unknown etiology in wuhan, China: potential for international spread via commercial air travel. *J. Trav. Med.* 2020; 27(2):008.
18. Lu. H, Stratton. C.W, Tang. Y.W. Outbreak of pneumonia of unknown etiology in wuhan China: the mystery and the miracle. *J. Med. Virol.* 2020; 92(4):401-402.
19. <https://www.who.int/docs/default-source/coronaviruse>.
20. American Academy of Pediatrics Subcommittee on Diagnosis and Management of Bronchiolitis: Diagnosis and management of bronchiolitis. *Pediatrics.* 2006; 118:1774-1793.
21. Martin J Vincent, Eric Bergeron, Suzanne Benjannet, Bobbie R Erickson, Pierre E Rollin, Thomas G Ksiazek, Nabil G Seidah and Stuart T Nicho. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology Journal.* 2005; 2:69.
22. Al-Ali KH, El-Badry AA. Anti-viral activity of two Labiate plants Naana Hassoi, Habak and Basil Rahan, of Al Madiah Almunawarah. *Journal of Medicine and Biomedical Sciences.* 2010; 1-7.
23. Kumar BNS, Swamy BNV, Swamy A and Murali A. A review of natural diuretics. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 2010; 1(4):615-63.
24. Prakash P, Gupta N. Therapeutic uses of *Ocimum sanctum* Linn. with a note on Eugenol and its pharmacological actions: A Short Review. *Indian Journal of Physiology and Pharmacology.* 2005; 49(2):125-131.
25. Hudson JB. The use of herbal extract in use control of influenza. *Journal of Medicinal Plants Research.* 2009; 3(13):1189-1195.
26. Alicia Martinez Lopez, Mirjana Persaud, Maritza Purav Chavez, Hongjie Zhang, Lijun Rong, Shufeng Liu, Tony T Wang, Stefan G. Sarfianos. Glycosylated diphyllin as a broad spectrum antiviral against zika virus. *The Lancet.* 2019; 47:269-283.
27. Jassim SAA, Naji MA. Novel antiviral agents: a medicinal plant perspective. *Journal of Applied Microbiology.* 2003; 95:412-427.
28. Asres K, Bucar F, Karting T, Witvrouw M, Pannecouque C, De Clercq E. Antiviral activity against human immunodeficiency virus type 1 (HIV-1) and type 2 (HIV-2) of ethnobotanically selected Ethiopian medicinal plants. *Phytotherapy Research.* 2001; 15:62-69.
29. Wajid Sarwar. Pharmacological and phytochemical studies on *Acacia modesta* Wall; A review”, *The Journal of Phytopharmacology.* 2016; 5(4):160-166.
30. Ahmad M, Sultana S, Fazl-i-Hadi S, Ben Hadda T, Rashid S, Zafar M, Yaseen G. An ethnobotanical study of medicinal plants in high mountainous region of Chail valley (District Swat-Pakistan). *J Ethnobiol Ethnomed.* 2014; 10:36.
31. Sher H, Hussain F. Ethnobotanical evaluation of some plant resources in Northern part of Pakistan. *Afr J Biotechnol.* 2009; 8(17):4066-76.
32. Alamgeer, Waqas Younis, Hira Asif, Amber Sharif, Humayun Riaz, Ishfaq Ali Bukhari, Asaad Mohamed Assir. Traditional medicinal plants used for respiratory disorders in Pakistan: a review of the ethno-medicinal and pharmacological evidence. *Chin Med.* 2018; 13:48
33. Weber, ND, Andersen DO, North JA, Murray BK, Lawson LD, Hughes BG. In vitro virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Med.* 1992; 58:417-423.
34. Casella S, Leonardi M, Melai B, Fratini F, Pistelli L. The Role of Diallyl Sulfides and Dipropyl Sulfides in the *In vitro* Antimicrobial Activity of the Essential Oil of Garlic, *Allium sativum* L., and Leek, *Allium porrum* L. *Phytother. Res.* 2012; 27:380-383.
35. Bryant AR. Zulu medicine and Medicine-men. Struik; Cape Town, South Africa, 1966.
36. Maria Carmen S Tan, Glenn G Oyong, Chien Chang Shen, Consolacion Y, Ragasa. Chemical Composition of *Andrographis paniculata* (Burm.f.) Nees. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 2016; 7(6):2405.
37. Manvi, Ganesh Prasad Garg. Evaluation of Pharmacognostical parameters and Hepatoprotective activity in *Bryonia alba* Linn. *J. Chem. Pharm. Res.* 2011; 3(6):99-109.
38. Li SY, Chen C, Zhang HQ, Guo HY, Wang H, Wang L. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antivir Res.* 2005; 67:18-23.
39. Laakso I, Seppanen-Laakso T, Hiltunen R, Ekundayo O. Composition of the essential oil of *Blumea lacera* DC. (Asteraceae) leaves from Nigeria. *Flavour Frag J.* 1989; 4:73-5.
40. Yaolan Li, Linda S.M.Ooi, Hua wang, Paul P.H. But and Vincent E.C.Ooi. Antiviral activities of medicinal herbs traditionally used in southern mainland China. *Phytother Res.* 2004; 18:718-22.
41. Yuan NN, Huang WH, Li YL, Cen YZ. Studies on the chemical constituents of *Blumea laciniata* Journal of Chinese medicinal materials. 2009; 32(7):1065-6.
42. Lami N, Kadota S, Kikuchi T. Constituents of the roots of *Boerhaavia diffusa* Linn. IV. Isolation and structure determination of boeravinones D, E and F. *Chemical and Pharmaceutical Bulletin.* 1992; 39(7):1863-1865.
43. Mitra R, Gupta RC. Punarnava: An ayurvedic drug of repute. In: *Applied Botany Abstracts.* Economic Botany Information Service, National Botanical Research Institute. Lucknow, Uttar Pradesh, India. 1997; 17(3):209-227.
44. Angonese MT, Moreira DL, Kaplan MAC. Perfil químico da familia acanthaceae. *Bol Mus Biol Mello Leitão* 1992; 1:3-6.
45. Edziri H, Mastouri M, Mahjoub MA, Ammar S, Mighri Z, Gutmann L and Aouni M. Antiviral activity of leaves extracts of *Marrubium alysson* L. *Journal of Medicinal Plants Research* 2011; 5(3):360-363.
46. Scaglia LFN, Retailleau P, Paolini J, Pannecouque C, Neyts J. Jatrophone diterpenes as inhibitors of chikungunya virus replication: Structure-activity relationship and discovery of a potent lead. *J Nat Prod.* 2014; 77:1505-1512.
47. Wilson B, Abraham G, Manju VS, Mathew M, Vimala B, Sundaresan S, Nambisan B. Antimicrobial activity of *Curcuma zedoaria* and *Curcuma malabarica* tubers. *J Ethnopharmacol.* 2005; 99:147-151.
48. Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chem.* 2005; 91:621-632.
49. Raut JS, Karuppaiyl SM. A status review on the medicinal properties of essential oils. *Ind. Crops Prod.* 2014; 62:250-264.
50. See DM, Broumand N, Sahl L, LES JG. In vitro effects of Echinacea and ginseng on natural killer and antibody-dependent cell cytotoxicity in healthy subjects and chronic fatigue syndrome or acquired immune deficiency syndrome patients. *Immunopharmacology.* 1997; 35:229-235.
51. Weber W, Taylor JA, Stoep AV, Weiss NS, Standish LJ, Calabrese C. *Echinacea purpurea* for prevention of upper respiratory tract infections in children. *J. Altern. Complement. Med.* 2005; 11:1021-1026.
52. Wang L., Jian S, Nan P, Zhong Y. Chemical composition of the essential oil of *Elephantopus scaber* from southern China. *Z Naturforsch C J Biosci.* 2004; 59(5-6):327-9.
53. Gopalakrishnan. Kalaiarasi T. Determination of biologically active constituents of the fruits of *Cucumis sativus* Linn. Using gc- ms analysis. *Int. J. Biol. Pharm. Res.* 2013; 4(7):523-527.
54. Lee Jia Ming, Adeline Chia Yoke Yin. Therapeutic Effects of *Glycyrrhizic Acid*. *Natural Product Communications.* 2013; 8(3):415-418.
55. Fiore C, Eisenhut M, Krausse R, Ragazzi E, Pellati D, Armanini D, Bielenberg J. Antiviral effects of *Glycyrrhiza* species. *Phytother Res.* 2008; 22(2):141-148.
56. Lau KM, Lee KM, Koon CM, Cheung CS, Lau CP, Ho HM, et al. Immunomodulatory and anti-SARS activities of *Houttuynia cordata*. *J Ethnopharmacol.* 2008; 118:79-85.
57. Chen L, Lin T, Zhang H, Su Y. Immune responses to foot-and-mouth disease DNA vaccines can be enhanced by coinjection with the *Isatis indigotica* extract. *Intervirology.* 2005; 48(4):207-212.
58. Heredia A, Davis C, Bamba D. Indirubin-3'-monoxime, a derivative of a Chinese antileukemia medicine, inhibits P-TEFb function and HIV-1 replication. *AIDS.* 2005; 19(18):2087-2095.
59. Hertel L, Chou S, Mocarski ES. Viral and cell cycle-regulated kinases in cytomegalovirus-induced pseudomitosis and replication. *PLoS Pathogens.* 2007; 3(1):6.
60. Hsuan SL, Chang SC, Wang SY. The cytotoxicity to leukemia cells and antiviral effects of *Isatis indigotica* extracts on pseudorabies virus. *J. Ethnopharmacol.* 2009; 123(1):61-67.
61. Lin CW, Tsai FJ, Tsai CH, Lai CC, Wan L, Ho TY, et al. Anti-SARS coronavirus 3C-like protease effects of *Isatis indigotica* root and plant-derived phenolic compounds. *Antivir Res.* 2005; 68:36-42.
62. Sunita Verma. Chemical constituents and pharmacological action of *Ocimum sanctum* (Indian holy basil-Tulsi). *The Journal of Phytopharmacology.* 2016; 5(5):205-207.
63. Shang X, Liabsuetrakul T, Sangsupawanich P, Xia X, He P, Cao H, McNeil E. Efficacy and safety of *Laggetta pterodonta* in children 3-24

- months with acute bronchiolitis: a randomized controlled trial. Clin Respir J. 2017; 11(3):296-304.
64. Li SY, Chen C, Zhang HQ, Guo HY, Wang H, Wang L. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. Antivir Res. 2005; 67:18-23.
 65. Dinda B, Debnath S, Arima S, Sato N, Harigaya Y. Chemical constituents from *Lasia spinose*, *Mussaenda* and *Wendlandia tinctoria*. J Indian Chem Soc. 2004; 81:73.
 66. Ooi LS, Ho WS, Ngai KL, Tian L Chan PKSun SS, Ooi VE. *Narcissus tazetta* lectin shows strong inhibitory effects against respiratory syncytial virus, influenza A (H1N1, H3N2, H5N1) and B viruses. J. Biosci. 2010; 35:95-103.
 67. Sai Durga Prasad S, Mallikarjuna K. Venumadhav R. Screening of Anti-Bacterial Activity of *Solanum trilobatum* Linn. Seed Extract against Dental Pathogens. Asian Journal of Plant Science and Research. 2015; 5(2):34-37.
 68. Foo LY. Amarulone, novel cyclic hydrolysable tannin from *Phyllanthus amarus*. Natural Product Letters. 1993; 3:45-52.
 69. Tan W, Jaganath I, Manikam I. Evaluation of antiviral activities of four local Malaysian *Phyllanthus* species against Herpes simplex viruses and possible antiviral target. International Journal of Medical Sciences. 2013; 10(13):1817-1892.
 70. Shanmugavelu M. Noikaluku Siddha parikaram. Part-I. Indian medicine-Homeopathy department. Government of India 2004.
 71. Ma SC, But PP, Ooi VE, He YH, Lee SH, Lee SF, Lin RC. Antiviral amentoflavone from *Selaginella sinensis*. Biol. Pharm. Bull.2001; 24:311-312.
 72. The Siddha Pharmacopoeia of India. Part-I. Ministry of health and welfare, Department of Ayurveda, Yoga and Naturopathy, Unani, Siddha and Homeopathy (AYUSH), Government of India. 2008.
 73. Agbor GA, Vinson JA, Oben JE, Ngogang JY. J. Herb.Pharmacotherap. 2007; 7:49-64.
 74. Reddy SV, Srinivas PV, Praveen B, Kishore KH, Raju BC, Murthy US, Rao JM. Phytomedicine: International J. Phytotherap. Phytopharmacol. 2004; 11:697-700.
 75. Li SY, Chen C, Zhang HQ, Guo HY, Wang H, Wang L. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. Antivir Res. 2005; 67:18-23.
 76. Hin, HS. Satsu H. Bae, MJ. Anti-inflammatory effect of chlorogenic acid on the IL-8 production in Caco-2 cells and the dextran sulphate sodium-induced colitis symptoms in C57BL/6 mice. Food Chem. 2015; 168:167-175.
 77. Bernardes WA, Lucarini R, Tozatti MG, Flauzino LG, Souza MG, Turatti IC, Silva ML A E, Martins CH, da Silva Filho AA, Cunha WR: Antibacterial activity of the essential oil from *Rosmarinus officinalis* and its major components against oral pathogens. Z Naturforsch C J Biosci. 2010; 65(9-10):588-93.
 78. Oluwatuyi M, Kaatz GW, Gibbons S. Antibacterial and resistance modifying activity of *Rosmarinus officinalis*. Phytochemistry. 2004; 65:3249-3254.
 79. Peng CH, Su JD, Chyau CC, Sung TY, Ho SS, Peng CC, Peng RY. Supercritical fluid extracts of rosemary leaves exhibit potent anti-inflammation and anti-tumor effects. Biosci Biotechnol Biochem. 2007; 71:2223-2232.
 80. Satoh T, Izumi M, Inukai Y, Tsutsumi Y, Nakayama N, Kosaka K, Shimojo Y, Kitajima C, Itoh K, Yokoi T, Shirasawa T: Carnosic acid protects neuronal HT22 Cells through activation of the antioxidant-responsive element in free carboxylic acid- and catechol hydroxyl moieties-depend manners. Neurosci Lett. 2008; 434:260-265.
 81. Li Y, Jiang R, Ooi LS, But PP, Ooi VE. Antiviral triterpenoids from the medicinal plant *Schefflera heptaphylla*. Phytother Res. 2007; 21(5):466-70.
 82. Hayashi K, Niwayama S, Hayashi T, Nago R, Ochiai H. In vitro and in vivo antiviral activity of scopadulcic acid B from *Scoparia dulcis*, Scrophulariaceae, against herpes simplex virus type 1. Antiviral Res. 1988; 9:345-354.
 83. Emam AM, Díaz Lanza AM, Matellano Fernández L, Faure R, Moussa AM. Biological activities of budlejasonin isolated from *Buddleja madagascariensis* and *Scrophularia scorodonia*. Pharmazie.1997; 52:76-77.
 84. Ma SC, But PP, Ooi VE, He YH, Lee SH, Lee SF, Lin RC. Antiviral amentoflavone from *Selaginella sinensis*. Biol Pharm Bull. 2001; 24: 311-312.
 85. Sahdeo Prasad, Amit K. Tyagi. Ginger and Its Constituents: Role in Prevention and Treatment of Gastrointestinal Cancer. Gastroenterology Research and Practice. 2015:0-11.
 86. Khare CP. Indian Medicinal Plants—An Illustrated Dictionary. First Indian Reprint, Springer (India) Pvt. Ltd. 2007: 717-718.
 87. Rastogi RP, Mehrotra BN. Compendium of Indian Medicinal Plants. 2nd Reprint, Central Drug Research Institute, Lucknow and National Institute of Science Communication, Council of Scientific and Industrial Research. 1998; 1:434-436.
 88. Qamar Uddin L, SamiullaV, Singh K, Jamil SS. Phytochemical and Pharmacological Profile of *Withania somnifera* Dunal: A Review. Journal of Applied Pharmaceutical Science. 2012; 02(01):170-175.
 89. Sidra Rehman, Usman A Ashfaq, Sana Riaz, Tariq Javed and Sheikh Riazuddin. Antiviral activity of Acacia nilotica against Hepatitis C Virus in liver infected cells. Virology Journal. 2011; 8:220.
 90. Neha Sharma. Efficacy of Garlic and Onion against virus. International Journal of Research in Pharmaceutical Sciences.2019; 10(4):3578-3586.
 91. Chien-Hsien, Chen, Tzu-Wei, Chou, Li-Hui, Cheng and Chin-Wen Ho. *In vitro* anti-adenoviral activity of five *Allium* plants, Journal of the Taiwan Institute of Chemical Engineers. 2011; 42(2):228-232.
 92. Luciana R, Tallini, Jean Paulo de, Andrade, Marcel Kaiser, Francesc Viladomat, Jerald J. Nair, José Angelo S, Zuanazzi, Jaume Bastida. Alkaloid Constituents of the Amaryllidaceae Plant *Amaryllis belladonna* L. Molecules. 20017; 22:1-12.
 93. Churiyah, Olivia Bunga pongtuluran, Elraderofaani, Tarwadi. Antiviral and Immunostimulant Activities of *Andrographis paniculata*, *Hayati*. Journal of Biosciences. 2015; 22(2):67-72.
 94. Efferth T, Romero MR, Wolf DG, Stamminger T, Marin JJ, Marschall M. Clinical Infectious Diseases. 2008; 47(6):804-811.
 95. Thomas Efferth. Beyond malaria: The inhibition of viruses by artemisinin-type compounds, Biotechnology Advances.2018; 36 (6): 1730-1737.
 96. Cecilia Perera, Thomas Effort. Antiviral Medicinal Herbs and Phytochemicals. J. Pharmacog.2012; 3(1):45-48.
 97. Pranati Nayak, Thirunavoukkarasu, M. A review of the plant *Boerhaavia diffusa*: its chemistry, pharmacology and therapeutical potential. J. Phytopharmacol. 2016; 5(2):83-92.
 98. Bochuan Yuan, Rui Yang, Yongsheng Ma, Shan Zhou, Xiaodong Zhang and Ying Liu. A systematic review of the active saikosaponins and extracts isolated from Radix Bupleuri and their applications, Pharmaceutical Biology. 2017; 55(1):620-635.
 99. Jun Xu, Zhao Xu, Wenming Zheng. A Review of the Antiviral Role of Green Tea Catechins. Molecules. 2017; 22(8):1337.
 100. Nogata Y, Sakamoto K, Shiratsuchi H, Ishii T, Yano M, Ohta H. Flavonoid composition of fruit tissues of citrus species. Bioscience Biotechnology & Biochemistry. 2006; 70(1):178-192.
 101. Qingjiao Liao, Zheng xu Qian, Rui Liu, Liwei An, Xulin Chen. Germacrone inhibits early stages of influenza virus infection, Antiviral Research. 2013; 100 (3):578-588.
 102. James Hudson, Selvarani Vimalanathan. Echinacea-A Source of Potent Antivirals for Respiratory Virus Infections. Pharmaceuticals. 2011; 4(7):1019-1031.
 103. Farha Arakkaveetil Kabeer, Remani Prathapan. Phytopharmacological Profile of *Elephantopus scaber*. Pharmacologia. 2014; 5(8):272-285.
 104. Andrea Derksen, Joachim Kuhn, Wali Hafezi, Jandirk Sendker, Christina Ehrhardt, Stephan Ludwig, Andreas Hensel. Antiviral activity of hydroalcoholic extract from *Eupatorium perfoliatum* L. against the attachment of influenza A virus. Journal of Ethnopharmacology. 2016; 188:144-152.
 105. Erik De Clercq. Antivirals and antiviral strategies. Nature Reviews Microbiology. 2004; 2:704-720.
 106. Cinatl J, Morgenstern B, Bauer G, Chandra P, Rabenau H, Doerr HW. Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated coronavirus. Lancet. 2003; 14(361):2045-2046.
 107. Lau KM, Lee KM, Koon CM, Cheung CS, Lau CP, Ho HM, et al. Immunomodulatory and anti-SARS activities of *Houttuynia cordata*. J Ethnopharmacol. 2008; 118:79-85.
 108. Zhuo Luo, Li-Fang Liu, Xiao-Hua Wang, Wen Li, Chong Jie, Huan Chen, Fan-Qin Wei, Dan-Hua Lu, Chang-Yu Yan, Bo Liu, Hiroshi Kurihara, Yi-Fang Li and Rong-Rong He. Epigoitrin, an Alkaloid From *Isatis indigotica*, Reduces H1N1 Infection in Stress-Induced Susceptible Model *in vivo* and *in vitro*, Frontiers of Pharmacology. 2019; 10(78):1-12.
 109. Rahul Chavan, Abhay Chowdhary. In Vitro Inhibitory Activity of *Justicia adhatoda* extracts against Influenza Virus Infection and Hemagglutination. Int. J. Pharm. Sci. Rev. Res. 2014; 25(2):231-236.
 110. Wenda Guan, Jing Li, Qiaolian Chen, Zhihong Jiang, Rongping Zhang, Xinhua Wang, Zifeng Yang and Xiping Pan. Pterodontic Acid Isolated from *Laggera pterodonta* Inhibits Viral Replication and Inflammation Induced by Influenza A Virus. Molecules. 2017; 22(10):1738.
 111. Shi-you Li, Cong Chen, Hai-qing Zhang, Hai-yan Guo, Hui Wang, Lin Wang, Xiang Zhang, Shi-neng Hua, Jun Yu, Pei-gen Xiao, Rong-song Li, Xuehai Tan. Identification of natural compounds with antiviral activities

- against SARS-associated coronavirus. *Antiviral Research*. 2005; 67:18–23.
112. Muhammad Mukhtar, Mohammad Arshad, Mahmood Ahmad, Roger J. Pomerantz, Brian Wigdahl, Zahida Parveen. Antiviral potentials of medicinal plants. *Virus Research*. 2008; 131(2):111–120.
113. Bjarne Gabrielsen, Thomas P. Monath, John W. Huggins, Deborah F. Kefauver, George R. Pettit, Grazyna Groszek, *et al.* Antiviral (RNA) activity of selected Amaryllidaceae isoquinoline constituents and synthesis of related substances. *Journal of Natural Products*. 1992; 55(11):1569–1581.
114. Ghoke SS, Sood R. Kumar N, Pateriya AK. Bhatia S, Mishra A, Dixit R, Singh VK, Desai DN, Kulkarni DD, Dimri U, Singh VP. Evaluation of antiviral activity of *Ocimum sanctum* and *Acacia arabica* leaves extracts against H9N2 virus using embryonated chicken egg model, *BMC Complementary and Alternative*. 2018; 18:174–186.
115. Kang EH, Kown TY, Oh GT, Park WF, Park SI, Park SK, Lee YI. Antiviral Medicinal Herbs and Phytochemicals. *Antiviral Research*. 2006; 72(2):100–106.
116. Jiang ZY, Liu WF, Zhang X, Luo J, Ma YB, Chen JJ. Anti-HBV active constituents from *Piper longum*. *Bioorganic and medicinal Chemistry Letters*. 2013; 23(7):2123–2127.
117. Priya NC, Saravana Kumari P. Antiviral Activities and Cytotoxicity Assay of Seed Extracts of *Piper longum* and *Piper nigrum* on Human Cell Lines. *International Journal of Pharmaceutical Sciences Review and Research*. 2017; 44(1):197–202.
118. Zorica Stojanović-Radić, Milica Pejić, Marina Dimitrijević, Ana Aleksić, Nanjangud V. Anil Kumar, Bahare Salehi, William C. Cho, Javad Sharifi-Rad. Piperine-A Major Principle of Black Pepper: A Review of Its Bioactivity and Studies. *Applied sciences Review*. 2019; 9:4270–4299.
119. Shi-you Li, Cong Chen, Hai-qing Zhang, Hai-yan Guo, Hui Wang, Lin Wang, Xiang Zhang, Shi-neng Hua, Jun Yu, Pei-gen Xiao, Rong-song Li, Xuehai Tan. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antiviral Research*. 2005; 67:18–23.
120. Han-Bo Shin, Myung-Soo, Choi Byeol Ryu. Antiviral activity of carnosic acid against respiratory syncytial virus. *Virology Journal*. 2013; 10(1):303.
121. Yaolan Li, Renwang Jiang, Linda SM. Ooi, Paul PH, But, Vincent EC Ooi. Antiviral triterpenoids from the medicinal plant *Schefflera heptaphylla*. *Phytotherapy Research*. 2007; 21(5):466–470.
122. Bermejo P, Abad MJ, Díaz AM, Fernández L, De Santos J, Sanchez S, Villaescusa L, Carrasco L, Irurzun A. Antiviral activity of seven iridoids, three saikosaponins and one phenylpropanoid glycoside extracted from *Bupleurum rigidum* and *Scrophularia scorodonia*. *Planta Medica*. 2002; 68(2):106–10.
123. Shuang Chung Ma, Paul Pui-Hay But, Vincent Eng Choon Ooi, Yue Hua He, Spencer Hon Sun Lee, Song Fung Lee, Rui Chou Lin. Antiviral Amentoflavone from *Selaginella sinensis*. *Biological & Pharmaceutical Bulletin*. 2001; 24(3):311–312.
124. Tariq Javed, Usman Ali Ashfaq, Sana Riaz, Sidra Rehman. *In vitro* antiviral activity of *Solanum nigrum* against Hepatitis C Virus. *Virology Journal*. 2011; 8(1):26–32.
125. Cai Z, Zhang G, Tang B, Liu Y, Fu X, Zhang X. Promising Anti-influenza Properties of Active Constituent of *Withania somnifera* Ayurvedic Herb in Targeting Neuraminidase of H1N1 Influenza: Computational Study. *Cell Biochemistry and Biophysics*. 2015; 72(3):727–39.
126. Abd El-Wahab A, H El-Adawi H, M El-Demellawy M. *In vitro* study of the antiviral activity of *Zingiber officinale*. *Planta Med*. 2009; 75(9):7–14.
127. Perera C, Efferth T. Antiviral medicinal herbs and phytochemicals. *Journal of Pharmacognosy*. 2012; 3(1):45–48.
128. Ventre K, Randolph AG. Ribavirin for respiratory syncytial virus infection of the lower respiratory tract in infants and young children. *Cochrane Database Syst. Rev*. 2007:CD000181.
129. Groneberg DA, Poutanen SM, Low DE, Lode H, Welte T, Zabel P. Treatment and vaccines for severe acute respiratory syndrome. *Lancet Infect. Dis*. 2005; 5:147–155.
130. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Med*. 2006; 3:343.
131. Arabi YM, Allothman A, Balkhy HH. Treatment of Middle East Respiratory Syndrome with a combination of lopinavir-ritonavir and interferon-beta1b (MIRACLE trial): study protocol for a randomized controlled trial. *Trials*. 2018; 19(1):81.
132. Arabi YM, Mandourah Y, Al-Hameed F, Sindi AA, Mekhlafi GAA. Corticosteroid Therapy for Critically Ill Patients with Middle East Respiratory Syndrome. *Am J Respir Crit Care Med*. 2020; 197:757–67.
133. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020; 395(10223):507–513.
134. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M. Remdesivir and Chloroquine Effectively Inhibit the Recently Emerged Novel Coronavirus (2019-nCoV) in Vitro. *Cell research*. 2020; 30:269–271.
135. Fernandes MJB, Barros AV, Melo MS, Simoni IC. Screening of Brazilian plants for antiviral activity against animal herpes viruses. *Journal of Medicinal Plants Research*. 2012; 6(12):2261–2265.
136. Simoni IC, Manha APS, Sciessere L, Hoe VMH, Takinami VH, Fernandes MJB. Evaluation of antiviral activity of Brazilian Cerrado plants against animal viruses. *Virus Review and Research*. 2007; 12:1–17.

HOW TO CITE THIS ARTICLE

Lakshmi Narayanan V, Anoop A. Antiviral efficacy of medicinal plants against respiratory viruses: Respiratory Syncytial Virus (RSV) and Coronavirus (CoV) / COVID 19. *J Phytopharmacol* 2020; 9(4):281–290.