

1987

Aphid Feeding Behavior: Relationship to Barley Yellow Dwarf Virus Resistance in *Agropyron* Species

Richard H. Shukle
USDA-ARS

David J. Lampe
Purdue University

Richard M. Lister
Purdue University

John E. Foster
University of Nebraska-Lincoln, john.foster@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>

 Part of the [Entomology Commons](#)

Shukle, Richard H.; Lampe, David J.; Lister, Richard M.; and Foster, John E., "Aphid Feeding Behavior: Relationship to Barley Yellow Dwarf Virus Resistance in *Agropyron* Species" (1987). *Faculty Publications: Department of Entomology*. 604.
<http://digitalcommons.unl.edu/entomologyfacpub/604>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Resistance

Aphid Feeding Behavior: Relationship to Barley Yellow Dwarf Virus Resistance in *Agropyron* Species

Richard H. Shukle, David J. Lampe, Richard M. Lister, and John E. Foster

First and fourth authors; research entomologists, USDA-ARS; second author, graduate research assistant, Department of Entomology; third author, professor, Department of Botany and Plant Pathology; Purdue University, West Lafayette, IN 47907.

We thank Dr. Eric S. Furgason and Richard G. Haywood for constructing the multichannel aphid feeding monitor used in this work. The technical assistance of Jo Anne McFtridge, Y. Adesina, and Donald R. Ross is also gratefully acknowledged.

Purdue University AES, Journal Paper 10,690.

Accepted for publication 21 October 1986 (submitted for electronic processing).

ABSTRACT

Shukle, R. H., Lampe, D. J., Lister, R. M., and Foster, J. E. 1987. Aphid feeding behavior: Relationship to barley yellow dwarf virus resistance in *Agropyron* species. *Phytopathology* 77: 725-729.

The infectibility of various *Agropyron* species (wheatgrasses) was examined with respect to three isolates of barley yellow dwarf virus (BYDV) by infesting them with appropriate vector aphids, followed by testing by enzyme-linked immunosorbent assay. Feeding behavior of the vectors *Rhopalosiphum padi* and *Sitobion avenae* was also electronically monitored to determine their ability to inoculate phloem. The results indicated that resistance to BYDV infection occurs in several *Agropyron*

species. For most species tested, resistance seemed due to failure in virus increase, but in some species a major constraint on infection was the inability of vectors to locate phloem. Two potential approaches to breeding for reduced BYDV in wheat by crossing with *Agropyron* species may thus be: incorporating factors reducing or preventing virus production and incorporating factors reducing the ability of vectors to inoculate plants successfully.

Additional key words: BYDV-resistant germ plasm, luteoviruses, wheat improvement.

The term barley yellow dwarf virus (BYDV) includes at least five variously interrelated types of luteoviruses (9,18) that are apparently restricted to gramineous plants. They cause yellow-dwarf diseases in cereals that affect the supply of wheat, oats, and barley throughout the world (6,18). Like the other luteoviruses,

they are obligatorily transmitted by aphids in a circulative manner and are restricted to phloem.

Two important vectors of BYDV are the English grain aphid, *Sitobion avenae* F., and the bird cherry oat aphid, *Rhopalosiphum padi* L. *S. avenae* transmits the MAV and PAV isolates (sensu Rochow, 17), whereas *R. padi* transmits the PAV and RPV isolates (17). Recent results in experiments with electronic monitoring of feeding behavior have provided direct evidence that penetration of phloem sieve elements is a prerequisite for BYDV transmission by these vectors (19, Lampe et al., *unpublished*). Infection with BYDV might therefore be avoided in cereal cultivars by resistance to phloem contact and ingestion.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. § 1734 solely to indicate this fact.

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 1987.

Currently, tolerance (sensu Cooper and Jones, 3) for BYDV infection occurs in wheat, *Triticum aestivum* L. em. Thell. (14,15), but there are no clear-cut examples of immunity or of the resistance (reduced virus production) that occur in some cultivars of barley (16). If such were available in wheat, or in species that can intercross with wheat, it could have considerable significance for BYDV resistance breeding programs.

In this regard, high levels of resistance (i.e., virus could not be detected by enzyme-linked immunosorbent assay [ELISA] in plants infested with viruliferous aphids) were recently reported with respect to two BYDV isolates, PAV and RMV, in several *Agropyron* species (23). There is, however, no information concerning such resistance with other isolates of BYDV, or whether it may actually reflect the inability of vectors to inoculate plants efficiently.

The objectives of the present study were: to determine if various *Agropyron* species show high levels of resistance to virus production with a PAV-like isolate of BYDV (P-PAV, 7) or with the MAV and RPV isolates of BYDV and to electronically monitor the feeding behavior of aphids on the *Agropyron* species to determine whether they are readily able to locate and ingest from phloem tissue.

MATERIALS AND METHODS

Insects and virus isolates. Viruliferous *R. padi* carrying the P-PAV or RPV isolates and *S. avenae* carrying the MAV isolate of BYDV, respectively, were obtained from cultures maintained on infected oats, *Avena sativa* L. 'Clintland 64', in isolated chambers at 21 C with a 16-hr photoperiod. Viruliferous aphids used to infest *Agropyron* species and oats and to monitor feeding behavior were apterous, virginoparous adults. The RPV and MAV isolates were subcultures maintained from cultures initially supplied by Dr. W. F. Rochow (24); the P-PAV isolate has been described (7).

Plants. Seedling plants of six wheatgrasses, *Agropyron intermedium* (Host) Beauv., *Agropyron elongatum* (Host) Beauv. 'Jose', *Agropyron sibiricum* (Willd.) Beauv., and *Agropyron smithii* Rydb. 'Arriba', 'Barton', and 'Rosanna' were grown from seed under greenhouse conditions beneath sodium lights (930 μ Einsteins/m²/sec) with a 16-hr photoperiod. Seeds of the *Agropyron* species were obtained from commercial sources. Plants were grown in sterile soil mix and irrigated with water containing a soluble NPK fertilizer (15-30-15). Tests for BYDV infectibility and aphid feeding behavior were conducted with seedling plants in the third-leaf stage (14–18 days after emergence).

Tests for infectibility with BYDV isolates. Plants were tested for infectibility with the BYDV isolates by infestation with 10 viruliferous, adult aphids (vide supra) per plant. Ten plants were used for each *Agropyron* species tested. Infested plants were maintained in isolation at 21 C (16-hr photoperiod) for 1 wk. Aphid survival was recorded and aphids were then killed by spraying with malathion. The plants were then grown for an additional 14 days under greenhouse conditions. Noninfested control plants and infested Clintland 64 oats were treated similarly. Tissue samples for ELISA consisted of each entire plant (including roots), washed clean and dried by blotting.

Extraction and ELISA. Plant samples were first pulverized in liquid nitrogen in a mortar. Potassium phosphate buffer (0.1 M, pH 7) containing 0.05% Tween-20 and 2% polyvinyl pyrrolidone (MW 40,000) was then added (1:6, w/v) along with a pinch of Carborundum, and the tissue ground further. The resulting extract was filtered through cheesecloth and stored for 1–2 wk at –76 C until assayed for virus. Virus in extracts was assayed by ELISA as described by Hammond et al (7) with the modifications of Skaria et al (24).

Monitoring aphid feeding behavior. Aphid feeding behavior was electronically monitored using a system similar to that described by McLean and Kinsey (10,11). In our system (19), a 12- μ m gold wire attached to the dorsum of the aphid with silver conducting paint serves as one electrode. After a 1-hr starvation period, the aphid is given access to the leaf of a plant and a 20 mV, 330 Hz signal is transmitted via the electrode attached to the aphid to a

ground implanted in the potting medium. During salivation and ingestion the aphid serves as a variable impedance and modulations of the signal are amplified and recorded.

This technique has been used to monitor the feeding behavior of several aphid species (2,13,19). Three basic waveform patterns are recorded using the system: an "S"-waveform corresponding to salivation and sheath-formation, an "X"-waveform corresponding to stylet penetration of sieve elements, and an "I" waveform corresponding to a continuous period of ingestion. Any combination of sequences of the three basic waveforms can occur during aphid feeding; however, an I-wave immediately preceded by an X-wave, or a series of X-waves, is considered to be diagnostic of phloem ingestion (11,19).

To record feeding behavior on *Agropyron* species, an individual aphid was placed on the second or third leaf of a seedling plant and feeding activities recorded for 180 min from the start of the first probe. All recordings were made in the laboratory at room temperature (22 C). For each *Agropyron* species, feeding behavior was recorded for six *S. avenae* and six *R. padi* on individual seedlings. The following events associated with aphid feeding were recorded: number of probes, salivation, number of phloem contacts, phloem ingestion, nonphloem ingestion, and nonprobing.

Histological association of stylet tip location. Association of stylet tip location with recorded waveforms was done histologically for *S. avenae* and *R. padi* feeding on Clintland 64 oats and *A. intermedium*. The procedure was similar to that described by McLean and Kinsey (11). At the occurrence of a waveform to be investigated, the aphid was anesthetized with CO₂. The wire connecting the aphid to the feeding monitor was then severed, and a leaf segment with the aphid in position was cut from the blade and placed into a Craff fixative containing 5% nicotine and 0.1% Triton X-100. This solution rapidly killed the anesthetized aphid. After 2 min, the leaf segment with the aphid in position was removed, trimmed to the desired size, and placed into a Craff fixative lacking the nicotine and Triton X-100. After fixation for 12–24 hr at 4 C, the tissue was dehydrated by passage through an ethanol series. The tissue was then embedded in Paraplast and 10- μ m sections cut. Throughout dehydration and embedding the aphid was left in position on the leaf segment. Sections were then stained with safranin and fast green, and the position of the aphid's stylets in the plant tissue was associated with the corresponding waveform.

RESULTS

ELISA test results. Aphids settled well on all the seedlings tested. Surviving apterae and newly deposited nymphs were noted on all of them 7 days after infestation. ELISA test results for the infested plants are summarized in Table 1. They indicated that in general, few infections had occurred among the *Agropyron* species tested. Exceptions to this were the high number of infections detected in *A. sibiricum* plants inoculated with the P-PAV isolate, and in *A. elongatum* 'Jose' inoculated with the RPV isolate. Among the *Agropyron* species inoculated with the MAV isolate, the only infections detected were in *A. elongatum* 'Jose' and *A. sibiricum* (one infection each). No infections with any of the isolates were detected in *A. intermedium* or in *A. smithii* 'Barton' and 'Rosanna'. Contrasting susceptibilities occurred to the different isolates. These were especially noticeable with *A. elongatum* 'Jose' and *A. sibiricum*.

Waveforms recorded during feeding. Typical waveforms recorded during feeding on Clintland 64 by *S. avenae* and *R. padi* are shown in Figure 1. These waveforms resemble those recorded for other species of aphids and host plants in that salivation (S), sieve element penetration (X), and ingestion (I) waveforms can be distinguished. In our system, the X-waveform corresponding to penetration of a sieve element generally is a sequence of M-shaped waves, although the shape and number of waves varies somewhat with the aphid species. The sequence of waveforms generated during probing and ingestion by *S. avenae* and *R. padi* was the same, except for the X-waveform generated upon penetration of a

TABLE 1. Enzyme-linked immunosorbent assay (ELISA) test results for seedlings of *Agropyron* species and Clintland 64 oats infested with aphids carrying the P-PAV, RPV, or MAV isolates of barley yellow dwarf virus

Host	ELISA test results								
	P-PAV isolate			RPV isolate			MAV isolate		
	No. positive ^a	Mean	(Range)	No. positive	Mean	(Range)	No. positive	Mean	(Range)
<i>A. intermedium</i>	0	—	(—)	0	—	(—)	0	—	(—)
<i>A. sibiricum</i>	9	0.27	(0.10–0.72)	4	0.13	(0.12–0.15)	1	0.14	(—)
<i>A. elongatum</i>									
cv. Jose	2	0.16	(0.11–0.20)	10	0.14	(0.08–0.24)	1	0.16	(—)
<i>A. smithii</i>									
cv. Arriba	1	0.10	(—)	2	0.11	(0.10–0.12)	0	—	(—)
cv. Barton	0	—	(—)	0	—	(—)	0	—	(—)
cv. Rosanna	0	—	(—)	0	—	(—)	0	—	(—)
<i>Avena sativa</i>									
cv. Clintland 64	5	0.90	(0.36–1.51)	4	0.37	(0.24–0.40)	6	1.66	(1.14–1.90)

^aELISA values exceeding twice those for healthy control values were rated as positive. Control values ranged from 0.03 to 0.04 for *Agropyron* spp. and 0.01 to 0.02 for Clintland 64. All tests were of 10 seedlings, except with Clintland 64, for which five, four, and six seedlings were tested with the PAV, RPV, and MAV isolates, respectively. All seedlings had surviving apterae and most had nymphs when inspected 7 days after infestation.

TABLE 2. Means of various electronically recorded events in feeding behavior of *Sitobion avenae* and *Rhopalosiphum padi* on different *Agropyron* species or Clintland 64 oats during 180 min of feeding

Host	Probes (no.)	Salivation (min)	Phloem ingestion (min)	Nonphloem ingestion (min)	Non-probing (min)
<i>S. avenae</i>					
<i>A. intermedium</i>	4 bc ^a	63 c	91 a	10 a	19 bc
<i>A. sibiricum</i>	3 c	65 c	94 a	10 a	11 c
<i>A. elongatum</i>					
(cv. Jose)	10 ab	126 a	8 cd	22 a	20 bc
<i>A. smithii</i>					
(cv. Arriba)	11 a	93 abc	2 d	17 a	56 ab
(cv. Barton)	12 a	78 bc	6 d	24 a	75 a
(cv. Rosanna)	6 abc	76 bc	44 bc	29 a	28 bc
Clintland 64 oats	8 abc	109 ab	51 ab	6 a	15 c
<i>R. padi</i>					
<i>A. intermedium</i>	6 a	58 a	41 b	49 ab	32 b
<i>A. sibiricum</i>	6 a	73 a	24 b	69 a	14 b
<i>A. elongatum</i>					
(cv. Jose)	6 a	72 a	67 ab	19 ab	21 b
<i>A. smithii</i>					
(cv. Arriba)	8 a	61 a	0 c	30 ab	88 a
(cv. Barton)	7 a	79 a	48 ab	14 ab	39 ab
(cv. Rosanna)	4 a	80 a	60 ab	18 ab	22 b
Clintland 64 oats	4 a	55 a	100 a	7 b	18 b

^aMeans ($N=6$) followed by different letters within columns for *S. avenae* and *R. padi*, respectively, are significantly different, $P<0.05$, Duncan's multiple range test with transformed scores ($\sqrt{X+1}$).

sieve element (Fig. 1). *S. avenae* generated two to six X-waves before commencing phloem ingestion, whereas *R. padi* generated only one. Additionally, the shape of the X-waveform varied slightly with the two aphid species.

Association of waveforms with stylet tip location. Histological preparations of six aphids of both species feeding on Clintland 64 oats showed that when the X-waveform or the S-X-I sequence were recorded, aphid stylets were invariably in contact with phloem. By contrast, in 12 preparations of aphids of both species where only the S-waveform or the S-I sequence were recorded, stylets were not in contact with phloem. It was sometimes possible to determine that stylet tips were inside a cell during the I-waveform (Fig. 2), thereby confirming that this waveform was recorded during ingestion, whereas S-waveforms were recorded during salivation.

Aphid feeding behavior. Means of various electronically recorded events in the probing/ingestion behavior of *S. avenae* and *R. padi* during 180 min of feeding on seedlings of the *Agropyron* species or Clintland 64 oats are shown in Table 2. Cultivars Arriba and Barton of *A. smithii* apparently resisted phloem contact by *S. avenae*. On both of these cultivars only two of the six aphids (33%) whose feeding behavior was monitored contacted phloem, and the

subsequent ingestion was for a short time (mean of 2 min for Arriba and 6 min for Barton). With cultivar Rosanna of *A. smithii*, however, all six *S. avenae* tested contacted phloem and ingested for relatively long periods (mean of 44 min).

A. elongatum 'Jose' also appeared resistant to phloem contact and ingestion by *S. avenae*, though not by *R. padi*. Only two of six *S. avenae* contacted phloem on this *Agropyron* species, and the subsequent ingestion of phloem sap was for a short time (mean of 8 min). *A. intermedium* and *A. sibiricum*, however, were both susceptible to phloem contact by *S. avenae*.

Within the species *A. smithii*, cultivar Arriba also seemed highly resistant to phloem contact by *R. padi* (i.e., none of the aphids tested contacted phloem). Phloem contact and ingestion was observed, however, with *A. smithii* cultivars Barton and Rosanna, as well as with the other *Agropyron* species tested (Table 2).

DISCUSSION

Sharma et al (23) reported that they were unable to detect the PAV and RMV isolates of BYDV by ELISA in plants of several species in the genus *Agropyron* infested with viruliferous aphids. Similarly, we have here identified several *Agropyron* species in which we could not detect P-PAV, RPV, or MAV isolates of BYDV by ELISA in a high proportion of plants infested with viruliferous aphids under conditions in which in this and many other experiments 100% of Clintland 64 oat plants become infected (Table 1). Interpretation of the results requires caution, but in general they clearly indicate considerable resistance (sensu Cooper and Jones, 3), if not immunity, to BYDV among the *Agropyron* species tested.

Monitoring the feeding behavior of *S. avenae* and *R. padi* (Table 2) provided direct evidence that the resistances to the P-PAV, RPV, and MAV isolates of BYDV apparent in those *Agropyron* species where aphids easily contacted and ingested from phloem were unlikely to result from inability of aphid vectors to inoculate the plants. Resistance in these plants would seem, therefore, to occur at the level of some virus-host interaction that prevents the virus from establishing or, perhaps, from replicating effectively. An example of this is illustrated by *A. intermedium*. For this species, ELISA indicated resistance to infection with the P-PAV, RPV, and MAV isolates (Table 1), but monitoring of feeding behavior (Table 2) showed that both *R. padi* and *S. avenae* contacted phloem easily and ingested for long periods. In contrast to this, resistance to phloem contact by *R. padi* and *S. avenae* seems likely to be a major component of the apparent resistance to BYDV observed with cultivar Arriba of *A. smithii* (Table 1), for which phloem contact was drastically reduced (Table 2). Though in tests for resistance with this cultivar a few plants gave positive ELISA values (Table 1), these may have resulted from occasional successful phloem contact by a few aphids.

A. smithii 'Barton' and *A. elongatum* 'Jose' demonstrated




Fig. 1. Typical waveforms recorded from (A) *Sitobion avenae* and (B) *Rhopalosiphum padi* feeding on Clintland 64 oats. Arrows indicate initiation of probes. S: salivation waveform, X: penetration of sieve element, I: continuous ingestion. The sequence S-X-I_p indicates ingestion from phloem, whereas the sequence S-I_{np} indicates ingestion from nonphloem tissue.




Fig. 2. Stylet tips located in a phloem cell from *Sitobion avenae* feeding on Clintland 64 oats. Aphid had generated an S-X-I waveform sequence. P: phloem cells, X: xylem vessel, ST: stylet tips, SS: salivary sheath, S: stylets within salivary sheath.

reduced phloem contact by *S. avenae*, but not by *R. padi*. Resistance to infection with the MAV isolate by *S. avenae* would therefore be expected with these cultivars, but ELISA also indicated this for the P-PAV isolate transmitted by *R. padi*. Contrast in resistance to phloem contact by *S. avenae* and *R. padi* raises the question of why one aphid species can contact phloem, while another has difficulty. Both aphid species probe these cultivars and ingest from tissues other than phloem, as they do with all of the *Agropyron* species tested. Thus, the resistance to phloem contact is not because aphids do not attempt to probe and feed. Perhaps it is due to differences in enzymes secreted during stylet penetration of plant tissues or in gustatory reactions that limit the ability of *S. avenae* to contact phloem, but not that of *R. padi*.

Resistance to phloem contact by *Schizaphis graminum* (Rondani) has been suggested to be associated with the chemical composition of intercellular pectins through which aphids probe (4,5,12). Resistance to location of phloem and ingestion of phloem sap by aphids could also be due to other factors such as lack of appropriate gustatory stimuli or the presence of repellents.

Our results suggest that "resistance" to the P-PAV, RPV, and MAV isolates of BYDV in the genus *Agropyron* can occur not only at the level of virus increase but also as a result of impaired vector efficiency due to failure to locate phloem. Because *Agropyron* species have been successfully crossed with wheat (21,22), there is a potential for introducing either or both of these kinds of "resistance" into wheat. *Agropyron* species have in fact been used successfully as sources of disease-resistant germ plasm for wheat improvement in the past (1,8,20,25).

LITERATURE CITED

1. Browder, L. E. 1980. A compendium of information about named genes for low reaction to *Puccinia recondita*. *Crop Sci.* 20:775-779.
2. Campbell, B. C., McLean, D. L., Kinsey, M. G., Jones, K. C., and Dreyer, D. L. 1982. Probing behavior of the greenbug (*Schizaphis graminum*, biotype C) on resistant and susceptible varieties of sorghum. *Entomol. Exp. Appl.* 31:140-156.
3. Cooper, J. I., and Jones, A. T. 1983. Responses of plants to viruses: Proposals for the use of terms. *Phytopathology* 73:127-128.
4. Dreyer, D. L., and Campbell, B. C. 1984. Association of the degree of methylation of intercellular pectin with plant resistance to aphids and with induction of aphid biotypes. *Experientia* 40:224-226.
5. Dreyer, D. L., Campbell, B. C., and Jones, K. C. 1984. Effect of bioregulator-treated sorghum on greenbug fecundity and feeding behavior: Implications for host-plant resistance. *Phytochemistry* 23:1593-1596.
6. Duffus, J. E. 1977. Aphids, viruses and the yellow plague. Pages 365-366 in: *Aphids as Virus Vectors*. K. F. Harris and E. Kurstak, eds. Academic Press, New York. 599 pp.
7. Hammond, J., Lister, R. M., and Foster, J. E. 1983. Purification, identity and some properties of an isolate of barley yellow dwarf virus from Indiana. *J. Gen. Virol.* 64:667-678.
8. Love, R. M., and Suneson, C. A. 1945. Cytogenetics of certain *Triticum-Agropyron* hybrids and their fertile derivatives. *Am. J. Bot.* 32:451-779.
9. Matthews, R. E. F. 1981. The groups of plant viruses. Pages 720-721 in: *Plant Virology*, 2nd ed. Academic Press, New York. 897 pp.
10. McLean, D. L., and Kinsey, M. C. 1964. A technique for electronically recording aphid feeding and salivation. *Nature (London)* 202:1358-1359.
11. McLean, D. L., and Kinsey, M. G. 1967. Probing behavior of the pea aphid *Acyrtosiphon pisum*. I. Definitive correlation of electronically recorded waveforms with aphid probing activities. *Ann. Entomol. Soc. Am.* 60:400-406.
12. Montllor, C. B., Campbell, B. C., and Mittler, T. E. 1983. Natural and induced differences in probing behavior of two biotypes of the greenbug, *Schizaphis graminum*, in relation to resistance in sorghum. *Entomol. Exp. Appl.* 34:99-106.
13. Nielson, M. W., and Don, H. 1974. Probing behavior of biotypes of the spotted alfalfa aphid on resistant and susceptible alfalfa clones. *Entomol. Exp. Appl.* 17:477-486.
14. Ohm, H. W., Patterson, F. L., Carrigan, L. L., Shaner, G. E., Foster, J. E., Finney, R. E., and Roberts, J. J. 1981. Registration of elmo common wheat germplasm. *Crop Sci.* 21:803.
15. Patterson, F. L., Ohm, H. W., Shaner, G. E., Finney, R. E., Gallun, R. L., Roberts, J. J., and Foster, J. E. 1982. Registration of Caldwell wheat. *Crop Sci.* 22:691-692.
16. Rasmusson, D. C., and Schaller, C. W. 1959. The inheritance of resistance in barley to the yellow-dwarf virus. *Agron. J.* 51:661-665.
17. Rochow, W. F. 1969. Biological properties of four isolates of barley yellow dwarf virus. *Phytopathology* 59:1580-1589.
18. Rochow, W. F., and Duffus, J. E. 1981. Luteoviruses and yellow diseases. Pages 147-170 in: *Handbook of Plant Virus Infections and Comparative Diagnosis*. E. Kurstak, ed. Elsevier North-Holland Biomedical Press, Amsterdam.
19. Scheller, H. V., and Shukle, R. H. 1986. Feeding behavior and transmission of barley yellow dwarf virus by *Sitobion avenae* on oats. *Entomol. Exp. Appl.* 40:189-195.
20. Schmidt, J. W., Sill, W. H., and Fellows, H. 1956. Range of reactions to wheat streak mosaic virus in hybrids derived from *Triticum vulgare* × *Agropyron elongatum*. *Agron. J.* 48:371-373.
21. Sharma, H. C., and Gill, B. S. 1983. Current status of wide hybridization in wheat. *Euphytica* 32:17-31.
22. Sharma, H. C., and Gill, B. S. 1983. New hybrids between *Agropyron* and wheat. II. Production, morphology and cytogenetic analysis of F1 hybrids and backcross derivatives. *Theor. Appl. Genet.* 66:111-121.
23. Sharma, H. C., Gill, B. S., and Uyemoto, J. K. 1984. High levels of resistance in *Agropyron* species to barley yellow dwarf and wheat streak mosaic viruses. *Phytopathol. Z.* 110:143-147.
24. Skaria, M., Lister, R. M., and Foster, J. E. 1984. Lack of barley yellow dwarf virus dosage effects on virus content in cereals. *Plant Dis.* 68:759-761.
25. Wells, D. O., Kota, R. S., Sandhu, H. B., Gardner, W. S., and Finney, K. F. 1982. Registration of one disomic substitution line and five translocation lines of winter wheat germplasm resistant to wheat streak mosaic virus. *Crop Sci.* 22:1277-1278.