

RESEARCH

Open Access



# Applications and common coupled fixed point results in ordered partial metric spaces

KPR Rao<sup>1</sup>, GNV Kishore<sup>2</sup>, Kenan Tas<sup>3</sup>, S Satyanaraya<sup>4\*</sup> and D Ram Prasad<sup>5</sup>

\*Correspondence:  
s.satyan1@gmail.com  
<sup>4</sup>Department of Computing, Adama  
Science and Technology University,  
Adama, Ethiopia  
Full list of author information is  
available at the end of the article

## Abstract

In this paper, we obtain a unique common coupled fixed point theorem by using  $(\psi, \alpha, \beta)$ -contraction in ordered partial metric spaces. We give an application to integral equations as well as homotopy theory. Also we furnish an example which supports our theorem.

**Keywords:** partial metric;  $w$ -compatible maps; coupled fixed point; mixed  $g$ -monotone property;  $\psi$ - $\alpha$ - $\beta$  contraction; homotopy theory

## 1 Introduction

The notion of a partial metric space (PMS) was introduced by Matthews [1] as a part of the study of denotational semantics of data flow networks. In fact, it is widely recognized that PMSs play an important role in constructing models in the theory of computation and domain theory in computer science (see *e.g.* [2–9]).

Matthews [1, 10], Oltra and Valero [11] and Altun *et al.* [12] proved some fixed point theorems in PMSs for a single map. For more work on fixed, common fixed point theorems in PMSs, we refer to [6, 13–27].

The notion of a coupled fixed point was introduced by Bhaskar and Lakshmikantham [28] and they studied some fixed point theorems in partially ordered metric spaces. Later some authors proved coupled fixed and coupled common fixed point theorems (see [16, 29–35]).

The aim of this paper is to study unique common coupled fixed point theorems of Jungck type maps by using a  $(\psi, \alpha, \beta)$ -contraction condition over partially ordered PMSs.

## 2 Preliminaries

First we recall some basic definitions and lemmas which play a crucial role in the theory of PMSs.

**Definition 2.1** (See [1, 10]) A partial metric on a non-empty set  $X$  is a function  $p : X \times X \rightarrow R^+$  such that, for all  $x, y, z \in X$ ,

$$(p_1) \quad x = y \Leftrightarrow p(x, x) = p(x, y) = p(y, y),$$

$$(p_2) \quad p(x, x) \leq p(x, y), p(y, y) \leq p(x, y),$$

- (p<sub>3</sub>)  $p(x, y) = p(y, x)$ ,
- (p<sub>4</sub>)  $p(x, y) \leq p(x, z) + p(z, y) - p(z, z)$ .

The pair  $(X, p)$  is called a PMS.

If  $p$  is a partial metric on  $X$ , then the function  $d_p : X \times X \rightarrow \mathbb{R}^+$ , given by

$$d_p(x, y) = 2p(x, y) - p(x, x) - p(y, y), \tag{1}$$

is a metric on  $X$ .

**Example 2.2** (See e.g. [10, 14, 20]) Consider  $X = [0, \infty)$  with  $p(x, y) = \max\{x, y\}$ . Then  $(X, p)$  is a PMS. It is clear that  $p$  is not a (usual) metric. Note that in this case  $d_p(x, y) = |x - y|$ .

**Example 2.3** (See [19]) Let  $X = \{[a, b] : a, b \in \mathbb{R}, a \leq b\}$  and define  $p([a, b], [c, d]) = \max\{b, d\} - \min\{a, c\}$ . Then  $(X, p)$  is a PMS.

Each partial metric  $p$  on  $X$  generates a  $T_0$  topology  $\tau_p$  on  $X$  which has as a base the family of open  $p$ -balls  $\{B_p(x, \varepsilon), x \in X, \varepsilon > 0\}$ , where  $B_p(x, \varepsilon) = \{y \in X : p(x, y) < p(x, x) + \varepsilon\}$  for all  $x \in X$  and  $\varepsilon > 0$ .

We now state some basic topological notions (such as convergence, completeness, continuity) on PMSs (see e.g. [1, 10, 12, 14, 20, 22]).

**Definition 2.4**

1. A sequence  $\{x_n\}$  in the PMS  $(X, p)$  converges to the limit  $x$  if and only if  $p(x, x) = \lim_{n \rightarrow \infty} p(x, x_n)$ .
2. A sequence  $\{x_n\}$  in the PMS  $(X, p)$  is called a Cauchy sequence if  $\lim_{n, m \rightarrow \infty} p(x_n, x_m)$  exists and is finite.
3. A PMS  $(X, p)$  is called complete if every Cauchy sequence  $\{x_n\}$  in  $X$  converges with respect to  $\tau_p$ , to a point  $x \in X$  such that  $p(x, x) = \lim_{n, m \rightarrow \infty} p(x_n, x_m)$ .
4. A mapping  $F : X \rightarrow X$  is said to be continuous at  $x_0 \in X$  if, for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $F(B_p(x_0, \delta)) \subseteq B_p(Fx_0, \epsilon)$ .

The following lemma is one of the basic results as regards PMS [1, 10, 12, 14, 20, 22].

**Lemma 2.5**

1. A sequence  $\{x_n\}$  is a Cauchy sequence in the PMS  $(X, p)$  if and only if it is a Cauchy sequence in the metric space  $(X, d_p)$ .
2. A PMS  $(X, p)$  is complete if and only if the metric space  $(X, d_p)$  is complete. Moreover,

$$\lim_{n \rightarrow \infty} d_p(x, x_n) = 0 \iff p(x, x) = \lim_{n \rightarrow \infty} p(x, x_n) = \lim_{n, m \rightarrow \infty} p(x_n, x_m). \tag{2}$$

Next, we give two simple lemmas which will be used in the proofs of our main results. For the proofs we refer [14].

**Lemma 2.6** Assume  $x_n \rightarrow z$  as  $n \rightarrow \infty$  in a PMS  $(X, p)$  such that  $p(z, z) = 0$ . Then  $\lim_{n \rightarrow \infty} p(x_n, y) = p(z, y)$  for every  $y \in X$ .

**Lemma 2.7** *Let  $(X, p)$  be a PMS. Then*

- (A) *if  $p(x, y) = 0$ , then  $x = y$ ,*
- (B) *if  $x \neq y$ , then  $p(x, y) > 0$ .*

**Remark 2.8** *If  $x = y$ ,  $p(x, y)$  may not be 0.*

**Definition 2.9** ([28]) *Let  $(X, \preceq)$  be a partially ordered set and  $F : X \times X \rightarrow X$ . Then the map  $F$  is said to have mixed monotone property if  $F(x, y)$  is monotone non-decreasing in  $x$  and monotone non-increasing in  $y$ ; that is, for any  $x, y \in X$ ,*

$$x_1 \preceq x_2 \text{ implies } F(x_1, y) \preceq F(x_2, y) \text{ for all } y \in X$$

and

$$y_1 \preceq y_2 \text{ implies } F(x, y_2) \preceq F(x, y_1) \text{ for all } x \in X.$$

**Definition 2.10** ([28]) *An element  $(x, y) \in X \times X$  is called a coupled fixed point of a mapping  $F : X \times X \rightarrow X$  if  $F(x, y) = x$  and  $F(y, x) = y$ .*

**Definition 2.11** ([30]) *An element  $(x, y) \in X \times X$  is called*

- (g<sub>1</sub>) *a coupled coincident point of mappings  $F : X \times X \rightarrow X$  and  $f : X \rightarrow X$  if  $fx = F(x, y)$  and  $fy = F(y, x)$ ,*
- (g<sub>2</sub>) *a common coupled fixed point of mappings  $F : X \times X \rightarrow X$  and  $f : X \rightarrow X$  if  $x = fx = F(x, y)$  and  $y = fy = F(y, x)$ .*

**Definition 2.12** ([30]) *The mappings  $F : X \times X \rightarrow X$  and  $f : X \rightarrow X$  are called  $w$ -compatible if  $f(F(x, y)) = F(fx, fy)$  and  $f(F(y, x)) = F(fy, fx)$  whenever  $fx = F(x, y)$  and  $fy = F(y, x)$ .*

Inspired by Definition 2.9, Lakshmikantham and Ćirić in [31] introduced the concept of a  $g$ -mixed monotone mapping.

**Definition 2.13** ([31]) *Let  $(X, \preceq)$  be a partially ordered set,  $F : X \times X \rightarrow X$  and  $g : X \rightarrow X$  be mappings. Then the map  $F$  is said to have a mixed  $g$ -monotone property if  $F(x, y)$  is monotone  $g$ -non-decreasing in  $x$  as well as monotone  $g$ -non-increasing in  $y$ ; that is, for any  $x, y \in X$ ,*

$$gx_1 \preceq gx_2 \text{ implies } F(x_1, y) \preceq F(x_2, y) \text{ for all } y \in X$$

and

$$gy_1 \preceq gy_2 \text{ implies } F(x, y_2) \preceq F(x, y_1) \text{ for all } x \in X.$$

Now we prove our main results.

### 3 Results and discussions

**Definition 3.1** Let  $(X, p)$  be a PMS, let  $F : X \times X \rightarrow X$  and  $g : X \rightarrow X$  be mappings. We say that  $F$  satisfies a  $(\psi, \alpha, \beta)$ -contraction with respect to  $g$  if there exist  $\psi, \alpha, \beta : [0, \infty) \rightarrow [0, \infty)$  satisfying the following:

- (3.1.1)  $\psi$  is continuous and monotonically non-decreasing,  $\alpha$  is continuous and  $\beta$  is lower semi continuous,
- (3.1.2)  $\psi(t) = 0$  if and only if  $t = 0, \alpha(0) = \beta(0) = 0,$
- (3.1.3)  $\psi(t) - \alpha(t) + \beta(t) > 0$  for  $t > 0,$
- (3.1.4)  $\psi(p(F(x, y), F(u, v))) \leq \alpha(M(x, y, u, v)) - \beta(M(x, y, u, v)), \forall x, y, u, v \in X, gx \leq gu, gy \geq gv$  and

$$M(x, y, u, v) = \max \left\{ p(gx, gu), p(gy, gv), p(gx, F(x, y)), p(gy, F(y, x)), p(gu, F(u, v)), p(gv, F(v, u)), \frac{p(gx, F(x, y))p(gy, F(y, x))}{1+p(gx, gu)+p(gy, gv)+p(F(x, y), F(u, v))}, \frac{p(gu, F(u, v))p(gv, F(v, u))}{1+p(gx, gu)+p(gy, gv)+p(F(x, y), F(u, v))} \right\}.$$

**Theorem 3.2** Let  $(X, \preceq)$  be a partially ordered set and  $p$  be a partial metric such that  $(X, p)$  is a PMS. Let  $F : X \times X \rightarrow X$  and  $g : X \rightarrow X$  be such that

- (3.2.1)  $F$  satisfies a  $(\psi, \alpha, \beta)$ -contraction with respect to  $g,$
- (3.2.2)  $F(X \times X) \subseteq g(X)$  and  $g(X)$  is a complete subspace of  $X,$
- (3.2.3)  $F$  has a mixed  $g$ -monotone property,
- (3.2.4) (a) if a non-decreasing sequence  $\{x_n\} \rightarrow x,$  then  $x_n \preceq x$  for all  $n,$   
 (b) if a non-increasing sequence  $\{y_n\} \rightarrow y,$  then  $y \preceq y_n$  for all  $n.$

If there exist  $x_0, y_0 \in X$  such that  $gx_0 \preceq F(x_0, y_0)$  and  $gy_0 \succeq F(y_0, x_0),$  then  $F$  and  $g$  have a coupled coincidence point in  $X \times X.$

*Proof* Let  $x_0, y_0 \in X$  be such that  $gx_0 \preceq F(x_0, y_0)$  and  $gy_0 \succeq F(y_0, x_0).$  Since  $F(X \times X) \subseteq g(X),$  we choose  $x_1, y_1 \in X$  such that

$$gx_0 \preceq F(x_0, y_0) = gx_1 \quad \text{and} \quad gy_0 \succeq F(y_0, x_0) = gy_1$$

and choose  $x_2, y_2 \in X$  such that

$$gx_2 = F(x_1, y_1) \quad \text{and} \quad gy_2 = F(y_1, x_1).$$

Since  $F$  has the mixed  $g$ -monotone property, we obtain

$$gx_0 \preceq gx_1 \preceq gx_2 \quad \text{and} \quad gy_0 \succeq gy_1 \succeq gy_2.$$

Continuing this process, we construct the sequences  $\{x_n\}$  and  $\{y_n\}$  in  $X$  such that

$$gx_{n+1} = F(x_n, y_n) \quad \text{and} \quad gy_{n+1} = F(y_n, x_n), \quad n = 0, 1, 2, \dots$$

with

$$\left. \begin{aligned} gx_0 &\preceq gx_1 \preceq gx_2 \preceq \dots & \text{and} \\ gy_0 &\succeq gy_1 \succeq gy_2 \succeq \dots \end{aligned} \right\} \tag{I}$$

Case (a): If  $gx_m = gx_{m+1}$  and  $gy_m = gy_{m+1}$  for some  $m$ , then  $(x_m, y_m)$  is a coupled coincidence point in  $X \times X$ .

Case (b): Assume  $gx_n \neq gx_{n+1}$  or  $gy_n \neq gy_{n+1}$  for all  $n$ .

Since  $gx_n \leq gx_{n+1}$  and  $gy_n \geq gy_{n+1}$ , from (3.2.1), we obtain

$$\begin{aligned} \psi(p(gx_n, gx_{n+1})) &= \psi(p(F(x_{n-1}, y_{n-1}), F(x_n, y_n))) \\ &\leq \alpha(M(x_{n-1}, y_{n-1}, x_n, y_n)) - \beta(M(x_{n-1}, y_{n-1}, x_n, y_n)), \\ M(x_{n-1}, y_{n-1}, x_n, y_n) &= \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), p(gx_{n-1}, gx_n), \\ p(gy_{n-1}, gy_n), p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}), \\ \frac{p(gx_{n-1}, gx_n)p(gy_{n-1}, gy_n)}{1+p(gx_{n-1}, gx_n)+p(gy_{n-1}, gy_n)+p(gx_n, gx_{n+1})}, \\ \frac{p(gx_n, gx_{n+1})p(gy_n, gy_{n+1})}{1+p(gx_{n-1}, gx_n)+p(gy_{n-1}, gy_n)+p(gx_n, gx_{n+1})} \end{array} \right\}. \end{aligned}$$

But

$$\frac{p(gx_{n-1}, gx_n)p(gy_{n-1}, gy_n)}{1 + p(gx_{n-1}, gx_n) + p(gy_{n-1}, gy_n) + p(gx_n, gx_{n+1})} \leq \max\{p(gx_{n-1}, gx_n), p(gx_n, gx_{n+1})\}$$

and

$$\frac{p(gx_n, gx_{n+1})p(gy_n, gy_{n+1})}{1 + p(gx_{n-1}, gx_n) + p(gy_{n-1}, gy_n) + p(gx_n, gx_{n+1})} \leq p(gy_n, gy_{n+1}).$$

Therefore

$$M(x_{n-1}, y_{n-1}, x_n, y_n) = \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\}.$$

Hence

$$\begin{aligned} \psi(p(gx_n, gx_{n+1})) &\leq \alpha \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right) \\ &\quad - \beta \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right). \end{aligned}$$

Similarly

$$\begin{aligned} \psi(p(gy_n, gy_{n+1})) &\leq \alpha \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right) \\ &\quad - \beta \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right). \end{aligned}$$

Put  $R_n = \max\{p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1})\}$ . Let us suppose that

$$R_n \neq 0 \quad \text{for all } n \geq 1. \tag{3}$$

Let, if possible, for some  $n$ ,  $R_{n-1} < R_n$ .

Now

$$\begin{aligned}
 \psi(R_n) &= \psi(\max\{p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1})\}) \\
 &= \max\{\psi(p(gx_n, gx_{n+1})), \psi(p(gy_n, gy_{n+1}))\} \\
 &\leq \alpha \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right) \\
 &\quad - \beta \left( \max \left\{ \begin{array}{l} p(gx_{n-1}, gx_n), p(gy_{n-1}, gy_n), \\ p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1}) \end{array} \right\} \right) \\
 &= \alpha(\max\{R_{n-1}, R_n\}) - \beta(\max\{R_{n-1}, R_n\}) \\
 &= \alpha(R_n) - \beta(R_n).
 \end{aligned}$$

From (3.1.2) and (3.1.3), it follows that  $R_n = 0$ , a contradiction.

Hence

$$R_n \leq R_{n-1}. \tag{4}$$

Thus  $\{R_n\}$  is a non-increasing sequence of non-negative real numbers and must converge to a real number  $r \geq 0$ .

Also

$$\psi(R_n) \leq \alpha(R_{n-1}) - \beta(R_{n-1}).$$

Letting  $n \rightarrow \infty$ , we get

$$\psi(r) \leq \alpha(r) - \beta(r).$$

From (3.1.2) and (3.1.3), we get  $r = 0$ . Thus

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \max\{p(gx_n, gx_{n+1}), p(gy_n, gy_{n+1})\} &= 0, \\
 \lim_{n \rightarrow \infty} p(gx_n, gx_{n+1}) = 0 &= \lim_{n \rightarrow \infty} p(gy_n, gy_{n+1}).
 \end{aligned} \tag{5}$$

Hence from  $(p_2)$ , we have

$$\lim_{n \rightarrow \infty} p(gx_n, gx_n) = 0 = \lim_{n \rightarrow \infty} p(gy_n, gy_n). \tag{6}$$

From (5) and (6) and by the definition of  $d_p$ , we get

$$\lim_{n \rightarrow \infty} d_p(gx_n, gx_{n+1}) = 0 = \lim_{n \rightarrow \infty} d_p(gy_n, gy_{n+1}). \tag{7}$$

Now we prove that  $\{gx_n\}$  and  $\{gy_n\}$  are Cauchy sequences.

To the contrary, suppose that  $\{gx_n\}$  or  $\{gy_n\}$  is not Cauchy.

This implies that  $d_p(gx_m, gx_n) \not\rightarrow 0$  or  $d_p(gy_m, gy_n) \not\rightarrow 0$  as  $n, m \rightarrow \infty$ .

Consequently

$$\max\{d_p(gx_m, gx_n), d_p(gy_m, gy_n)\} \not\rightarrow 0 \text{ as } n, m \rightarrow \infty.$$

Then there exist an  $\epsilon > 0$  and monotone increasing sequences of natural numbers  $\{m_k\}$  and  $\{n_k\}$  such that  $n_k > m_k > k$ . We have

$$\max\{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \geq \epsilon \tag{8}$$

and

$$\max\{d_p(gx_{m_k}, gx_{n_k-1}), d_p(gy_{m_k}, gy_{n_k-1})\} < \epsilon. \tag{9}$$

From (8) and (9), we have

$$\begin{aligned} \epsilon &\leq \max\{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \\ &\leq \max\{d_p(gx_{m_k}, gx_{n_k-1}), d_p(gy_{m_k}, gy_{n_k-1})\} \\ &\quad + \max\{d_p(gx_{n_k-1}, gx_{n_k}), d_p(gy_{n_k-1}, gy_{n_k})\} \\ &< \epsilon + \max\{d_p(gx_{n_k-1}, gx_{n_k}), d_p(gy_{n_k-1}, gy_{n_k})\}. \end{aligned}$$

Letting  $k \rightarrow \infty$  and using (7), we get

$$\lim_{k \rightarrow \infty} \max\{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} = \epsilon. \tag{10}$$

By the definition of  $d_p$  and using (6) we get

$$\lim_{k \rightarrow \infty} \max\{p(gx_{m_k}, gx_{n_k}), p(gy_{m_k}, gy_{n_k})\} = \frac{\epsilon}{2}. \tag{11}$$

From (8), we have

$$\begin{aligned} \epsilon &\leq \max\{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \\ &\leq \max\{d_p(gx_{m_k}, gx_{m_k-1}), d_p(gy_{m_k}, gy_{m_k-1})\} \\ &\quad + \max\{d_p(gx_{m_k-1}, gx_{n_k}), d_p(gy_{m_k-1}, gy_{n_k})\} \\ &\leq 2 \max\{d_p(gx_{m_k}, gx_{m_k-1}), d_p(gy_{m_k}, gy_{m_k-1})\} \\ &\quad + \max\{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\}. \end{aligned} \tag{12}$$

Letting  $k \rightarrow \infty$ , using (7), (10) and (12), we get

$$\lim_{k \rightarrow \infty} \max\{d_p(gx_{m_k-1}, gx_{n_k}), d_p(gy_{m_k-1}, gy_{n_k})\} = \epsilon. \tag{13}$$

Hence, we get

$$\lim_{k \rightarrow \infty} \max\{p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k})\} = \frac{\epsilon}{2}. \tag{14}$$

From (9), we have

$$\begin{aligned}
 \epsilon &\leq \max \{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \\
 &\leq \max \{d_p(gx_{m_k}, gx_{m_k-1}), d_p(gy_{m_k}, gy_{m_k-1})\} \\
 &\quad + \max \{d_p(gx_{m_k-1}, gx_{n_k+1}), d_p(gy_{m_k-1}, gy_{n_k+1})\} \\
 &\quad + \max \{d_p(gx_{n_k+1}, gx_{n_k}), d_p(gy_{n_k+1}, gy_{n_k})\} \\
 &\leq 2 \max \{d_p(gx_{m_k}, gx_{m_k-1}), d_p(gy_{m_k}, gy_{m_k-1})\} \\
 &\quad + \max \{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \\
 &\quad + 2 \max \{d_p(gx_{n_k}, gx_{n_k+1}), d_p(gy_{n_k}, gy_{n_k+1})\}.
 \end{aligned}
 \tag{15}$$

Letting  $k \rightarrow \infty$ , using (7), (10) and (15), we get

$$\lim_{k \rightarrow \infty} \max \{d_p(gx_{m_k-1}, gx_{n_k+1}), d_p(gy_{m_k-1}, gy_{n_k+1})\} = \epsilon.
 \tag{16}$$

Hence, we have

$$\lim_{k \rightarrow \infty} \max \{p(gx_{m_k-1}, gx_{n_k+1}), p(gy_{m_k-1}, gy_{n_k+1})\} = \frac{\epsilon}{2}.
 \tag{17}$$

Now from (8), we have

$$\begin{aligned}
 \epsilon &\leq \max \{d_p(gx_{m_k}, gx_{n_k}), d_p(gy_{m_k}, gy_{n_k})\} \\
 &\leq \max \{d_p(gx_{m_k}, gx_{n_k+1}), d_p(gy_{m_k}, gy_{n_k+1})\} \\
 &\quad + \max \{d_p(gx_{n_k+1}, gx_{n_k}), d_p(gy_{n_k+1}, gy_{n_k})\}.
 \end{aligned}$$

Letting  $k \rightarrow \infty$  and using (7), we obtain

$$\begin{aligned}
 \epsilon &\leq \lim_{k \rightarrow \infty} \max \{d_p(gx_{m_k}, gx_{n_k+1}), d_p(gy_{m_k}, gy_{n_k+1})\} + 0 \\
 &\leq \lim_{k \rightarrow \infty} \max \left\{ \begin{aligned} &2p(gx_{m_k}, gx_{n_k+1}) - p(gx_{m_k}, gx_{m_k}) - p(gx_{n_k+1}, gx_{n_k+1}), \\ &2p(gy_{m_k}, gy_{n_k+1}) - p(gy_{m_k}, gy_{m_k}) - p(gy_{n_k+1}, gy_{n_k+1}) \end{aligned} \right\} \\
 &= 2 \lim_{k \rightarrow \infty} \max \{p(gx_{m_k}, gx_{n_k+1}), p(gy_{m_k}, gy_{n_k+1})\}, \text{ from (6).}
 \end{aligned}$$

Thus,

$$\frac{\epsilon}{2} \leq \lim_{k \rightarrow \infty} \max \{p(gx_{m_k}, gx_{n_k+1}), p(gy_{m_k}, gy_{n_k+1})\}.$$

By the properties of  $\psi$ ,

$$\begin{aligned}
 \psi\left(\frac{\epsilon}{2}\right) &\leq \lim_{k \rightarrow \infty} \psi(\max \{p(gx_{m_k}, gx_{n_k+1}), p(gy_{m_k}, gy_{n_k+1})\}) \\
 &= \lim_{k \rightarrow \infty} \max \{\psi(p(gx_{m_k}, gx_{n_k+1})), \psi(p(gy_{m_k}, gy_{n_k+1}))\}.
 \end{aligned}
 \tag{18}$$



Now

$$\begin{aligned} \psi(p(gx_{m_k}, gx_{n_k+1})) &= \psi(p(F(x_{m_k-1}, y_{m_k-1}), F(x_{n_k}, y_{n_k}))) \\ &\leq \alpha(M(x_{m_k-1}, y_{m_k-1}, x_{n_k}, y_{n_k})) - \beta(M(x_{m_k-1}, y_{m_k-1}, x_{n_k}, y_{n_k})) \\ &= \alpha \left( \max \left\{ \begin{aligned} &p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k}), p(gx_{m_k-1}, gx_{m_k}), \\ &p(gy_{m_k-1}, gy_{m_k}), p(gx_{n_k}, gx_{n_k+1}), p(gy_{n_k}, gy_{n_k+1}), \\ &\frac{p(gx_{m_k-1}, gx_{m_k})p(gy_{m_k-1}, gy_{m_k})}{1+p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k})+p(gx_{m_k}, gx_{n_k+1})}, \\ &\frac{p(gx_{n_k}, gx_{n_k+1})p(gy_{n_k}, gy_{n_k+1})}{1+p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k})+p(gx_{m_k}, gx_{n_k+1})} \end{aligned} \right\} \right) \\ &\quad - \beta \left( \max \left\{ \begin{aligned} &p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k}), p(gx_{m_k-1}, gx_{m_k}), \\ &p(gy_{m_k-1}, gy_{m_k}), p(gx_{n_k}, gx_{n_k+1}), p(gy_{n_k}, gy_{n_k+1}), \\ &\frac{p(gx_{m_k-1}, gx_{m_k})p(gy_{m_k-1}, gy_{m_k})}{1+p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k})+p(gx_{m_k}, gx_{n_k+1})}, \\ &\frac{p(gx_{n_k}, gx_{n_k+1})p(gy_{n_k}, gy_{n_k+1})}{1+p(gx_{m_k-1}, gx_{n_k}), p(gy_{m_k-1}, gy_{n_k})+p(gx_{m_k}, gx_{n_k+1})} \end{aligned} \right\} \right). \end{aligned}$$

Letting  $k \rightarrow \infty$ , we have

$$\lim_{k \rightarrow \infty} \psi(p(gx_{m_k}, gx_{n_k+1})) \leq \alpha\left(\frac{\epsilon}{2}\right) - \beta\left(\frac{\epsilon}{2}\right).$$

Similarly, we obtain

$$\lim_{k \rightarrow \infty} \psi(p(gy_{m_k}, gy_{n_k+1})) \leq \alpha\left(\frac{\epsilon}{2}\right) - \beta\left(\frac{\epsilon}{2}\right).$$

Hence from (18), we have

$$\psi\left(\frac{\epsilon}{2}\right) \leq \alpha\left(\frac{\epsilon}{2}\right) - \beta\left(\frac{\epsilon}{2}\right).$$

From (3.1.2) and (3.1.3), we get  $\frac{\epsilon}{2} = 0$ , a contradiction.

Hence  $\{gx_n\}$  and  $\{gy_n\}$  are Cauchy sequences in the metric space  $(X, d_p)$ .

Hence we have  $\lim_{n,m \rightarrow \infty} d_p(gx_n, gx_m) = 0 = \lim_{n,m \rightarrow \infty} d_p(gy_n, gy_m)$ .

Now from the definition of  $d_p$  and from (6), we have

$$\lim_{n \rightarrow \infty} p(gx_n, gx_m) = 0 = \lim_{n \rightarrow \infty} p(gy_n, gy_m). \tag{19}$$

Suppose  $g(X)$  is a complete subspace of  $X$ .

Since  $\{gx_n\}$  and  $\{gy_n\}$  are Cauchy sequences in a complete metric space  $(g(X), d_p)$ . Then  $\{gx_n\}$  and  $\{gy_n\}$  converges to some  $u$  and  $v$  in  $g(X)$  respectively. Thus

$$\lim_{n \rightarrow \infty} d_p(gx_n, u) = 0$$

and

$$\lim_{n \rightarrow \infty} d_p(gy_n, v) = 0$$

for some  $u$  and  $v$  in  $g(X)$ .

Since  $u, v \in g(X)$ , there exist  $x, y \in X$  such that  $u = gx$  and  $v = gy$ .

Since  $\{gx_n\}$  and  $\{gy_n\}$  are Cauchy sequences,  $gx_n \rightarrow u, gy_n \rightarrow v, gx_{n+1} \rightarrow u$  and  $gy_{n+1} \rightarrow v$ .

From Lemma 2.5(2) and (19), we obtain

$$p(u, u) = \lim_{n \rightarrow \infty} p(gx_n, u) = p(v, v) = \lim_{n \rightarrow \infty} p(gy_n, v) = 0. \tag{20}$$

Now we prove that  $\lim_{n \rightarrow \infty} p(F(x, y), gx_n) = p(F(x, y), u)$ .

By definition of  $d_p$ ,

$$d_p(F(x, y), gx_n) = 2p(F(x, y), gx_n) - p(F(x, y), F(x, y)) - p(gx_n, gx_n).$$

Letting  $n \rightarrow \infty$ , we have

$$d_p(F(x, y), u) = 2 \lim_{n \rightarrow \infty} p(F(x, y), gx_n) - p(F(x, y), F(x, y)) - 0, \text{ from (6).}$$

By definition of  $d_p$  and (19), we have

$$\lim_{n \rightarrow \infty} p(F(x, y), gx_n) = p(F(x, y), u).$$

Similarly,  $\lim_{n \rightarrow \infty} p(F(y, x), gy_n) = p(F(y, x), v)$ .

From  $(p_4)$ , we have

$$\begin{aligned} p(u, F(x, y)) &\leq p(u, gx_{n+1}) + p(gx_{n+1}, F(x, y)) - p(gx_{n+1}, gx_{n+1}) \\ &= p(u, gx_{n+1}) + p(gx_{n+1}, F(x, y)). \end{aligned}$$

Letting  $n \rightarrow \infty$ , we have

$$p(u, F(x, y)) \leq 0 + \lim_{n \rightarrow \infty} p(F(x_n, y_n), F(x, y)).$$

Also from (3.2.4), we get  $gx_n \leq gx$  and  $gy_n \geq gy$ . Since  $\psi$  is a continuous and non-decreasing function, we get

$$\begin{aligned} \psi(p(u, F(x, y))) &\leq \lim_{n \rightarrow \infty} \psi(p(F(x_n, y_n), F(x, y))) \\ &\leq \lim_{n \rightarrow \infty} [\alpha(M(x_n, y_n, x, y)) - \beta(M(x_n, y_n, x, y))], \\ M(x_n, y_n, x, y) &= \max \left\{ \begin{array}{l} p(gx_n, u), p(gy_n, v), p(gx_n, gx_{n+1}), \\ p(gy_n, gy_{n+1}), p(u, F(x, y)), p(v, F(y, x)), \\ \frac{p(gx_n, gx_{n+1})p(gy_n, gy_{n+1})}{1+p(gx_n, u)+p(gy_n, v)+p(gx_{n+1}, F(x, y))}, \\ \frac{p(u, F(x, y))p(v, F(y, x))}{1+p(gx_n, u)+p(gy_n, v)+p(gx_{n+1}, F(x, y))} \end{array} \right\} \\ &\rightarrow \max\{p(u, F(x, y)), p(v, F(y, x))\} \text{ as } n \rightarrow \infty. \end{aligned}$$

Therefore

$$\psi(p(u, F(x, y))) \leq \alpha \left( \max \left\{ \begin{array}{l} p(u, F(x, y)), \\ p(v, F(y, x)) \end{array} \right\} \right) - \beta \left( \max \left\{ \begin{array}{l} p(u, F(x, y)), \\ p(v, F(y, x)) \end{array} \right\} \right).$$

Similarly,

$$\psi(p(v, F(y, x))) \leq \alpha \left( \max \left\{ \begin{matrix} p(u, F(x, y)), \\ p(v, F(y, x)) \end{matrix} \right\} \right) - \beta \left( \max \left\{ \begin{matrix} p(u, F(x, y)), \\ p(v, F(y, x)) \end{matrix} \right\} \right).$$

Hence

$$\begin{aligned} & \psi(\max\{p(u, F(x, y)), p(v, F(y, x))\}) \\ &= \max\{\psi(p(u, F(x, y))), \psi(p(v, F(y, x)))\} \\ &\leq \alpha \left( \max \left\{ \begin{matrix} p(u, F(x, y)), \\ p(v, F(y, x)) \end{matrix} \right\} \right) - \beta \left( \max \left\{ \begin{matrix} p(u, F(x, y)), \\ p(v, F(y, x)) \end{matrix} \right\} \right). \end{aligned}$$

It follows that  $\max\{p(u, F(x, y)), p(v, F(y, x))\} = 0$ . So  $F(x, y) = u$  and  $F(y, x) = v$ .

Hence  $F(x, y) = gx = u$  and  $F(y, x) = gy = v$ .

Hence  $F$  and  $g$  have a coincidence point in  $X \times X$ . □

**Theorem 3.3** *In addition to the hypothesis of Theorem 3.2, we suppose that for every  $(x, y), (x^1, y^1) \in X \times X$  there exists  $(u, v) \in X \times X$  such that  $(F(u, v), F(v, u))$  is comparable to  $(F(x, y), F(y, x))$  and  $(F(x^1, y^1), F(y^1, x^1))$ . If  $(x, y)$  and  $(x^1, y^1)$  are coupled coincidence points of  $F$  and  $g$ , then*

$$\begin{aligned} F(x, y) = gx = gx^1 = F(x^1, y^1) \quad \text{and} \\ T(y, x) = gy = gy^1 = F(y^1, x^1). \end{aligned}$$

Moreover, if  $(F, g)$  is  $w$ -compatible, then  $F$  and  $g$  have a unique common coupled fixed point in  $X \times X$ .

*Proof* The proof follows from Theorem 3.2 and the definition of comparability. □

**Theorem 3.4** *Let  $(X, \leq)$  be a partially ordered set and  $p$  be a partial metric such that  $(X, p)$  is a complete PMS. Let  $F : X \times X \rightarrow X$  be such that*

$$(3.4.1) \quad \psi(p(F(x, y), F(u, v))) \leq \alpha(\max\{p(x, u), p(y, v)\}) - \beta(\max\{p(x, u), p(y, v)\}),$$

$\forall x, y, u, v \in X, x \leq u$  and  $y \geq v$ , where  $\psi, \alpha$  and  $\beta$  are defined in Definition 3.1 and

- (3.4.2) (a) *If a non-decreasing sequence  $\{x_n\} \rightarrow x$ , then  $x_n \leq x$  for all  $n$ , and*  
 (b) *if a non-increasing sequence  $\{y_n\} \rightarrow y$ , then  $y \leq y_n$  for all  $n$ .*

*If there exist  $x_0, y_0 \in X$  such that  $x_0 \leq F(x_0, y_0)$  and  $y_0 \geq F(y_0, x_0)$ , then  $F$  has a unique coupled fixed point in  $X \times X$ .*

**Example 3.5** Let  $X = [0, 1]$ , let  $\leq$  be partially ordered on  $X$  by

$$x \leq y \iff x \geq y.$$

The mapping  $F : X \times X \rightarrow X$  defined by  $F(x, y) = \frac{x^2 + y^2}{8(x + y + 1)}$  and  $p : X \times X \rightarrow [0, \infty)$  by  $p(x, y) = \max\{x, y\}$  is a complete partial metric on  $X$ . Define  $\psi, \alpha, \beta : [0, \infty) \rightarrow [0, \infty)$  by  $\psi(t) = t$ ,

$\alpha(t) = \frac{t}{2}$  and  $\beta(t) = \frac{t}{4}$ . We have

$$\begin{aligned}
 p(F(x, y), F(u, v)) &= \max \left\{ \frac{x^2 + y^2}{8(x + y + 1)}, \frac{u^2 + v^2}{8(u + v + 1)} \right\} \\
 &= \frac{1}{4} \left[ \max \left\{ \frac{x^2}{x + y + 1}, \frac{u^2}{u + v + 1} \right\} + \max \left\{ \frac{y^2}{x + y + 1}, \frac{v^2}{u + v + 1} \right\} \right] \\
 &\leq \frac{1}{8} \left[ \max \left\{ \frac{x^2}{x + 1}, \frac{u^2}{u + 1} \right\} + \max \left\{ \frac{y^2}{y + 1}, \frac{v^2}{v + 1} \right\} \right] \\
 &\leq \frac{1}{8} \left[ \max \left\{ \frac{x}{x + 1}, \frac{u}{u + 1} \right\} + \max \left\{ \frac{y}{y + 1}, \frac{v}{v + 1} \right\} \right] \\
 &\leq \frac{1}{8} [\max\{x, u\} + \max\{y, v\}] \\
 &= \frac{1}{8} [p(x, u) + p(y, v)] \\
 &\leq \frac{1}{4} \max\{p(x, u), p(y, v)\} \\
 &= \alpha(\max\{p(x, u), p(y, v)\}) - \beta(\max\{p(x, u), p(y, v)\}).
 \end{aligned}$$

Hence all conditions of Theorem 3.4 hold. From Theorem 3.4,  $(0, 0)$  is a unique coupled fixed point of  $F$  in  $X \times X$ .

### 3.1 Application to integral equations

In this section, we study the existence of a unique solution to an initial value problem, as an application to Theorem 3.4.

Consider the initial value problem

$$\begin{aligned}
 x^1(t) &= f(t, x(t), x(t)), \quad t \in I = [0, 1], \\
 x(0) &= x_0,
 \end{aligned} \tag{21}$$

where  $f : I \times [\frac{x_0}{4}, \infty) \times [\frac{x_0}{4}, \infty) \rightarrow [\frac{x_0}{4}, \infty)$  and  $x_0 \in \mathbb{R}$ .

**Theorem 3.6** Consider the initial value problem (21) with  $f \in C(I \times [\frac{x_0}{4}, \infty) \times [\frac{x_0}{4}, \infty))$  and

$$\int_0^t f(s, x(s), y(s)) \, ds \leq \max \left\{ \frac{1}{4} \int_0^t f(s, x(s), x(s)) \, ds - \frac{9x_0}{16}, \frac{1}{4} \int_0^t f(s, y(s), y(s)) \, ds - \frac{9x_0}{16} \right\}.$$

Then there exists a unique solution in  $C(I, [\frac{x_0}{4}, \infty))$  for the initial value problem (21).

*Proof* The integral equation corresponding to initial value problem (21) is

$$x(t) = x_0 + \int_0^t f(s, x(s), x(s)) \, ds. \tag{22}$$

Let  $X = C(I, [\frac{x_0}{4}, \infty))$  and  $p(x, y) = \max\{x - \frac{x_0}{4}, y - \frac{x_0}{4}\}$  for  $x, y \in X$ . Define  $\psi, \alpha, \beta : [0, \infty) \rightarrow [0, \infty)$  by  $\psi(t) = t, \alpha(t) = \frac{1}{2}t$  and  $\beta(t) = \frac{1}{4}t$ . Define  $F : X \times X \rightarrow X$  by

$$F(x, y)(t) = x_0 + \int_0^t f(s, x(s), y(s)) ds.$$

Now

$$\begin{aligned} & p(F(x, y)(t), F(u, v)(t)) \\ &= \max\left\{F(x, y) - \frac{x_0}{4}, F(u, v) - \frac{x_0}{4}\right\} \\ &= \max\left\{\frac{3x_0}{4} + \int_0^t f(s, x(s), y(s)) ds, \frac{3x_0}{4} + \int_0^t f(s, u(s), v(s)) ds\right\} \\ &\leq \max\left\{\frac{3x_0}{4} + \max\left\{\frac{1}{4} \int_0^t f(s, x(s), x(s)) ds - \frac{9x_0}{16}, \frac{1}{4} \int_0^t f(s, y(s), y(s)) ds - \frac{9x_0}{16}\right\}, \right. \\ &\quad \left.\frac{3x_0}{4} + \max\left\{\frac{1}{4} \int_0^t f(s, u(s), u(s)) ds - \frac{9x_0}{16}, \frac{1}{4} \int_0^t f(s, v(s), v(s)) ds - \frac{9x_0}{16}\right\}\right\} \\ &= \max\left\{\max\left\{\frac{x(t)}{4} - \frac{x_0}{16}, \frac{y(t)}{4} - \frac{x_0}{16}\right\}, \max\left\{\frac{u(t)}{4} - \frac{x_0}{16}, \frac{v(t)}{4} - \frac{x_0}{16}\right\}\right\} \\ &= \frac{1}{4} \max\left\{\max\left\{x(t) - \frac{x_0}{4}, u(t) - \frac{x_0}{4}\right\}, \max\left\{y(t) - \frac{x_0}{4}, v(t) - \frac{x_0}{4}\right\}\right\} \\ &= \frac{1}{4} \max\{p(x, u), p(y, v)\} \\ &= \alpha(\max\{p(x, u), p(y, v)\}) - \beta(\max\{p(x, u), p(y, v)\}). \end{aligned}$$

Thus  $F$  satisfies the condition (3.4.1) of Theorem 3.4. From Theorem 3.4, we conclude that  $F$  has a unique coupled fixed point  $(x, y)$  with  $x = y$ . In particular  $x(t)$  is the unique solution of the integral equation (22). □

### 3.2 Application to homotopy

In this section, we study the existence of a unique solution to homotopy theory.

**Theorem 3.7** *Let  $(X, p)$  be a complete PMS,  $U$  be an open subset of  $X$  and  $\bar{U}$  be a closed subset of  $X$  such that  $U \subseteq \bar{U}$ . Suppose  $H : \bar{U} \times \bar{U} \times [0, 1] \rightarrow X$  is an operator such that the following conditions are satisfied:*

- (i)  $x \neq H(x, y, \lambda)$  and  $y \neq H(y, x, \lambda)$  for each  $x, y \in \partial U$  and  $\lambda \in [0, 1]$  (here  $\partial U$  denotes the boundary of  $U$  in  $X$ ),
- (ii)  $\psi(p(H(x, y, \lambda), H(u, v, \lambda))) \leq \alpha(\max\{p(x, y), p(u, v)\}) - \beta(\max\{p(x, y), p(u, v)\}) \forall x, y \in \bar{U}$  and  $\lambda \in [0, 1]$ , where  $\psi, \alpha : [0, \infty) \rightarrow [0, \infty)$  is continuous and non-decreasing and  $\beta : [0, \infty) \rightarrow [0, \infty)$  is lower semi continuous with  $\psi(t) - \alpha(t) + \beta(t) > 0$  for  $t > 0$ ,
- (iii) there exists  $M \geq 0$  such that

$$p(H(x, y, \lambda), H(x, y, \mu)) \leq M|\lambda - \mu|$$

for every  $x \in \bar{U}$  and  $\lambda, \mu \in [0, 1]$ .

Then  $H(\cdot, 0)$  has a coupled fixed point if and only if  $H(\cdot, 1)$  has a coupled fixed point.

*Proof* Consider the set

$$A = \{ \lambda \in [0, 1] : (x, y) = H(x, y, \lambda) \text{ for some } x, y \in U \}.$$

Since  $H(\cdot, 0)$  has a coupled fixed point in  $U$ , we have  $0 \in A$ , so that  $A$  is a non-empty set.

We will show that  $A$  is both open and closed in  $[0, 1]$  so by the connectedness we have  $A = [0, 1]$ .

As a result,  $H(\cdot, 1)$  has a fixed point in  $U$ . First we show that  $A$  is closed in  $[0, 1]$ .

To see this let  $\{\lambda_n\}_{n=1}^\infty \subseteq A$  with  $\lambda_n \rightarrow \lambda \in [0, 1]$  as  $n \rightarrow \infty$ .

We must show that  $\lambda \in A$ .

Since  $\lambda_n \in A$  for  $n = 1, 2, 3, \dots$ , there exist  $x_n, y_n \in U$  with  $u_n = (x_n, y_n) = H(x_n, y_n, \lambda_n)$ . Consider

$$\begin{aligned} p(x_n, x_{n+1}) &= p(H(x_n, y_n, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_{n+1})) \\ &\leq p(H(x_n, y_n, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_n)) \\ &\quad + p(H(x_{n+1}, y_{n+1}, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_{n+1})) \\ &\quad - p(H(x_{n+1}, y_{n+1}, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_n)) \\ &\leq p(H(x_n, y_n, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_n)) + M|\lambda_n - \lambda_{n+1}|. \end{aligned}$$

Letting  $n \rightarrow \infty$ , we get

$$\lim_{n \rightarrow \infty} p(x_n, x_{n+1}) \leq \lim_{n \rightarrow \infty} p(H(x_n, y_n, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_n)) + 0.$$

Since  $\psi$  is continuous and non-decreasing we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \psi(p(x_n, x_{n+1})) &\leq \lim_{n \rightarrow \infty} \psi(p(H(x_n, y_n, \lambda_n), H(x_{n+1}, y_{n+1}, \lambda_n))) \\ &\leq \lim_{n \rightarrow \infty} [\alpha(\max\{p(x_n, x_{n+1}), p(y_n, y_{n+1})\}) - \beta(\max\{p(x_n, x_{n+1}), p(y_n, y_{n+1})\})]. \end{aligned}$$

Similarly

$$\begin{aligned} \lim_{n \rightarrow \infty} \psi(p(y_n, y_{n+1})) &\leq \lim_{n \rightarrow \infty} [\alpha(\max\{p(x_n, x_{n+1}), p(y_n, y_{n+1})\}) - \beta(\max\{p(x_n, x_{n+1}), p(y_n, y_{n+1})\})]. \end{aligned}$$

It follows that

$$\lim_{n \rightarrow \infty} p(x_n, x_{n+1}) = 0 = \lim_{n \rightarrow \infty} p(y_n, y_{n+1}). \tag{23}$$

From  $(p_2)$ ,

$$\lim_{n \rightarrow \infty} p(x_n, x_n) = 0 = \lim_{n \rightarrow \infty} p(y_n, y_n). \tag{24}$$

By the definition of  $d_p$ , we obtain

$$\lim_{n \rightarrow \infty} d_p(x_n, x_{n+1}) = 0 = \lim_{n \rightarrow \infty} d_p(y_n, y_{n+1}). \tag{25}$$

Now we prove that  $\{x_n\}$  and  $\{y_n\}$  are Cauchy sequences in  $(X, d_p)$ . Contrary to this hypothesis, suppose that  $\{x_n\}$  or  $\{y_n\}$  is not Cauchy.

There exists an  $\epsilon > 0$  and a monotone increasing sequence of natural numbers  $\{m_k\}$  and  $\{n_k\}$  such that  $n_k > m_k$ ,

$$\max\{d_p(x_{m_k}, x_{n_k}), d_p(y_{m_k}, y_{n_k})\} \geq \epsilon \tag{26}$$

and

$$\max\{d_p(x_{m_k}, x_{n_k-1}), d_p(y_{m_k}, y_{n_k-1})\} < \epsilon. \tag{27}$$

From (26) and (27), we obtain

$$\begin{aligned} \epsilon &\leq \max\{d_p(x_{m_k}, x_{n_k}), d_p(y_{m_k}, y_{n_k})\} \\ &\leq \max\{d_p(x_{m_k}, x_{n_k-1}), d_p(y_{m_k}, y_{n_k-1})\} \\ &\quad + \max\{d_p(x_{n_k-1}, x_{n_k}), d_p(y_{n_k-1}, y_{n_k})\} \\ &< \epsilon + \max\{d_p(x_{n_k-1}, x_{n_k}), d_p(y_{n_k-1}, y_{n_k})\}. \end{aligned}$$

Letting  $k \rightarrow \infty$  and then using (25), we get

$$\lim_{k \rightarrow \infty} \max\{d_p(x_{m_k}, x_{n_k}), d_p(y_{m_k}, y_{n_k})\} = \epsilon. \tag{28}$$

Hence from the definition of  $d_p$  and from (24), we get

$$\lim_{k \rightarrow \infty} \max\{p(x_{m_k}, x_{n_k}), p(y_{m_k}, y_{n_k})\} = \frac{\epsilon}{2}. \tag{29}$$

Letting  $k \rightarrow \infty$  and then using (28) and (25) in

$$|d_p(x_{m_k}, x_{n_k+1}) - d_p(x_{m_k}, x_{n_k})| \leq d_p(x_{n_k+1}, x_{n_k}),$$

we get

$$\lim_{k \rightarrow \infty} d_p(x_{n_k+1}, x_{m_k}) = \epsilon. \tag{30}$$

Hence, we have

$$\lim_{k \rightarrow \infty} p(x_{n_k+1}, x_{m_k}) = \frac{\epsilon}{2}. \tag{31}$$

Similarly

$$\lim_{k \rightarrow \infty} p(y_{n_k+1}, y_{m_k}) = \frac{\epsilon}{2}. \tag{32}$$

Consider

$$\begin{aligned} p(x_{m_k}, x_{n_k+1}) &= p(H(x_{m_k}, y_{m_k}, \lambda_{m_k}), H(x_{n_k+1}, y_{n_k+1}, \lambda_{n_k+1})) \\ &\leq p(H(x_{m_k}, y_{m_k}, \lambda_{m_k}), H(x_{m_k}, y_{m_k}, \lambda_{n_k+1})) \\ &\quad + p(H(x_{m_k}, y_{m_k}, \lambda_{n_k+1}), H(x_{n_k+1}, y_{n_k+1}, \lambda_{n_k+1})) \\ &\quad - p(H(x_{m_k}, y_{m_k}, \lambda_{n_k+1}), H(x_{m_k}, y_{m_k}, \lambda_{n_k+1})) \\ &\leq M|\lambda_{m_k} - \lambda_{n_k+1}| + p(H(x_{m_k}, y_{m_k}, \lambda_{n_k+1}), H(x_{n_k+1}, y_{n_k+1}, \lambda_{n_k+1})). \end{aligned}$$

Since  $\{\lambda_n\}$  is Cauchy, letting  $k \rightarrow \infty$  in the above, we get

$$\frac{\epsilon}{2} \leq \lim_{k \rightarrow \infty} p(H(x_{m_k}, y_{m_k}, \lambda_{n_k+1}), H(x_{n_k+1}, y_{n_k+1}, \lambda_{n_k+1})).$$

Since  $\psi$  is continuous and non-decreasing we obtain

$$\begin{aligned} \psi\left(\frac{\epsilon}{2}\right) &\leq \lim_{k \rightarrow \infty} \psi(p(H(x_{m_k}, y_{m_k}, \lambda_{n_k+1}), H(x_{n_k+1}, y_{n_k+1}, \lambda_{n_k+1}))) \\ &\leq \lim_{k \rightarrow \infty} [\alpha(\max\{p(x_{m_k}, x_{n_k+1}), p(y_{m_k}, y_{n_k+1})\}) \\ &\quad - \beta(\max\{p(x_{m_k}, x_{n_k+1}), p(y_{m_k}, y_{n_k+1})\})] \\ &= \alpha\left(\frac{\epsilon}{2}\right) - \beta\left(\frac{\epsilon}{2}\right). \end{aligned}$$

It follows that  $\epsilon \leq 0$ , which is a contradiction.

Hence  $\{x_n\}$  and  $\{y_n\}$  are Cauchy sequences in  $(X, d_p)$  and

$$\lim_{n,m \rightarrow \infty} d_p(x_n, x_m) = 0 = \lim_{n,m \rightarrow \infty} d_p(y_n, y_m).$$

By the definition of  $d_p$  and (24), we get  $\lim_{n,m \rightarrow \infty} p(x_n, x_m) = 0 = \lim_{n,m \rightarrow \infty} p(y_n, y_m)$ .

From Lemma 2.5, we conclude (a)  $\{x_n\}$  and  $\{y_n\}$  are Cauchy sequences in  $(X, p)$ .

Since  $(X, p)$  is complete, from Lemma 2.5(b), we conclude there exist  $u, v \in U$  with

$$p(u, u) = \lim_{n \rightarrow \infty} p(x_n, u) = \lim_{n \rightarrow \infty} p(x_{n+1}, u) = \lim_{n,m \rightarrow \infty} p(x_n, x_m) = 0, \tag{33}$$

$$p(v, v) = \lim_{n \rightarrow \infty} p(x_n, v) = \lim_{n \rightarrow \infty} p(x_{n+1}, v) = \lim_{n,m \rightarrow \infty} p(y_n, y_m) = 0. \tag{34}$$

From Lemma 2.6, we get  $\lim_{n \rightarrow \infty} p(x_n, H(u, v, \lambda)) = p(u, H(u, v, \lambda))$ .

Now,

$$\begin{aligned} p(x_n, H(u, v, \lambda)) &= p(H(x_n, y_n, \lambda_n), H(u, v, \lambda)) \\ &\leq p(H(x_n, y_n, \lambda_n), H(x_n, y_n, \lambda)) + p(H(x_n, y_n, \lambda), H(u, v, \lambda)) \end{aligned}$$



$$\begin{aligned}
 & -p(H(x_n, y_n, \lambda), H(x_n, y_n, \lambda)) \\
 & \leq M|\lambda_n - \lambda| + p(H(x_n, y_n, \lambda), H(u, v, \lambda)).
 \end{aligned}$$

Letting  $n \rightarrow \infty$ , we obtain

$$p(u, H(u, v, \lambda)) \leq \lim_{n \rightarrow \infty} p(H(x_n, y_n, \lambda), H(u, v, \lambda)).$$

Since  $\psi$  is continuous and non-decreasing, we obtain

$$\begin{aligned}
 \psi(p(u, H(u, v, \lambda))) & \leq \lim_{n \rightarrow \infty} \psi(p(H(x_n, y_n, \lambda), H(u, v, \lambda))) \\
 & \leq \lim_{n \rightarrow \infty} [\alpha(\max\{p(x_n, u), p(y_n, v)\}) - \beta(\max\{p(x_n, u), p(y_n, v)\})] \\
 & = 0.
 \end{aligned}$$

It follows that  $p(u, H(u, v, \lambda)) = 0$ . Thus  $u = H(u, v, \lambda)$ . Similarly  $v = H(v, u, \lambda)$ .

Thus  $\lambda \in A$ . Hence  $A$  is closed in  $[0, 1]$ .

Let  $\lambda_0 \in A$ . Then there exist  $x_0, y_0 \in U$  with  $x_0 = H(x_0, y_0, \lambda_0)$ .

Since  $U$  is open, there exists  $r > 0$  such that  $B_p(x_0, r) \subseteq U$ .

Choose  $\lambda \in (\lambda_0 - \epsilon, \lambda_0 + \epsilon)$  such that  $|\lambda - \lambda_0| \leq \frac{1}{M^n} < \epsilon$ .

Then  $x \in B_p(x_0, r) = \{x \in X / p(x, x_0) \leq r + p(x_0, x_0)\}$ . We have

$$\begin{aligned}
 p(H(x, y, \lambda), x_0) & = p(H(x, y, \lambda), H(x_0, y_0, \lambda_0)) \\
 & \leq p(H(x, y, \lambda), H(x, y, \lambda_0)) + p(H(x, y, \lambda_0), H(x_0, y_0, \lambda_0)) \\
 & \quad - p(H(x, y, \lambda_0), H(x, y, \lambda_0)) \\
 & \leq M|\lambda - \lambda_0| + p(H(x, y, \lambda_0), H(x_0, y_0, \lambda_0)) \\
 & \leq \frac{1}{M^{n-1}} + p(H(x, y, \lambda_0), H(x_0, y_0, \lambda_0)).
 \end{aligned}$$

Letting  $n \rightarrow \infty$ , we obtain

$$p(H(x, y, \lambda), x_0) \leq p(H(x, y, \lambda_0), H(x_0, y_0, \lambda_0)).$$

Since  $\psi$  is continuous and non-decreasing, we have

$$\begin{aligned}
 \psi(p(H(x, y, \lambda), x_0)) & \leq \psi(p(H(x, y, \lambda_0), H(x_0, y_0, \lambda_0))) \\
 & \leq \alpha(\max\{p(x, x_0), p(y, y_0)\}) \\
 & \quad - \phi(\max\{p(x, x_0), p(y, y_0)\}).
 \end{aligned}$$

Similarly

$$\begin{aligned}
 & \psi(p(H(y, x, \lambda), y_0)) \\
 & \leq \alpha(\max\{p(x, x_0), p(y, y_0)\}) - \phi(\max\{p(x, x_0), p(y, y_0)\}).
 \end{aligned}$$

Thus

$$\begin{aligned} &\psi(\max\{p(H(x, y, \lambda), x_0), p(H(y, x, \lambda), y_0)\}) \\ &\leq \alpha(\max\{p(x, x_0), p(y, y_0)\}) - \phi(\max\{p(x, x_0), p(y, y_0)\}) \\ &\leq \psi(\max\{p(x, x_0), p(y, y_0)\}). \end{aligned}$$

Since  $\psi$  is non-decreasing, we have

$$\begin{aligned} \max\{p(H(x, y, \lambda), x_0), p(H(y, x, \lambda), y_0)\} &\leq \max\{p(x, x_0), p(y, y_0)\} \\ &\leq \max\{r + p(x_0, x_0), r + p(y_0, y_0)\}. \end{aligned}$$

Thus for each fixed  $\lambda \in (\lambda_0 - \epsilon, \lambda_0 + \epsilon)$ ,  $H(\cdot, \lambda) : \overline{B_p(x_0, r)} \rightarrow \overline{B_p(x_0, r)}$ .

Since also (ii) holds and  $\psi$  and  $\alpha$  are continuous and non-decreasing and  $\beta$  is continuous with  $\psi(t) - \alpha(t) + \beta(t) > 0$  for  $t > 0$ , all conditions of Theorem 3.4 are satisfied.

Thus we deduce that  $H(\cdot, \lambda)$  has a coupled fixed point in  $\overline{U}$ . But this coupled fixed point must be in  $U$  since (i) holds.

Thus  $\lambda \in A$  for any  $\lambda \in (\lambda_0 - \epsilon, \lambda_0 + \epsilon)$ .

Hence  $(\lambda_0 - \epsilon, \lambda_0 + \epsilon) \subseteq A$  and therefore  $A$  is open in  $[0, 1]$ .

For the reverse implication, we use the same strategy. □

**Corollary 3.8** *Let  $(X, p)$  be a complete PMS,  $U$  be an open subset of  $X$  and  $H : \overline{U} \times \overline{U} \times [0, 1] \rightarrow X$  with the following properties:*

- (1)  $x \neq H(x, y, t)$  and  $y \neq H(y, x, t)$  for each  $x, y \in \partial U$  and each  $\lambda \in [0, 1]$  (here  $\partial U$  denotes the boundary of  $U$  in  $X$ ),
- (2) there exist  $x, y \in \overline{U}$  and  $\lambda \in [0, 1], L \in [0, 1]$ , such that

$$p(H(x, y, \lambda), H(u, v, \mu)) \leq L \max\{p(x, u), p(y, v)\},$$

- (3) there exists  $M \geq 0$ , such that

$$p(H(x, \lambda), H(x, \mu)) \leq M \cdot |\lambda - \mu|$$

for all  $x \in \overline{U}$  and  $\lambda, \mu \in [0, 1]$ .

If  $H(\cdot, 0)$  has a fixed point in  $U$ , then  $H(\cdot, 1)$  has a fixed point in  $U$ .

*Proof* The proof follows by taking  $\psi(x) = x, \phi(x) = x - Lx$  with  $L \in [0, 1]$  in Theorem 3.7. □

#### 4 Conclusions

In this paper we conclude some applications on homotopy theory and integral equations by using coupled fixed point theorems in ordered PMSs.

##### Competing interests

The authors declare that they have no competing interests.

##### Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

**Author details**

<sup>1</sup>Department of Mathematics, Acharya Nagarjuna University, Nagarjuna Nagar, Guntur, Andhra Pradesh 522 510, India. <sup>2</sup>Department of Mathematics, K L University, Vaddeswaram, Guntur, Andhra Pradesh 522 502, India. <sup>3</sup>Department of Mathematics and Computer Science, Cankaya University, Ankara, Turkey. <sup>4</sup>Department of Computing, Adama Science and Technology University, Adama, Ethiopia. <sup>5</sup>Department of Mathematics, Nallamallareddy Engineering College, Divya Nagar, Hyderabad, Telangana 500088, India.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 31 March 2017 Accepted: 24 August 2017 Published online: 01 December 2017

**References**

1. Matthews, SG: Partial metric topology. Research Report 212, Department of Computer Science, University of Warwick (1992)
2. Abodayeh, K, Mlaiki, N, Abdeljawad, T, Shatanawi, W: Relations between partial metric spaces and M-metric spaces, Caristi Kirk's theorem in M-metric type spaces. *J. Math. Anal.* **7**(3), 1-12 (2016)
3. Heckmann, R: Approximation of metric spaces by partial metric spaces. *Appl. Categ. Struct.* **7**(1-2), 71-83 (1999)
4. Kopperman, R, Matthews, SG, Pajoohesh, H: What do partial metrics represent? In: *Spatial Representation: Discrete vs. Continuous Computational Models*. Dagstuhl Seminar Proceedings, No. 04351, Internationales Begegnungs- und Forschungszentrum für Informatik (IBFI), Schloss Dagstuhl, Germany (2005)
5. Künzi, HPA, Pajoohesh, H, Schellekens, MP: Partial quasi-metrics. *Theor. Comput. Sci.* **365**(3), 237-246 (2006)
6. O'Neill, SJ: Two topologies are better than one. Technical report, University of Warwick, Coventry, UK (1995). <http://www.dcs.warwick.ac.uk/reports/283.html>
7. Schellekens, M: The Smyth completion: a common foundation for denotational semantics and complexity analysis. *Electron. Notes Theor. Comput. Sci.* **1**, 535-556 (1995). doi:10.1016/S1571-0661(04)00029-5
8. Schellekens, MP: A characterization of partial metrizable spaces: domains are quantifiable. *Theor. Comput. Sci.* **305**(1-3), 409-432 (2003). doi:10.1016/S0304-3975(02)00705-3
9. Waszkiewicz, P: Partial metrizable continuous posets. *Math. Struct. Comput. Sci.* **16**(2), 359-372 (2006)
10. Matthews, SG: Partial metric topology. In: *Proceedings of the 8th Summer Conference on General Topology and Applications*. Annals of the New York Academy of Sciences, vol. 728, pp. 183-197 (1994)
11. Oltra, S, Valero, O: Banach's fixed point theorem for partial metric spaces. *Rend. Ist. Mat. Univ. Trieste* **36**(1-2), 17-26 (2004)
12. Altun, I, Sola, F, Simsek, H: Generalized contractions on partial metric spaces. *Topol. Appl.* **157**(18), 2778-2785 (2010)
13. Abbas, M, Aydi, H, Radenović, S: Fixed point of T-Hardy-Rogers contractive mappings in ordered partial metric spaces. *Int. J. Math. Math. Sci.* **2012**, Article ID 313675 (2012)
14. Abdeljawad, T, Karapinar, E, Tas, K: Existence and uniqueness of a common fixed point on partial metric spaces. *Appl. Math. Lett.* **24**(11), 1894-1899 (2011)
15. Altun, I, Erduran, A: Fixed point theorems for monotone mappings on partial metric spaces. *Fixed Point Theory Appl.* **2011**, Article ID 508730 (2011). doi:10.1155/2011/508730
16. Aydi, H: Some coupled fixed point results on partial metric spaces. *Int. J. Math. Math. Sci.* **2011**, Article ID 647091 (2011). doi:10.1155/2011/647091
17. Aydi, H: Fixed point results for weakly contractive mappings in ordered partial metric spaces. *J. Adv. Math. Stud.* **4**(2), 1-12 (2011)
18. Čojbašić, V, Radenović, S, Chauhan, S: Common fixed point of generalized weakly contractive maps in 0-complete partial metric spaces. *Acta Math. Sci.* **34B**(4), 1345-1356 (2014)
19. Ilić, D, Pavlović, V, Rakočević, V: Some new extensions of Banach's contraction principle to partial metric spaces. *Appl. Math. Lett.* **24**(8), 1326-1330 (2011). doi:10.1016/j.aml.2011.02.025
20. Karapinar, E, Erhan, IM: Fixed point theorems for operators on partial metric spaces. *Appl. Math. Lett.* **24**(11), 1900-1904 (2011)
21. Karapinar, E: Weak  $\phi$ -contraction on partial contraction and existence of fixed points in partially ordered sets. *Math. Aeterna* **1**(4), 237-244 (2011)
22. Karapinar, E: Generalizations of Caristi Kirk's theorem on partial metric spaces. *Fixed Point Theory Appl.* **2011**, 4 (2011)
23. Rao, KPR, Kishore, GNV: A unique common fixed point theorem for four maps under  $\psi$ - $\phi$  contractive condition in partial metric spaces. *Bull. Math. Anal. Appl.* **3**(3), 56-63 (2011)
24. Radenović, S: Classical fixed point results in 0-complete partial metric spaces via cyclic-type extension. *Bull. Allahabad Math. Soc.* **31**(Part 1), 39-55 (2016)
25. Shukla, S, Radenović, S: Some common fixed point theorems for  $F$ -contraction type mappings in 0-complete partial metric spaces. *J. Math.* **2013**, Article ID 878730 (2013)
26. Valero, O: On Banach fixed point theorems for partial metric spaces. *Appl. Gen. Topol.* **6**(2), 229-240 (2005)
27. Vetro, F, Radenović, S: Nonlinear quasi-contractions of Ćirić type in partial metric spaces. *Appl. Math. Comput.* **219**, 1594-1600 (2012)
28. Bhaskar, TG, Lakshmikantham, V: Fixed point theorems in partially ordered metric spaces and applications. *Nonlinear Anal.* **65**, 1379-1393 (2006)
29. Abdeljawad, T: Coupled fixed point theorems for partially contractive type mappings. *Fixed Point Theory Appl.* **2012**, 148 (2012)
30. Abbas, M, Alikhan, M, Radenović, S: Common coupled fixed point theorems in cone metric spaces for  $w$ -compatible mappings. *Appl. Math. Comput.* **217**(1), 195-202 (2010)
31. Lakshmikantham, V, Ćirić, L: Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces. *Nonlinear Anal.* **70**(12), 4341-4349 (2009). doi:10.1016/j.na.2008.09.020

32. Rao, KPR, Kishore, GNV, Van Luong, N: A unique common coupled fixed point theorem for four maps under  $\psi$ - $\phi$  contractive condition in partial metric spaces. *CUBO* **14**(03), 115-127 (2012)
33. Rao, KPR, Kishore, GNV, Raju, VCC: A coupled fixed point theorem for two pairs of  $w$ -compatible maps using altering distance function in partial metric space. *J. Adv. Res. Pure Math.* **4**(4), 96-114 (2012)
34. Radenović, S: Some coupled coincidence points results of monotone mappings in partially ordered metric spaces. *Int. J. Anal. Appl.* **5**(2), 174-184 (2014)
35. Shatanawi, W, Samet, B, Abbas, M: Coupled fixed point theorems for mixed monotone mappings in ordered partial metric spaces. *Math. Comput. Model.* **55**(3-4), 680-687 (2012)

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](http://springeropen.com)

---