

APPROXIMATING COMMON FIXED POINTS OF FINITE FAMILY OF ASYMPTOTICALLY NONEXPANSIVE NON-SELF MAPPINGS

G. S. Saluja*

Abstract

Let K be a nonempty closed convex nonexpansive retract of a real uniformly convex Banach space E with P as a nonexpansive retraction. Let $T_1, T_2, \dots, T_N: K \rightarrow E$ be N asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{\alpha_n^i\}$, $\{\beta_n^i\}$ and $\{\gamma_n^i\}$ are sequences in $[0, 1]$ with $\alpha_n^i + \beta_n^i + \gamma_n^i = 1$ for all $i = 1, 2, \dots, N$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ iteratively by (6), where $\{u_n^i\}$ are bounded sequences in K with $\sum_{n=1}^{\infty} u_n^i < \infty$. (i) If the dual E^* of E has the *Kadec-Klee* property, then $\{x_n\}$ converges weakly to a common fixed point $x^* \in F$; (ii) if $\{T_1, T_2, \dots, T_N\}$ satisfies condition (B), then $\{x_n\}$ converges strongly to a common fixed point $x^* \in F$.

1 Introduction and preliminaries

Let K be a nonempty closed convex subset of a Banach space E . A self mapping $T: K \rightarrow K$ is called asymptotically nonexpansive if there exists a sequence $\{u_n\} \subset [0, \infty)$; $u_n \rightarrow 0$ as $n \rightarrow \infty$ such that for all $x, y \in K$, the following inequality holds:

$$\|T^n x - T^n y\| \leq (1 + u_n) \|x - y\|, \quad \forall n \geq 1. \quad (1)$$

T is called uniformly L -Lipschitzian if there exists a constant $L > 0$ such that for

2000 *Mathematics Subject Classifications.* 47H10.

Key words and Phrases. Asymptotically nonexpansive nonself mapping, common fixed point, multi step iterative sequences with errors for nonself mappings, real uniformly convex Banach space.

Received: December 17, 2007

Communicated by Vladimir Rakočević

all $x, y \in K$,

$$\|T^n x - T^n y\| \leq L \|x - y\|, \quad \forall n \geq 1. \quad (2)$$

The class of asymptotically nonexpansive maps was introduced by Goebel and Kirk [12] as an important generalization of the class of nonexpansive maps (i.e., mappings $T: K \rightarrow K$ such that $\|Tx - Ty\| \leq \|x - y\|, \forall x, y \in K$) who proved that if K is a nonempty closed convex subset of a real uniformly convex Banach space and T is an asymptotically nonexpansive self-mapping of K , then T has a fixed point.

Iterative techniques for approximating fixed points of nonexpansive mappings and asymptotically nonexpansive mappings have been studied by various authors (see e.g., [24], [2, 3], [5], [21], [17], [6], [23], [1], [10, 11], [7], [13, 14, 15, 16]) using the Mann iteration method (see e.g., [27]) or the Ishikawa iteration method (see e.g., [20]).

In 1978, Bose [19] proved that if K is a bounded closed convex nonempty subset of a uniformly convex Banach space E satisfying Opial's [29] condition and $T: K \rightarrow K$ is an asymptotically nonexpansive mapping, then the sequence $\{T^n x\}$ converges weakly to a fixed point of T provided T is asymptotically regular at $x \in K$, i.e., $\lim_{n \rightarrow \infty} \|T^n x - T^{n+1} x\| = 0$. Passty [6] and also Xu [8] proved that the requirement that E satisfies Opial's condition can be replaced by the condition that E has a Frechet differentiable norm. Furthermore, Tan and Xu [13, 14] later proved that the asymptotic regularity of T can be weakened to the weakly asymptotic regularity of T at x , i.e., $\omega - \lim_{n \rightarrow \infty} (T^n x - T^{n+1} x) = 0$.

In [10, 11], Schu introduced a modified Mann process to approximate fixed points of asymptotically nonexpansive self-maps defined on nonempty closed convex and bounded subsets of a Hilbert space H .

In 1994, Rhoades [1] extended the Schu's result to uniformly convex Banach space using a modified Ishikawa iteration method.

In all the above results, the operator T remains a self-mapping of a nonempty closed convex subset K of a uniformly convex Banach space. If, however, the domain of T , $D(T)$ is a proper subset of E (and this is the case in several applications), and T maps $D(T)$ into E , then the iteration processes of Mann and Ishikawa studied by these authors; and their modifications introduced by Schu may fail to be well defined.

In 2003, Chidume et al [4] studied the iterative scheme defined by

$$x_1 \in K, \quad x_{n+1} = P((1 - \alpha_n)x_n + \alpha_n T(PT)^{n-1}x_n), \quad n \geq 1, \quad (3)$$

in the framework of uniformly convex Banach space, where K is a closed convex nonexpansive retract of a real uniformly convex Banach space E with P as a nonexpansive retract. $T: K \rightarrow E$ is an asymptotically nonexpansive nonself map with sequence $\{k_n\} \subset [1, \infty)$, $k_n \rightarrow 1$ as $n \rightarrow \infty$. $\{\alpha_n\}_{n=1}^{\infty}$ is a real sequence in $[0, 1]$ satisfying the condition $\epsilon \leq \alpha_n \leq 1 - \epsilon$ for all $n \geq 1$ and for some $\epsilon > 0$. They proved strong and weak convergence theorems for asymptotically nonexpansive nonself maps.

In 2005, Shahzad [18] studied the sequence $\{x_n\}$ defined by

$$x_1 \in K, \quad x_{n+1} = P((1 - \alpha_n)x_n + \alpha_n TP[(1 - \beta_n)x_n + \beta_n Tx_n]), \quad (4)$$

where K is a nonempty closed convex nonexpansive retract of a real uniformly convex Banach space E with P as a nonexpansive retraction. He proved weak and strong convergence theorems for nonself nonexpansive mappings in Banach spaces. Recently, Su and Qin [28] studied the sequence $\{x_n\}$ defined by

$$x_1 \in K,$$

$$\begin{aligned} z_n &= P(\alpha_n'' T(PT)^{n-1} x_n + (1 - \alpha_n'') x_n), \\ y_n &= P(\alpha_n' T(PT)^{n-1} z_n + (1 - \alpha_n') x_n), \\ x_{n+1} &= P(\alpha_n T(PT)^{n-1} y_n + (1 - \alpha_n) x_n), \end{aligned} \quad (5)$$

where $\{\alpha_n\}$, $\{\alpha_n'\}$ and $\{\alpha_n''\}$ are real sequences in $(0, 1)$ and K is a nonempty closed convex nonexpansive retract of a uniformly convex Banach space E with P as a nonexpansive retraction. They proved weak and strong convergence theorems for asymptotically nonexpansive nonself mappings in uniformly convex Banach space. Motivated by Su and Qin [28] and some others, the purpose of this paper is to construct a multi step iterative scheme with errors for approximating common fixed point of a finite family of asymptotically nonexpansive nonself mappings (when such a fixed point exists) and to prove weak and strong convergence theorems for such maps.

Let K be a nonempty closed convex subset of a real uniformly convex Banach space E . In this paper, the following iteration scheme is studied:

$$\begin{aligned} x_n^1 &= P(\alpha_n^1 T_1(PT_1)^{n-1} x_n + \beta_n^1 x_n + \gamma_n^1 u_n^1) \\ x_n^2 &= P(\alpha_n^2 T_2(PT_2)^{n-1} x_n^1 + \beta_n^2 x_n^1 + \gamma_n^2 u_n^2) \end{aligned}$$

on K if there exists a nondecreasing function $f: [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f(r) > 0$ for all $r \in (0, \infty)$ such that for all $x \in K$, $\|x - Tx\| \geq f(d(x, F))$ where $d(x, F(T)) = \inf\{\|x - p\| : p \in F(T)\}$.

(ii) A family $\{T_1, T_2, \dots, T_N\}$ of N self mappings on K with $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$ is said to satisfy condition (B) on K if there exist f and d as in (i) such that $\max_{1 \leq i \leq N} \{\|x - T_i x\|\} \geq f(d(x, F))$ for all $x \in K$.

Note that condition (B) reduces to condition (A), when $T_i = T$ for all $i = 1, 2, \dots, N$.

In order to prove our main results, we will make use of the following lemmas:

Lemma 1.1 (Tan and Xu [15]): Let $\{a_n\}_{n=1}^\infty$, $\{\beta_n\}_{n=1}^\infty$ and $\{r_n\}_{n=1}^\infty$ be sequences of nonnegative real numbers satisfying

$$a_{n+1} \leq (1 + r_n)a_n + \beta_n, \quad \forall n \in \mathbb{N}.$$

If $\sum_{n=1}^\infty r_n < \infty$, $\sum_{n=1}^\infty \beta_n < \infty$. Then

(i) $\lim_{n \rightarrow \infty} a_n$ exists.

(ii) If $\liminf_{n \rightarrow \infty} a_n = 0$, then $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 1.2 (see [8]): Let $p > 1$ and $R > 1$ be two fixed numbers and E a Banach space. Then E is uniformly convex if and only if there exists a continuous, strictly increasing and convex function $g: [0, \infty) \rightarrow [0, \infty)$ with $g(0) = 0$ such that $\|\lambda x + (1 - \lambda)y\|^p \leq \lambda \|x\|^p + (1 - \lambda) \|y\|^p - W_p(\lambda)g(\|x - y\|)$ for all $x, y \in B_R(0) = \{x \in E : \|x\| \leq R\}$, and $\lambda \in [0, 1]$, where $W_p(\lambda) = \lambda(1 - \lambda)^p + \lambda^p(1 - \lambda)$.

Lemma 1.3 (demiclosed principle for nonselfmap [4]): Let E be a uniformly convex Banach space, K a nonempty closed convex subset of E . Let $T: K \rightarrow E$ be an asymptotically nonexpansive mapping with $\{k_n\} \subset [1, \infty)$ and $k_n \rightarrow 1$ as $n \rightarrow \infty$. Then $I - T$ is demiclosed with respect to zero.

Lemma 1.4 (see [9]): Let E be a real reflexive Banach space such that its dual E^* has the *Kadec-Klee* property. Let $\{x_n\}$ be a bounded sequence in E and $x^*, y^* \in w_w(x_n)$; here $w_w(x_n)$ denotes the weak w -limit set of $\{x_n\}$. Suppose $\lim_{n \rightarrow \infty} \|tx_n + (1 - t)x^* - y^*\|$ exists for all $t \in [0, 1]$. Then $x^* = y^*$.

2 Main results

Definition 2.1 (see [4]): Let E be a real normed linear space, K a nonempty subset of E . Let $P: E \rightarrow K$ be the nonexpansive retraction of E onto K . A map

$T: K \rightarrow E$ is said to be asymptotically nonexpansive if there exists a sequence $\{u_n\} \subset [0, \infty)$; $u_n \rightarrow 0$ as $n \rightarrow \infty$ such that for all $x, y \in K$, the following inequality holds:

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq (1 + u_n) \|x - y\|, \quad \forall n \geq 1. \quad (7)$$

T is called uniformly L -Lipschitzian if there exists a constant $L > 0$ such that for all $x, y \in K$

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq L \|x - y\|, \quad \forall n \geq 1. \quad (8)$$

Lemma 2.2: Let E be a real uniformly convex Banach space and K be a nonempty closed convex subset which is also a nonexpansive retract of E . Let $T_1, T_2, \dots, T_N: K \rightarrow K$ be N asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{\alpha_n^i\}$, $\{\beta_n^i\}$ and $\{\gamma_n^i\}$ are sequences in $[0, 1]$ with $\alpha_n^i + \beta_n^i + \gamma_n^i = 1$ for all $i = 1, 2, \dots, N$. From arbitrary $x_1 \in K$, define the sequence $\{x_n\}$ iteratively by (6), where $\{u_n^i\}$ are bounded sequences in K with $\sum_{n=1}^{\infty} u_n^i < \infty$. Then

(i) $\|x_{n+1} - x^*\| = \|x_n^N - x^*\| \leq (1 + b_n^{N-1}) \|x_n - x^*\| + d_n^{N-1}$, for all $n \geq 1$, $x^* \in F$ and for some sequences $\{b_n^i\}$ and $\{d_n^i\}$ for all $i = 1, 2, \dots, N$ of numbers such that $\sum_{n=1}^{\infty} b_n^i < \infty$ and $\sum_{n=1}^{\infty} d_n^i < \infty$.

(ii) There exists a constant $M > 0$ such that $\|x_{n+m} - x^*\| \leq M \|x_n - x^*\|$ for all $n, m \geq 1$ and $x^* \in F$.

Proof: (i) Let $x^* \in F$, then from (6) we have

$$\begin{aligned} \|x_n^1 - x^*\| &= \|P(\alpha_n^1 T_1 (PT_1)^{n-1} x_n + \beta_n^1 x_n + \gamma_n^1 u_n^1) - Px^*\| \\ &\leq \alpha_n^1 \|T_1 (PT_1)^{n-1} x_n - x^*\| + \beta_n^1 \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\| \\ &\leq \alpha_n^1 (1 + r_n^1) \|x_n - x^*\| + \beta_n^1 \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\| \\ &\leq (1 - \beta_n^1) (1 + r_n^1) \|x_n - x^*\| + \beta_n^1 (1 + r_n^1) \|x_n - x^*\| \\ &\quad + \gamma_n^1 \|u_n^1 - x^*\| \\ &\leq (1 + r_n^1) \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\| \\ &\leq (1 + r_n^1) \|x_n - x^*\| + d_n^0 \end{aligned}$$

where $d_n^0 = \gamma_n^1 \|u_n^1 - x^*\|$. Since $\sum_{n=1}^{\infty} \gamma_n^1 < \infty$, then $\sum_{n=1}^{\infty} d_n^0 < \infty$. Next, we note that

$$\begin{aligned} \|x_n^2 - x^*\| &= \|P(\alpha_n^2 T_2 (PT_2)^{n-1} x_n^1 + \beta_n^2 x_n + \gamma_n^2 u_n^2) - Px^*\| \\ &\leq \alpha_n^2 \|T_2 (PT_2)^{n-1} x_n^1 - x^*\| + \beta_n^2 \|x_n - x^*\| + \gamma_n^2 \|u_n^2 - x^*\| \\ &\leq \alpha_n^2 (1 + r_n^2) \|x_n^1 - x^*\| + \beta_n^2 \|x_n - x^*\| + \gamma_n^2 \|u_n^2 - x^*\| \end{aligned}$$

$$\begin{aligned}
&\leq \alpha_n^2(1+r_n^2)[(1+r_n^1)\|x_n-x^*\|+d_n^0]+\beta_n^2\|x_n-x^*\| \\
&\quad +\gamma_n^2\|u_n^2-x^*\| \\
&\leq [(1+r_n^1)(1+r_n^2)\alpha_n^2+\beta_n^2]\|x_n-x^*\|+\alpha_n^2(1+r_n^2)d_n^0 \\
&\quad +\gamma_n^2\|u_n^2-x^*\| \\
&\leq (\alpha_n^2+\beta_n^2)(1+r_n^1)(1+r_n^2)\|x_n-x^*\|+\alpha_n^2(1+r_n^2)d_n^0 \\
&\quad +\gamma_n^2\|u_n^2-x^*\| \\
&\leq (1+r_n^1+r_n^2+r_n^1r_n^2)\|x_n-x^*\|+\alpha_n^2(1+r_n^2)d_n^0+\gamma_n^2\|u_n^2-x^*\| \\
&\leq (1+b_n^1)\|x_n-x^*\|+d_n^1
\end{aligned}$$

where $d_n^1 = \alpha_n^2(1+r_n^2)d_n^0 + \gamma_n^2\|u_n^2-x^*\|$ and $b_n^1 = (1+r_n^1+r_n^2+r_n^1r_n^2)$. Since $\sum_{n=1}^{\infty} d_n^0 < \infty$, $\sum_{n=1}^{\infty} \gamma_n^2 < \infty$, $\sum_{n=1}^{\infty} r_n^i < \infty$ for $i = 1, 2$, and so $\sum_{n=1}^{\infty} d_n^1 < \infty$ and $\sum_{n=1}^{\infty} b_n^1 < \infty$. Similarly, we have

$$\begin{aligned}
&\|x_n^3-x^*\| = \|P(\alpha_n^3T_3(PT_3)^{n-1}x_n^2+\beta_n^3x_n+\gamma_n^3u_n^3)-Px^*\| \\
&\leq \alpha_n^3\|T_3(PT_3)^{n-1}x_n^2-x^*\|+\beta_n^3\|x_n-x^*\|+\gamma_n^3\|u_n^3-x^*\| \\
&\leq \alpha_n^3(1+r_n^3)\|x_n^2-x^*\|+\beta_n^3\|x_n-x^*\|+\gamma_n^3\|u_n^3-x^*\| \\
&\leq \alpha_n^3(1+r_n^3)[(1+b_n^1)\|x_n-x^*\|+d_n^1]+\beta_n^3\|x_n-x^*\| \\
&\quad +\gamma_n^3\|u_n^3-x^*\| \\
&\leq [\alpha_n^3(1+r_n^3)(1+b_n^1)+\beta_n^3]\|x_n-x^*\|+\alpha_n^3(1+r_n^3)d_n^1 \\
&\quad +\gamma_n^3\|u_n^3-x^*\| \\
&\leq (\alpha_n^3+\beta_n^3)(1+b_n^1)(1+r_n^3)\|x_n-x^*\|+\alpha_n^3(1+r_n^3)d_n^1 \\
&\quad +\gamma_n^3\|u_n^3-x^*\| \\
&\leq (1+b_n^1)(1+r_n^3)\|x_n-x^*\|+d_n^2 \\
&\leq (1+b_n^2)\|x_n-x^*\|+d_n^2
\end{aligned}$$

where $b_n^2 = b_n^1+r_n^3+b_n^1r_n^3$ and $d_n^2 = \alpha_n^3(1+r_n^3)d_n^1+\gamma_n^3\|u_n^3-x^*\|$. Since $\sum_{n=1}^{\infty} b_n^1 < \infty$, $\sum_{n=1}^{\infty} r_n^3 < \infty$, $\sum_{n=1}^{\infty} d_n^1 < \infty$ and $\sum_{n=1}^{\infty} \gamma_n^3 < \infty$, so $\sum_{n=1}^{\infty} b_n^2 < \infty$ and $\sum_{n=1}^{\infty} d_n^2 < \infty$.

By continuing the above process, there exists a nondecreasing sequences $\{d_n^{l-1}\}$ and $\{b_n^{l-1}\}$ such that $\sum_{n=1}^{\infty} d_n^{l-1} < \infty$ and $\sum_{n=1}^{\infty} b_n^{l-1} < \infty$ and

$$\|x_n^i-x^*\| \leq (1+b_n^{i-1})\|x_n-x^*\|+d_n^{i-1}, \quad \forall n \geq 1, \quad \forall i = 1, 2, \dots, N.$$

Thus

$$\|x_{n+1}-x^*\| = \|x_n^N-x^*\| \leq (1+b_n^{N-1})\|x_n-x^*\|+d_n^{N-1}, \quad \forall n \in N.$$

This completes the proof of (i).

(ii) Since $1 + x \leq e^x$ for all $x > 0$. Then from (i) it can be obtained that

$$\begin{aligned}
& \|x_{n+m} - x^*\| \leq (1 + b_{n+m-1}^{N-1}) \|x_{n+m-1} - x^*\| + d_{n+m-1}^{N-1} \\
& \leq e^{b_{n+m-1}^{N-1}} \|x_{n+m-1} - x^*\| + d_{n+m-1}^{N-1} \\
& = e^{(b_{n+m-1}^{N-1} + b_{n+m-2}^{N-1})} \|x_{n+m-2} - x^*\| + e^{b_{n+m-1}^{N-1}} d_{n+m-2}^{N-1} + d_{n+m-1}^{N-1} \\
& = e^{(b_{n+m-1}^{N-1} + b_{n+m-2}^{N-1})} \|x_{n+m-2} - x^*\| + e^{b_{n+m-1}^{N-1}} (d_{n+m-1}^{N-1} + d_{n+m-2}^{N-1}) \\
& = \dots\dots\dots \\
& = \dots\dots\dots \\
& = e^{\sum_{k=n}^{n+m-1} b_k^{N-1}} \|x_n - x^*\| + e^{\sum_{k=n}^{n+m-1} b_k^{N-1}} \cdot \sum_{k=n}^{n+m-1} d_k^{N-1} \\
& = M \cdot \|x_n - x^*\| + M \cdot \sum_{k=n}^{n+m-1} d_k^{N-1}, \text{ where } M = e^{\sum_{k=n}^{\infty} b_k^{N-1}}
\end{aligned}$$

This completes the proof of (ii).

Lemma 2.3: Let E be a normed linear space and K be a nonempty closed and convex subset which is also a nonexpansive retract of E . Let $T_1, T_2, \dots, T_N: K \rightarrow K$ be N uniformly L -Lipschitzian mappings. Let $\{x_n\}$ be the sequence defined by (6) with sequences $\{u_n^i\}$ in K for all $i = 1, 2, \dots, N$ and $\{\alpha_n^i\}, \{\beta_n^i\}$ and $\{\gamma_n^i\}$ are sequences in $[0, 1]$ satisfying $\alpha_n^i + \beta_n^i + \gamma_n^i = 1$ for all $i = 1, 2, \dots, N$. Set $c_n^i = \|x_n - T_i(PT_i)^{n-1}x_n\|$ for all $i = 1, 2, \dots, N$. If $\lim_{n \rightarrow \infty} \|x_n - T_i(PT_i)^{n-1}x_n\| = 0$, $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$, then $\lim_{n \rightarrow \infty} \|x_n - T_i x_n\| = 0$.

Proof: Since T_i is uniformly L -Lipschitzian for all $i = 1, 2, \dots, N$, we have

$$\begin{aligned}
\|x_{n+1} - T_i x_{n+1}\| & \leq \|x_{n+1} - T_i(PT_i)^n x_{n+1}\| + \|T_i(PT_i)^n x_{n+1} - T_i x_{n+1}\| \\
& \leq c_{n+1}^i + L \|T_i(PT_i)^{n-1} x_{n+1} - x_{n+1}\| \\
& \leq c_{n+1}^i + L \{ \|x_{n+1} - x_n\| + \|x_n - T_i(PT_i)^{n-1} x_n\| \\
& \quad + \|T_i(PT_i)^{n-1} x_n - T_i(PT_i)^{n-1} x_{n+1}\| \} \\
& \leq c_{n+1}^i + L \{ \|x_{n+1} - x_n\| + c_n^i + L \|x_{n+1} - x_n\| \} \\
& \leq c_{n+1}^i + L(L+1) \|x_{n+1} - x_n\| + Lc_n^i \rightarrow 0 \text{ as } n \rightarrow \infty
\end{aligned}$$

This completes the proof.

Remark 2.4: If we put $P = I$ (identity map) in Lemma 2.2, then it generalizes the corresponding Lemma of Schu [11] for one mapping. Further, if $F = \bigcap_{i=1}^N F(T_i) \neq \phi$ and $\lim_{n \rightarrow \infty} \|x_n - T_i(PT_i)^{n-1}x_n\| = 0$ for all $i = 1, 2, \dots, N$, then we have $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$.

Theorem 2.5: Let E be a real uniformly convex Banach space and K be a nonempty closed convex subset which is also a nonexpansive retract of E . Let $T_1, T_2, \dots,$

$T_N: K \rightarrow K$ be N uniformly continuous asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{x_n\}$ be the sequence defined by (6) with $\sum_{n=1}^{\infty} \gamma_n^i < \infty$ and $\{\alpha_n^i\} \subseteq [\varepsilon, 1 - \varepsilon]$ for all $i = 1, 2, \dots, N$, for some $\varepsilon \in (0, 1)$. Then $\|x_n - T_i x_n\| = 0$ for all $i = 1, 2, \dots, N$.

Proof: Let $x^* \in F = \bigcap_{i=1}^N F(T_i)$. Then by Lemma 2.1 (i) and Lemma 1.1, $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists. Let $\lim_{n \rightarrow \infty} \|x_n - x^*\| = a$. If $a = 0$, then by the continuity of each T_i the conclusion follows. Now suppose that $a > 0$. Firstly, we are now to show that $\lim_{n \rightarrow \infty} \|T_N(PT_N)^{n-1}x_n - x_n\| = 0$. Since $\{x_n\}$ and $\{u_n^i\}$ are bounded for all $i = 1, 2, \dots, N$, there exists $R > 0$ such that $x_n - x^* + \gamma_n^i(u_n^i - x_n), T_i(PT_i)^{n-1}x_n^{i-1} - x^* + \gamma_n^i(u_n^i - x_n) \in B_R(0)$ for all $n \geq 1$ and for all $i = 1, 2, \dots, N$. Using Lemma 1.2, we have

$$\begin{aligned}
\|x_n^N - x^*\|^2 &= \|P(\alpha_n^N T_N(PT_N)^{n-1}x_n^{N-1} + \beta_n^N x_n + \gamma_n^N u_n^N) - Px^*\|^2 \\
&= \|\alpha_n^N T_N(PT_N)^{n-1}x_n^{N-1} + \beta_n^N x_n + \gamma_n^N u_n^N - x^*\|^2 \\
&= \|\alpha_n^N (T_N(PT_N)^{n-1}x_n^{N-1} - x^* + \gamma_n^N (u_n^N - x_n)) \\
&\quad + (1 - \alpha_n^N)(x_n - x^* + \gamma_n^N (u_n^N - x_n))\|^2 \\
&\leq \alpha_n^N \|T_N(PT_N)^{n-1}x_n^{N-1} - x^* + \gamma_n^N (u_n^N - x_n)\|^2 + (1 - \alpha_n^N) \\
&\quad \|x_n - x^* + \gamma_n^N (u_n^N - x_n)\|^2 \\
&\quad - W_2(\alpha_n^N)g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) \\
&\leq \alpha_n^N (\|T_N(PT_N)^{n-1}x_n^{N-1} - x^*\| + \gamma_n^N \|u_n^N - x_n\|)^2 + (1 - \alpha_n^N) \\
&\quad (\|x_n - x^*\| + \gamma_n^N \|u_n^N - x_n\|)^2 \\
&\quad - W_2(\alpha_n^N)g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) \\
&\leq \alpha_n^N [(1 + b_n^{N-2})\|x_n - x^*\| + d_n^{N-2} + \gamma_n^N \|u_n^N - x_n\|]^2 + (1 - \alpha_n^N) \\
&\quad [(1 + b_n^{N-2})\|x_n - x^*\| + d_n^{N-2} + \gamma_n^N \|u_n^N - x_n\|]^2 \\
&\quad - W_2(\alpha_n^N)g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) \\
&\leq [(1 + b_n^{N-2})\|x_n - x^*\| + d_n^{N-2} + \gamma_n^N \|u_n^N - x_n\|]^2 \\
&\quad - W_2(\alpha_n^N)g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) \\
&\leq [\|x_n - x^*\| + \lambda_n^{N-2}]^2 - W_2(\alpha_n^N)g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|)
\end{aligned} \tag{9}$$

where $\lambda_n^{N-2} = d_n^{N-2} + \gamma_n^N \|u_n^N - x_n\|$. Observe that $\varepsilon^3 \leq W_2(\alpha_n^N)$ now (9) implies that $\varepsilon^3 g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 + \rho_n^{N-2}$, where $\rho_n^{N-2} = 2\lambda_n^{N-2} + (\lambda_n^{N-2})^2$. Since $\sum_{n=1}^{\infty} d_n^{N-2} < \infty$ and $\sum_{n=1}^{\infty} \gamma_n^{N-2} < \infty$, we get $\sum_{n=1}^{\infty} \rho_n^{N-2} < \infty$. This implies that $\lim_{n \rightarrow \infty} g(\|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\|) = 0$. Since g is strictly increasing and continuous at 0, it follows that

$$\lim_{n \rightarrow \infty} \|T_N(PT_N)^{n-1}x_n^{N-1} - x_n\| = 0.$$

Since $T_N, \forall N$ is asymptotically nonexpansive, note that

$$\begin{aligned} \|x_n - x^*\| &\leq \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| + \|T_N(PT_N)^{n-1}x_n^{N-1} - x^*\| \\ &= \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| + (1 + r_n^N) \|x_n^{N-1} - x^*\| \end{aligned}$$

for all $n \geq 1$. Thus $a = \lim_{n \rightarrow \infty} \|x_n - x^*\| \leq \liminf_{n \rightarrow \infty} \|x_n^{N-1} - x^*\| \leq \limsup_{n \rightarrow \infty} \|x_n^{N-1} - x^*\| \leq a$ and therefore $\lim_{n \rightarrow \infty} \|x_n^{N-1} - x^*\| = a$. Using the same argument in the proof above, we have

$$\begin{aligned} &\|x_n^{N-1} - x^*\|^2 \\ &\leq \alpha_n^{N-1} \|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x^* + \gamma_n^{N-1}(u_n^{N-1} - x_n)\|^2 + (1 - \alpha_n^{N-1}) \\ &\quad \|x_n - x^* + \gamma_n^{N-1}(u_n^{N-1} - x_n)\|^2 \\ &\quad - W_2(\alpha_n^{N-1})g(\|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\|) \\ &\leq \alpha_n^{N-1} [(1 + b_n^{N-3}) \|x_n - x^*\| + d_n^{N-3} + \gamma_n^{N-1} \|u_n^{N-1} - x_n\|]^2 + (1 - \alpha_n^{N-1}) \\ &\quad [(1 + b_n^{N-3}) \|x_n - x^*\| + d_n^{N-3} + \gamma_n^{N-1} \|u_n^{N-1} - x_n\|]^2 \\ &\quad - W_2(\alpha_n^{N-1})g(\|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\|) \\ &\leq [(1 + b_n^{N-3}) \|x_n - x^*\| + d_n^{N-3} + \gamma_n^{N-1} \|u_n^{N-1} - x_n\|]^2 \\ &\quad - W_2(\alpha_n^{N-1})g(\|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\|) \\ &\leq [\|x_n - x^*\| + \lambda_n^{N-3}]^2 \\ &\quad - W_2(\alpha_n^{N-1})g(\|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\|) \quad (10) \end{aligned}$$

where $\lambda_n^{N-3} = d_n^{N-3} + \gamma_n^{N-1} \|u_n^{N-1} - x_n\|$.

This implies that

$$\varepsilon^3 g(\|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\|) \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 + \rho_n^{N-3},$$

where $\rho_n^{N-3} = 2\lambda_n^{N-3} + (\lambda_n^{N-3})^2$

and therefore $\lim_{n \rightarrow \infty} \|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\| = 0$.

Thus, we have

$$\begin{aligned} &\|x_n - T_N(PT_N)^{n-1}x_n\| \\ &\leq \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| + \|T_N(PT_N)^{n-1}x_n^{N-1} - T_N(PT_N)^{n-1}x_n\| \end{aligned}$$

$$\begin{aligned}
&\leq \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| + (1 + r_n^N) \|x_n^{N-1} - x_n\| \\
&\leq \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| \\
&\quad + (1 + r_n^N) \|\alpha_n^{N-1}T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} + \beta_n^{N-1}x_n + \gamma_n^{N-1}u_n^{N-1} - x_n\| \\
&\leq \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| \\
&\quad + (1 + r_n^N)[\alpha_n^{N-1} \|T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2} - x_n\| + \gamma_n^{N-1} \|u_n^{N-1} - x_n\|]
\end{aligned}$$

since

$$\lim_{n \rightarrow \infty} \|x_n - T_N(PT_N)^{n-1}x_n^{N-1}\| = 0, \quad \lim_{n \rightarrow \infty} \|x_n - T_{N-1}(PT_{N-1})^{n-1}x_n^{N-2}\| = 0$$

and $\sum_{n=1}^{\infty} \gamma_n^{N-1} < \infty$, $\sum_{n=1}^{\infty} r_n^N < \infty$, it follows that

$$\lim_{n \rightarrow \infty} \|x_n - T_N(PT_N)^{n-1}x_n\| = 0.$$

Similarly, by using the same argument as in the proof above we have

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \|x_n - T_{N-2}(PT_{N-2})^{n-1}x_n^{N-3}\| = \lim_{n \rightarrow \infty} \|x_n - T_{N-3}(PT_{N-3})^{n-1}x_n^{N-4}\| \\
&=, \dots, = \lim_{n \rightarrow \infty} \|x_n - T_2(PT_2)^{n-1}x_n^1\| = 0.
\end{aligned}$$

This implies that

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \|x_n - T_{N-1}(PT_{N-1})^{n-1}x_n\| = \lim_{n \rightarrow \infty} \|x_n - T_{N-2}(PT_{N-2})^{n-1}x_n\| \\
&=, \dots, = \lim_{n \rightarrow \infty} \|x_n - T_3(PT_3)^{n-1}x_n\| = 0.
\end{aligned}$$

It remains to show that

$$\lim_{n \rightarrow \infty} \|x_n - T_1(PT_1)^{n-1}x_n\| = 0, \quad \lim_{n \rightarrow \infty} \|x_n - T_2(PT_2)^{n-1}x_n\| = 0.$$

Note that

$$\begin{aligned}
&\|x_n^1 - x^*\|^2 \\
&\leq \alpha_n^1 (\|T_1(PT_1)^{n-1}x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|)^2 + (1 - \alpha_n^1) \\
&\quad (\|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|)^2 - W_2(\alpha_n^1)g(\|T_1(PT_1)^{n-1}x_n - x_n\|) \\
&\leq \alpha_n^1 [(1 + r_n^1) \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|]^2 + (1 - \alpha_n^1) \\
&\quad [(1 + r_n^1) \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|]^2 - W_2(\alpha_n^1)g(\|T_1(PT_1)^{n-1}x_n - x_n\|) \\
&\leq [(1 + r_n^1) \|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|]^2 \\
&\quad - W_2(\alpha_n^1)g(\|T_1(PT_1)^n x_{n-1} - x_n\|) \\
&\leq [\|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|]^2 - W_2(\alpha_n^1)g(\|T_1(PT_1)^{n-1}x_n - x_n\|)
\end{aligned}$$

Thus, we have $\varepsilon^3 g(\|T_1(PT_1)^{n-1}x_n - x_n\|) \leq [\|x_n - x^*\| + \gamma_n^1 \|u_n^1 - x^*\|]^2 - \|x_n^1 - x^*\|^2$ and therefore $\lim_{n \rightarrow \infty} \|T_1(PT_1)^{n-1}x_n - x_n\| = 0$.

Since

$$\begin{aligned} & \|x_n - T_2(PT_2)^{n-1}x_n\| \\ & \leq \|x_n - T_2(PT_2)^{n-1}x_n^1\| + \|T_2(PT_2)^{n-1}x_n^1 - T_2(PT_2)^{n-1}x_n\| \\ & \leq \|x_n - T_2(PT_2)^{n-1}x_n^1\| + (1 + r_n^2) \|x_n^1 - x_n\| \\ & \leq \|x_n - T_2(PT_2)^{n-1}x_n^1\| \\ & \quad + (1 + r_n^2) \|\alpha_n^1 T_1(PT_1)^{n-1}x_n + \beta_n^1 x_n + \gamma_n^1 u_n^1 - x_n\| \\ & \leq \|x_n - T_2(PT_2)^{n-1}x_n^1\| \\ & \quad + (1 + r_n^2) [\alpha_n^1 \|T_1(PT_1)^{n-1}x_n - x_n\| + \gamma_n^1 \|u_n^1 - x_n\|] \end{aligned}$$

it implies that $\lim_{n \rightarrow \infty} \|T_2(PT_2)^{n-1}x_n - x_n\| = 0$. Therefore

$$\lim_{n \rightarrow \infty} \|T_i(PT_i)^{n-1}x_n - x_n\| = 0$$

for all $i = 1, 2, \dots, N$.

On the other hand, by Remark 2.3, it is clear that $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$. Therefore, by Lemma 2.2, we can conclude that $\|x_n - T_i x_n\| = 0$ as $n \rightarrow \infty$. This completes the proof.

Theorem 2.6: Let E be a real uniformly convex Banach space and K be a nonempty closed convex subset which is also a nonexpansive retract of E . Let $T_1, T_2, \dots, T_N: K \rightarrow K$ be N asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{\alpha_n^i\}$ be the sequence defined by (6) with $\sum_{n=1}^{\infty} \gamma_n^i < \infty$ and $\{\alpha_n^i\} \subseteq [\varepsilon, 1 - \varepsilon]$ for all $i = 1, 2, \dots, N$, for some $\varepsilon \in (0, 1)$. Then for all $u, v \in F$, the limit

$$\lim_{n \rightarrow \infty} \|tx_n + (1-t)u - v\|$$

exists for all $t \in [0, 1]$.

Proof: By Lemma 2.2(i) and by Lemma 1.1, we have $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for all $x^* \in F$. This implies that $\{x_n\}$ is bounded. Observe that there exists $R > 0$ such that $\{x_n\} \subset C = B_R(0) \cap K$, where $B_R(0) = \{x \in E : \|x\| \leq R\}$. Then C is a nonempty closed convex bounded subset of E . Let $a_n(t) = \|tx_n + (1-t)u - v\|$. Then $\lim_{n \rightarrow \infty} a_n(0) = \|u - v\|$ and from Lemma 2.2(i) and 1.1, $\lim_{n \rightarrow \infty} a_n(1) = \lim_{n \rightarrow \infty} \|x_n - v\|$ exists. Without loss of generality, we may assume that $\lim_{n \rightarrow \infty} \|x_n - u\| = r > 0$ and $t \in (0, 1)$.

For any $n \geq 1$ and for all $i = 1, 2, \dots, N$, we define $A_n^i: C \rightarrow C$ by

$$A_n^1 = P(\alpha_n^1 T_1(PT_1)^{n-1} + \beta_n^1 I + \gamma_n^1 u_n^1)$$

$$\begin{aligned}
 A_n^2 &= P(\alpha_n^2 T_2 (PT_2)^{n-1} A_n^1 + \beta_n^2 I + \gamma_n^2 u_n^2) \\
 \dots &= \dots\dots\dots \\
 \dots &= \dots\dots\dots \\
 A_n^N &= P(\alpha_n^N T_N (PT_N)^{n-1} A_n^{N-1} + \beta_n^N I + \gamma_n^N u_n^N)
 \end{aligned}$$

Thus for all $x, y \in K$, we have

$$\begin{aligned}
 &\|A_n^i x - A_n^i y\| \\
 &= \|P(\alpha_n^i T_i (PT_i)^{n-1} A_n^{i-1} x + \beta_n^i x + \gamma_n^i u_n^i) \\
 &\quad - P(\alpha_n^i T_i (PT_i)^{n-1} A_n^{i-1} y + \beta_n^i y + \gamma_n^i u_n^i)\| \\
 &\leq \|(\alpha_n^i T_i (PT_i)^{n-1} A_n^{i-1} x + \beta_n^i x + \gamma_n^i u_n^i) - (\alpha_n^i T_i (PT_i)^{n-1} A_n^{i-1} y + \beta_n^i y + \gamma_n^i u_n^i)\| \\
 &\leq \alpha_n^i \|T_i (PT_i)^{n-1} A_n^{i-1} x \\
 &\quad - T_i (PT_i)^{n-1} A_n^{i-1} y\| + \beta_n^i \|x - y\| \\
 &\leq \alpha_n^i (1 + r_n^i) \|A_n^{i-1} x - A_n^{i-1} y\| + \beta_n^i \|x - y\| \\
 &\leq (1 + r_n^i) \|A_n^{i-1} x - A_n^{i-1} y\| + \|x - y\| \\
 &\leq k_n^i \|A_n^{i-1} x - A_n^{i-1} y\| + \|x - y\|
 \end{aligned}$$

where $k_n^i = (1 + r_n^i)$ for all $i = 2, 3, \dots, N$. Since $\sum_{n=1}^\infty r_n^i < \infty$ for all $i = 1, 2, \dots, N$, then $\prod_{i=1}^N k_n^i < \infty$, and

$$\begin{aligned}
 \|A_n^1 x - A_n^1 y\| &\leq \alpha_n^1 (1 + r_n^1) \|x - y\| + \beta_n^1 \|x - y\| \\
 &\leq [\alpha_n^1 (1 + r_n^1) + \beta_n^1] \|x - y\| \\
 &\leq [\alpha_n^1 + \beta_n^1 + \alpha_n^1 r_n^1] \|x - y\| \\
 &\leq (1 + r_n^1) \|x - y\| = k_n^1 \|x - y\|.
 \end{aligned}$$

This implies that

$$\begin{aligned}
 \|A_n^N x - A_n^N y\| &\leq \alpha_n^N \|T_N (PT_N)^{n-1} A_n^{N-1} x - T_N (PT_N)^{n-1} A_n^{N-1} y\| + \beta_n^N \|x - y\| \\
 &\leq \alpha_n^N (1 + r_n^N) \|A_n^{N-1} x - A_n^{N-1} y\| + \beta_n^N \|x - y\| \\
 &\leq (1 + r_n^N) \|A_n^{N-1} x - A_n^{N-1} y\| + \|x - y\| \\
 &\leq k_n^N \|A_n^{N-1} x - A_n^{N-1} y\| + \|x - y\|, \text{ where } k_n^N = (1 + r_n^N) \\
 &\leq k_n^N [k_n^{N-1} \|A_n^{N-2} x - A_n^{N-2} y\| + \|x - y\|] + \|x - y\| \\
 &\leq k_n^N k_n^{N-1} \|A_n^{N-2} x - A_n^{N-2} y\| + (1 + k_n^N) \|x - y\| \\
 &\leq \dots\dots\dots \\
 &\leq \dots\dots\dots
 \end{aligned}$$

$$\begin{aligned}
&\leq k_n^N k_n^{N-1} \dots k_n^2 \|A_n^1 x - A_n^1 y\| + [1 + k_n^N + k_n^N k_n^{N-1} + \\
&\quad \dots + k_n^N k_n^{N-1} \dots k_n^2] \|x - y\| \\
&\leq k_n^N k_n^{N-1} \dots k_n^2 k_n^1 \|x - y\| + [1 + k_n^N + k_n^N k_n^{N-1} + \\
&\quad \dots + k_n^N k_n^{N-1} \dots k_n^2] \|x - y\| \\
&\leq \left(\prod_{i=1}^N k_n^i \right) \|x - y\| + [1 + \left(\prod_{i=N}^N k_n^i \right) + \left(\prod_{i=N-1}^N k_n^i \right) + \\
&\quad \dots + \left(\prod_{i=2}^N k_n^i \right)] \|x - y\| \\
&\leq \left[\prod_{i=1}^N k_n^i + \prod_{i=2}^N k_n^i + \dots + \prod_{i=N}^N k_n^i + 1 \right] \|x - y\| \\
&\leq [\delta_n^1 + \delta_n^2 + \dots + \delta_n^N + 1] \|x - y\|, \text{ where } \delta_n^j = \prod_{i=j}^N k_n^i \\
&\leq \left(\sum_{p=0}^N \delta_n^p \right) \|x - y\| \\
&\leq \|x - y\|
\end{aligned}$$

that is

$$\|A_n^N x - A_n^N y\| \leq \|x - y\|.$$

Set $S_{n,m} = A_{n+m-1}^N A_{n+m-2}^N \dots A_n^N$, $m \geq 1$, and

$$b_{n,m} = \|S_{n,m}(tx_n + (1-t)u) - (tS_{n,m}x_n + (1-t)S_{n,m}u)\|.$$

It is easy to see that $A_n^N x_n = x_{n+1}$, $S_{n,m}x_n = x_{n+m}$ and $\|S_{n,m}x - S_{n,m}y\| \leq \|x - y\|$.

We show first that, for any $x^* \in F$, $\|S_{n,m}x^* - x^*\| \rightarrow 0$ uniformly for all $m \geq 1$ as $n \rightarrow \infty$. Indeed, for any $x^* \in F$, we have

$$\begin{aligned}
\|A_n^i x^* - x^*\| &\leq \alpha_n^i \|T_i (PT_i)^{n-1} A_n^{i-1} x^* - x^*\| + \gamma_n^i \|u_n^i - x^*\| \\
&\leq \alpha_n^i (1 + r_n^i) \|A_n^{i-1} x^* - x^*\| + \gamma_n^i \|u_n^i - x^*\| \\
&\leq (1 + r_n^i) \|A_n^{i-1} x^* - x^*\| + \|u_n^i - x^*\| \\
&\leq k_n^i \|A_n^{i-1} x^* - x^*\| + \|u_n^i - x^*\|, \text{ where } k_n^i = (1 + r_n^i)
\end{aligned}$$

for all $i = 2, 3, \dots, N$, and $k_n^i = (1 + r_n^i)$ for all $i = 1, 2, \dots, N$ and

$$\begin{aligned}
\|A_n^1 x^* - x^*\| &\leq \gamma_n^1 \|u_n^1 - x^*\| \\
&\leq \|u_n^1 - x^*\|.
\end{aligned}$$

Therefore

$$\begin{aligned}
\|A_n^N x^* - x^*\| &\leq \alpha_n^N \|T_N(PT_N)^{n-1} A_n^{N-1} x^* - x^*\| + \gamma_n^N \|u_n^N - x^*\| \\
&\leq \alpha_n^N (1 + r_n^N) \|A_n^{N-1} x^* - x^*\| + \gamma_n^N \|u_n^N - x^*\| \\
&\leq (1 + r_n^N) \|A_n^{N-1} x^* - x^*\| + \|u_n^N - x^*\| \\
&\leq k_n^N \|A_n^{N-1} x^* - x^*\| + \|u_n^N - x^*\| \\
&\leq \delta_n^1 \|u_n^1 - x^*\| + \delta_n^2 \|u_n^2 - x^*\| + \dots \\
&\quad + \delta_n^N \|u_n^{N-1} - x^*\| + \|u_n^N - x^*\| \\
&\leq M \sum_{p=0}^N \delta_n^p
\end{aligned}$$

for all $n \geq 1$, where $M = \max\{\sup_{n \geq 1}\{\|u_n^1 - x^*\|\}, \dots, \sup_{n \geq 1}\{\|u_n^N - x^*\|\}\}$ and $\delta_n^j = \prod_{i=j}^N k_n^i$ such that $\sum_{n=1}^{\infty} \delta_n^j < \infty$. Hence

$$\begin{aligned}
\|S_{n,m} x^* - x^*\| &= \|A_{n+m-1}^N A_{n+m-2}^N \dots A_n^N x^* - x^*\| \\
&\leq \|A_{n+m-1}^N A_{n+m-2}^N \dots A_n^N x^* - A_{n+m-1}^N A_{n+m-2}^N \dots A_{n+1}^N x^*\| \\
&\quad + \|A_{n+m-1}^N A_{n+m-2}^N \dots A_{n+1}^N x^* - A_{n+m-1}^N A_{n+m-2}^N \dots A_{n+2}^N x^*\| \\
&\quad + \dots + \|A_{n+m-1}^N x^* - x^*\| \\
&\leq \|A_n^N x^* - x^*\| + \|A_{n+1}^N x^* - x^*\| + \dots + \|A_{n+m-1}^N x^* - x^*\| \\
&\leq M \sum_{p=0}^N \delta_n^p + M \sum_{p=0}^N \delta_{n+1}^p + \dots + M \sum_{p=0}^N \delta_{n+m-1}^p \\
&\leq M \sum_{l=n}^{n+m-1} \left(\sum_{p=0}^N \delta_l^p \right) = \xi_n^{x^*}
\end{aligned}$$

since $\sum_{l=1}^{\infty} \delta_l^p < \infty$, we have $\xi_n^{x^*} \rightarrow 0$ as $n \rightarrow \infty$ and hence $\|S_{n,m} x^* - x^*\| \rightarrow 0$ as $n \rightarrow \infty$. Observe that

$$\begin{aligned}
a_{n+m}(t) &= \|tS_{n,m}x_n + (1-t)u - v\| \\
&\leq \|tS_{n,m}x_n + (1-t)u - S_{n,m}(tx_n + (1-t)u)\| \\
&\quad + \|S_{n,m}(tx_n + (1-t)u) - v\| \\
&\leq \|tS_{n,m}x_n + (1-t)S_{n,m}u - S_{n,m}(tx_n + (1-t)u) + (1-t)(u - S_{n,m}u)\| \\
&\quad + \|S_{n,m}(tx_n + (1-t)u) - v\| \\
&\leq b_{n,m} + (1-t) \|u - S_{n,m}u\| + \|S_{n,m}(tx_n + (1-t)u) - v\| \\
&\leq b_{n,m} + \|S_{n,m}(tx_n + (1-t)u) - S_{n,m}v\| + \|S_{n,m}v - v\| \\
&\quad + (1-t) \|u - S_{n,m}u\| \\
&\leq b_{n,m} + \|tx_n + (1-t)u - v\| + \|S_{n,m}v - v\| + (1-t) \|u - S_{n,m}u\| \\
&\leq b_{n,m} + a_n(t) + \xi_n^v + (1-t)\xi_n^u
\end{aligned} \tag{*}$$

By using [9], Theorem 2.3], we have

$$\begin{aligned} b_{n,m} &\leq \varphi^{-1}(\|x_n - u\| - \|S_{n,m}x_n - S_{n,m}u\|) \\ &\leq \varphi^{-1}(\|x_n - u\| - \|x_{n+m} - u + u - S_{n,m}u\|) \\ &\leq \varphi^{-1}(\|x_n - u\| - (\|x_{n+m} - u\| - \|S_{n,m}u - u\|)) \end{aligned}$$

and so the sequence $\{b_{n,m}\}$ converges uniformly to 0 as $n \rightarrow \infty$ for all $m \geq 1$. Thus, fixing n and letting $m \rightarrow \infty$ in (*), we have

$$\limsup_{m \rightarrow \infty} a_{n+m}(t) \leq \varphi^{-1}(\|x_n - u\| - (\lim_{m \rightarrow \infty} \|x_m - u\| - \xi_n^u)) + a_n(t) + \xi_n^v + (1-t)\xi_n^u$$

and again letting $n \rightarrow \infty$

$$\limsup_{n \rightarrow \infty} a_n(t) \leq \varphi^{-1}(0) + \liminf_{n \rightarrow \infty} a_n(t) + 0 + 0 = \liminf_{n \rightarrow \infty} a_n(t).$$

This shows that $\lim_{n \rightarrow \infty} a_n(t)$ exists, that is,

$$\lim_{n \rightarrow \infty} \|tx_n + (1-t)u - v\|$$

exists for all $t \in [0, 1]$. This completes the proof.

Theorem 2.7: Let E be a real uniformly convex Banach space such that its dual E^* has the *Kadec-Klee* property and K a nonempty closed convex subset of E which is also a nonexpansive retract of E . Let $T_1, T_2, \dots, T_N: K \rightarrow K$ be N asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{x_n\}$ be the sequence defined by (6) with $\sum_{n=1}^{\infty} \gamma_n^i < \infty$ and $\{\alpha_n^i\} \subseteq [\varepsilon, 1 - \varepsilon]$ for all $i = 1, 2, \dots, N$, for some $\varepsilon \in (0, 1)$. Then $\{x_n\}$ converges weakly to some common fixed point $x^* \in F$.

Proof: By Lemma 2.2 (i) and 1.1, we have $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for all $x^* \in F$. This implies that $\{x_n\}$ is bounded. Since E is reflexive, there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ converges weakly to some $x^* \in K$. By Theorem 2.5, we have $\lim_{j \rightarrow \infty} \|x_{n_j} - T_i x_{n_j}\| = 0$ for all $i = 1, 2, \dots, N$. Now Lemma 1.3 guarantees that $I - T_i$ is demiclosed at zero for all $i = 1, 2, \dots, N$. This implies that $T_i x^* = x^*$ for all $i = 1, 2, \dots, N$, hence this means that $x^* \in F$. It remains to show that $\{x_n\}$ converges weakly to x^* . Suppose $\{x_{n_i}\}$ is another subsequence of $\{x_n\}$ converges weakly to some y^* . Then $y^* \in K$ and so $x^*, y^* \in \omega_w(x_n) \cap F$. By Theorem 2.6, the limit

$$\lim_{n \rightarrow \infty} \|tx_n + (1-t)x^* - y^*\|$$

exists for all $t \in [0, 1]$. By Lemma 1.4, we have $x^* = y^*$. As a result, $\omega_w(x_n) \cap F$ is a singleton, and so $\{x_n\}$ converges weakly to some common fixed point $x^* \in F$.

This completes the proof.

Theorem 2.8: Let E be a real uniformly convex Banach space and K be a nonempty closed convex subset which is also a nonexpansive retract of E . Let $T_1, T_2, \dots, T_N: K \rightarrow K$ be N asymptotically nonexpansive nonself mappings with sequences $\{r_n^i\}$ such that $\sum_{n=1}^{\infty} r_n^i < \infty$, for all $1 \leq i \leq N$ and $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Suppose $\{T_1, T_2, \dots, T_N\}$ satisfies condition (B). Let $\{x_n\}$ be the sequence defined by (6) with $\sum_{n=1}^{\infty} \gamma_n^i < \infty$ and $\{\alpha_n^i\} \subseteq [\varepsilon, 1 - \varepsilon]$ for all $i = 1, 2, \dots, N$, for some $\varepsilon \in (0, 1)$. Then $\{x_n\}$ converges strongly to a common fixed point of the mappings $\{T_1, T_2, \dots, T_N\}$.

Proof: From Lemma 2.2(i) and 1.1 we see that $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for all $x^* \in F = \bigcap_{i=1}^N F(T_i)$. Let $\lim_{n \rightarrow \infty} \|x_n - x^*\| = a$ for some $a \geq 0$. Without loss of generality, if $a = 0$, then there is nothing to prove. Assume that $a > 0$, as proved in Lemma 2.2(i), we have

$$\|x_{n+1} - x^*\| \leq (1 + b_n^{N-1}) \|x_n - x^*\| + d_n^{N-1}, \quad \forall n \in N$$

where $\{b_n^i\}_{n=1}^{\infty}$ and $\{d_n^i\}_{n=1}^{\infty}$ for all $i = 1, 2, \dots, N$ are nonnegative real sequences such that $\sum_{n=1}^{\infty} b_n^i < \infty$ and $\sum_{n=1}^{\infty} d_n^i < \infty$ for all $i = 1, 2, \dots, N$. This gives that

$$d(x_{n+1}, F) \leq (1 + b_n^{N-1})d(x_n, F) + d_n^{N-1}, \quad \forall n \in N.$$

Applying Lemma 1.1 to the above inequality, we obtained that $\lim_{n \rightarrow \infty} d(x_n, F)$ exists. Also by Theorem 2.5, $\lim_{n \rightarrow \infty} \|x_n - T_i x_n\| = 0$ for all $i = 1, 2, \dots, N$. Since $\{T_1, T_2, \dots, T_N\}$ satisfies condition (B), we conclude that $\lim_{n \rightarrow \infty} d(x_n, F) = 0$.

Next we show that $\{x_n\}$ is a Cauchy sequence. Since $\lim_{n \rightarrow \infty} d(x_n, F) = 0$, given any $\varepsilon > 0$, there exists a natural number n_0 such that $d(x_n, F) < \frac{\varepsilon}{3}$ for all $n \geq n_0$. So we can find $p^* \in F$ such that $\|x_{n_0} - p^*\| < \frac{\varepsilon}{2}$. For all $n \geq n_0$ and $m \geq 1$, we have

$$\begin{aligned} \|x_{n+m} - x_n\| &\leq \|x_{n+m} - p^*\| + \|x_n - p^*\| \\ &\leq \|x_{n_0} - p^*\| + \|x_n - p^*\| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

This shows that $\{x_n\}$ is a Cauchy sequence and so is convergent since E is complete. Let $\lim_{n \rightarrow \infty} x_n = q^*$. Then $q^* \in K$. It remains to show that $q^* \in F$. Let $\varepsilon_1 > 0$ be given. Then there exists a natural number n_1 such that $\|x_n - q^*\| < \frac{\varepsilon_1}{4}$ for all $n \geq n_1$. Since $\lim_{n \rightarrow \infty} d(x_n, F) = 0$, there exists a natural number $n_2 \geq n_1$ such that for all $n \geq n_2$ we have $d(x_n, F) < \frac{\varepsilon_1}{5}$ and in particular, we have $d(x_{n_2}, F) < \frac{\varepsilon_1}{5}$. Therefore, there exists $w^* \in F$ such that $\|x_{n_2} - w^*\| < \frac{\varepsilon_1}{4}$. For any $i \in I$ and

$n \geq n_2$, we have

$$\begin{aligned}
 \|T_i q^* - q^*\| &\leq \|T_i q^* - w^*\| + \|w^* - q^*\| \\
 &\leq 2\|q^* - w^*\| \\
 &\leq 2(\|q^* - x_{n_2}\| + \|x_{n_2} - w^*\|) \\
 &< 2\left(\frac{\varepsilon_1}{4} + \frac{\varepsilon_1}{4}\right) \\
 &< \varepsilon_1.
 \end{aligned}$$

This implies that $T_i q^* = q^*$. Hence $q^* \in F(T_i)$ for all $i \in I$ and so $q^* \in F = \bigcap_{i=1}^N F(T_i)$. Thus $\{x_n\}$ converges strongly to a common fixed point of the mappings $\{T_1, T_2, \dots, T_N\}$. This completes the proof.

Remark 2.9: Theorem 2.5 to 2.8 extend the corresponding results of Su and Qin [28] to the case of multistep iterative sequences with errors for a finite family of asymptotically nonexpansive nonself mappings and also they extend many known results.

Remark 2.10: Our results also extend the corresponding results of Plubtieng and Ungchittarakool [22] to more general class of nonexpansive nonself mappings.

Remark 2.11: Our results also extend the corresponding results of Shahzad [18] to the case of multistep iterative sequences with errors for a finite family of more general class of nonexpansive mappings.

References

- [1] B.E. Rhoades, *Fixed point iteration for certain nonlinear mappings*, J. Math. Anal. Appl. 183(1994), 118-120.
- [2] C.E. Chidume, *On the approximation of fixed points of nonexpansive mappings*, Houston J. Math. 7(1981), 345-355.
- [3] C.E. Chidume, *Nonexpansive mappings, generalizations and iterative algorithms*. In: Agarwal R.P., O'Reagan D.eds. Nonlinear Analysis and Application. To V. Lakshmikantham on his 80th Birthday (Research Monograph), Dordrecht: Kluwer Academic Publishers, pp. 383-430.
- [4] C.E. Chidume; E.U. Ofoedu and H. Zegeye, *Strong and weak convergence theorems for asymptotically nonexpansive mappings*, J. of Mathematical Analysis and Applications, 280(2003), 364-374.
- [5] C.E. Chidume; N. Shahzad and H. Zegeye, *Strong convergence theorems for nonexpansive mappings in arbitrary Banach spaces*, Nonlinear Anal. Submitted.

- [6] G.B. Passty, *Construction of fixed points for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. 84(1982), 212-216.
- [7] H.F. Senter and W.G. Dotson, *Approximating fixed points of nonexpansive mappings*, Proc. Amer. Math. Soc. 44(1974), 375-380.
- [8] H.K. Xu, *Existence and convergence for fixed points of mappings of asymptotically nonexpansive type*, Nonlinear Analysis, 16(1991), 1139-1146.
- [9] J. Garcia Falset; W. Kaczor; T. Kuczumow and S. Reich, *Weak convergence theorems for asymptotically nonexpansive mappings and semigroups*, Nonlinear Analysis. Theory, Methods and Applications 43(2001), no.3, 377-401.
- [10] J. Schu, *Iterative construction of fixed points of asymptotically nonexpansive mappings*, J. Math. Anal. Appl. 158(1991), 407-413.
- [11] J. Schu, *Weak and strong convergence theorems to fixed points of asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. 43(1991), 153-159.
- [12] K. Goebel and W.A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. 35(1972), 171-174.
- [13] K.K. Tan and H.K. Xu, *A nonlinear ergodic theorem for asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. 45(1992), 25-36.
- [14] K.K. Tan and H.K. Xu, *The nonlinear ergodic theorem for asymptotically nonexpansive mapping in Banach spaces*, Proc. Amer. Math. Soc. 114(1992), 399-404.
- [15] K.K. Tan and H.K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. 178(1993), 301-308.
- [16] K.K. Tan and H.K. Xu, *Fixed point iteration processes for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. 122(1994), 733-739.
- [17] M.O. Osilike and S.C. Aniagbosor, *Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings*, Math. Comput. Modelling 32(2000), 1181-1191.
- [18] N. Shahzad, *Approximating fixed points of non-self nonexpansive mappings in Banach spaces*, Nonlinear Analysis 61(2005), no.6, 1031-1039.
- [19] S.C. Bose, *Weak convergence to a fixed point of an asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. 68(1978), 305-308.
- [20] S. Ishikawa, *Fixed points by a new iteration method*, Proc. Amer. Math. Soc. 44(1974), 147-150.
- [21] S. Ishikawa, *Fixed points and iteration of a nonexpansive mapping in a Banach space*, Proc. Amer. Math. Soc. 59(1976), 65-71.

- [22] S. Plubtieng and K. Ungchittarakool, *Weak and strong convergence of finite family with errors of nonexpansive non-self mappings*, Fixed Point Theory and Applications, Volume 2006, Article ID 81493, pages 1-12.
- [23] S. Reich, *Weak convergence theorems for nonexpansive mappings in Banach spaces*, J. Math. Anal. Appl. 67(1979), 274-276.
- [24] S.S. Chang; Y.J. Cho and H. Zhou, *Demi-closed principle and weak convergence problems for asymptotically nonexpansive mappings*, J. Korean Math. Soc. 38(6) (2001), 1245-1260.
- [25] W. Kaczor, *Weak convergence of almost orbits of asymptotically nonexpansive commutative semigroups*, J. Mathematical Analysis and Applications 272(2002), no.2, 565-574.
- [26] W.J. Davis and P. Enflo, *Contractive projections on l_p spaces*, Analysis at Urbana, Vol.I (Urbana, IL, 1986-1987), London Math. Soc. Lecture Note Ser., vol. 137, Cambridge University Press, Cambridge, 1989, pp. 151-161.
- [27] W.R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. 4(1953), 506-510.
- [28] Yongfu Su and Xiaolong Qin, *Approximating fixed points of non-self asymptotically nonexpansive mappings in Banach spaces*, J. Applied Mathematics and Stochastic Analysis, Volume 2006, Article ID 21961, pages 1-13.
- [29] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. 73(1967), 591-597.

Department of Mathematics & I.T., Govt. College of Science, Raipur (C.G.),
India

E-mail: saluja.1963@rediffmail.com