

Approximation algorithms for TSP with neighborhoods in the plane

Adrian Dumitrescu *

University of Wisconsin–Milwaukee
Milwaukee, WI 53201-0784
ad@cs.uwm.edu

Joseph S. B. Mitchell †

Stony Brook University
Stony Brook, NY 11794-3600
jsbm@ams.sunysb.edu

Abstract

In the Euclidean TSP with neighborhoods (TSPN), we are given a collection of n regions (*neighborhoods*) and we seek a shortest tour that visits each region. As a generalization of the classical Euclidean TSP, TSPN is also NP-hard. In this paper, we present new approximation results for the TSPN, including (1) a constant-factor approximation algorithm for the case of arbitrary connected neighborhoods having comparable diameters; and (2) a PTAS for the important special case of disjoint unit disk neighborhoods (or nearly disjoint, nearly-unit disks). Our methods also yield improved approximation ratios for various special classes of neighborhoods, which have previously been studied. Further, we give a linear-time $O(1)$ -approximation algorithm for the case of neighborhoods that are (infinite) straight lines.

1 Introduction

A salesman wants to meet a set of n potential buyers. Each buyer specifies a connected region in the plane, his *neighborhood*, within which he is willing to meet the salesman. For example, the neighborhoods may be disks centered at buyers' locations, and the radius of each disk specifies the maximum distance a buyer is willing to travel to the meeting place. The salesman wants to find a tour of shortest length that visits all of buyers' neighborhoods and finally returns to its initial departure point. A variant of the problem, which we will address in this paper, is that in which no departure point is specified, and only a tour of the neighborhoods is to be found. This problem, which is known as the *TSP with neighborhoods* (TSPN), is a generalization of the classic Euclidean Traveling Salesman Problem (TSP), in which the regions, or “neighborhoods,” are single points, and consequently is NP-hard [9, 21].

Related Work. The TSP has a long and rich history of research in combinatorial optimization. It has been studied extensively in many forms, including geometric instances; see [4, 13, 14, 20, 24]. The problem is known to be NP-hard, even for points in the Euclidean plane [9, 21]. It has recently been shown that the geometric instances of the TSP, including the Euclidean TSP, have a polynomial-time approximation scheme, as developed by Arora [2] and Mitchell [17], and later improved by Rao and Smith [23].

*This work was done while the author was a visiting faculty member at Stony Brook University.

†Partially supported by grants from HRL Laboratories, NASA Ames, the National Science Foundation (CCR-9732220), Northrop-Grumman Corporation, Sandia National Labs, and Sun Microsystems.

Arkin and Hassin [1] were the first to study approximation algorithms for the geometric TSPN. They gave $O(1)$ -approximation algorithms for several special cases, including parallel segments of equal length, translates of a convex region, translates of a connected region, and more generally, for regions which have diameter segments that are *parallel* to a common direction, and the ratio between the longest and the shortest diameter is bounded by a constant.

For the general case of connected polygonal regions, Mata and Mitchell [15] obtained an $O(\log n)$ -approximation algorithm, based on “guillotine rectangular subdivisions”, with time bound $O(N^5)$, where N is the total complexity of the n regions. Gudmundsson and Levcopoulos [10] have recently obtained a faster method, which, for any fixed $\epsilon > 0$, is guaranteed to perform at least one of the following tasks¹: (1) it outputs a tour of length at most $O(\log n)$ times optimum in time $O(n \log n + N)$; (2) it outputs a tour of length at most $(1 + \epsilon)$ times optimum in time $O(N^3)$. So far, no polynomial-time approximation algorithm is known for general connected regions.

Recently it was shown that TSPN is APX-hard and cannot be approximated within a factor of 1.000374 unless $P=NP$ [5, 11]. In fact, the inapproximability factor for the vertex cover problem in graphs with degree bounded by 5, as stated in [7], and on which the result in [11] is based, implies a factor larger than that.

In the time since this paper first appeared, it has been shown by de Berg et. al [5] that the TSPN has an $O(1)$ -approximation algorithm in the case that the regions are connected and *fat*. Also, Schwartz and Safra [25] have improved the lower bounds of [5, 11] on the hardness of approximation of several variants of the TSPN problem, and Jonsson [12] has given an $O(n)$ time $\sqrt{2}$ -approximation algorithm for the case of regions that are lines in the plane; our previous $O(n)$ time $\pi/2$ -approximation algorithm appears in Section 5.

Summary of Our Results. In this paper, we obtain several approximation results on the geometric TSPN, including:

- (1) We extend the approaches initiated in [1] and obtain the first $O(1)$ -approximation algorithm for the TSPN having connected regions of the same or similar diameter. This solves among others, the open problem posed in [1], to provide a constant-factor approximation algorithm for TSPN on segments of the same length and *arbitrary* orientation.
- (2) We give a polynomial-time approximation scheme (PTAS) for the case of disjoint unit disks or the case of nearly disjoint disks of nearly the same size. The algorithm is based on applying the m -guillotine method with a new area-based charging scheme. The fact that there is a PTAS for the case in which the neighborhoods are “nice,” with no point lying in more than a constant number of neighborhoods, should be contrasted with the fact that the TSPN on arbitrary regions is APX-hard. The construction in the proof of [11] utilizes “skinny” neighborhoods, which intersect each other extensively.
- (3) We also give modest improvements on earlier approximation bounds in [1] for the cases of parallel segments of equal length, translates of a convex region, and translates of a connected region.
- (4) We present simple algorithms which achieve a constant-factor guarantee for the case of equal disks and for the case of infinite straight lines.

¹One does not know in advance which one will be accomplished.

Preliminaries. The input to our algorithms will be a set \mathcal{R} of n regions, each of which is a simply-connected, closed subset of the plane, \mathbb{R}^2 , bounded by a finite union of arcs of constant-degree algebraic curves (degenerate regions are simply points). Since the regions are assumed to be closed, they include the points that lie on the curves that form their boundary. The assumption that regions are *simply* connected means that each region has no “holes.”

Ideally, each region is a subset of the plane lying inside a simple, closed, continuous curve, together with the curve itself. However, we can only deal with regions that are computer-representable, hence the above definition.

Examples of allowable regions include simple polygons, whose boundaries are unions of a finite number of straight line segments, circular disks, regions bounded by straight segments and circular arcs, infinite straight lines, etc. We let N denote the total number of arcs specifying all n regions in \mathcal{R} , i.e. the total combinatorial complexity of the input.

A *tour* (or *circuit*) T , is a closed continuous curve that visits each region of \mathcal{R} . The *length* of tour T , denoted $|T|$, is the Euclidean length of the curve T . To avoid ambiguity, the size of a finite set X is denoted by $\#X$. In the *TSP with neighborhoods* (TSPN) problem, our goal is to compute a tour whose length is guaranteed to be close to the shortest possible length of a tour. We let T^* denote any optimal tour and let $L^* = |T^*|$ denote its length. An algorithm that outputs a tour whose length is guaranteed to be at most $c \cdot L^*$ is said to be a *c-approximation algorithm* and to have an *approximation ratio* of c . A family of $(1 + \epsilon)$ -approximation algorithms, parameterized by $\epsilon > 0$, and each running in polynomial time for fixed ϵ , is said to be a *polynomial-time approximation scheme* (PTAS).

Outline of the Paper. In Section 2 we use some simple packing arguments to yield approximation algorithms for the TSPN for equal-size disks. Section 3 presents a PTAS for the TSPN for equal-size disks. In Section 4 we give an approximation algorithm for the TSPN for regions having the same diameter. Finally, in Section 5, we give an approximation algorithm for the case of regions that are infinite straight lines. We conclude with a short list of open problems for future research.

2 Equal-Size Disks

We begin by giving some simple arguments and corresponding algorithms that achieve a constant approximation ratio for TSPN on a set \mathcal{R} of n disks of the same size. Without loss of generality, we assume that all disks have unit radius. Our results carry over naturally to disks of nearly the same size, with corresponding changes in the approximation factor.

First, we consider the case of disjoint unit disks. The algorithm is simple and natural: using known PTAS results for TSP on points [2, 17, 23], compute, in time $O(n \log n)$, a $(1 + \epsilon)$ -approximate tour, $T = T_C$, of the center points of the n disks. We refer to it as the *center tour*. Clearly T_C is a valid region tour. We claim that

Proposition 1 *Given set of n disjoint unit disks, one can compute a tour, T , whose length satisfies*

$$|T| \leq \left(1 + \frac{8}{\pi}\right)|T^*| + 8(1 + \epsilon),$$

where T^* is an optimal tour. The running time is dominated by that of computing a $(1 + \epsilon)$ -approximate tour of n points.

Proof. Put $L^* = |T^*|$. Since T^* visits all disks, the area A_{L^*} swept by a disk of radius 2, whose center moves along T^* , covers all of the unit disks \mathcal{R} . This area is bounded as follows

$$\pi n \leq A_{L^*} \leq 4L^* + 4\pi.$$

Thus, $n \leq 4 + \frac{4L^*}{\pi}$. A center tour of length at most

$$L^* + 2n \leq L^* + 2\left(4 + \frac{4L^*}{\pi}\right) = \left(1 + \frac{8}{\pi}\right)L^* + 8$$

can be obtained by going along T^* and making a detour of length at most 2 to visit the center of a disk at each point where T^* first visits a disk. Hence, the length of the computed tour is bounded as claimed. \square

For large n , the approximation ratio above is $(1 + \frac{8}{\pi})(1 + \epsilon) \leq 3.55$; for small constant values of n , the problem can be solved exactly using brute force. We note that for any algorithm that outputs a tour on the center points, we cannot expect an approximation ratio smaller than 2. To see this, consider a large square, and place almost touching unit disks along its perimeter, both on its inside and on its outside. All of the disks touch the perimeter, which is also an optimal disk tour except at the four corners of the square. The length of the disk center tour is roughly two times the perimeter of the square.

When the disks are nearly of the same size, so that the ratio between the maximum and minimum radius is bounded by a constant $k > 1$, for large n , the approximation ratio is about $1 + \frac{8k^2}{\pi}$.

Next we consider the case in which the disks can overlap. First, we compute a maximal independent (pairwise-disjoint) set I of disks. Next, we compute T_I , a $(1 + \epsilon)$ -approximate tour of the center points of disks in I . Finally, we output a tour T obtained by following the tour T_I , taking detours around the boundaries of each of the disks in I , as illustrated in Figure 1. More specifically, we select an arbitrary disk D_0 and one of the intersection points, s , between D_0 and T_I . We start at point s and go clockwise along T_I . Whenever the boundary of a disk of I is encountered, we follow clockwise around the boundary of the disk until we encounter again the tour T_I . When we finally reach s , we continue clockwise around D_0 to T_I and then continue counterclockwise around T_I , again taking detours clockwise around the disks of I that we encounter along the way. The tour T finally ends when we return the second time to s . In this way, our tour T traverses the boundary of each disk of I exactly once and therefore visits all of the disks that are not in I as well. We remark that this method of constructing a feasible tour of the disks results in a slightly better worst-case ratio than another natural strategy for extending T_I to a full disk tour: While traversing T_I , each time one encounters the boundary of a disk $D \in I$, traverse the entire circumference of D exactly once, and then go directly to the point where the tour T_I exits D and continue along T_I .

Denote by T^* an optimal disk tour, of length L^* , and by T_I^* an optimal disk tour of the subset $I \subseteq \mathcal{R}$ of disks. A constant approximation ratio can be derived from the following three inequalities:

$$|T| \leq \pi|T_I| + 2\pi, \tag{1}$$

$$|T_I| \leq \left(1 + \frac{8}{\pi}\right)|T_I^*| + 8)(1 + \epsilon), \tag{2}$$

$$|T_I^*| \leq L^*. \tag{3}$$




Figure 1: Construction of the tour T from the tour T_I on the center points of a maximal independent set, I , of disks. Starting from s , we follow the thick solid tour, back to s , and then follow the thick dashed tour back around to s again.

The third inequality follows from the fact that $I \subseteq \mathcal{R}$, and the second from the case of disjoint unit disks considered above. To check the first inequality, decompose T_I into $\#I$ parts, assuming $\#I \geq 2$, one for each disk in I , by cutting each segment between two consecutive centers in the middle. Let $x + y = d \geq 2$ be the length of one of these parts of tour T_I , corresponding to a disk D , where x, y are the lengths of the two segments of T_I adjacent to the center of D . Write a_1, a_2 for the arc lengths of D when its boundary is traversed by T ; we have $a_1 + a_2 = 2\pi$. Writing the ratio of the length of the corresponding part of T to the length, d , of this part of T_I , we get

$$\frac{(x - 1 + a_1 + y - 1) + (x - 1 + a_2 + y - 1)}{d} = \frac{2(d + \pi - 2)}{d} \leq \pi,$$

the maximum being attained when $d = 2$. When $\#I = 1$, we have $|T_I| = 0, |T| = 2\pi$. Thus (1) is satisfied in all cases. Putting (1), (2), (3) together, we get

Proposition 2 *Given set of n unit disks, possibly overlapping, one can compute a tour, T , whose length satisfies*

$$|T| \leq ((\pi + 8)|T^*| + 8\pi)(1 + \epsilon) + 2\pi,$$

where T^* is an optimal tour. The running time is dominated by that of computing a $(1 + \epsilon)$ -approximate tour of n points.

For large n , the approximation ratio is $(\pi + 8)(1 + \epsilon) \leq 11.15$. We note that the approximation ratio we have obtained with this approach for disjoint unit disks, 3.55, (resp., 11.15 for unit disks) is better (resp., weaker) than $\sqrt{3^2 + 7^2} \approx 7.62$, the approximation ratio that will be given at the end of Section 4 for translates of a convex region, which applies, of course, to the case of unit disks.

3 A PTAS for Disjoint Equal Disks

In this section we present a polynomial-time approximation scheme for the TSPN in the case of regions that are equal-size disks or nearly equal-size disks.

Given the powerful methods that have been developed to obtain PTAS’s for various geometric optimization problems, such as the Euclidean TSP, it is natural to suspect that these same techniques may apply to the TSPN. Indeed, one may expect that TSPN should have a PTAS based on applying existing methods. However, we know now, from the recent APX-hardness result of [11], that this cannot be. What goes wrong?

The basic issue we must address in order to apply these techniques is to be able to write a recursion to solve an appropriate “succinct” subproblem with dynamic programming. What is the subproblem “responsible” for solving? For problems involving *points*, the subproblem can be made responsible for constructing some kind of inexpensive network on the points *inside* the subproblem defined by a rectangle, and to interconnect this network with the boundary in some nicely controlled way, e.g., with only a constant complexity of connection, in the case of *m*-guillotine methods. The problem with *regions* is that they can cross subproblem boundaries. Then, we do not know if the subproblem is responsible to visit the region, or if the region is visited *outside* the subproblem. We cannot afford to enumerate the subset of regions that cross the boundary for which the subproblem is responsible – there are too many such subsets, leading to too many subproblems. Thus, we need a new idea.

Our approach is to employ a new type of structural result, based on the general method of *m*-guillotine subdivisions. In particular, we show how to transform an optimal tour into one of a special class of tours that recursively has a special *m*-guillotine structure, permitting us to have a succinct, constant-size specification of the subset of regions, crossing the subproblem boundary, for which the subproblem is “responsible” in that it must visit these regions on its interior. In order to bound the increase in tour length in performing this transformation, we must “charge off” the added tour length to some small fraction of the length of the optimal tour, just as is done in proving the bounds for the *m*-guillotine PTAS method for TSP. In order to do this charging, we must assume some special structure on the class of neighborhoods in the TSPN, e.g., that the regions $\mathcal{D} = \{D_1, \dots, D_n\}$ are pairwise-disjoint, equal-size disks, or have a similar structure allowing us to relate tour length to area.

Here, we show how the approach applies to disjoint disks having equal radii, δ ; generalizations are readily made to the case of nearly equal radii, with a constant upper bound on the ratio of radii, and to the case of “modestly overlapping,” in which the disks become pairwise-disjoint if all of them are decreased in size by a constant factor while keeping the center points the same.

We begin with some definitions, largely following the notation of [17]. Let G be an embedding of a planar graph, and let L denote the total Euclidean length of its edges, E . We can assume without loss of generality that G is restricted to the unit square, B ; i.e., $E \subset \text{int}(B)$.

Consider an axis-aligned rectangle W , a *window*, with $W \subseteq B$. Rectangle W will correspond to a subproblem in a dynamic programming algorithm. Let ℓ be an axis-parallel line intersecting W . We refer to ℓ as a *cut* and assume, without loss of generality, that ℓ is vertical.

The intersection, $\ell \cap (E \cap \text{int}(W))$, of a cut ℓ with $E \cap \text{int}(W)$ consists of a possibly empty set of subsegments of ℓ . These subsegments are possibly singleton points. Let ξ be the number of endpoints of such subsegments along ℓ , and let the points be denoted by p_1, \dots, p_ξ , in order of decreasing y -coordinate along ℓ . For a positive integer m , we define the *m*-span, $\sigma_m(\ell)$, of ℓ with respect to W as follows. If $\xi \leq 2(m-1)$, then $\sigma_m(\ell) = \emptyset$; otherwise, $\sigma_m(\ell)$ is defined to be the line segment, $p_m p_{\xi-m+1}$, joining the m th endpoint, p_m , with the m th-from-the-last endpoints, $p_{\xi-m+1}$. Refer to Figure 2. Note that the segment $p_m p_{\xi-m+1}$ may be of zero length in case $p_m = p_{\xi-m+1}$.

The intersection, $\ell \cap \mathcal{D} \cap W$ of ℓ with the disks that intersect W consists of a possibly empty set of $\xi_{\mathcal{D}} \leq |\mathcal{D} \cap W|$ subsegments of ℓ , one for each disk that is intersected by ℓ in W . Let these disks




Figure 2: Definition of m -span: Here, the 3-span, $\sigma_3(\ell)$ of ℓ with respect to the window $W \subset B$ is highlighted with a thick shaded vertical segment.




Figure 3: Definition of m -disk-span: Here, the 3-disk-span, $\sigma_{3,\mathcal{D}}(\ell)$ of ℓ with respect to the window $W \subset B$ is highlighted with a thick shaded vertical segment.

be $D_1, D_2, \dots, D_{\xi_{\mathcal{D}}}$, in order of decreasing y -coordinate. For a positive integer m , we define the m -disk-span, $\sigma_{m, \mathcal{D}}(\ell)$, of ℓ with respect to W as follows. If $\xi_{\mathcal{D}} \leq 2m$, then $\sigma_{m, \mathcal{D}}(\ell) = \emptyset$; otherwise, $\sigma_{m, \mathcal{D}}(\ell)$ is defined to be the possibly zero-length line segment joining the bottom endpoint of $\ell \cap D_m$ with the top endpoint of $\ell \cap D_{\xi_{\mathcal{D}}-m+1}$. Refer to Figure 3.

Line ℓ is an m -good cut with respect to W if $\sigma_m(\ell) \subseteq E$ and $\sigma_{m, \mathcal{D}}(\ell) \subseteq E$. In particular, if $\xi \leq 2(m-1)$ and $\xi_{\mathcal{D}} \leq 2m$, then ℓ is trivially an m -good cut, since both the m -span and the m -disk-span are empty in this case.

We now say that E satisfies the m -guillotine property with respect to window W if either (1) W does not fully contain any disk; or (2) there exists an m -good cut, ℓ , that splits W into W_1 and W_2 , and, recursively, E satisfies the m -guillotine property with respect to both W_1 and W_2 .

Theorem 1 *Let G be an embedded connected planar graph, with edge set E , of total length L , and let \mathcal{D} be a given set of pairwise-disjoint equal-radius disks (of radius δ) each of which intersects E . Assume that E and \mathcal{D} are contained in the unit square B . Then, for any positive integer m , there exists a planar graph G' that satisfies the m -guillotine property with respect to B and has an edge set $E' \supseteq E$ of length*

$$L' \leq \left(1 + \frac{\sqrt{2} + 16/\pi}{m}\right) L + \frac{16\delta}{m}.$$

Proof. We convert G into a new graph G' by adding to E a new set of horizontal/vertical edges whose total length is at most $\frac{\sqrt{2}+16/\pi}{m}L + \frac{16\delta}{m}$. The construction is recursive: at each stage, we show that there exists a cut, ℓ , with respect to the current window W (which initially is the unit square B), such that we can afford to add both the m -span and the m -disk-span to E .

We say that a point p on a cut ℓ is m -dark with respect to ℓ and W if, along $\ell^\perp \cap \text{int}(W)$, there are at least m edges of E intersected by ℓ^\perp on each side of p , where ℓ^\perp is the line through p and perpendicular to ℓ .² We say that a subsegment of ℓ is m -dark (with respect to W) if all points of the segment are m -dark with respect to ℓ and W .

The important property of m -dark points along ℓ is the following: Assume, without loss of generality, that ℓ is horizontal. We consider any line segment that lies along an edge of E to have a *top* side and a *bottom* side; the top is the side that can be seen from above, from a point with $y = +\infty$. Then, if all points on subsegment pq of ℓ are m -dark, we can charge the length of pq off to the bottoms of the first m subsegments, $E^+ \subseteq E$, of edges that lie above pq , and the tops of the first m subsegments, $E^- \subseteq E$, of edges that lie below pq , since we know that there are at least m edges “blocking” pq from the top/bottom of W . We charge pq ’s length half to E^+ , charging each of the m levels of E^+ from below, with $\frac{1}{2m}$ units of charge, and half to E^- , charging each of the m levels of E^- from above, with $\frac{1}{2m}$ units of charge. We refer to this type of charge as the “red” charge.

We say that a point p on a cut ℓ is m -disk-dark with respect to ℓ and W if, along $\ell^\perp \cap \text{int}(W)$, there are at least m disks of \mathcal{D} that have a nonempty intersection with $\ell^\perp \cap W$ on each side of p . Here, again, ℓ^\perp is the line through p and perpendicular to ℓ . We say that a subsegment of ℓ is m -disk-dark with respect to W if all points of the segment are m -disk-dark with respect to ℓ and W . The *chargeable* length within W of a cut ℓ is defined to be the sum of the lengths of its m -dark portion and its m -disk-dark portion. Refer to Figure 4.

²We can think of the edges E as being “walls” that are not very effective at blocking light — light can go through $m-1$ walls, but is stopped when it hits the m th wall; then, p on a line ℓ is m -dark if p is not illuminated when light is shone in from the boundary of W , along the direction of ℓ^\perp .




Figure 4: Definition of m -disk-dark: Here, the points that are 2-disk-dark with respect to ℓ are those four subsegments of ℓ that lie within the shaded regions, which comprise the set $R_x^{(2,D)}$ of points of W that are 2-disk-dark with respect to horizontal cuts.

The important property of m -disk-dark points along horizontal ℓ is the following: If all points on subsegment pq of ℓ are m -disk-dark, then we can charge the length of pq off to the bottoms of the first m disks that lie above pq , and the tops of the first m disks that lie below pq , since we know that there are at least m disks “blocking” pq from the top/bottom of W . We charge pq ’s length half upwards, charging the bottoms of each of the m “levels” of disks with $\frac{1}{2m}$ units of charge, and half downwards, charging the tops of each of the m “levels” of disks with $\frac{1}{2m}$ units of charge. We refer to this type of charge as the “blue” charge.

A cut ℓ is *favorable* if its chargeable length within W is *at least as large as* the sum of the lengths of the m -span and the m -disk-span. We do not attempt to take advantage of the fact that there may be overlap among the m -span and the m -disk-span, nor that there may be portions of these two segments that are already part of E ; an argument taking advantage of these facts may improve slightly the constants in some of our bounds.

The existence of a favorable cut is guaranteed by the following key lemma, whose proof is similar to that of the key lemma in [17]:

Lemma 1 *For any G and any window W , there is a favorable cut.*

Proof. We show that there must be a favorable cut that is either horizontal or vertical.

Let $f(x)$ denote the “cost” of the vertical line, ℓ_x , passing through the point $(x, 0)$, where “cost” means the sum of the lengths of the m -span and the m -disk-span for ℓ_x . Then,

$$f(x) = |\sigma_m(\ell_x)| + |\sigma_{m,D}(\ell_x)|.$$

Thus,

$$A_x = \int_0^1 f(x)dx = A_x^{(m)} + A_x^{(m,D)} = \int_0^1 |\sigma_m(\ell_x)|dx + \int_0^1 |\sigma_{m,D}(\ell_x)|dx,$$

where $A_x^{(m)} = \int_0^1 |\sigma_m(\ell_x)|dx$ is the area of the x -monotone region $R_x^{(m)}$ of points of B that are m -dark with respect to horizontal cuts, and $A_x^{(m,D)} = \int_0^1 |\sigma_{m,D}(\ell_x)|dx$ is the area of the x -monotone

region $R_x^{(m,\mathcal{D})}$ of points of B that are m -disk-dark with respect to horizontal cuts. Refer to Figure 4. Similarly, define $g(y)$ to be the cost of the horizontal line through y , and let $A_y = \int_0^1 g(y)dy$.

Assume, without loss of generality, that $A_x \geq A_y$. We claim that there exists a horizontal favorable cut; i.e., we claim that there exists a horizontal cut, ℓ , such that its chargeable length is at least as large as the cost of ℓ , meaning that the length of its m -dark portion plus its m -disk-dark portion is at least $|\sigma_m(\ell)| + |\sigma_{m,\mathcal{D}}(\ell)|$. To see this, note that A_x can be computed by switching the order of integration, “slicing” the regions $R_x^{(m)}$ and $R_x^{(m,\mathcal{D})}$ horizontally, rather than vertically; i.e., $A_x = \int_0^1 h(y)dy = \int_0^1 h_m(y)dy + \int_0^1 h_{m,\mathcal{D}}(y)dy$, where $h_m(y)$ is the m -dark length of the horizontal line through y , $h_{m,\mathcal{D}}(y)$ is the length of the intersection of $R_x^{(m,\mathcal{D})}$ with a horizontal line through y , and $h(y)$ is the chargeable length of the horizontal line through y . In other words, $h_m(y)$ (resp., $h_{m,\mathcal{D}}(y)$) is the length of the m -dark (resp., m -disk-dark) portion of the horizontal line through y . Thus, since $A_x \geq A_y$, we get that $\int_0^1 h(y)dy \geq \int_0^1 g(y)dy \geq 0$. Thus, it cannot be that for all values of $y \in [0, 1]$, $h(y) < g(y)$, so there exists a $y = y^*$ for which $h(y^*) \geq g(y^*)$. The horizontal line through this y^* is a cut satisfying the claim of the lemma.

If, instead, we had $A_x \leq A_y$, then we would get a *vertical* cut satisfying the claim. \square

Now that we know there must be a favorable cut, ℓ , we can charge off the cost of the m -span and the m -disk-span of ℓ , making “red” charge on the bottoms (resp., tops) of segments of E that lie above (resp., below) m -dark points of ℓ , and making “blue” charge on the bottoms (resp., tops) of disks that lie above (resp., below) m -disk-dark points of ℓ . We then recurse on each side of the cut, in the two new windows.

After a portion of E has been charged *red* on one side, due to a cut ℓ , it will be within m levels of the boundary of the windows on either side of ℓ , and, hence, within m levels of the boundary of any future windows, found deeper in the recursion, that contain the portion. Thus, no portion of E will ever be charged *red* more than once from each side, in each of the two directions, horizontal or vertical, so no portion of E will ever pay more than $\sqrt{2}/m$ times its length in red charge. We charge at the rate of $\frac{1}{2m}$ per unit length of the perimeter of the segment’s axis-aligned bounding box, and the worst case is achieved for a segment of slope ± 1 . Thus, the total red charge is at most $\frac{\sqrt{2}}{m}L$.

Similarly, no disk will ever have its boundary charged *blue* more than once from each of the two directions, horizontal or vertical. Since it is charged at the rate of $\frac{1}{2m}$ per unit length of its axis-aligned bounding box, whose perimeter is 8δ , we get a total blue charge of at most $\frac{4\delta}{m}n$. We now appeal to the lower bound from the previous section, which was based on an area argument. Here, for radius δ disks, that argument shows that $\pi\delta^2n \leq A_L \leq 4\delta L + (2\delta)^2\pi$, from which we get $n \leq \frac{4}{\pi\delta}L + 4$. Note that this area argument uses the fact that E is connected. Thus, the total blue charge is at most $\frac{16}{\pi m}L + \frac{16\delta}{m}$.

It is also important to note that we are always charging red portions of the original edge set E : the new edges added are never themselves charged, since they lie on window boundaries and cannot therefore serve to make a portion of some future cut m -dark.

Overall, then, the total increase in length caused by adding the m -spans and m -disk-spans along favorable cuts is bounded by

$$\frac{16 + \pi\sqrt{2}}{\pi m}L + \frac{16\delta}{m}.$$

Our goal in adding the m -disk-span is to obtain a succinct representation of which disks that straddle the boundary of a window are visited within the window and which are visited outside

the window. The m -disk-span segment visits all but a constant, $O(m)$, number of the disks on the corresponding side of the window. There is, however, one remaining issue with respect to the m -disk-span segments: We need to argue that we can “afford,” within our charging scheme, to *connect* the m -disk-span to the input edge set, E . This is because, in our dynamic programming optimization, we will find a shortest possible planar graph with the m -guillotine property that obeys certain connectivity constraints, as well as other properties that guarantee that the graph has an Eulerian subgraph spanning all disks. The optimal graph that we compute uses the m -disk-span segments to visit the corresponding disks on the boundaries of windows that define subproblems.

Remark. (*Clarification added after journal publication.*) We had previously phrased the above as “find a shortest possible connected planar graph with the m -guillotine property”; we have rephrased to be “find a shortest possible planar graph with the m -guillotine property that obeys certain connectivity constraints”.

In particular, we add connections to E from the endpoints of the m -disk-span to the point of E in the corresponding disk that is closest to the endpoint. We know that this connection is of length at most 2δ per endpoint of the m -disk-span, since the disks are of diameter 2δ . In total, this adds only 4δ to the length of the m -disk-span. Assuming the m -disk-span stabs at least three or more disks, its length is at least $\Omega(\delta)$, implying that we can charge off this extra 4δ for the connections in the same way that we charge off the m -disk-spans themselves. If, on the other hand, the m -disk-span stabs only one or two disks, then we can afford to skip the addition of the m -disk-span altogether, and just keep track in the dynamic program of the necessary information for these couple extra disks, specifying whether they are to be visited within the window or not.

Remark. (*Clarification added after journal publication; thanks to Sophie Spirkl for her inquiry and input.*) We argue above that we can afford to add connections to E , since the length added is proportional to the bridge length. Our dynamic programming algorithm computes a minimum-length planar network with the m -guillotine property that obeys certain connectivity constraints, so that all of the network is connected except (possibly) the m -disk-spans, which may not be connected to the rest of the (connected) network. The objective function in the dynamic program requires that we minimize the total length of the network, counting the lengths of the bridges that serve as m -disk-spans a constant number of times. Then, since we know we can afford to add length proportional to the total m -disk-span lengths, we know we can add the connections mentioned above, resulting in an overall connected network (and can make it Eulerian with appropriate doubling of the bridge segments, in the usual way, as mentioned again below), appropriately close to the optimal length. \square

Corollary 1 *The TSPN for a set of disjoint equal-size disks has a PTAS. The same is true for a set of nearly disjoint, nearly equal-size disks, for which there is a constant upper bound on the ratio of largest to smallest radius and there is a constant factor such that the disks become disjoint if their radii are each multiplied by the factor while keeping the center points the same.*

Proof. We consider only the case of disjoint disks each of radius δ ; the generalization to nearly disjoint, nearly equal-size disks is straightforward.

We impose a regular m -by- m grid on each disk; let \mathcal{G} denote the resulting set of $O(m^2n)$ grid points.

Consider an optimal tour, OPT , of length L^* . Now, OPT is a simple polygon. We can perturb the vertices of OPT so that each lies at a grid point in \mathcal{G} , resulting in a new tour, OPT' ,

visiting every disk, whose length is at most $L^* + O(n\delta/m)$. Using the fact, from the previous section, that $n \leq \frac{4}{\pi\delta}L^* + 4$, and assuming that $n \geq 8$, we get that the length of OPT' is at most $L^* + O(\frac{4}{\pi m}L^* + \frac{4\delta}{m}) \leq L^*(1 + O(\frac{8}{\pi m}))$. If $n < 8$, then we can solve the problem in constant time by brute force.

Theorem 1 implies that we can convert OPT' , which consists of some set E of edges, into a planar graph, OPT'' , obeying the m -guillotine property, while not increasing the total length by too much. In particular, the length of OPT'' is at most

$$(1 + \frac{\sqrt{2} + 16/\pi}{m})L^*(1 + O(\frac{8}{\pi m})) + \frac{16\delta}{m} \leq (1 + O(1/m))L^*.$$

We now apply a dynamic programming algorithm, running in $O(n^{O(m)})$ time, to compute a minimum-length planar graph having a prescribed set of properties: (1) it satisfies the m -guillotine property, which is necessary for the dynamic program to have the claimed efficiency; (2) it visits at least one grid point of \mathcal{G} in each region D_i ; and (3) it contains an Eulerian subgraph that spans the disks. This third condition that allows us to extract a tour in the end. We only outline here the dynamic programming algorithm; the details are very similar to those of [17], with the modification to account for the m -disk-span.

A subproblem is defined by a rectangle W whose coordinates are among those of the grid points \mathcal{G} , together with a constant amount ($O(m)$) of information about how the solution to the subproblem interacts across the boundary of W with the solution outside of W . This information includes the following:

- (a) For each of the four sides of W , we specify a “bridge” segment and at most $2m$ other segments with endpoints among \mathcal{G} that cross the side; this is done exactly as in the case of the Euclidean TSP on points, as in [17].
- (b) For each of the four sides of W , we specify a “disk bridge” segment corresponding to the m -disk-span, and, for each of at most $2m$ disks that are not intersected by the disk bridge segment, we specify in a single bit whether the disk is to be visited within the subproblem or not; if not, it is visited outside the window W .
- (c) We specify a required “connection pattern” within W . In particular, we indicate which subsets of the $O(m)$ specified edges crossing the boundary of W are required to be connected within W . This, again, is done exactly as is detailed for the Euclidean TSP on point sets in [17].

In order to end up with a graph having an Eulerian subgraph spanning the disks, we use the same trick as done in [17]: we “double” the bridge segments and the disk bridge segments, and then require that the number of connections on each side of a bridge segment satisfy a parity condition. Exactly as in [17], this allows us to extract a tour from the planar graph that results from the dynamic programming algorithm, which gives a shortest possible graph that obeys the specified conditions.

The result is that in polynomial time ($O(n^{O(m)})$) one can compute a shortest possible graph, from a special class of such graphs, and this graph spans the regions \mathcal{D} . Theorem 1 guarantees that the length of the resulting graph is very close, within factor $1 + O(1/m)$, to the length of an optimal solution to the TSPN. (We also know from the remarks previously, that we can afford to add connections to assure that the m -disk-spans are connected to the rest of the (connected) network.) Thus, once we extract a tour from the Eulerian subgraph, we have the desired $(1 + \epsilon)$ -approximation solution, where $\epsilon = O(1/m)$.

Finally, we mention that the running time of $O(n^{O(m)})$ can be improved to $O(n^C)$, for a constant C independent of m , using the notion of grid-rounded guillotine subdivisions, as in [18, 19]; here, the dependence on the constant m is exponential in the multiplicative constant concealed by the big-Oh of $O(n^C)$. We suspect that the techniques of Rao and Smith [23], based on Arora’s method [2], can be used to improve the time bound to $O(n \log n)$, while possibly also addressing the problem in higher dimensions; we leave this for future work. \square

4 Connected Regions of the Same Diameter

In this section we give a constant-factor approximation algorithm for the TSPN problem that applies in the case that all regions have the same diameter or nearly the same diameter.

The *diameter* of a region, δ , is the distance between two points in the region that are farthest apart. Without loss of generality, we assume that all regions have unit diameter, $\delta = 1$. The general method we use is to carefully select a representative point in each region, and then compute an almost optimal tour on these representative points. This approach was initiated in [1]. We also employ the following tool:

Lemma 2 (Combination Lemma) [1] *Given regions that can be partitioned into two types, and constants c_1, c_2 bounding the error ratios with which we can approximate optimal tours on regions of types 1 and 2, then we can approximate the optimal tour on all regions with an error ratio bounded by $c_1 + c_2 + 2$.*

As noted in [1], repeated application of this lemma may be useful: if there are k different region types, with individual approximation bounds of c_1, \dots, c_k , the resulting ratio is bounded by $c_1 + \dots + c_k + 2(k - 1)$. Now we describe the algorithm, which is similar in many aspects to the one in [1].

In each region, compute a unit-length diameter segment; in case of multiple such segments, select one arbitrarily. Computing a diameter segment of a region can be done efficiently, in time linear in the complexity of the region. Classify the regions into two types: (1) those for which the selected diameter is *almost horizontal*, by which we mean its slope is between -45° and 45° ; (2) those for which the selected diameter is *almost vertical*, by which we mean all the others. Use algorithm **A** (below) for each of these two region types. We will prove that a constant ratio is achievable for each class, after a suitable transformation is applied to regions of type (2). We then apply the Combination Lemma to obtain a constant-factor approximation for all regions.

Algorithm **A** gives a constant-factor approximation of the optimal tour for regions of type (1). Regions of type (2) are readily handled by rotating them by 90° to obtain type (1) regions. When the diameters are nearly the same, so that the ratio of the largest to the smallest is bounded by a constant, we can still get an approximation algorithm with a constant ratio; however since this ratio is rather large even for same diameters, we omit the calculations.

A set of lines is a *cover* of a set of regions if each region is intersected by at least one line from the covering set. We refer to such a set of lines as *covering lines*.

Algorithm A. Input: a set of n regions of type (1).




Figure 5: Illustration of Step 1 in Algorithm A

Step 1. Construct a cover of the regions by a minimum number of vertical lines. This procedure works in a greedy fashion, namely the leftmost line is as far right as possible, so that it is a right tangent of some region. To obtain this cover, the intervals of projection on the x -axis of all regions are computed and a greedy cover of this set of intervals is found. After removing all intervals covered by a previous line, and there are still uncovered intervals, another covering line is repeatedly added to the cover. At the same time, a representative point for each region is arbitrarily selected on the corresponding covering line and inside the region, e.g., the topmost boundary point of intersection between the region and its covering line. In this way n representative points, one per region, are selected. An illustration of this procedure appears in Figure 5. An important remark is that the representative points are not necessarily on the selected diameters, since a diameter may not be entirely contained in its region.

Assuming that we have only regions of type (1), the greedy cover has the effect of obtaining a large enough horizontal distance between any two consecutive covering lines. We remark that the greedy covering algorithm for a set of closed intervals on a line is known to output a cover of minimum size, a property which carries over to our vertical line cover.

Step 2. Proceed according to the following three cases.

Case 1: The greedy cover contains one covering line.

Compute a smallest perimeter axis-aligned rectangle Q that intersects all regions, where Q is considered a two-dimensional domain. Let w and h denote Q 's width and height respectively. Consider Q to be a graph with four vertices and four edges (its four sides). Let e_1 and e_2 be the vertical segments of height h that partition Q into three equal-width parts. Add to the graph Q a double edge corresponding to e_1 and a doubled edge corresponding to e_2 . The

resulting graph G is an Eulerian multigraph, since all node degrees are even, with 8 vertices and 12 edges. Output any Euler tour T of G . While such a tour is not, in general, the shortest possible tour visiting the regions, it suffices for our purposes of approximation.

Case 2: The greedy cover contains two covering lines.

Move (if possible) the rightmost vertical covering line to the left as much as possible (while still covering all regions). Recompute the representative points obtained in this way. Set D to be the distance between the two covering lines (clearly $D > 0$).

Case 2.1: $D \geq 3$.

Compute an axis-aligned rectangle Q of width $w = D$, with its vertical sides along the two covering lines, and of minimal height, which includes all representative points (on the two covering lines). Let h denote Q 's height. Output the tour T that is the perimeter of Q .

Case 2.2: $D \leq 3$ (similar to Case 1).

Compute a smallest perimeter axis-aligned rectangle Q that touches (intersects) all regions, where Q is considered to be a two-dimensional domain. Let w and h denote Q 's width and height respectively. Note that $w \leq 5$. Let e_1, \dots, e_7 be the vertical segments of height h that partition Q into eight equal-width parts. Consider the edges of Q , together with doubled copies of the edges e_1, \dots, e_7 , to define an Eulerian multigraph, G , having 18 vertices and 32 edges. Output any Euler tour T of G .

Case 3: The greedy cover contains at least three covering lines.

Compute T , a $(1 + \epsilon)$ -approximate tour of the representative points as the output tour.

If the regions are simple polygons, then a minimum-perimeter touching rectangle Q is determined by at most four contact points with region boundary arcs. A brute force procedure examining all possible k -tuples, $k \leq 4$, of such arcs computes Q in $O(N^5)$ time. The total running time of the approximation algorithm is either bounded by the complexity of the above step or by the complexity of computing $(1 + \epsilon)$ -approximate tours on n points, depending on the size of the greedy cover.

Theorem 2 *Given a set \mathcal{R} of n connected regions of the same diameter in the plane, a $O(1)$ -approximation of an optimal tour can be computed in polynomial time.*

Proof. Let OPT be an optimal region tour. We address each of the cases we distinguished in the previous algorithm. We will use repeatedly the following simple fact (see [1]): For positive a, b, w, h the following inequality holds

$$aw + bh \leq \sqrt{a^2 + b^2} \sqrt{w^2 + h^2}. \quad (4)$$

Case 1. Write $diag(Q)$ for the diagonal of the rectangle Q . We first argue that T visits all regions. Since all regions are covered by a unique covering line, they lie in a vertical strip of width $\leq 2\delta = 2$. So $w \leq 2$. The horizontal projection of each type (1) region (on the x -axis) is at least $1/\sqrt{2}$. Hence, each region is intersected either by the boundary of Q or by one of the two vertical segments inside Q , since these segments partition Q into three subrectangles each of width at most $2/3 < 1/\sqrt{2}$.

(Each region that is not intersected by the perimeter of Q lies entirely inside Q). Consequently, T is a valid region tour. We first give a lower bound on $|OPT|$. Since the length of a tour touching all four sides of a rectangle is at least twice the length of the diagonal of the rectangle (see [1]),

$$|OPT| \geq 2diag(Q) = 2\sqrt{w^2 + h^2}.$$

The length of T is

$$|T| = 2w + 6h = 2(w + 3h) \leq 2\sqrt{10}\sqrt{w^2 + h^2} \leq \sqrt{10}|OPT|.$$

Case 2.1: $D \geq 3$. We distinguish two sub-cases.

Case 2.1.a: $h \leq 2$. Recall that $w = D \geq 3$.

$$|OPT| \geq 2(w - 2),$$

$$|T| = 2w + 2h \leq 2w + 4 \leq 10(w - 2) \leq 5|OPT|.$$

Case 2.1.b: $h \geq 2$. Since each region has unit diameter, the optimal tour may lie inside Q at distance at most one from its boundary, so it must touch all four sides of some rectangle Q' having width $\geq w - 2$ and height $\geq h - 2$. Hence

$$|OPT| \geq 2\sqrt{(w - 2)^2 + (h - 2)^2}.$$

Since $8w + 8h \geq 40$, we have

$$|T| = 2w + 2h \leq 10((w - 2) + (h - 2)) \leq 10\sqrt{2}\sqrt{(w - 2)^2 + (h - 2)^2} \leq 5\sqrt{2}|OPT|.$$

Case 2.2: $D \leq 3$. The horizontal projection of each region is at least $1/\sqrt{2}$ and the rectangle Q , of width $w \leq 5$, is partitioned into 8 subrectangles, each of width $\leq 5/8$, by the vertical segments e_1, \dots, e_7 . Thus, since $1/\sqrt{2} > 5/8$, T visits all regions. (As in Case 1, each region that is not intersected by the perimeter of Q lies entirely inside Q ; hence, it is intersected by one of the seven vertical segments.) A similar calculation yields

$$|OPT| \geq 2diag(Q) = 2\sqrt{w^2 + h^2}.$$

$$|T| = 2w + 16h \leq 2\sqrt{1^2 + 8^2}\sqrt{w^2 + h^2} \leq 2(8.1)\sqrt{w^2 + h^2} \leq 8.1|OPT|.$$

Case 3. Partition the optimal tour OPT into blocks OPT_i , with $i \geq 1$. OPT_1 starts at an arbitrary point of intersection of OPT with the leftmost covering line, and ends at the last intersection of OPT with the second from the left covering line, before OPT intersects a different covering line. Notice that OPT does not cross to the left of the leftmost covering line. In general, the blocks OPT_i are determined by the last point of intersection of OPT with a covering line, before OPT crosses a different covering line. In Figure 5, the hypothetical optimum tour is partitioned into six blocks AB, BC, CD, DE, EF, FA . For example, OPT crosses the second vertical line twice before crossing the third vertical line, and B is the last point of intersection with the second line. Consider the bounding box Q of OPT_i , the smallest perimeter axis-aligned rectangle which includes OPT_i . Write w for its width and h for its height. There are two cases to consider.

Case 3.1: OPT_i intersects regions stabbed by two consecutive covering lines only, l_1, l_2 say, at distance w_1 . This implies that OPT_i lies between these two covering lines, so $w = w_1$. Without

loss of generality, OPT_i touches the lower side of Q , at some point C , before it touches the upper side of Q , at some point D . Since the horizontal projection of each region is at least $1/\sqrt{2}$, we have $w_1 \geq 1/\sqrt{2}$. Let $0 \leq a, b \leq h$ specify the starting and ending points A and B of OPT_i ; see Figure 6. Namely, a is the vertical distance between A , the start point of OPT_i , and the lower horizontal side of Q , and b is the vertical distance between B , the end point of OPT_i , and the upper horizontal side of Q . By considering the reflections of OPT_i with respect to the two horizontal sides of Q , we get that $|OPT_i|$ is bounded from below by the length of the polygonal line $A'CDB'$:

$$|OPT_i| \geq |A'CDB'| \geq \sqrt{(h+a+b)^2 + w_1^2}. \quad (5)$$




Figure 6: Case 3.1: 2 lines.

We show that there exists a partial tour T_i (a path) of the representative points of all the regions OPT_i visits, of length bounded by $c|OPT_i|$ for some positive constant c , where T_i starts at A and ends at B . Take $T_i = ADEFGB$, where the points D, E, F, G are on the lines l_1, l_2 at unit distance from the corners of Q ; see Figure 6.

$$\begin{aligned} |T_i| &\leq (a+1) + (1+h+1) + (w_1) + (1+h+1) + (1+h-b) \leq 3(h+a+b) + (w_1+6) \\ &\leq 3(h+a+b) + (6\sqrt{2}+1)w_1 \leq \sqrt{3^2 + (6\sqrt{2}+1)^2} \sqrt{(h+a+b)^2 + w_1^2} < 9.95|OPT_i|. \end{aligned} \quad (6)$$

Put T_i together to get a tour T of the representative points, having length smaller than $9.95|OPT|$. Then, for $\epsilon \leq 0.05$, the $(1+\epsilon)$ approximate tour of representatives, which the algorithm actually computes, has length at most $10|OPT|$.

Case 3.2: OPT_i intersects regions stabbed by three consecutive covering lines only, l_1, l_2 and l_3 . Denote by w_1 (resp., w_2) the horizontal distance between l_1 and l_2 (resp., l_2 and l_3). This implies that OPT_i lies between these three covering lines, but it does not touch the rightmost line l_3 . We can assume that OPT_i starts at A and ends at B , and as in the previous case, that it touches the lower side of Q , at some point C , before it touches the upper side of Q , at some point D . Let l' be the supporting line of the right side of Q , and w' be the horizontal distance between l_2 and l' . We have $w_1, w_2 \geq 1/\sqrt{2}$ and $w' \geq \max(0, w_2 - 1)$. The case when OPT_i touches the upper side of Q before it touches l' is shown in Figure 7; the other case is similar and we get the same bound on $|OPT_i|$. Let a, b, A, B be as before. We distinguish two sub-cases:

Case 3.2.a: $w_2 \leq 1$. The lower bound on OPT_i we have used earlier is still valid.

$$|OPT_i| \geq \sqrt{(h+a+b)^2 + w_1^2}.$$

We show a partial tour T_i of the representative points of all the regions OPT_i visits. Take $T_i = ADEFBGHIFB$; see Figure 7. Points E, F and I (resp., D, G and H) are on the lines l_1, l_2 and l_3 at unit vertical distance above the upper side (resp., below the lower side) of Q .

$$\begin{aligned} |T_i| &\leq (a+1) + (1+h+1) + w_1 + (1+h+1) + w_2 + (1+h+1) + w_2 + (1+b) \\ &\leq 3(h+a+b) + (w_1 + 2w_2 + 8) \leq 3(h+a+b) + (10\sqrt{2} + 1)w_1 \\ &\leq \sqrt{3^2 + (10\sqrt{2} + 1)^2} \sqrt{(h+a+b)^2 + w_1^2} < 15.45|OPT_i|. \end{aligned} \quad (7)$$

Put T_i together to get a tour T of the representative points, having length smaller than $15.45|OPT|$. Then, for $\epsilon \leq 0.05$, the $(1 + \epsilon)$ -approximate tour of representatives has length at most $15.5|OPT|$.




Figure 7: Case 3.2: 3 lines.

Case 3.2.b: $w_2 \geq 1$. We use a different lower bound on OPT_i .

$$|OPT_i| \geq \sqrt{(h+a+b)^2 + (w_1 + 2w_2 - 2)^2}, \quad (8)$$

which we get by considering the reflections of OPT_i with respect to the two horizontal sides of Q and with respect to l' ; see Figure 7. Take $T_i = ADEFBGHIFB$ as in Case 3.2.a.

$$\begin{aligned} |T_i| &\leq 3(h+a+b) + (w_1 + 2w_2 + 8) \leq 3(h+a+b) + (10\sqrt{2} + 1)(w_1 + 2w_2 - 2) \\ &\leq \sqrt{3^2 + (10\sqrt{2} + 1)^2} \sqrt{(h+a+b)^2 + (w_1 + 2w_2 - 2)^2} < 15.45|OPT_i|. \end{aligned} \quad (9)$$

Put T_i together to get a tour T of the representative points, having length smaller than $15.45|OPT|$. Then, for $\epsilon \leq 0.05$, the $(1 + \epsilon)$ -approximate tour of representatives has length at most $15.5|OPT|$. The overall approximation ratio of the algorithm, derived from the Combination Lemma is $15.5 + 15.5 + 2 = 33$. \square

Comparison with [1]. We point out some of the similarities and differences between the techniques of Arkin and Hassin [1] and our algorithm and its analysis, which is based on theirs. First, in [1], three different algorithms are presented, for parallel equal-length segments, translates of a convex region, and translates of an arbitrary connected region. The second two algorithms are refinements of the first, and representative points are chosen differently in each case. We presented here a single algorithm that works for all regions of type (1). Our consideration of cases 1 and 2 are slightly different from the corresponding cases of [1], which allows us to handle a larger class of inputs, namely regions of type (1); e.g. in case 2, [1] distinguishes between the subcases $D \leq 1$ and $D \geq 1$, while we distinguish between $D \leq 3$ and $D \geq 3$, and the cases are treated slightly differently. Finally, the analysis of the algorithms is based on similar ideas, e.g. to divide the optimal tours into blocks; our analysis differs in being able to address a single algorithm and a larger class of inputs.

Some Special Cases

We note that our calculations of the approximation ratio for connected regions of a same diameter, give improved bounds for three cases addressed in [1]:

1. parallel equal segments, from $3\sqrt{2} + 1$, to $3\sqrt{2}$,
2. translates of a convex region, from $\sqrt{3^2 + 7^2} + 1$, to $\sqrt{3^2 + 7^2}$,
3. translates of a connected region, from $\sqrt{3^2 + 11^2} + 1$, to $\sqrt{3^2 + 11^2}$.

The reason why these cases can be improved, and the new approximation ratios have similar expressions with the old ones, is that our algorithm **A** is similar to the algorithms in [1] for those cases.

We exemplify here the case of parallel equal segments and omit details for the rest. The algorithm computes a greedy cover of the segments, assumed to be of unit length, using vertical lines. Then it proceeds according to the cardinality of the cover. Cases 1 and 2 are treated in [1], and the ratio is $\sqrt{2}$; this is not the bottleneck case. In Case 1 (one covering line), an optimal tour is easy to obtain. In Case 2 (two covering lines), a smallest aligned rectangle which touches all segments is the output tour. In Case 3 (three or more covering lines) the algorithm computes an almost optimal tour of the representative points, as algorithm **A** does. Its analysis is divided into two sub-cases, as in the proof of Theorem 2. In the first sub-case (OPT_i intersects segments covered by two consecutive covering lines only), the lower bound in Equation (5) on $|OPT_i|$ is valid. The upper bound in Equation (6) on $|T_i|$ is adjusted by dropping the constant term equal to +6. Then

$$|T_i| \leq 3(h + a + b) + w_1 \leq \sqrt{3^2 + 1^2}|OPT_i| = \sqrt{10}|OPT_i|.$$

In the second sub-case (OPT_i intersects segments covered by three consecutive covering lines only), the lower bound in Equation (8) on $|OPT_i|$ is valid. The upper bound in Equation (9) on $|T_i|$ is adjusted by dropping the constant term equal to +8. We also have that $w_1, w_2 > 1$. Then

$$|T_i| < 3(h + a + b) + w_1 + 2w_2 \leq 3(h + a + b) + 3(w_1 + 2w_2 - 2) \leq 3\sqrt{2}|OPT_i|.$$

The overall approximation ratio obtained for parallel equal segments is $3\sqrt{2}$.

5 Lines

We consider now the case in which the n regions defining the TSPN instance are infinite straight lines in the plane. It is interesting that this case allows for an exact solution in polynomial time:

Proposition 3 *Given a set L of n infinite straight lines in the plane, a shortest tour that visits L can be computed in polynomial time.*

Proof. We convert the problem to an instance of the *watchman route problem* in a simple polygon. A *watchman route* in a polygon is a tour inside the polygon, such that every point in the polygon is visible from some point along the tour. The watchman route problem asks for a watchman route of minimum length [3]. The problem is known to have an $O(n^5)$ algorithm; see [26], as well as [8]. Let B be a rectangle that contains all of the vertices of the arrangement of L . At any one of the two points of intersection between a line $l_i \in L$ and the boundary of B , we extend a very narrow “spike” outward from that point, along l_i , for some fixed distance. Let P be the simple polygon having $3n + 4$ vertices that is the union of B and these n spikes, as illustrated in Figure 8. We make the observation that a tour T visits all of the lines in L if and only if it sees all of the polygon P , as required by the watchman route problem. Consequently, we can solve the TSPN on the set of lines L by solving the watchman route problem on polygon P . \square




Figure 8: Proof of Proposition 3.

Given the high running time of the watchman route algorithms, it is of interest to consider more efficient algorithms that may approximate the optimal solution. To this end, we now present a *linear-time* constant-factor approximation algorithm.

Let $L = \{l_1, \dots, l_n\}$ be the input set of n lines. A *minimum touching circle (disk)* is a circle of minimum radius which intersects all of the lines in L . The algorithm computes and outputs C_L , a minimum touching circle for L . We will show that this provides a tour of length at most $\frac{\pi}{2}|OPT|$, where, as usual, OPT denotes an optimal tour. First we argue about the approximation ratio, and leave for later the presentation of the algorithm. To start, we assume for simplicity that no two lines are parallel, though this assumption will be later removed.

Observation 1 *The optimal tour of a set of lines is a (possibly degenerate) convex polygon P .*

Proof. It is easy to see that an optimal tour must be polygonal, consisting of a finite union of straight line segments. If an optimal tour were a non-convex polygon P , we obtain a contradiction to its optimality since the boundary of the convex hull of P , which is shorter than P , also visits all of the lines L , since P does. \square

Observation 2 C_L is determined by 3 lines in L , i.e. it is the inscribed circle in the triangle Δ formed by these 3 lines.

We distinguish two cases:

Case 1. Δ is an acute triangle. It is well known that for an acute triangle, the minimum perimeter inscribed triangle (having a vertex on each side of the triangle) is its *pedal* triangle, whose vertices are the feet of the altitudes of the given triangle (see e.g. [22]). So in this case, the optimal tour OPT_Δ , which visits the 3 lines of Δ is its pedal triangle; we denote by y its perimeter. Clearly y is a lower bound on $|OPT|$: $|OPT| \geq |OPT_\Delta| = y$. Denote by s the semi-perimeter of Δ , by R the radius of its circumscribed circle, and by r the radius of its inscribed circle.

Fact 1 For an acute triangle Δ , $s > 2R$.

Proof. If A, B, C are the angles of Δ , this is equivalent to

$$R(\sin A + \sin B + \sin C) > 2R.$$

After simplification with R , this is a well known inequality in the geometry of an acute triangle ([6], page 18). \square

Fact 2 For an acute triangle Δ , $y = \frac{2rs}{R}$.

A proof of this equality can be found in [6], page 86.

Claim 1 For an acute triangle Δ , $r < \frac{y}{4}$.

Proof. Putting the above together, we get $r = \frac{Ry}{2s} < \frac{Ry}{4R} = \frac{y}{4}$. \square

The length of the output tour is bounded as follows

$$|C_L| = 2\pi r < 2\pi \frac{y}{4} = \frac{\pi}{2} y \leq \frac{\pi}{2} |OPT|.$$

Case 2. Δ is an obtuse triangle. In this case, $|OPT_\Delta| = 2h$, where h is the length of the altitude corresponding to the obtuse angle, say A . Clearly $2h$ is a lower bound on $|OPT|$: $|OPT| \geq |OPT_\Delta| = 2h$.

Fact 3 For any triangle Δ , $h > 2r$.

Proof. Let S denote the area of the triangle with side lengths a , b , and c . Then, we know that $S = ah/2$, by our definition of the altitude h . We also know from elementary geometry that $S = (a + b + c)r/2$, recalling that r is the radius of the inscribed circle of the triangle. Thus, $ah = (a + b + c)r$, from which we obtain

$$\frac{h}{r} = \frac{a + b + c}{a} > \frac{a + a}{a} = 2,$$

using the triangle inequality. □

Using this inequality, we have $|OPT_\Delta| = 2h > 4r$. The length of the output tour is bounded as follows

$$|C_L| = 2\pi r = \frac{\pi}{2}(4r) < \frac{\pi}{2}|OPT_\Delta| \leq \frac{\pi}{2}|OPT|.$$

Thus, in both cases, the approximation ratio is $\frac{\pi}{2} \leq 1.58$.

When C_L is determined by 3 lines in L of which two are parallel at distance h , we say that they form a generalized triangle Δ . Clearly $2h$ is a lower bound on $|OPT|$: $|OPT| \geq |OPT_\Delta| = 2h$. We also have $r = h/2$, where r is the radius of C_L , thus

$$|C_L| = 2\pi r = \pi h \leq \frac{\pi}{2}|OPT|.$$

We now describe the algorithm for computing C_L . The distance $d(p, l)$ from a point p of coordinates (x_0, y_0) to a line l of equation $ax + by + c = 0$ is

$$d(p, l) = \frac{|ax_0 + by_0 + c|}{\sqrt{a^2 + b^2}}.$$

Let the lines in L have the equations

$$(l_i :) \quad a_i x + b_i y + c_i = 0, \quad i = 1, \dots, n.$$

Finding a minimum touching circle amounts to finding the center coordinates (x, y) , and a minimum radius z , s.t.

$$\frac{|a_i x + b_i y + c_i|}{\sqrt{a_i^2 + b_i^2}} \leq z, \quad i = 1, \dots, n.$$

This is equivalent to solving the following 3-dimensional linear program

$$\min z \quad \text{subject to} \quad \left\{ \frac{a_i x + b_i y + c_i}{\sqrt{a_i^2 + b_i^2}} \leq z, \quad \frac{-a_i x - b_i y - c_i}{\sqrt{a_i^2 + b_i^2}} \leq z, \quad i = 1, \dots, n \right\},$$

which takes $O(n)$ time [16]. Consequently, we have proved

Theorem 3 *Given a set L of n (infinite straight) lines in the plane, a $\frac{\pi}{2}$ -approximate tour that visits L can be computed in $O(n)$ time.*

6 Conclusion

Several open problems remain, including

- (1) Is there a constant-factor approximation algorithm for arbitrary connected regions in the plane? What if the regions are disconnected? (giving us a geometric version of a “one-of-a-set TSP”)
- (2) What approximation bounds can be obtained in higher dimensions? Our packing arguments for disjoint disks lift to higher dimensions, but our other methods do not readily generalize. A particularly intriguing special case is the generalization of the case of infinite straight lines: What can be said in 3-space for the TSPN on a set of lines or of planes?
- (3) Is there a PTAS for general pairwise-disjoint regions in the plane?

Acknowledgements

We thank Estie Arkin and Michael Bender for several useful discussions on the TSPN problem. We thank the anonymous referees for their detailed comments and suggestions, which greatly improved the paper.

References

- [1] E. M. Arkin and R. Hassin, Approximation algorithms for the geometric covering salesman problem, *Discrete Appl. Math.*, **55**:197–218, 1994.
- [2] S. Arora, Nearly linear time approximation schemes for Euclidean TSP and other geometric problems, *J. of the ACM*, **45**(5):1–30, 1998.
- [3] T. Asano, S. K. Ghosh, and T. C. Shermer, Visibility in the plane, In *Handbook of Computational Geometry* (J.-R. Sack and J. Urrutia, editors), pages 829–876. Elsevier Science Publishers B.V. North-Holland, Amsterdam, 2000.
- [4] J. L. Bentley, Fast algorithms for geometric traveling salesman problems, *ORSA J. on Comput.*, **4**(4):387–411, 1992.
- [5] M. de Berg, J. Gudmundsson, M. J. Katz, C. Levcopoulos, M. H. Overmars, and A. F. van der Stappen, TSP with neighborhoods of varying size, *Proc. 10th Annual European Symposium on Algorithms*, to appear, September, 2002.
- [6] O. Bottema, R. Ž. Djordjević, R. Janić, D. S. Mitrinović and P. M. Vasić, *Geometric Inequalities*, Wolters-Noordhoff, Groningen, 1969.
- [7] P. Berman and M. Karpinski, On some tighter inapproximability results, Technical Report TR98-065, ECCO, 1998.
- [8] S. Carlsson, H. Jonsson and B. J. Nilsson, Finding the shortest watchman route in a simple polygon, *Discrete Comput. Geom.* **22**(3):377–402, 1999.
- [9] M. R. Garey and D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, W. H. Freeman, New York, NY, 1979.

- [10] J. Gudmundsson and C. Levcopoulos, A fast approximation algorithm for TSP with neighborhoods, *Nordic J. of Comput.*, **6**:469–488, 1999.
- [11] J. Gudmundsson and C. Levcopoulos, Hardness result for TSP with neighborhoods, Technical Report LU-CS-TR:2000-216, Department of Computer Science, Lund University, Sweden, 2000.
- [12] H. Jonsson, The traveling salesman problem for lines in the plane, *Inform. Process. Lett.*, **82**(3):137–142, 2002.
- [13] M. Jünger, G. Reinelt and G. Rinaldi, The traveling salesman problem, In *Network Models, Handbook of Operations Research/Management Science* (M. O. Ball, T. L. Magnanti, C. L. Monma, and G. L. Nemhauser, editors), pages 225–330, Elsevier Science, Amsterdam, 1995.
- [14] E. L. Lawler, J. K. Lenstra, A. H. G. Rinnooy Kan and D. B. Shmoys, editors, *The Traveling Salesman Problem*. John Wiley & Sons, New York, NY, 1985.
- [15] C. Mata and J. S. B. Mitchell, Approximation algorithms for geometric tour and network design problems, *Proc. 11th Annu. ACM Sympos. Comput. Geom.*, pages 360–369, 1995.
- [16] N. Megiddo, Linear programming in linear time when the dimension is fixed, *J. of ACM*, **31**:114–127, 1984.
- [17] J. S. B. Mitchell, Guillotine subdivisions approximate polygonal subdivisions: A simple polynomial-time approximation scheme for geometric TSP, k -MST, and related problems, *SIAM J. on Comput.*, **28**(4):1298–1309, 1999.
- [18] J. S. B. Mitchell. Guillotine subdivisions approximate polygonal subdivisions: Part III – Faster polynomial-time approximation schemes for geometric network optimization. Manuscript, University at Stony Brook, 1997.
- [19] J. S. B. Mitchell. Approximation algorithms for geometric optimization problems. In *Proc. Ninth Canadian Conference on Computational Geometry*, Queen’s University, Kingston, Canada, August 11-14, 1997, pp. 229–232.
- [20] J. S. B. Mitchell, Geometric shortest paths and network optimization, In *Handbook of Computational Geometry* (J.-R. Sack and J. Urrutia, editors), pages 633–701. Elsevier Science Publishers B.V. North-Holland, Amsterdam, 2000.
- [21] C. H. Papadimitriou, The Euclidean traveling salesman problem is NP-complete, *Theoret. Comput. Sci.*, **4**:237–244, 1977.
- [22] H. Rademacher and O. Toeplitz, *The Enjoyment of Mathematics*, Princeton University Press, 1957; a translation from *Von Zahlen und Figuren*, Springer, Berlin, 1933.
- [23] S. B. Rao and W. D. Smith, Approximating geometrical graphs via “spanners” and “banyans”, *Proc. 30th Annu. ACM Sympos. Theory Comput.*, 540–550, 1998.
- [24] G. Reinelt, Fast heuristics for large geometric traveling salesman problems, *ORSA J. Comput.*, **4**:206–217, 1992.

- [25] O. Schwartz and S. Safra, On the complexity of approximating TSP with neighborhoods and related problems, Manuscript (submitted), July, 2002.
- [26] X. Tan, Fast computation of shortest watchman routes in simple polygons, *Inform. Process. Lett.*, **77**(1):27–33, 2001.