

APPROXIMATION OF JENSEN MEASURES BY IMAGE MEASURES UNDER HOLOMORPHIC FUNCTIONS AND APPLICATIONS

SHANGQUAN BU AND WALTER SCHACHERMAYER

ABSTRACT. We show that Jensen measures defined on \mathbb{C}^n or more generally on a complex Banach space X can be approximated by the image of Lebesgue measure on the torus under X -valued polynomials defined on \mathbb{C} . We give similar characterizations for Jensen measures in terms of analytic martingales and Hardy martingales. The results are applied to approximate plurisubharmonic martingales by Hardy martingales, which enables us to give a characterization of the analytic Radon-Nikodym property of Banach spaces in terms of convergence of plurisubharmonic martingales, thus solving a problem of G. A. Edgar.

0. INTRODUCTION

Let X be a Banach space and μ a Radon probability measure on X with first moment (i.e., $\int_X \|x\| d\mu(x) < \infty$). It is well known that there is a unique $x_0 \in X$, called the *barycenter* of μ verifying

$$f(x_0) \leq \int_X f(x) d\mu(x),$$

for every real-valued convex Lipschitz function on X . We then call μ a *Choquet measure* for x_0 .

Recall the following easy folklore result (compare [E3] for further results in this context; unexplained notation will be defined below):

Theorem (0). *Let μ be a Radon probability measure with first moment on a Banach space X and $x_0 \in X$. The following are equivalent:*

- (i) μ is a Choquet measure with barycenter x_0 .
- (ii) There is a Bochner integrable function $f : [0, 1] \rightarrow X$ with expectation $\mathbb{E}(f) = \int_0^1 f(x) d\lambda(x) = x_0$ and such that the image measure $f(\lambda)$ equals μ .
- (iii) μ can be approximated in the narrow topology with respect to the class of Lipschitz functions on X by the final distribution $D_n(\mathbb{P})$ of a finite dyadic martingale $(D_i)_{i=0}^n$ defined on a probability space $(\Omega, \Sigma, \mathbb{P})$ and starting at x_0 , i.e., $D_0 \equiv x_0$.
- (iv) μ equals the final distribution $M_n(\mathbb{P})$ of a finite martingale $(M_i)_{i=0}^n$ defined on a probability space $(\Omega, \Sigma, \mathbb{P})$ and starting at x_0 , i.e., $M_0 \equiv x_0$.

Received by the editors January 11, 1989 and, in revised form, February 7, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46B20, 32F05, 60G46.

Key words and phrases. Plurisubharmonic martingales, Hardy martingales, Jensen measures, the analytic Radon-Nikodym property.

We have stated the rather easy Theorem (0) as it is the “real” analogue of the “complex” Theorem (A) below, where the term Choquet measure is replaced by Jensen measure, integrable function by holomorphic function, dyadic martingale by analytic martingale and the term martingale in (iv) by Hardy martingale (unexplained notions will be defined in the subsequent section):

Theorem (A). *Let μ be a Radon probability measure with first moment on a complex Banach space X and $x_0 \in X$. Equip $\mathcal{M}^1(X)$ with the weak topology induced by $\text{Lip}(X)$. The following are equivalent:*

(i) μ is a Jensen measure on X with barycenter x_0 .

(ii) μ is in the closure of the subset

$$\mathbf{P} = \{g(\lambda) : g : \mathbb{C} \rightarrow X \text{ is a polynomial with } g(0) = x_0\}$$

of $\mathcal{M}^1(X)$.

(iii) μ is in the closure of the subset

$$\mathbf{A} = \{F_n(\lambda^n) : (F_i)_{i=0}^n \text{ is an analytic martingale with } F_0 \equiv x_0\}$$

of $\mathcal{M}^1(X)$.

(iv) μ is in the closure of the subset

$$\mathbf{H} = \{F_n(\lambda^n) : (F_i)_{i=0}^n \text{ is a Hardy martingale with } F_0 \equiv x_0\}$$

of $\mathcal{M}^1(X)$.

Remark. The implication (ii) \Rightarrow (i), (iii) \Rightarrow (i) and (iv) \Rightarrow (i) are easily seen to be true. The equivalence of (ii) and (iii) has essentially been proved by G. A. Edgar [E2] while the equivalence of (ii) and (iv) follows from a theorem of N. Ghoussoub and B. Maurey [G-M, Theorem 4.1].

The decisive new information given by Theorem (A) is how to approximate an abstract Jensen measure by *analytic objects* as in (ii), (iii) or (iv). Note that the only *analytic concept* appearing in the definition of Jensen measures is that of plurisubharmonic functions, against which μ is tested via inequality (1) below.

We are afraid that the reader interested in several complex variables might be turned off by the infinite-dimensional setting and the concepts of analytic and Hardy martingales. We therefore formulate another version of Theorem (A), which is more in the spirit of several complex variables and we give a proof which does not rely on the concepts of analytic and Hardy martingales. However, the basic idea of the proof is the same as in Theorem (A).

Theorem (B). *Let U be a domain in \mathbb{C}^n and μ a probability measure with compact support in U . Then μ is a Jensen measure on U with barycenter $x_0 \in U$ if and only if μ can be approximated by image measure $g(\lambda)$, where g is polynomials $g : \mathbb{C} \rightarrow \mathbb{C}^n$, $g(\overline{\mathbb{D}}) \subseteq U$, $g(0) = x_0$ in the following sense: For every finite set $\{f_1, f_2, \dots, f_m\}$ of continuous functions on U and $\varepsilon > 0$ there is g as above such that for $1 \leq i \leq m$,*

$$|\langle f_i, \mu \rangle - \langle f_i, g(\lambda) \rangle| = \left| \int_U f_i(x) d\mu(x) - \int_0^1 f_i \circ g(e^{2\pi i\theta}) d\theta \right| < \varepsilon.$$

Let us come back again to the context of Banach spaces: it was proved by G. A. Edgar [E2] that convergence of X -valued L^1 -bounded analytic martingales characterizes the analytic Radon-Nikodym property of X introduced by

A. Bukhvalov and A. Danilevich [Bu-Da] and the corresponding theorem for Hardy martingales has been observed by D. J. H. Garling [Gar, Theorem 3].

Note that the definition of analytic martingales as well as Hardy martingales refers to a special representation of these martingales, namely that they are defined on the probability space $(\mathbb{T}^{\mathbb{N}}, \lambda^{\mathbb{N}})$. This is somehow unsatisfactory from a probabilistic point of view. G. A. Edgar has introduced the representation free concept of plurisubharmonic martingales (Definition I.8. below).

One easily verifies that a Hardy martingale (and therefore an analytic martingale) is a plurisubharmonic martingale [Gar, Theorem 1]. Theorem (C), which is the second main result of this paper, gives a kind of converse.

Theorem (C). *Let X be a complex Banach space, $(M_n)_{n=0}^{\infty}$ an X -valued plurisubharmonic martingale and $(\varepsilon_n)_{n=1}^{\infty}$ positive numbers. Then there is a representation $(F_n)_{n=0}^{\infty}$ of $(M_n)_{n=0}^{\infty}$ defined on $(\mathbb{T}^{\mathbb{N}}, \lambda^{\mathbb{N}})$ such that F_n depends only on the first n coordinates of $\mathbb{T}^{\mathbb{N}}$ (and may therefore be identified with a function on \mathbb{T}^n) and a Hardy martingale $(G_n)_{n=0}^{\infty}$ such that, for every $n \in \mathbb{N}$,*

$$\|(F_n - F_{n-1}) - (G_n - G_{n-1})\|_{L^1(\mathbb{T}^n, X)} < \varepsilon_n.$$

The solution to the problem of G. A. Edgar [E2] now follows immediately from Theorem (C). This result has also been proved by N. Ghoussoub and B. Maurey [G-M] by different methods.

Corollary (D). *A Banach space X has the analytic Radon-Nikodym property if and only if L^1 -bounded X -valued plurisubharmonic martingales converge almost surely.*

We now describe the organization of this paper.

In §1 we gather the necessary definitions and notations. In §2 we prove Theorem (A). The proof turns out to be surprisingly simple and uses the Hahn-Banach theorem in a crucial way.

In §3 we prove Theorem (B), which is formulated in the local setting (i.e., for domains in \mathbb{C}^n) and we therefore also prove some technical results.

In §4 we prepare the tools needed for Theorem (C): A more precise and parametrised version of Theorem (A) is proved (Proposition IV.2) and we have to use some techniques from measure theory (disintegration of measures, measurable selections).

In §5 we then prove Theorem (C) and Corollary (D). We also note an application of Theorem (C) to Analytic Martingale Transform spaces introduced by D. J. H. Garling [Gar], extending a result of Xu [X].

For unexplained notation we refer to [L-T] for the Banach space concepts and to [Ra] or [K] for the concepts of several complex variables.

Acknowledgement. We thank B. Maurey for substantial help in establishing a proof of Corollary (D) and we also thank N. Ghoussoub, P. Müller and E. Perkins for their advice in the preparation of this paper.

I. DEFINITIONS AND NOTATIONS

Throughout this paper, \mathbb{T} will denote the torus $\{e^{2\pi i\theta} : 0 \leq \theta < 1\}$ which we shall freely identify with a subset of \mathbb{C} or with $[0, 1[$. The open (resp. closed) disc of \mathbb{C} will be denoted by \mathbb{D} (resp. $\overline{\mathbb{D}}$). Normalized Lebesgue measure on \mathbb{T} as well as on $[0, 1]$ will always be denoted by λ .

X will denote a complex Banach space; we shall consider Radon probability measures on X (see e.g., [Schw2]). As every Radon probability measure μ is supported by a separable subspace of X we shall assume throughout the paper without loss of generality that X is separable, hence the set of Radon probability measures on X coincides with the set of all probability measures defined on the Borel σ -field generated by the metric topology of X [Schw2]. If μ is a measure with first moment on X (i.e., $\int \|x\| d\mu(x) < \infty$) and $\phi : X \rightarrow \mathbb{R}$ is a Lipschitz function, then we may form

$$\langle \phi, \mu \rangle = \int_X \phi(x) d\mu(x).$$

If U is a domain in \mathbb{C}^n , μ is a Radon measure on U and $\phi : U \rightarrow \mathbb{R}$ is a measurable function, we shall also denote the scalar product as above if the right term makes sense.

If U is a domain in \mathbb{C} , a function $g : U \rightarrow X$ is called analytic (or holomorphic) if, for every $x^* \in X^*$, $x^* \circ g$ is analytic. A function $g : \mathbb{C} \rightarrow X$ of the form $g(z) = \sum_{n=0}^N x_n z^n$ with $x_n \in X$ and $N \in \mathbb{N}$ will be called an X -valued polynomial on \mathbb{C} . Note that, if U contains $\overline{\mathbb{D}}$ and $g : U \rightarrow X$ is analytic then we can approximate g by X -valued polynomials uniformly on $\overline{\mathbb{D}}$ (see e.g., [Ch]).

We shall denote for $1 \leq p \leq \infty$ by $L^p(\mathbb{T}, X)$ the space of Bochner integrable functions $f : \mathbb{T} \rightarrow X$ equipped with the norm

$$\|f\|_p = \left(\int_0^1 \|f(e^{2\pi i\theta})\|^p d\theta \right)^{1/p},$$

for $1 \leq p < \infty$ and for $p = \infty$

$$\|f\|_\infty = \operatorname{ess\,sup}_{t \in \mathbb{T}} \|f(t)\|,$$

and by $H_0^p(\mathbb{T}, X)$ the subspace of $L^p(\mathbb{T}, X)$ formed by the elements g verifying, for every $k \geq 0$,

$$\int_0^1 e^{2k\pi i\theta} f(e^{2\pi i\theta}) d\theta = 0.$$

We shall identify elements $f \in H_0^p(\mathbb{T}, X)$ with functions on $\overline{\mathbb{D}}$, i.e., the analytic extension of f to $\overline{\mathbb{D}}$ obtained via the Poisson kernel.

Denote by $\mathcal{M}^1(X)$ the space of finite measures on X with first moment, i.e., $\int_X \|x\| d|\mu|(x) < \infty$, and by $\operatorname{Lip}(X)$ the space of Lipschitz functions on X . The scalar product $\langle \cdot, \cdot \rangle$ defined above places these spaces in duality and we shall equip $\mathcal{M}^1(X)$ with the weak topology induced by $\operatorname{Lip}(X)$.

If $(\Omega, \Sigma, \mathbb{P})$ is a probability space, (Ω', Σ') a measure space and $F : \Omega \rightarrow \Omega'$ a measurable map, we denote by $F(\mathbb{P})$ the image measure of \mathbb{P} under F which is defined, for $A \in \Sigma'$, by

$$F(\mathbb{P})(A) = \mathbb{P}(F^{-1}(A)).$$

Definition I.1. If $(\mu_\alpha)_{\alpha \in I}$ is a net of probability measures on a polish space (E, d) , we shall say that $(\mu_\alpha)_{\alpha \in I}$ converges narrowly or in the narrow topology

to a probability measure μ if, for every bounded continuous function $f : E \rightarrow \mathbb{R}$,

$$\lim_{\alpha} \langle f, \mu_{\alpha} \rangle = \langle f, \mu \rangle.$$

Definition I.2 [E1]. Let U be a domain in X . A function $\phi : U \rightarrow \mathbb{R} \cup \{-\infty\}$ is called *plurisubharmonic on U* if ϕ is upper semicontinuous and if for every $x, y \in X$ such that $\{x + \mathbb{D}y\} \subseteq U$

$$\phi(x) \leq \int_0^1 \phi(x + e^{2\pi i\theta}y) d\theta.$$

Definition I.3 (compare [E1]). Let X be a complex Banach space and μ a probability measure on X with first moment. We say that μ is a *Jensen measure on X with barycenter $x_0 \in X$* if, for every Lipschitz plurisubharmonic function, ϕ on X

$$(1) \quad \phi(x_0) \leq \int_X \phi(y) d\mu(y).$$

Remark I.4. First note that there are more plurisubharmonic functions than convex Lipschitz functions on X and therefore fewer Jensen measures than Choquet measures on X . For a general account on Jensen measures we refer to [Gam].

Classically Jensen measures are supposed to have compact support. In view of the application to L^1 -bounded martingales (Theorem (C) below) we place ourselves into the more general context of measures with first moment; hence we have to restrict ourselves to require inequality (1) only for Lipschitz plurisubharmonic functions and not arbitrary plurisubharmonic functions to avoid integrability problems. However in the context of measures with compact support on a domain U of \mathbb{C}^n it will be more natural to adopt the subsequent concept:

Definition I.5. Let U be a domain in \mathbb{C}^n and μ a probability measure with compact support K in U . We say that μ is a *Jensen measure on U with barycenter $x_0 \in U$* if, for every plurisubharmonic function $\varphi : U \rightarrow \mathbb{R} \cup \{-\infty\}$ we have

$$\varphi(x_0) \leq \int_U \varphi(x) d\mu(x).$$

Remark I.6. Note that by the upper semicontinuity of φ the integral on the right-hand side is well defined (with values in $\mathbb{R} \cup \{-\infty\}$). The definition is more in the classical spirit of Jensen measures and does not refer to Lipschitz functions as Definition I.3 above. We shall show in Proposition III.4 below that these two definitions are consistent.

Definition I.7. For a Banach space X , a sequence of functions $(F_n)_{n=0}^{\infty}$, $F_0 \equiv x_0$, $F_n \in L^1(\mathbb{T}^n, \lambda^n, X)$ (where \mathbb{T}^0 is a one point space) is called an *X -valued*

(a) *analytic martingale* ([Bo-Da], see also [D-G-T] and [E1]) if, for $n \in \mathbb{N}$ and $(\theta_1, \theta_2, \dots, \theta_{n-1}) \in \mathbb{T}^{n-1}$

$$F_n(\theta_1, \theta_2, \dots, \theta_n) - F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1}) = f_n(\theta_1, \theta_2, \dots, \theta_{n-1})e^{2\pi i\theta_n};$$

(b) *Hardy martingale* ([Gar], see also [G-M]) if, for every $n \in \mathbb{N}$ and $(\theta_1, \theta_2, \dots, \theta_{n-1}) \in \mathbb{T}^{n-1}$, the function

$$\theta_n \rightarrow d_n(\theta_1, \theta_2, \dots, \theta_{n-1}, \theta_n),$$

defined by

$$\begin{aligned} & d_n(\theta_1, \theta_2, \dots, \theta_{n-1}, \theta_n) \\ &= F_n(\theta_1, \theta_2, \dots, \theta_{n-1}, \theta_n) - F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1}), \end{aligned}$$

is in $H_0^1(\mathbb{T}, X)$.

If U is a domain in X we call $(F_n)_{n=1}^\infty$ a U -valued analytic (resp. Hardy) martingale if in addition to the above requirements, for every $n \in \mathbb{N}$ and $(\theta_1, \theta_2, \dots, \theta_{n-1}) \in \mathbb{T}^{n-1}$,

$$\begin{aligned} & F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1}) + f_n(\theta_1, \theta_2, \dots, \theta_{n-1})re^{2\pi i\theta_n} \in U, \\ & \text{(resp. } F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1}) + d_n(\theta_1, \theta_2, \dots, \theta_{n-1}, re^{2\pi i\theta_n}) \in U), \end{aligned}$$

for every $0 \leq r \leq 1$ and $\theta_n \in \mathbb{T}$.

Obviously analytic martingales are Hardy martingales. The term *martingale* is justified as one may identify $(F_n)_{n=0}^\infty$ in an obvious way with a stochastic process on $(\mathbb{T}^\mathbb{N}, \lambda^\mathbb{N})$ equipped with its natural filtration $(\Sigma_n)_{n=0}^\infty$ which is readily verified to be a martingale (see [Gar]).

Definition 1.8 (compare [E1]). An X -valued martingale $(M_n)_{n=0}^\infty$, defined on a probability space $(\Omega, \Sigma, \mathbb{P})$ and such that $M_0 \equiv x_0$ is called a *plurisubharmonic martingale* if, for every Lipschitz plurisubharmonic function ϕ on X , the stochastic process $(\phi \circ M_n)_{n=0}^\infty$ is a submartingale.

One can easily observe that Hardy martingales (whence, in particular, analytic martingales) are plurisubharmonic martingales.

To end this section let us point out the easy implications among the above concepts: Let $g \in H_0^1(\mathbb{T}, X)$ and identify g with a function g on \mathbb{D} which is analytic in \mathbb{D} . Then the image measure $g(\lambda)$ is a Jensen measure on X with barycenter 0. Indeed, if $\phi : X \rightarrow \mathbb{R}$ is a Lipschitz plurisubharmonic function then $\phi \circ g$ is subharmonic on \mathbb{D} and the radial limits exist almost surely; therefore

$$\phi(0) = \phi \circ g(0) \leq \int_0^1 \phi \circ g(e^{2\pi i\theta}) d\theta = \int_X \phi(x) d(g(\lambda))(x).$$

Similarly one verifies that for a finite Hardy martingale on X (whence, in particular, for a finite analytic martingale) $(F_i)_{i=0}^n$, with $F_0 \equiv x_0$, the image measure $F_n(\lambda^n)$ is a Jensen measure on X with barycenter x_0 (compare [Gar, Theorem 1]).

This remark takes care of the easy implications of Theorem (A) above. In the next section we shall prove that the reverse implications also hold true.

II. THE PROOF OF THEOREM (A)

Recall the characterization of the plurisubharmonic hull of a function which has been proved by G. A. Edgar [E1, Lemma 2.1]. We give a version formulated for domains (compare also Proposition III.4 below) and use a slightly weaker hypothesis than in [E1]:

Proposition II.1. *Let $U \subseteq X$ be a domain and $f : U \rightarrow \mathbb{R} \cup \{-\infty\}$ an upper semicontinuous function. Define $f_0 = f$ and for $n \geq 1$*

$$f_n(x) = \inf \left\{ \int_0^1 f_{n-1}(x + e^{2\pi i\theta} y) d\theta \right\}$$

there the inf is taken over all $y \in X$ such that $\{x + \overline{\mathbb{D}}y\} \subseteq U$. Then $(f_n)_{n=0}^\infty$ decreases pointwise to the largest plurisubharmonic function \hat{f} on U dominated by f .

Proof. It is obvious that $(f_n)_{n=0}^\infty$ decreases. We verify inductively that f_n is upper semicontinuous: $f_0 = f$ is upper semicontinuous. Suppose f_{n-1} is upper semicontinuous and let $(x_k)_{k=0}^\infty$ in U be such that $\lim_{k \rightarrow \infty} x_k = x_0$. If $y_0 \in X$ is such that $\{x_0 + \overline{\mathbb{D}}y_0\} \subseteq U$ then there is k_0 such that $\{x_k + \overline{\mathbb{D}}y_0\} \subseteq U$ for $k \geq k_0$. The upper semicontinuous function f_{n-1} is bounded above on the relatively compact set $\bigcup_{k=k_0}^\infty \{x_k + \overline{\mathbb{D}}y_0\}$ and, for every $z \in \overline{\mathbb{D}}$,

$$f_{n-1}(x_0 + zy_0) \geq \limsup_{k \rightarrow \infty} f_{n-1}(x_k + zy_0).$$

Hence we obtain from Fatou's lemma that, for every $y_0 \in X$ verifying $\{x_0 + \overline{\mathbb{D}}y_0\} \subseteq U$,

$$\begin{aligned} \int_0^1 f_{n-1}(x_0 + e^{2\pi i\theta} y_0) d\theta &\geq \limsup_{k \rightarrow \infty} \int_0^1 f_{n-1}(x_k + e^{2\pi i\theta} y_0) d\theta \\ &\geq \limsup_{k \rightarrow \infty} f_n(x_k), \end{aligned}$$

and therefore

$$f_n(x_0) \geq \limsup_{k \rightarrow \infty} f_n(x_k).$$

This shows that each f_n and therefore \hat{f} is upper semicontinuous.

For every plurisubharmonic function ϕ on U , $\phi \leq f$, we have $\phi \leq f_n$ for every $n \in \mathbb{N}$. Indeed, clearly $\phi \leq f_0$ and suppose that $\phi \leq f_{n-1}$. Then for every $x_0 \in U$ and $y_0 \in X$ such that $\{x_0 + \overline{\mathbb{D}}y_0\} \subseteq U$

$$\int_0^1 f_{n-1}(x_0 + e^{2\pi i\theta} y_0) d\theta \geq \int_0^1 \phi(x_0 + e^{2\pi i\theta} y_0) d\theta \geq \phi(x_0),$$

whence $f_n(x_0) \geq \phi(x_0)$, which gives the inductive step. Hence we conclude that $\hat{f} \geq \phi$ for every plurisubharmonic function ϕ on U dominated by f .

Finally we have to show the mean value inequality for \hat{f} which follows from the Beppo Levi's monotone convergence theorem: for $x_0 \in U$, $y_0 \in X$, with $\{x_0 + \overline{\mathbb{D}}y_0\} \subseteq U$

$$\begin{aligned} \hat{f}(x_0) &= \lim_{n \rightarrow \infty} f_n(x_0) \leq \lim_{n \rightarrow \infty} \int_0^1 f_{n-1}(x_0 + e^{2\pi i\theta} y_0) d\theta \\ &= \int_0^1 \hat{f}(x_0 + e^{2\pi i\theta} y_0) d\theta. \end{aligned}$$

The proof is complete. \square

Remark II.2. As noted by G. A. Edgar [E1] one also may write the definition of f_n in the following way:

$$f_n(x) = \inf\{\mathbb{E}(f(F_n)) : (F_i)_{i=0}^n \text{ is a } U\text{-valued analytic martingale with } F_0 \equiv x\}.$$

Proof of Theorem (A).

(i) \Rightarrow (iii) We first show that the set

$$\mathbf{A} = \{F_n(\lambda^n) : (F_i)_{i=0}^n \text{ is an } X\text{-valued analytic martingale with } F_0 \equiv x_0\},$$

is a convex subset of $\mathcal{M}^1(X)$: let $(F'_i)_{i=0}^n$ and $(F''_i)_{i=0}^m$ be two analytic martingales as above. We may assume $n = m$. Define now an analytic martingale $(F_i)_{i=0}^{n+1}$ by letting $F_0 \equiv F_1 \equiv x_0$ and for $1 \leq i \leq n$,

$$F_{i+1}(\theta_1, \theta_2, \dots, \theta_{i+1}) = \begin{cases} F'_i(\theta_2, \dots, \theta_{i+1}) & \text{if } 0 \leq \theta_1 < \frac{1}{2}, \\ F''_i(\theta_2, \dots, \theta_{i+1}) & \text{if } \frac{1}{2} \leq \theta_1 < 1. \end{cases}$$

Clearly

$$F_{n+1}(\lambda^{n+1}) = \{F'_n(\lambda^n) + F''_n(\lambda^n)\}/2,$$

thus showing the convexity of \mathbf{A} .

If the conclusion of (iii) were false then, by the Hahn-Banach theorem we could find a Lipschitz function f on X and reals $\alpha < \beta$ such that

$$\langle f, \mu \rangle = \int_X f d\mu \leq \alpha,$$

while

$$\langle f, \nu \rangle = \int_X f d\nu \geq \beta,$$

for every $\nu \in \mathbf{A}$. Note that the last line can be rewritten as

$$\int_{\mathbb{T}^n} f \circ F_n d\lambda^n \geq \beta,$$

for every X -valued analytic martingale starting at x_0 , hence by the preceding remark

$$\hat{f}(x_0) \geq \beta.$$

This gives the desired contradiction as we obtain the absurd inequality

$$\alpha \geq \langle f, \mu \rangle \geq \langle \hat{f}, \mu \rangle \geq \hat{f}(x_0) \geq \beta.$$

Note that \hat{f} is Lipschitz by a remark of N. Ghoussoub and B. Maurey (see [G-M]), so no integrability problems arise.

(iii) \Rightarrow (iv) Obvious.

(iv) \Rightarrow (ii) Let $(F_i)_{i=0}^n$ be an X -valued Hardy martingale, $F_0 \equiv x_0$. By [G-M, Theorem 6.1] we may find for $\varepsilon > 0$, a function $g^\varepsilon: V_\varepsilon \rightarrow X$, which is analytic in a neighbourhood V_ε of \mathbb{D} , $g^\varepsilon(0) = x_0$ and a continuous surjection $\pi: \mathbb{T} \rightarrow \mathbb{T}^n$ such that $\pi(\lambda) = \lambda^n$ and

$$\int_0^1 \|F_n(\pi(e^{2\pi i\theta})) - g^\varepsilon(e^{2\pi i\theta})\| d\theta < \varepsilon/2.$$

Find an X -valued polynomial p^ε such that, for $|z| \leq 1$,

$$\|p^\varepsilon(z) - g^\varepsilon(z)\| < \varepsilon/2.$$

If $f : X \rightarrow \mathbb{R}$ is a Lipschitz function with Lipschitz constant M , then

$$\begin{aligned} |\langle f, F_n(\lambda^n) \rangle - \langle f, p^\varepsilon(\lambda) \rangle| &= \left| \int_0^1 (f \circ F_n(\pi(e^{2\pi i\theta})) - f \circ p^\varepsilon(e^{2\pi i\theta})) d\theta \right| \\ &\leq M \int_0^1 \|F_n(\pi(e^{2\pi i\theta})) - p^\varepsilon(e^{2\pi i\theta})\| d\theta < M\varepsilon, \end{aligned}$$

which readily shows that $F_n(\lambda^n)$ is in the closure of the subset \mathbf{P} in (ii).

(ii) \Rightarrow (i) This implication follows from the argument in the first section and the observation that, for a net $(\mu_\alpha)_{\alpha \in I}$ of Jensen measures on X with barycenter x_0 converging in $\mathcal{M}^1(X)$, the limit μ again is a Jensen measure on X with barycenter x_0 . \square

III. THE PROOF OF THEOREM (B)

We now turn to the setting of Theorem (B) for which we shall give a self-contained proof. We start with an easy but crucial example.

Example III.1. Consider $X = \mathbb{C}$ and denote by λ the Lebesgue measure on the torus $\mathbb{T} \subseteq \mathbb{C}$ and by δ_0 the Dirac measure at 0. The measure $\mu = (\lambda + \delta_0)/2$ clearly is a Jensen measure on \mathbb{C} with barycenter 0.

However there is no polynomial $p : \mathbb{C} \rightarrow \mathbb{C}$ such that $p(\lambda) = \mu$. Indeed such a polynomial would have to equal zero on a subset of \mathbb{T} of measure 1/2 and therefore have to be identically zero, a contradiction. In fact, it is well known that any not identically zero function f in $H^\infty(\mathbb{T})$ equals zero at most on a subset of \mathbb{T} of measure 0 (compare the proof of Lemma III.2 below), hence μ is not the image measure $f(\lambda)$ of any f in $H^\infty(\mathbb{T})$ either.

This shows that the set \mathbf{P} appearing in Theorem (A) fails to be convex (contrary to the set \mathbf{A}). Note however that it follows from Theorem (A) that the closure of \mathbf{P} equals the set of Jensen measures on X and therefore is convex.

We shall need the following result related to outer functions:

Lemma III.2. Let $A \subseteq \mathbb{T}$ be compact with measure $\lambda(A) = \alpha$, G be an open neighbourhood of A in $\overline{\mathbb{D}}$, and $\varepsilon > 0$. There is a sequence $(p_n)_{n=1}^\infty$ of \mathbb{C} -valued polynomials on \mathbb{C} , mapping \mathbb{D} into \mathbb{D} such that

- (i) $p_n(0) = 0$ for every $n \in \mathbb{N}$.
- (ii) $|p_n(z)| < \varepsilon$ for $z \in \overline{\mathbb{D}} \setminus G$ and $n \in \mathbb{N}$.
- (iii) $(p_n(\lambda \upharpoonright_A))_{n=1}^\infty$ converges narrowly to $\alpha\lambda$.
- (iv) $(p_n(\lambda \upharpoonright_{\mathbb{T} \setminus A}))_{n=1}^\infty$ converges narrowly to $(1 - \alpha)\delta_0$.

In particular $(p_n(\lambda))_{n=1}^\infty$ converges narrowly to $\alpha\lambda + (1 - \alpha)\delta_0$.

Proof. We may suppose $0 < \alpha < 1$. Fix $n \in \mathbb{N}$ and define h_n on \mathbb{T} by

$$h_n(e^{2\pi i\theta}) = \begin{cases} 0 & \text{if } e^{2\pi i\theta} \in A, \\ -n & \text{if } e^{2\pi i\theta} \in \mathbb{T} \setminus A, \end{cases}$$

and extend h_n via the Poisson kernel to a function on $\overline{\mathbb{D}}$ which is harmonic on \mathbb{D} . Let \hat{h}_n be its harmonic conjugate normalized by $\hat{h}_n(0) = 0$ and let

$$g_n = \exp(h_n + i\hat{h}_n),$$

which is a function in $H^\infty(\mathbb{T})$. We have

$$g_n(0) = \exp(h_n(0)) = e^{-\alpha n},$$

and

- (1) $e^{-n} < |g_n(z)| < 1$ for $z \in \mathbb{D}$,
- (2) $|g_n(z)| = 1$ for $z \in A$,
- (3) $|g_n(z)| = e^{-n}$ for $z \in \mathbb{T} \setminus A$.

The last two lines imply that any cluster point of the image measures $(g_n(\lambda))_{n=1}^\infty$ in the narrow topology is necessarily of the form $\alpha\nu + (1 - \alpha)\delta_0$ where ν is a measure supported by the torus \mathbb{T} . We shall show that ν necessarily equals Lebesgue measure. We thank B. Maurey for providing the following proof which is simpler than the original one.

As g_n is analytic on \mathbb{D} for $n \geq 1$, for every harmonic function f defined on \mathbb{C} , $f \circ g_n$ is harmonic on \mathbb{D} . Thus for every $k \geq 1$,

$$\begin{aligned} \alpha \int_0^1 e^{2\pi k i \theta} d\nu(\theta) &= \lim_{n \rightarrow \infty} \int_0^1 g_n^k(e^{2\pi i \theta}) d\theta \\ &= \lim_{n \rightarrow \infty} e^{-\alpha k n} = 0, \end{aligned}$$

the last line using the harmonicity of the function $z \rightarrow z^k$. As ν is a positive measure and for all $k \geq 1$

$$\int_0^1 e^{2\pi k i \theta} d\nu(\theta) = 0,$$

we conclude that ν is Lebesgue measure. Hence

$$\lim_{n \rightarrow \infty} g_n(\lambda \upharpoonright_{\mathbb{T} \setminus A}) = (1 - \alpha)\delta_0, \quad \lim_{n \rightarrow \infty} g_n(\lambda \upharpoonright_A) = \alpha\lambda,$$

with respect to the narrow topology.

We still have to approximate the holomorphic function g_n by appropriate polynomials p_n . First note that it follows by the same argument as above that for every $k \in \mathbb{N}$, the sequence $(g_n^k)_{n=1}^\infty$ is a sequence in the unit ball of $H^\infty(\mathbb{T})$ such that $(g_n^k(\lambda \upharpoonright_{\mathbb{T} \setminus A}))_{n=1}^\infty$ converges narrowly to $(1 - \alpha)\delta_0$ and $(g_n^k(\lambda \upharpoonright_A))_{n=1}^\infty$ converges narrowly to $\alpha\lambda$. Let G_1 be a neighbourhood of A in \mathbb{D} which is relatively compact in G . As $(|g_n^k(z)|)_{k=1}^\infty$ converges to zero uniformly in $z \in \mathbb{D} \setminus G_1$ and $n \in \mathbb{N}$ as $k \rightarrow \infty$, we can find $k \in \mathbb{N}$ such that

$$|g_n^k(z)| < \varepsilon/2 \quad \text{for } n \in \mathbb{N}, z \in \mathbb{D} \setminus G_1.$$

Next note that $\lim_{r \rightarrow 1, r < 1} g_n^k(rz) = g_n^k(z)$ for almost all $z \in \mathbb{T}$. It follows quickly that we may find a sequence $(r_n)_{n=1}^\infty$ in $]0, 1[$ tending sufficiently fast to 1 such that the functions $q_n(z) = g_n^k(r_n z)$ verify

- (ii) $|q_n(z)| < \varepsilon/2$ for $z \in \mathbb{D} \setminus G_1$.
- (iii) $(q_n(\lambda \upharpoonright_A))_{n=1}^\infty$ converges narrowly to $\alpha\lambda$.
- (iv) $(q_n(\lambda \upharpoonright_{\mathbb{T} \setminus A}))_{n=1}^\infty$ converges narrowly to $(1 - \alpha)\delta_0$.

Finally find appropriate polynomials $(p_n)_{n=1}^\infty$ approximating $(q_n)_{n=1}^\infty$ uniformly on \mathbb{D} such that the conclusion of the lemma holds true. \square

We now give an analogue of Proposition II.1 in terms of holomorphic functions instead of analytic martingales. Apparently this result is well known in

the theory of several complex variables but as we could not find an explicit reference in the literature we include a proof for the sake of completeness, which we formulate in the more general setting of Banach spaces.

Proposition III.3. *Let U be a domain in a Banach space X and $f : U \rightarrow \mathbb{R}$ be bounded and uniformly continuous. Define for $x \in U$,*

$$\psi(x) = \inf \int_0^1 f \circ g(e^{2\pi i\theta}) d\theta,$$

where the inf is taken over all X -valued polynomials g such that $g(0) = x$ and $g(\overline{\mathbb{D}}) \subseteq U$. Then ψ equals the largest plurisubharmonic function \hat{f} on U dominated by f .

Proof. ψ is upper semicontinuous: indeed, let $(x_k)_{k=1}^\infty$ in U converge to $x_0 \in U$ and choose, for $\varepsilon > 0$, a polynomial $g : \mathbb{C} \rightarrow X$, $g(0) = x_0$, $g(\overline{\mathbb{D}}) \subseteq U$ such that

$$\psi(x_0) > \int_0^1 f \circ g(e^{2\pi i\theta}) d\theta - \varepsilon.$$

There is $k_0 \in \mathbb{N}$, such that for $k \geq k_0$, $\{g(\overline{\mathbb{D}}) - x_0 + x_k\} \subseteq U$ and by Lebesgue's theorem on dominated convergence

$$\begin{aligned} \limsup_{k \rightarrow \infty} \psi(x_k) &\leq \limsup_{k \geq \infty, k \rightarrow k_0} \int_0^1 f(g(e^{2\pi i\theta}) - x_0 + x_k) d\theta \\ &\leq \int_0^1 f(g(e^{2\pi i\theta})) d\theta < \psi(x_0) + \varepsilon, \end{aligned}$$

which shows that ψ is upper semicontinuous.

For every plurisubharmonic function ϕ on U with $\phi \leq f$ we have $\phi \leq \psi$ as for every polynomial $g : \mathbb{C} \rightarrow X$ with $g(\overline{\mathbb{D}}) \subseteq U$, $g(0) = x_0$ we have

$$\phi(x_0) \leq \int_0^1 \phi \circ g(e^{2\pi i\theta}) d\theta \leq \int_0^1 f \circ g(e^{2\pi i\theta}) d\theta,$$

whence $\phi(x_0) \leq \psi(x_0)$.

We still have to show the mean value inequality for ψ . Let $x_0 \in U$ and $y_0 \in X$, $y_0 \neq 0$, such that $\{x_0 + \overline{\mathbb{D}}y_0\} \subseteq U$. We may assume $x_0 = 0$. Fix $\varepsilon > 0$ and $M > 0$ such that $|f|$ is bounded by M and such that, for $x, y \in U$, $\|x - y\| < \varepsilon/M$ implies $|f(x) - f(y)| < \varepsilon$.

Let \mathcal{D} be a countable subset of the space $\text{Pol}_0(\mathbb{C}, X)$ of X -valued polynomials g , $g(0) = 0$, which is dense with respect to the topology of uniform convergence on $\overline{\mathbb{D}}$.

For $\theta \in \mathbb{T}$ we may find $g_\theta \in \mathcal{D}$ such that

$$\{g_\theta(\overline{\mathbb{D}}) + e^{2\pi i\theta}y_0\} \subseteq U,$$

and

$$\psi(e^{2\pi i\theta}y_0) > \int_0^1 f(g_\theta(e^{2\pi it}) + e^{2\pi i\theta}y_0) dt - \varepsilon.$$

Choose finitely many $(\theta_j)_{j=1}^m$ in $[0, 1[$, compact subsets $(A_j)_{j=1}^m$ of \mathbb{T} and open subsets $(G_j)_{j=1}^m$ of $\overline{\mathbb{D}}$ such that

(i) $e^{2\pi i\theta_j} \in A_j \subseteq G_j$, $(G_j)_{j=1}^m$ are pairwise disjoint and

$$\text{diam}(G_j) < \varepsilon/2M\|y_0\| \quad \text{for } 1 \leq j \leq m;$$

(ii) $\sum_{j=1}^m \lambda(A_j) > 1 - \varepsilon/M$;

(iii) $|\psi(e^{2\pi i\theta} y_0) - \psi(y_0 e^{2\pi i\theta_j})| < \varepsilon$ for $1 \leq j \leq m$ and $e^{2\pi i\theta} \in A_j$;

(iv) $\{g_{\theta_j}(\overline{\mathbb{D}}) + zy_0\} \subseteq U$ for $1 \leq j \leq m$ and $z \in \overline{G_j}$;

which is easily seen to be possible. Define $g_j = g_{\theta_j}$ for $1 \leq j \leq m$.

Let $L \geq 1$ be a Lipschitz constant for the functions $(g_j)_{j=1}^m$ on $\overline{\mathbb{D}}$ and find $0 < \delta < \varepsilon/2M$ such that, for K denoting the compact set

$$K = \{\overline{\mathbb{D}}y_0\} \cup \left\{ \bigcup_{j=1}^m \bigcup_{z \in \overline{G_j}} \{g_j(\overline{\mathbb{D}}) + zy_0\} \right\},$$

the set

$$K_\delta = \{x \in X : \text{dist}_{\|\cdot\|}(x, K) \leq \delta\},$$

is contained in U . Apply Lemma III.3 to find, for $1 \leq j \leq m$, sequences $(p_{j,n})_{n=1}^\infty$ of \mathbb{C} -valued polynomials mapping \mathbb{D} into \mathbb{D} , such that for $1 \leq j \leq m$,

(i) $p_{j,n}(0) = 0$, for $n \in \mathbb{N}$;

(ii) $|p_{j,n}(z)| < \delta/mL$ for $z \in \overline{\mathbb{D}} \setminus G_j$ and $n \in \mathbb{N}$;

(iii) $(p_{j,n}(\lambda 1_{A_j}))_{n=1}^\infty$ converges narrowly to $\lambda(A_j)\lambda$.

Define, for $n \in \mathbb{N}$, the polynomial $h_n : \mathbb{C} \rightarrow X$ by

$$h_n(z) = zy_0 + \sum_{j=1}^m g_j \circ p_{j,n}(z).$$

Let us check that $h_n(\overline{\mathbb{D}}) \subseteq K_\delta \subseteq U$. Indeed, if $z \in \overline{\mathbb{D}}$, $z \notin \bigcup_{j=1}^m G_j$, then, for $1 \leq j \leq m$, $|p_{j,n}(z)| < \delta/mL$, whence

$$\left\| \sum_{j=1}^m g_j \circ p_{j,n}(z) \right\| < mL(\delta/mL) = \delta,$$

and therefore $h_n(z) \in K_\delta$.

If $z \in G_{j_0}$, for some $1 \leq j_0 \leq m$, then

$$h_n(z) = zy_0 + g_{j_0}(p_{j_0,n}(z)) + \sum_{j \neq j_0} g_j \circ p_{j,n}(z),$$

whence

$$\|h_n(z) - (g_{j_0}(p_{j_0,n}(z)) - zy_0)\| = \left\| \sum_{j \neq j_0} g_j \circ p_{j,n}(z) \right\| < \delta,$$

and therefore again $h_n(z) \in K_\delta$.

Finally we may estimate, for $n \in \mathbb{N}$,

$$\begin{aligned} & \int_0^1 f \circ h_n(e^{2\pi i\theta}) d\theta \\ & \leq \sum_{j=1}^m \int_{A_j} f \left\{ e^{2\pi i\theta} y_0 + g_j \circ p_{j,n}(e^{2\pi i\theta}) + \sum_{k \neq j} g_k \circ p_{k,n}(e^{2\pi i\theta}) \right\} d\theta + \varepsilon \\ & \leq \sum_{j=1}^m \int_{A_j} f \{ \{g_j \circ p_{j,n}(e^{2\pi i\theta}) + y_0 e^{2\pi i\theta_j}\} + k_j(\theta) \} d\theta + \varepsilon, \end{aligned}$$

where

$$k_j(\theta) = \sum_{k \neq j} g_k \circ p_{k,n}(e^{2\pi i\theta}) + (e^{2\pi i\theta} - e^{2\pi i\theta_j})y_0,$$

so that, for $\theta \in A_j$, $\|k_j(\theta)\| < \varepsilon/2M + \delta < \varepsilon/M$.

By the equicontinuity assumption on f we therefore can estimate

$$\int_0^1 f \circ h_n(e^{2\pi i\theta}) d\theta \leq \sum_{j=1}^m \int_{A_j} f(g_j \circ p_{j,n}(e^{2\pi i\theta}) + y_0 e^{2\pi i\theta_j}) d\theta + 2\varepsilon,$$

passing to $n \rightarrow \infty$ we obtain

$$\begin{aligned} \psi(0) & \leq \lim_{n \rightarrow \infty} \sum_{j=1}^m \int_{A_j} f(g_j \circ p_{j,n}(e^{2\pi i\theta}) + y_0 e^{2\pi i\theta_j}) d\theta + 2\varepsilon \\ & = \sum_{j=1}^m \lambda(A_j) \int_0^1 f(g_j(e^{2\pi i\theta}) + y_0 e^{2\pi i\theta_j}) d\theta + 2\varepsilon \\ & \leq \sum_{j=1}^m \lambda(A_j) \psi(y_0 e^{2\pi i\theta_j}) + 3\varepsilon \\ & \leq \sum_{j=1}^m \int_{A_j} \psi(e^{2\pi i\theta} y_0) d\theta + 4\varepsilon \\ & \leq \int_0^1 \psi(e^{2\pi i\theta} y_0) d\theta + 5\varepsilon, \end{aligned}$$

the last line using the fact that $\psi \geq -M$. As $\varepsilon > 0$ is arbitrary we obtain the mean value inequality for ψ and finish the proof of III.4. \square

Proof of Theorem (B). Fix μ, K, U and x_0 as in Theorem (B). We may assume $x_0 = 0$. We shall first prove Theorem (B) with respect to the class of bounded uniformly continuous functions on U . Set

$$\mathbf{P} = \{g(\lambda) : g : \mathbb{C} \rightarrow \mathbb{C}^n \text{ a polynomial, } g(0) = 0, g(\overline{\mathbb{D}}) \subseteq U\},$$

which we consider as a subset of $\mathcal{M}^c(U)$ the space of Radon measures with compact support in U . This space is in duality with the space $C^{\text{ucb}}(U)$ of bounded uniformly continuous functions on U and we equip $\mathcal{M}^c(U)$ with the weak topology induced by $C^{\text{ucb}}(U)$.

We now show that the closure of \mathbf{P} is convex. Let $(g_j)_{j=1}^m$ be \mathbb{C}^n -valued polynomials, $g_j(0) = 0, g_j(\overline{\mathbb{D}}) \subseteq U$ for $1 \leq j \leq m$ and $(c_j)_{j=1}^m$ positive scalars, $\sum_{j=1}^m c_j = 1$.

Fix $\varepsilon > 0$ and find disjoint compact sets $(A_j)_{j=1}^m$ of \mathbb{T} of measure $\lambda(A_j) = (1 - \varepsilon)c_j$ and disjoint neighbourhoods $(G_j)_{j=1}^m$ of $(A_j)_{j=1}^m$ in $\overline{\mathbb{D}}$. Fix a norm $\|\cdot\|$ on \mathbb{C}^n , find $\delta > 0$ such that $\min_{1 \leq j \leq m} \text{dist}_{\|\cdot\|}(g_j(\overline{\mathbb{D}}), \mathbb{C}^n \setminus U) > \delta$ and let L be a Lipschitz constant for $(g_j)_{j=1}^m$ on $\overline{\mathbb{D}}$.

Now apply Lemma III.2 to find sequences $(p_{j,k})_{k=1}^\infty$ of \mathbb{C} -valued polynomials such that, for $1 \leq j \leq m$,

- (i) $p_{j,k}(0) = 0$ for $k \in \mathbb{N}$;
- (ii) $|p_{j,k}(z)| < \delta/Lm$ for $z \in \overline{\mathbb{D}} \setminus G_j$ and $k \in \mathbb{N}$;
- (iii) $(p_{j,k}(\lambda \upharpoonright_{A_j}))_{k=1}^\infty$ converges narrowly to $(1 - \varepsilon)c_j \lambda$;
- (iv) $(p_{j,k}(\lambda \upharpoonright_{\mathbb{T} \setminus A_j}))_{k=1}^\infty$ converges narrowly to $(1 - (1 - \varepsilon)c_j)\delta_0$. Then

$$h_k = \sum_{j=1}^m g_j \circ p_{j,k}$$

is a \mathbb{C}^k -valued polynomial, $h_k(0) = 0$ and $h_k(\overline{\mathbb{D}})$ is contained in U . For every $f \in C^{\text{ucb}}(U)$ we have

$$\lim_{k \rightarrow \infty} \langle f, h_k(\lambda) \rangle = (1 - \varepsilon) \left\langle f, \sum_{j=1}^m c_j g_j(\lambda) \right\rangle,$$

which readily shows that $\sum_{j=1}^m c_j g_j(\lambda)$ is in the closure of P .

Hence similarly as in the proof of Theorem (A) we now are in a position to apply the Hahn-Banach theorem: If μ were not in the closure of \mathbf{P} , then we could find $f \in C^{\text{ucb}}(U)$ and $\alpha < \beta$ such that

$$\langle f, \mu \rangle = \int_U f d\mu \leq \alpha, \quad \langle f, \nu \rangle = \int_U f d\nu \geq \beta,$$

for every $\nu \in A$. The last line can be rewritten as

$$\int_{\mathbb{T}} f \circ g d\lambda \geq \beta$$

for every polynomial $g : \mathbb{C} \rightarrow \mathbb{C}^n$, $g(0) = 0$, $g(\overline{\mathbb{D}}) \subseteq U$ whence Proposition III.3 implies that $\hat{f}(0) \geq \beta$ and we arrive at the desired contradiction:

$$\alpha \geq \langle f, \mu \rangle \geq \langle \hat{f}, \mu \rangle \geq \hat{f}(0) \geq \beta.$$

Now for $\varepsilon > 0$, f_1, f_2, \dots, f_m continuous functions on U bounded by $M > 0$, there is $\delta > 0$ such that $K + \overline{B}(0, \delta) \subseteq U$, there are g_1, g_2, \dots, g_m uniformly continuous functions on U bounded by M such that, for $1 \leq i \leq m$, f_i and g_i coincide on $K + \overline{B}(0, \delta)$. Note that there is a uniformly continuous function h on U with $0 \leq h \leq 1$, and such that h equals 1 on $U \setminus \{K + \overline{B}(0, \delta)\}$ and equals 0 on K . Applying the conclusion of Theorem (B) for $\varepsilon > 0$, g_1, g_2, \dots, g_m and h , we can find a polynomial $p : \mathbb{C} \rightarrow \mathbb{C}^n$, $p(0) = 0$, $p(\overline{\mathbb{D}}) \subseteq U$, such that

$$\begin{aligned} |\langle h, \mu \rangle - \langle h, p(\lambda) \rangle| &< \varepsilon, \\ |\langle g_i, \mu \rangle - \langle g_i, p(\lambda) \rangle| &< \varepsilon \quad \text{for } 1 \leq i \leq m. \end{aligned}$$

Note that μ is supported by K and h vanishes on K , so $|\langle h, p(\lambda) \rangle| < \varepsilon$. Note also that $0 \leq h \leq 1$ and h equals 1 on $U \setminus \{K + \overline{B}(0, \delta)\}$, we have then

$p(\lambda)\{U \setminus \{K + \overline{B}(0, \delta)\}\} \leq \varepsilon$, whence

$$|\langle f_i, \mu \rangle - \langle f_i, p(\lambda) \rangle| \leq |\langle g_i, \mu \rangle - \langle g_i, p(\lambda) \rangle| + 2Mp(\lambda)\{U \setminus \{K + \overline{B}(0, \delta)\}\} \leq \varepsilon + 2M\varepsilon \quad \text{for } 1 \leq i \leq m.$$

This shows that the conclusion of Theorem (B) holds true for f_1, f_2, \dots, f_m .
 \square

To end this section we show the compatibility of Definition I.3 and Definition I.5 above.

Proposition III.4. *Let μ be a probability measure on \mathbb{C}^n with compact support. If the inequality*

$$(1) \quad \varphi(0) \leq \int_{\mathbb{C}^n} \varphi(x) d\mu(x)$$

holds true for all Lipschitz plurisubharmonic functions φ on \mathbb{C}^n , then (1) holds true for all plurisubharmonic functions φ on \mathbb{C}^n .

We shall say that a function $f : \mathbb{C}^n \rightarrow \mathbb{R}$ is locally Lipschitz if, for every bounded open subset U of \mathbb{C}^n , $f|_U$ is a Lipschitz function on U . The proof of Proposition III.4 will rely on the following result:

Proposition III.5. *Let φ be a locally Lipschitz plurisubharmonic function on \mathbb{C}^n . Then φ can be approximated by Lipschitz plurisubharmonic functions uniformly on compact subsets of \mathbb{C}^n .*

Admitting Proposition III.5 it is fairly standard to deduce Proposition III.4:

Proof of Proposition III.4. Fix a plurisubharmonic function φ on \mathbb{C}^n . It follows from the argument in [Ra, Theorem II.4.12] that there is a decreasing sequence $(\varphi_j)_{j=1}^\infty$ of plurisubharmonic C^∞ -functions on \mathbb{C}^n decreasing pointwise to φ . In particular each φ_j is locally Lipschitz, hence we may find by Proposition III.5 a sequence $(\psi_j)_{j=1}^\infty$ of Lipschitz plurisubharmonic functions on \mathbb{C}^n such that $|\varphi_j - \psi_j| \leq 1/j$ on $\text{supp}(\mu) \cup \{0\}$.

Assuming the validity of inequality (1) for every ψ_j we obtain

$$\varphi_j(0) \leq \int_{\mathbb{C}^n} \varphi_j(x) d\mu(x) + 2/j,$$

whence by the Beppo-Levi theorem

$$\varphi(0) \leq \int_{\mathbb{C}^n} \varphi(x) d\mu(x). \quad \square$$

Proof of Proposition III.5. Fix a norm $\|\cdot\|$ on $X = \mathbb{C}^n$ and a locally Lipschitz plurisubharmonic function φ on X . We shall approximate φ uniformly on $B_1 = \{x : \|x\| \leq 1\}$ by Lipschitz plurisubharmonic functions ψ on X . We may assume that $\varphi|_{B_1} \geq 0$ and we may find $k_0 \in \mathbb{N}$ such that $\varphi|_{B_1}$ obeys a Lipschitz constant less than k_0 .

For $k \geq k_0$, let φ_k be the largest function on \mathbb{C}^n satisfying a Lipschitz constant k and coinciding with φ on B_1 , and denote by $\hat{\varphi}_k$ the plurisubharmonic envelope of φ_k . The sequence $(\hat{\varphi}_k)_{k=k_0}^\infty$ is an increasing sequence of Lipschitz plurisubharmonic functions on \mathbb{C}^n [G-M, Lemma II.1] such that, for every $k \geq k_0$,

$$\hat{\varphi}_k|_{B_1} \leq \varphi|_{B_1}.$$

By Dini's theorem it will suffice to prove that $(\hat{\varphi}_k)_{k=k_0}^\infty$ tends pointwise to φ on B_1 .

Let us assume to the contrary that there is $x_0 \in B_1$ and $\alpha < \beta$ such that $\hat{\varphi}_k(x_0) < \alpha$ for all $k \geq k_0$ while $\varphi(x_0) = \beta$ and let us work towards a contradiction. For $k \geq k_0$ we can find, by Proposition III.3, a polynomial $p_k : \mathbb{C} \rightarrow X$, $p_k(0) = x_0$ and such that

$$(2) \quad \mathbb{E}(\varphi_k \circ p_k) = \int_0^1 \varphi_k(p_k(e^{2i\pi\theta})) d\theta < \alpha.$$

Hence for every $\varepsilon > 0$

$$(3) \quad \lim_{k \rightarrow \infty} \mathbb{E} \|p_k \mathbb{1}_{\{\|p_k\| > 1+\varepsilon\}}\| = 0.$$

Indeed, for every $M \in \mathbb{N}$ there is $k_1 \geq k_0$ such that for $k \geq k_1$ and $x \in X$, $\|x\| > 1 + \varepsilon$, $\varphi_k(x) > M\|x\|$, hence

$$\lim_{k \rightarrow \infty} \mathbb{E} \|p_k \mathbb{1}_{\{\|p_k\| > 1+\varepsilon\}}\| \leq \lim_{k \rightarrow \infty} M^{-1} \mathbb{E}(\varphi_k \circ p_k) < M^{-1}\alpha,$$

which proves (3). In particular the sequence $\{\|p_k\|\}_{k=k_0}^\infty$ is uniformly integrable in $L^1(\mathbb{T}, \lambda)$.

Denote by $(W_t)_{t \leq \tau}$ Brownian motion on \mathbb{C} modelled on some probability space $(\Omega, \Sigma, \mathbb{P})$, with $W_0 \equiv 0$ and stopped at the first time τ when $|W_\tau| = 1$. For $k \geq k_0$ denote by τ_k the stopping time

$$\tau_k(\omega) = \tau(\omega) \wedge \inf\{t : \|p_k \circ W_t(\omega)\| \geq 2\},$$

and let $A_k = \{\omega \in \Omega : \tau_k(\omega) < \tau(\omega)\}$. We claim that

$$(4) \quad \lim_{k \rightarrow \infty} \mathbb{P}(A_k) = 0.$$

Indeed, the process $\{(\|p_k \circ W_t\| - 3/2)^+\}_{t \leq \tau}$ is a submartingale whence

$$\begin{aligned} \mathbb{P}(A_k)/2 &= \mathbb{E}((\|p_k \circ W_{\tau_k}\| - 3/2)^+ \mathbb{1}_{A_k}) \\ &\leq \mathbb{E}((\|p_k \circ W_\tau\| - 3/2)^+ \mathbb{1}_{A_k}) \\ &\leq \mathbb{E}((\|p_k \circ W_\tau\| - 3/2)^+) \\ &\leq \mathbb{E}(\|p_k \mathbb{1}_{\{\|p_k\| > 3/2\}}\|), \end{aligned}$$

whence (4) follows from (3). Now let R_k (resp. S_k) be the X -valued random variable $p_k \circ W_\tau$ (resp. $p_k \circ W_{\tau_k}$) defined on $(\Omega, \Sigma, \mathbb{P})$. The sequence $(R_k)_{k=k_0}^\infty$ has the same law as the sequence $(p_k)_{k=k_0}^\infty$ of random variables on (\mathbb{T}, λ) , so the sequence $(R_k)_{k=1}^\infty$ is uniformly integrable in $L^1(\mathbb{P}, X)$. Clearly $\|S_k(\omega)\|$ is bounded by 2 for $k \geq k_0$ and $\omega \in \Omega$. We may estimate

$$\lim_{k \rightarrow \infty} \mathbb{E}(\|R_k - S_k\|) \leq \lim_{k \rightarrow \infty} (\mathbb{E}(\|R_k \mathbb{1}_{A_k}\|) + \mathbb{E}(\|S_k \mathbb{1}_{A_k}\|)) = 0,$$

whence for every Lipschitz function f on \mathbb{C}^n

$$(5) \quad \limsup_{k \rightarrow \infty} \mathbb{E}(f \circ R_k) = \limsup_{k \rightarrow \infty} \mathbb{E}(f \circ S_k).$$

The function φ is plurisubharmonic whence for $k \geq k_0$ the process $(\varphi \circ p_k \circ W_t)_{t \leq \tau}$ is a submartingale and therefore

$$\beta = \varphi(x_0) \leq \mathbb{E}(\varphi \circ p_k \circ W_{\tau_k}) = \mathbb{E}(\varphi \circ S_k).$$

On the other hand let $m \geq k_0$ be big enough such that $\varphi_m|_{B_2} \geq \varphi|_{B_2}$. Then we may apply (5) and (1) to obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \mathbb{E}(\varphi \circ S_k) &\leq \limsup_{k \rightarrow \infty} \mathbb{E}(\varphi_m \circ S_k) \\ &\leq \limsup_{k \rightarrow \infty} \mathbb{E}(\varphi_m \circ R_k) \\ &\leq \limsup_{k \rightarrow \infty} \mathbb{E}(\varphi_k \circ R_k) \\ &= \limsup_{k \rightarrow \infty} \mathbb{E}(\varphi_k \circ p_k) \leq \alpha, \end{aligned}$$

a contradiction finishing the proof of Proposition III.5. \square

IV. A VARIATION OF THEOREM (A)

In order to prove Theorem (C) we shall need a refinement of Theorem (A). The proof involves standard but cumbersome measure theoretical arguments and will be given in this section.

Proposition IV.1. *Let μ be a Jensen measure on a complex Banach space X with barycenter x_0 and $\varepsilon > 0$. Then there is a polynomial $g : \mathbb{C} \rightarrow X$ and a measurable function $f \in L^1(\mathbb{T}, X)$ such that*

- (i) $f(\lambda) = \mu$,
- (ii) $g(0) = x_0$,
- (iii) $\|f - g\|_1 = \int_0^1 \|f(t) - g(e^{2\pi it})\| d\lambda(t) < \varepsilon$.

Before proving Proposition IV.1 we deduce a parametrized version, which will be precisely what we need:

Proposition IV.2. *Let (E, d) be a polish space equipped with its Borel σ -algebra Σ and let ρ be a probability measure on Σ . Let $(\mu_z)_{z \in E}$ be a family of Jensen measures on X with barycenter $(x_z)_{z \in E}$ depending measurably on z in the sense that, for every $\phi \in \text{Lip}(X)$,*

$$z \rightarrow \langle \phi, \mu_z \rangle = \int_X \phi(x) d\mu_z(x),$$

is Borel measurable. Then, for $\varepsilon > 0$, there is a ρ -measurable function

$$\begin{aligned} (F, G) : X &\rightarrow L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X), \\ z &\rightarrow (f_z, g_z) \end{aligned}$$

such that, for every $z \in X$, g_z is the restriction to \mathbb{T} of a polynomial on \mathbb{C} and

- (i) $f_z(\lambda) = \mu_z$,
- (ii) $g_z(0) = x_z$,
- (iii) $\|f_z - g_z\| < \varepsilon$.

Proof. Define \mathbf{M} to be the subset of $X \times L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X)$

$$\mathbf{M} = \{(z, f, g) : g \text{ is the restriction to } \mathbb{T} \text{ of a polynomial on } \mathbb{C}, f(\lambda) = \mu_z, g(0) = \mathbb{E}(f), \text{ and } \|f - g\| < \varepsilon\}.$$

The set \mathbf{M} is Borel in $X \times L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X)$. Indeed, let $(\phi_n)_{n=4}^\infty$ be a sequence of Lipschitz functions on X separating the points of $\mathcal{M}^1(X)$ and let

$$\begin{aligned} \mathbf{M}_1 &= \{(z, f, g) : g \text{ is the restriction to } \mathbb{T} \text{ of a polynomial on } \mathbb{C}\}, \\ \mathbf{M}_2 &= \{(z, f, g) : \|f - g\| < \varepsilon\}, \\ \mathbf{M}_3 &= \{(z, f, g) : g(0) = \mathbb{E}(f)\}, \\ \mathbf{M}_n &= \{(z, f, g) : \langle \phi_n, f(\lambda) \rangle = \langle \phi_n, \mu_z \rangle\} \text{ for } n \geq 4. \end{aligned}$$

One can easily check that the above sets are Borel in $X \times L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X)$, hence $\mathbf{M} = \bigcap_{n=1}^\infty \mathbf{M}_n$ is so too.

By Proposition IV.1 the natural projection from M to X is onto hence by a measurable selection theorem (see, e.g., [Co, Theorem 8.5.3] or [H-J, Theorem 9.5]) we can find a ρ -measurable selection

$$(F, G) : X \rightarrow L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X),$$

such that, for every $z \in X$, $(z, (F, G)(z))$ is in \mathbf{M} . This means that the function (F, G) satisfies our requirements. \square

Let us now start to prove Proposition IV.1. Recall first the elementary fact that, if μ is a positive measure on a polish space (E, d) of mass $\|\mu\| = \mu(E) = \alpha$ and A is a subset of $[0, 1]$ of measure $\lambda(A) = \alpha$ then there is a Borel measurable function $f : A \rightarrow E$ such that $f(\lambda \upharpoonright_A) = \mu$. The next lemma builds on this observation.

Lemma IV.3. *Let μ be a Radon probability measure on a polish space (E, d) , $g : [0, 1] \rightarrow E$ Lebesgue measurable and $\varepsilon, \delta > 0$. Suppose further that there are disjoint sets $(A_i)_{i=1}^n$ of $[0, 1]$ and disjoint open sets $(B_i)_{i=1}^n$ in E such that, letting $\alpha_i = \lambda(A_i)$ and $\beta_i = \mu(B_i)$ for $1 \leq i \leq n$*

- (i) $\sum_{i=1}^n \beta_i > 1 - \delta/2$;
- (ii) $\alpha_i > \beta_i - \delta/2n$ for $1 \leq i \leq n$;
- (iii) $\text{diam}(B_i) = \sup\{d(x, y) : x, y \in B_i\} < \varepsilon$ for $1 \leq i \leq n$;
- (iv) $g(t) \in B_i$ for $t \in A_i$ and $1 \leq i \leq n$.

Then there is a Borel measurable function $f : [0, 1] \rightarrow E$ with $f(\lambda) = \mu$ and a measurable subset $C \subseteq [0, 1]$ of measure $\lambda(C) > 1 - \delta$ such that $\|f(t) - g(t)\| < \varepsilon$ on C .

Proof. For $1 \leq i \leq n$, let $\gamma_i = \min(\alpha_i, \beta_i)$, find Borel measurable subsets C_i of A_i of measure $\lambda(C_i) = \gamma_i$ and let $\mu_i = (\gamma_i/\beta_i)\mu \upharpoonright_{B_i}$, so that the mass of the measures μ_i equals $\|\mu_i\| = \gamma_i$. Note that $\sum_{i=1}^n \gamma_i > 1 - \delta$. Let $C = \bigcup_{i=1}^n C_i$, $C_0 = [0, 1] \setminus C$, and $\mu_0 = \mu - \sum_{i=1}^n \mu_i$.

For $i = 0, 1, \dots, n$ apply the preceding remark to find measurable functions

$$f_0 : C_0 \rightarrow E, \quad f_i : C_i \rightarrow B_i \text{ for } 1 \leq i \leq n,$$

such that

$$f_i(\lambda \upharpoonright_{C_i}) = \mu_i \text{ for } 0 \leq i \leq n.$$

The function $f = \sum_{i=0}^n f_i \upharpoonright_{C_i}$ satisfies the requirements. \square

Proof of Proposition IV.1. As μ has a first moment we may find $\delta > 0$ such that, for $B \subseteq X$, $\mu(B) < \delta$ we have

$$(1) \quad \int_B \|x\| d\mu(x) < \varepsilon.$$

Next find disjoint compact sets $(K_i)_{i=1}^n$ in X of diameter less than ε such that

$$(2) \quad \sum_{i=1}^n \mu(K_i) > 1 - \delta/2,$$

and open disjoint neighbourhoods $(B_i)_{i=1}^n$ of $(K_i)_{i=1}^n$ of diameter less than ε such that $\mu(B_i \setminus K_i) < \delta/4n$. Now choose Lipschitz functions $(\phi_i)_{i=1}^n$ from X to $[0, 1]$ such that ϕ_i equals 1 on K_i and vanishes outside of B_i .

By Theorem (A) we may find a polynomial $g : \mathbb{C} \rightarrow X$ with $g(0) = x_0 = \text{bary}(\mu)$ such that

$$(3) \quad |\langle \phi_i, \mu \rangle - \langle \phi_i, g(\lambda) \rangle| < \delta/4n \quad \text{for } 1 \leq i \leq n,$$

$$\left| \int_X \|x\| d\mu(x) - \int_X \|x\| d(g(\lambda))(x) \right| < \varepsilon.$$

For $A_i = g^{-1}(B_i)$ let us check that the requirements of Lemma IV.3 are satisfied. The only condition which is not obvious is (ii) which for $1 \leq i \leq n$ follows from

$$\begin{aligned} \lambda(A_i) &\geq \langle \phi_i, g(\lambda) \rangle > \langle \phi_i, \mu \rangle - \delta/4n \\ &\geq \mu(K_i) - \delta/4n \geq \mu(B_i) - \delta/2n. \end{aligned}$$

Hence we may find f and C as in Lemma IV.3.

Note that it follows from (1) and the relation $\mu = f(\lambda)$ that for any subset $A \subseteq \mathbb{T}$ of measure $\lambda(A) < \delta$ we have

$$\int_A \|f(t)\| d\lambda(t) < \varepsilon.$$

In order to estimate $\|f - g\|_1$ the crucial point is to control the L^1 -mass of g on $C_0 = \mathbb{T} \setminus C$:

$$\begin{aligned} \int_{C_0} \|g(t)\| d\lambda(t) &= \int_{\mathbb{T}} \|g(t)\| d\lambda(t) - \int_C \|g(t)\| d\lambda(t) \\ &\leq \int_{\mathbb{T}} \|f(t)\| d\lambda(t) - \int_C \|f(t)\| d\lambda(t) + \int_C \|f(t) - g(t)\| d\lambda(t) + \varepsilon \\ &< \int_{C_0} \|f(t)\| d\lambda(t) + 2\varepsilon < 3\varepsilon, \end{aligned}$$

where in the second line we have used (3).

We therefore can estimate

$$\begin{aligned} \|f - g\|_1 &< \int_C \|f(t) - g(t)\| d\lambda(t) + \int_{C_0} \|f(t)\| d\lambda(t) + \int_{C_0} \|g(t)\| d\lambda(t) \\ &< \varepsilon + \varepsilon + 3\varepsilon = 5\varepsilon, \end{aligned}$$

thus finishing the proof of Proposition IV.1. \square

V. THE PROOF OF THEOREM (C)

Let $(M_n)_{n=1}^\infty$ be an X -valued stochastic process defined on a probability space $(\Omega, \Sigma, \mathbb{P})$. Note that there is a standard representation of this process obtained in the following way. Let

$$\vec{M} : \Omega \rightarrow X^\mathbb{N}, \quad \omega \rightarrow (M_n(\omega))_{n=1}^\infty,$$

denote by μ the image measure $\vec{M}(\mathbb{P})$, by $\pi_n : X^{\mathbb{N}} \rightarrow X$ the projection onto the n th coordinate and by $\Pi_n : X^{\mathbb{N}} \rightarrow X^n$ the projection onto the n first coordinates. Equipping $X^{\mathbb{N}}$ with the Borel σ -algebra Σ and letting Σ_n the σ -algebra generated by Π_n , the process $(\pi_n)_{i=1}^{\infty}$ on the probability space $(X^{\mathbb{N}}, \Sigma, \mu)$ is adapted to $(\Sigma_n)_{i=1}^{\infty}$ and has the same law as $(M_n)_{i=1}^{\infty}$.

From now on we suppose that $(M_n)_{i=1}^n$ is a martingale and we fix its standard representation, i.e., the measure μ on $X^{\mathbb{N}}$. For $n \in \mathbb{N}$ we denote by X_n the n th coordinate of $X^{\mathbb{N}}$, by μ_n the image measure $\mu_n = \pi_n(\lambda)$ on X_n , and by ρ_n the image measure $\rho_n = \Pi_n(\lambda)$ on X^n . X^0 will be identified with the one point set $\{0\}$ and ρ_0 with the unique probability measure on X^0 . The elements $(x_1, x_2, \dots, x_n) \in X^n$ will occasionally be denoted by y_n .

By a well-known disintegration theorem (see, e.g., [Schw, Theorem 5.44]) we can, for $n \in \mathbb{N}$, disintegrate ρ_n with respect to its marginal ρ_{n-1} , i.e., we can find a measurable map

$$F_n : X^{n-1} \rightarrow \mathcal{M}^1(X), \quad y_{n-1} \rightarrow \mu_{n, y_{n-1}},$$

such that

$$(1) \quad \rho_n = \int_{X^{n-1}} \mu_{n, y_{n-1}} d\rho_{n-1}(y_{n-1}).$$

Measurability here means that, for every $f \in \text{Lip}(X)$ the map

$$\langle f, F_n(\cdot) \rangle : X^{n-1} \rightarrow \mathbb{R}, \quad y_{n-1} \rightarrow \langle f, \mu_{n, y_{n-1}} \rangle,$$

is Borel measurable and formula (1) is a symbolic way of writing that, for every $f \in \text{Lip}(X^n)$ we have

$$\langle f, \rho_n \rangle = \int_{X^{n-1}} \int_{X_n} f(y_{n-1}, x_n) d\mu_{n, y_{n-1}}(x_n) d\rho_{n-1}(y_{n-1}).$$

Proposition V.1. *Let $(M_n)_{n=1}^{\infty}$ be a martingale with values in a complex Banach space X and let the probability measure μ on $X^{\mathbb{N}}$ be its standard representation. Letting $M_0 \equiv \mathbb{E}(M_1)$ then using the above notation, $(M_n)_{n=1}^{\infty}$ is a plurisubharmonic martingale if and only if μ_1 and, for $n \geq 2$ and ρ_{n-1} almost all $y_{n-1} \in X^{n-1}$, $\mu_{n, y_{n-1}}$ are Jensen measures on X .*

Proof. First note that, for $n \geq 2$ and for ρ_{n-1} almost all $y_{n-1} = (x_1, x_2, \dots, x_{n-1})$ the barycenter of $\mu_{n, y_{n-1}}$ written symbolically as

$$\text{bary}(\mu_{n, y_{n-1}}) = \int_{X_n} x_n d\mu_{n, y_{n-1}}(x_n),$$

equals x_{n-1} . Indeed let $(x_i^*)_{i=1}^{\infty}$ be a sequence in X^* separating points of X ; we have to show that, for $i \in \mathbb{N}$ and for ρ_{n-1} almost all $y_{n-1} = (x_1, \dots, x_{n-1})$

$$(1) \quad \langle x_{n-1}, x_i^* \rangle = \int_{X_n} \langle x_n, x_i^* \rangle d\mu_{n, y_{n-1}}(x_n).$$

As by assumption $(x_i^* \circ \pi_n)_{n=1}^{\infty}$ is a martingale on $(X^{\mathbb{N}}, \mu)$ with respect to the filtration $(\Sigma_n)_{n=1}^{\infty}$ we have for every Borel subset $A \subseteq X^{n-1}$ and every $i \in \mathbb{N}$

$$\int_A \langle x_{n-1}, x_i^* \rangle d\rho_{n-1}(y_{n-1}) = \int_A \int_{X_n} \langle x_n, x_i^* \rangle d\mu_{n, y_{n-1}}(x_n) d\rho_{n-1}(y_{n-1}),$$

which readily gives (1).

If now μ_1 and, for $n \geq 2$ and ρ_{n-1} almost all $y_{n-1} \in X^{n-1}$ the measure $\mu_{n,y_{n-1}}$ is Jensen on X , then it is straightforward to check that $(M_n)_{n=0}^\infty$ is a plurisubharmonic martingale.

Conversely, if $(M_n)_{n=1}^\infty$ is a plurisubharmonic martingale then it follows immediately that μ_1 is a Jensen measure on X . Fix $n \geq 2$; we have to show that, for ρ_{n-1} almost all $y_{n-1} = (x_1, x_2, \dots, x_{n-1}) \in X^{n-1}$ we have that for all Lipschitz plurisubharmonic functions ϕ on X

$$\phi(x_{n-1}) \leq \int_{X_n} \phi(x_n) d\mu_{n,y_{n-1}}(x_n).$$

Let $(\phi_n)_{n=1}^\infty$ be a sequence in the set $\text{PSH}_1(X)$ of Lipschitz plurisubharmonic functions on X with Lipschitz constant less than 1 that vanish at the origin such that $(\phi_n)_{n=1}^\infty$ is dense with respect to the topology of pointwise convergence on X . We have to show that the function Φ on X^{n-1} defined by

$$\Phi(y_{n-1}) = \inf_{\phi \in \text{PSH}_1(X)} \left\{ \int_{X_n} \phi(x_n) d\mu_{n,y_{n-1}}(x_n) - \phi(x_{n-1}) \right\}$$

is not strictly negative on a set of ρ_{n-1} -positive measure. By applying the subsequent Lemma V.2 to the measure $\mu_{n,y_{n-1}} - \delta_{\{x_{n-1}\}}$ we conclude that

$$\Phi(y_{n-1}) = \inf_{i \in \mathbb{N}} \left\{ \int_{X_n} \phi_i(x_n) d\mu_{n,y_{n-1}}(x_n) - \phi_i(x_{n-1}) \right\},$$

and it will therefore suffice to show that, for every $i \in \mathbb{N}$,

$$\Phi_i(y_{n-1}) = \int_{X_n} \phi_i(x_n) d\mu_{n,y_{n-1}}(x_n) - \phi_i(x_{n-1})$$

is greater than or equal to zero ρ_{n-1} -almost surely. If this were not the case, we could find a Borel set $A \subseteq X^{n-1}$ such that

$$\int_A \Phi_i(y_{n-1}) d\rho_{n-1}(y_{n-1}) < 0,$$

which is contradictory to the assumption that $(\phi_i \circ \pi_n)_{n=1}^\infty$ is a submartingale on $(X^\mathbb{N}, \mu)$ in view of

$$\int_A \int_{X_n} \phi_i(x_n) d\mu_{n,y_{n-1}}(x_n) d\rho_{n-1}(y_{n-1}) < \int_A \phi_i(x_{n-1}) d\rho_{n-1}(y_{n-1}). \quad \square$$

We have used the subsequent lemma whose proof is left to the reader:

Lemma V.2. *Let μ be a finite signed measure on X such that the absolute value $|\mu|$ has a first moment. Let C be a set of functions with Lipschitz constant bounded by 1 and \bar{C} its closure in the topology of pointwise convergence on X . Then*

$$\sup_{\phi \in C} \langle \phi, \mu \rangle = \sup_{\phi \in \bar{C}} \langle \phi, \mu \rangle.$$

Proof of Theorem (C). Let $(M_n)_{n=0}^\infty$ be an X -valued plurisubharmonic martingale, $M_0 \equiv x_0$. Let μ be its standard representation on $X^\mathbb{N}$ and, using the above notation, $(\mu_n)_{n=1}^\infty$ and $(\rho_n)_{n=0}^\infty$ the marginals of μ on X_n and X^n respectively. We proceed by induction on $n \in \mathbb{N}$.

Let $F_0 \equiv G_0 \equiv x_0$. For $n = 1$ we infer from Proposition V.1 that μ_1 is a Jensen measure on X with barycenter x_0 . We may apply Proposition IV.1

to find $F_1 \in L^1(\mathbb{T}, X)$ and an X -valued polynomial G_1 such that $G_1(0) = x_0$, $F_1(\lambda) = \mu_1$ and

$$\|(F_1 - F_0) - (G_1 - G_0)\|_1 < \varepsilon_1.$$

Suppose that we have defined $(F_i)_{i=0}^{n-1}$ and $(G_i)_{i=0}^{n-1}$, such that $(G_i)_{i=0}^{n-1}$ is a Hardy martingale,

$$\|(F_i - F_{i-1}) - (G_i - G_{i-1})\|_1 < \varepsilon_1 \quad \text{for } 1 \leq i \leq n-1,$$

and such that denoting by

$$\vec{F}_{n-1} : \mathbb{T}^{n-1} \rightarrow X^{n-1},$$

$$\vec{F}_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1}) = (F_1(\theta_1), F_2(\theta_1, \theta_2), \dots, F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})),$$

we have

$$(1) \quad \vec{F}_{n-1}(\lambda^{n-1}) = \rho_{n-1},$$

which means that $(F_i)_{i=0}^{n-1}$ is a representation of $(M_i)_{i=0}^{n-1}$.

Consider the disintegration of ρ_n discussed above,

$$(2) \quad \rho_n = \int_{X^{n-1}} \mu_{n, y_{n-1}} d\rho_{n-1}(y_{n-1}).$$

We may suppose by Proposition V.1 that, for every $y_{n-1} \in X^{n-1}$, $\mu_{n, y_{n-1}}$ is a Jensen measure on X with barycenter x_{n-1} , where $y_{n-1} = (x_1, x_2, \dots, x_{n-1})$. We now can apply Proposition IV.2 for $(E, \rho) = (\mathbb{T}^{n-1}, \lambda^{n-1})$ to find λ^{n-1} -measurable functions

$$(F, G) : \mathbb{T}^{n-1} \rightarrow L^1(\mathbb{T}, X) \times L^1(\mathbb{T}, X),$$

$$(\theta_1, \theta_2, \dots, \theta_{n-1}) \rightarrow \{f_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\theta_n), g_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\theta_n)\}$$

such that, for every $(\theta_1, \theta_2, \dots, \theta_{n-1}) \in \mathbb{T}^{n-1}$, $g_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\cdot)$ is the restriction to \mathbb{T} of an X -valued polynomial defined on \mathbb{C} verifying

$$g_{\theta_1, \theta_2, \dots, \theta_{n-1}}(0) = F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})$$

such that

$$(3) \quad f_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\lambda) = \mu_{n, \vec{F}_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})}$$

and

$$\|f_{\theta_1, \theta_2, \dots, \theta_{n-1}} - g_{\theta_1, \theta_2, \dots, \theta_{n-1}}\|_1 < \varepsilon_n.$$

Now let

$$F_n(\theta_1, \theta_2, \dots, \theta_n) = f_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\theta_n)$$

and

$$G_n(\theta_1, \theta_2, \dots, \theta_n) = G_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})$$

$$+ (g_{\theta_1, \theta_2, \dots, \theta_{n-1}}(\theta_n) - F_{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})),$$

then (1) holds true with $(n-1)$ replaced by n in view of (2) and (3). Clearly $(G_i)_{i=0}^n$ is a Hardy martingale and we can estimate

$$\|(F_n - F_{n-1}) - (G_n - G_{n-1})\|_1$$

$$= \int_{\mathbb{T}^{n-1}} \|f_{\theta_1, \theta_2, \dots, \theta_{n-1}} - g_{\theta_1, \theta_2, \dots, \theta_{n-1}}\|_1 d\lambda^{n-1}(\theta_1, \theta_2, \dots, \theta_{n-1})$$

$$< \varepsilon_n.$$

This proves Theorem (C). \square

With Theorem (C) at our disposition we now can harvest Corollary (D) without any further effort:

Proof of Corollary (D). Let $(M_n)_{n=0}^\infty$ be an L^1 -bounded plurisubharmonic martingale (i.e., $\sup_n \|M_n\|_1 < \infty$). Apply Theorem (C) to find a representation $(F_n)_{n=0}^\infty$ of $(M_n)_{n=0}^\infty$ on $\mathbb{T}^{\mathbb{N}}$ such that F_n depends only on the first n coordinates and a Hardy martingale $(G_n)_{n=0}^\infty$ such that, for $n \in \mathbb{N}$,

$$\|(F_n - F_{n-1}) - (G_n - G_{n-1})\|_1 < 2^{-n}.$$

Clearly the process $(F_n - G_n)_{n=0}^\infty$ converges almost surely by the Borel-Cantelli lemma and $(G_n)_{n=0}^\infty$ is L^1 -bounded.

If X has the analytic Radon-Nikodym property then $(G_n)_{n=0}^\infty$ converges almost surely [Gar] and therefore $(F_n)_{n=0}^\infty$ does so too.

Conversely if L^1 -bounded plurisubharmonic martingales converge almost surely then in particular L^1 -bounded analytic martingales do so, whence by Edgar's theorem [E2] X has the analytic Radon-Nikodym property. \square

To end this section we give an application of Theorem (C). We say that a complex Banach space X is an AMT (resp. HMT resp. PSH-MT) space if for some $0 < p < \infty$ there is a constant C_p such that for every X -valued analytic (resp. Hardy, resp. plurisubharmonic) martingale $(M_n)_{n=0}^N$ and for every predictable process $(V_n)_{n=1}^N$ bounded in absolute value by 1 we have the following estimate on the martingale transform:

$$(1) \quad \left\| \sum_{n=1}^N V_n (M_n - M_{n-1}) \right\|_p \leq C_p \|M_n\|_p.$$

We refer to [Gar] for a discussion of these concepts. For example D. J. H. Garling proved that $L^1([0, 1])$ is an HMP space [Gar, Theorem 10] while G. Pisier [P] has shown that the space of trace class operators on l^2 fails to be an AMT space. Let us also note that a martingale transform of an analytic (resp. Hardy, resp. plurisubharmonic) martingale is an analytic (resp. Hardy, resp. plurisubharmonic) martingale.

It has been shown by Xu [X] that the concepts of AMT and HMT spaces coincide. Xu's result combined with Theorem (C) gives the following result:

Theorem V.3. *The concepts of AMT, HMT and PSH-MT spaces coincide.*

REFERENCES

- [Bo-Da] J. Bourgain and W. J. Davis, *Martingale transforms and complex uniform convexity*, Trans. Amer. Math. Soc. **294** (1986), 501–515.
- [Bu-Da] A. V. Bukhvalov and A. A. Danilevich, *Boundary properties of analytic and harmonic functions with values in Banach space*, Mat. Zametki **31** (1982), no. 2, 203–214; English transl. in Math. Notes **31** (1982), 104–110.
- [Ch] S. B. Chae, *Holomorphy and calculus in normed spaces*, Pure Appl. Math., Dekker, New York, 1985.
- [Co] D. Cohn, *Measure theory*, Birkhäuser, Stuttgart, 1980.
- [D-G-T] W. J. Davis, D. H. J. Garling, and N. Tomczak-Jaegermann, *The complex convexity of complex quasi-normed linear spaces*, J. Funct. Anal. **55** (1984), 110–150.

- [Dow] P. N. Dowling, *Representation operators and the analytic Radon-Nikodym property in Banach spaces*, Proc. Roy. Irish. Acad. **85A** (1985), 143–150.
- [Du] R. Durrett, *Brownian motion and martingales in analysis*, Wadsworth, Belmont, Calif., 1984.
- [E1] G. A. Edgar, *Complex martingale convergence*, Lecture Notes in Math., vol. 1116, Springer-Verlag, Berlin and New York, 1985, pp. 38–59.
- [E2] —, *Analytic martingale convergence*, J. Funct. Anal. **69** (1986), 268–280.
- [E3] —, *Extremal integral representations*, J. Funct. Anal. **23** (1976), 145–161.
- [Gam] T. W. Gamelin, *Uniform algebras and Jensen measures*, Lecture Notes Series, vol. 32, Cambridge Univ. Press, Cambridge, 1978.
- [Gar] D. J. H. Garling, *On martingales with values in a complex Banach space*, Proc. Cambridge Philos. Soc. **104** (1988), 399–406.
- [G-M] N. Ghoussoub and B. Maurey, *Plurisubharmonic martingales and barriers in complex quasi-Banach spaces*, Preprint, 1988.
- [H-J] J. Hoffmann-Jorgensen, *The theory of analytic spaces*, Lecture Notes Ser. in Math., Aarhus Univ., Aarhus, 1970.
- [K] S. G. Krantz, *Function theory of several complex variables*, Pure and Appl. Math. Ser., Wiley, New York, 1982.
- [L-T] J. Lindenstrauss and L. Trafriri, *Classical Banach spaces. I*, Ergebnisse der Math. 92, Springer, 1977.
- [P] G. Pisier, Personal communication.
- [Ra] M. Range, *Holomorphic functions and integrable representations in several complex variables*, Springer-Verlag, New York, 1986.
- [Schw1] L. Schwartz, *Disintegration of measures*, Tata Inst. Fund. Res. Stud. Math., Bombay, 1976.
- [Schw2] —, *Radon measures on arbitrary topological spaces and cylindrical measures*, Tata Inst. Fund. Res., Oxford Univ. Press, Oxford, 1973.
- [X] Q. H. Xu, Personal communication.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PARIS VII, PARIS, FRANCE

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WUHAN, WUHAN, CHINA

INSTITUT FÜR MATHEMATIK, JOHANNES KEPLER UNIVERSITÄT, A-4040 LINZ, AUSTRIA