

Approximation of Stochastic Partial Differential Equations by a Kernel-based Collocation Method

Igor Cialenco, Gregory E. Fasshauer and Qi Ye*

igor@math.iit.edu, fasshauer@iit.edu, qye3@iit.edu

Department of Applied Mathematics
Illinois Institute of Technology
10 West 32nd Str, Bld E1, Room 208
Chicago, Illinois, USA 60616

First Circulated: August 21, 2011

This Version: May 11, 2012

Forthcoming in *International Journal of Computer Mathematics*

Abstract

In this paper we present the theoretical framework needed to justify the use of a kernel-based collocation method (meshfree approximation method) to estimate the solution of high-dimensional stochastic partial differential equations (SPDEs). Using an implicit time stepping scheme, we transform stochastic parabolic equations into stochastic elliptic equations. Our main attention is concentrated on the numerical solution of the elliptic equations at each time step. The estimator of the solution of the elliptic equations is given as a linear combination of reproducing kernels derived from the differential and boundary operators of the SPDE centered at collocation points to be chosen by the user. The random expansion coefficients are computed by solving a random system of linear equations. Numerical experiments demonstrate the feasibility of the method.

Keywords: kernel-based collocation; numerical solutions; stochastic partial differential equation; reproducing kernel; Matérn function; Gaussian process.

Mathematics Subject Classification (2010): 46E22 65D05 60G15 60H15 65N35.

*Corresponding author

1 Introduction

Stochastic partial differential equations (SPDEs) frequently arise from applications in areas such as physics, engineering and finance. However, in many cases it is difficult to derive an explicit form of their solution. Moreover, current numerical algorithms often show limited success for high-dimensional problems or in complex domains – even for deterministic partial differential equations. The *kernel-based* approximation method (*meshfree* approximation method [4, 11, 21]) is a relatively new numerical tool for the solutions of high-dimensional problems. In this paper we apply – to our knowledge for the first time – such a kernel-based collocation method to construct numerical estimators for stochastic partial differential equations. Galerkin-type approximation methods based on the eigenvalues and eigenfunctions of the underlying differential operator are currently very popular for the numerical solution of SPDEs [7, 17, 14]. With the kernel-based meshfree collocation method introduced here explicit knowledge of these eigenvalues and eigenfunctions is not required since the kernels can be directly obtained as Green kernels of the differential operators [10, 9]. Stochastic collocation methods using a polynomial basis are also frequently found in the literature [2, 18]. These methods usually require the collocation points to lie on a regular grid. In our method the collocation points can be placed at rather arbitrarily scattered locations. This allows for the use of either deterministic or random designs such as, e.g., uniform or Sobol’ points, but also for adaptively chosen locations. In this paper we do not study the design aspect of our algorithm, but reserve this important aspect for future work. Another advantage of using a meshfree method is its ability – also not explicitly pursued here – to deal with problems on a complex domain $\mathcal{D} \subset \mathbb{R}^d$, $d \geq 1$, by using appropriately placed collocation points. Another advantage of this method is its high efficiency, in the sense that once certain matrices are inverted and factored we can compute, essentially for free, the value of the approximated solution at any point in the spatial domain and at any event from sample space. In particular the method is suitable for simulation of a large number of sample paths of the solution. In this article we present only a general framework for this new numerical method and prove weak convergence of the corresponding schemes. We conclude the paper with a numerical implementation of this method applied to a one-dimensional stochastic heat equation with Dirichlet boundary conditions driven by an additive space-time white noise (colored in space). Much more details, as well as some of the aspects just mentioned, will be discussed in Qi Ye’s Ph.D. thesis [23] or in future publications.

1.1 The method in a nutshell

Assume that \mathcal{D} is a *regular* open bounded domain in \mathbb{R}^d (see Appendix B), and let \mathcal{H} be a separable Hilbert space of functions defined on \mathcal{D} . Also, let $(\Omega_W, \mathcal{F}_W, \{\mathcal{F}_t\}, \mathbb{P}_W)$ be a stochastic basis with the usual assumptions. We consider the following parabolic Itô

equation

$$\begin{cases} dU_t = \mathcal{A}U_t dt + \sigma dW_t, & \text{in } \mathcal{D}, \quad 0 < t < T, \\ BU_t = 0, & \text{on } \partial\mathcal{D}, \\ U_0 = u_0, \end{cases} \quad (1.1)$$

where \mathcal{A} is a linear elliptic operator in \mathcal{H} , B is a boundary operator for Dirichlet or Neumann boundary conditions, $u_0 \in \mathcal{H}$, and W is a Wiener process in \mathcal{H} , with mean zero and spatial covariance function R given by $\mathbb{E}(W(t, \mathbf{x})W(s, \mathbf{y})) = \min\{t, s\}R(\mathbf{x}, \mathbf{y})$, $\mathbf{x}, \mathbf{y} \in \mathcal{D}$, $t, s > 0$, and $\sigma > 0$ (see for instance [5]).

We assume that equation (1.1) has a unique solution $U \in L_2(\Omega_W \times (0, T); \mathcal{H})$.

The proposed numerical method for solving a general SPDE of the form (1.1) can be described as follows:

- S1) Discretize (1.1) in time by the implicit Euler scheme at equally spaced time points $0 = t_0 < t_1 < \dots < t_n = T$,

$$U_{t_j} - U_{t_{j-1}} = \mathcal{A}U_{t_j} \delta t + \sigma \delta W_j, \quad j = 1, \dots, n, \quad (1.2)$$

where $\delta t := t_j - t_{j-1}$ and $\delta W_j := W_{t_j} - W_{t_{j-1}}$.

- S2) Since it follows from (1.2) and the definition of Brownian motion that the noise increment δW_j at each time instance t_j is independent from the solution $U_{t_{j-1}}$ at the previous step, we simulate the Gaussian field with covariance structure $R(\mathbf{x}, \mathbf{y})$ at a finite collection of predetermined *collocation points*

$$X_{\mathcal{D}} := \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathcal{D}, \quad X_{\partial\mathcal{D}} := \{\mathbf{x}_{N+1}, \dots, \mathbf{x}_{N+M}\} \subset \partial\mathcal{D}.$$

- S3) Let the differential operator $P := I - \delta t \mathcal{A}$, and the noise term $\xi := \sigma \delta W_j$. Since ξ is a Gaussian field with $\mathbb{E}(\xi_{\mathbf{x}}) = 0$ and $\text{Cov}(\xi_{\mathbf{x}} \xi_{\mathbf{y}}) = \sigma^2 \delta t R(\mathbf{x}, \mathbf{y})$, equation (1.2) together with the corresponding boundary condition becomes an elliptic SPDE of the form

$$\begin{cases} Pu = f + \xi, & \text{in } \mathcal{D}, \\ Bu = 0, & \text{on } \partial\mathcal{D}, \end{cases} \quad (1.3)$$

where $u := U_{t_j}$ is seen as an unknown part and $f := U_{t_{j-1}}$ and ξ are viewed as given parts. We will solve for u using a kernel-based collocation method written as

$$u(\mathbf{x}) \approx \hat{u}(\mathbf{x}) := \sum_{k=1}^N c_k P_2 K^*(\mathbf{x}, \mathbf{x}_k) + \sum_{k=1}^M c_{N+k} B_2 K^*(\mathbf{x}, \mathbf{x}_{N+k}), \quad \mathbf{x} \in \mathcal{D}, \quad (1.4)$$

where K is a *reproducing kernel* and the integral-type kernels K^* , $P_2 K^*$, $B_2 K^*$ are defined in Lemmas 2.1 and B.2. The unknown random coefficients $\mathbf{c} := (c_1, \dots, c_{N+M})^T$ are obtained by solving a random system of linear equations (with constant deterministic system matrix and different random right-hand side) at each time step. Details are provided in Section 3.

S4) Repeat S2 and S3 for all $j = 1, \dots, n$.

Obviously, many other – potentially better – time stepping schemes could be applied here. However, as mentioned earlier, we focus mainly on step S3 and are for the time being content with using the implicit Euler scheme. Naturally, the rate of convergence of the above numerical scheme depends on the size of the time step δt , and the *fill distance* $h_X := \sup_{\mathbf{x} \in \mathcal{D}} \min_{\mathbf{x}_k \in X_{\mathcal{D}} \cup X_{\partial \mathcal{D}}} \|\mathbf{x} - \mathbf{x}_k\|_2$ of the collocation points. We support this statement empirically in Section 4. We should mention that even for deterministic time-dependent PDEs to find the exact rates of convergence of kernel-based methods is a delicate and nontrivial question, only recently solved in [12]. We will address this question in the case of SPDEs in future works.

The fundamental building block of our mesh-free method is the reproducing kernel $K : \mathcal{D} \times \mathcal{D} \rightarrow \mathbb{R}$ and its reproducing-kernel Hilbert space $H_K(\mathcal{D})$ (see Appendix A for more details). By the very nature of such a kernel-based method, the approximate solution U_{t_j} , $j = 1, \dots, n$, must live in $H_K(\mathcal{D})$. Thus, we make the following standing assumption throughout the paper:

The solution U of the original equation (1.1) lies in a Hilbert space \mathcal{H} which can be embedded in the reproducing kernel Hilbert space $H_K(\mathcal{D})$.

Usually, \mathcal{H} is a Sobolev space $\mathcal{H}^m(\mathcal{D})$, for some positive m . In this case it is possible to choose an appropriate kernel K such that the above embedding assumption is satisfied. For a general discussion of existence and uniqueness of the solution of problem (1.1) see, e.g., [19, 6, 5].

2 Reproducing-kernel collocation method for Gaussian processes

In this section we briefly review the standard kernel-based approximation method for high-dimensional interpolation problems. However, since we will later be interested in solving a stochastic PDE, we present the following material mostly from the stochastic point of view. For further discussion of this method we refer the reader to the recent survey papers [20, 8] and references therein.

Assume that the function space $H_K(\mathcal{D})$ is a reproducing-kernel Hilbert space with norm $\|\cdot\|_{K, \mathcal{D}}$ and its reproducing kernel $K \in C(\overline{\mathcal{D}} \times \overline{\mathcal{D}})$ is symmetric positive definite (see Appendix A.1). Given the data values $\{y_j\}_{j=1}^N \subset \mathbb{R}$ at the collocation points $X_{\mathcal{D}} := \{\mathbf{x}_j\}_{j=1}^N \subset \mathcal{D}$ of an unknown function $u \in H_K(\mathcal{D})$, i.e.,

$$y_j = u(\mathbf{x}_j), \quad \mathbf{x}_j = (x_{1,j}, \dots, x_{d,j}) \in \mathcal{D} \subset \mathbb{R}^d, \quad j = 1, \dots, N,$$

the goal is to find *an optimal estimator* from $H_K(\mathcal{D})$ that interpolates these data.

Definition 2.1 ([3, Definition 3.28]). A stochastic process $S : \mathcal{D} \times \Omega \rightarrow \mathbb{R}$ is said to be *Gaussian* with mean $\mu : \mathcal{D} \rightarrow \mathbb{R}$ and covariance kernel $\Phi : \mathcal{D} \times \mathcal{D} \rightarrow \mathbb{R}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ if, for any pairwise distinct points $X_{\mathcal{D}} := \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathcal{D}$, the random vector $\mathbf{S} := (S_{\mathbf{x}_1}, \dots, S_{\mathbf{x}_N})^T$ is a multi-normal random variable on $(\Omega, \mathcal{F}, \mathbb{P})$ with mean $\boldsymbol{\mu}$ and covariance matrix Φ , i.e., $\mathbf{S} \sim \mathcal{N}(\boldsymbol{\mu}, \Phi)$, where $\boldsymbol{\mu} := (\mu(\mathbf{x}_1), \dots, \mu(\mathbf{x}_N))^T$ and $\Phi := (\Phi(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}$.

2.1 Data fitting problems via deterministic interpolation and simple kriging

In the *deterministic formulation* of kernel interpolation we solve an optimization problem by minimizing the reproducing-kernel norm subject to interpolation constraints, i.e.,

$$\hat{u}_K = \arg \min_{u \in \mathbf{H}_K(\mathcal{D})} \{\|u\|_{K,\mathcal{D}} \text{ s.t. } u(\mathbf{x}_j) = y_j, j = 1, \dots, N\}.$$

In this case, the minimum norm interpolant (also called the collocation solution) $\hat{u}_K(\mathbf{x})$ is a linear combination of “shifts” of the reproducing kernel K ,

$$\hat{u}_K(\mathbf{x}) := \sum_{k=1}^N c_k K(\mathbf{x}, \mathbf{x}_k), \quad \mathbf{x} \in \mathcal{D}, \quad (2.1)$$

where the coefficients $\mathbf{c} := (c_1, \dots, c_N)^T$ are obtained by solving the following system of linear equations

$$\mathbf{K}\mathbf{c} = \mathbf{y}_0, \quad (2.2)$$

with $\mathbf{K} := (K(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}$ and $\mathbf{y}_0 := (y_1, \dots, y_N)^T$.

For simple kriging, i.e., in the *stochastic formulation*, we let S be a Gaussian process with mean 0 and covariance kernel K on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Kriging is based on the modeling assumption that u is a realization of the Gaussian field S . The data values y_1, \dots, y_N are then realizations of the random variables $S_{\mathbf{x}_1}, \dots, S_{\mathbf{x}_N}$. The optimal unbiased predictor of $S_{\mathbf{x}}$ based on \mathbf{S} is equal to

$$\hat{U}_{\mathbf{x}} := \sum_{k=1}^N c_k(\mathbf{x}) S_{\mathbf{x}_k} = \arg \min_{U \in \text{span}\{S_{\mathbf{x}_j}\}_{j=1}^N} \mathbb{E}|U - S_{\mathbf{x}}|^2,$$

where the coefficients $\mathbf{c}(\mathbf{x}) := (c_1(\mathbf{x}), \dots, c_N(\mathbf{x}))^T$ are given by

$$\mathbf{c}(\mathbf{x}) = \mathbf{K}^{-1} \mathbf{k}(\mathbf{x})$$

with $\mathbf{k}(\mathbf{x}) := (K(\mathbf{x}, \mathbf{x}_1), \dots, K(\mathbf{x}, \mathbf{x}_N))^T$ and the same matrix \mathbf{K} as above. We can also compute that

$$\mathbb{E}(\hat{U}_{\mathbf{x}} | S_{\mathbf{x}_1} = y_1, \dots, S_{\mathbf{x}_N} = y_N) = \hat{u}_K(\mathbf{x}).$$

Note that, in the kriging approach we consider only the values of the stochastic process S at the collocation points, and view the obtained vector as a random variable. However, if we

view S as a real function, then $\mathbb{P}(S \in \mathbf{H}_K(\mathcal{D})) = 0$ by [16, Theorem 7.3]. A simple example for this fact is given by the scalar Brownian motion defined in the domain $\mathcal{D} := (0, 1)$ (see, e.g., [10, Example 5.1]). This means that it is difficult to apply the kriging formulation to PDE problems. Next we will introduce a new stochastic data fitting approach that will subsequently allow us to perform kernel-based collocation for stochastic PDEs.

2.2 Data fitting problems via a new stochastic approach

From now on we will view the separable reproducing-kernel Hilbert space $\mathbf{H}_K(\mathcal{D})$ as a sample space and its Borel σ -field $\mathcal{B}(\mathbf{H}_K(\mathcal{D}))$ as a σ -algebra to set up the probability spaces so that the stochastic process $S_{\mathbf{x}}(\omega) := \omega(\mathbf{x})$ is Gaussian. We use the techniques of [13, 16] to verify Lemma 2.1, which is a restatement of [16, Theorem 7.2]. This theoretical result is a generalized form of Wiener measure defined on the measurable space $(\mathbf{C}[0, \infty), \mathcal{B}(\mathbf{C}[0, \infty)))$, called canonical space, such that the coordinate mapping process $W_x(\omega) := \omega(x)$ is a Brownian motion (see, for instance, [15], Chapter 2).

Lemma 2.1. *Let the positive definite kernel $K \in \mathbf{C}(\overline{\mathcal{D}} \times \overline{\mathcal{D}})$ be the reproducing kernel of the reproducing-kernel Hilbert space $\mathbf{H}_K(\mathcal{D})$. Given a function $\mu \in \mathbf{H}_K(\mathcal{D})$ there exists a probability measure \mathbb{P}^μ defined on $(\Omega_K, \mathcal{F}_K) := (\mathbf{H}_K(\mathcal{D}), \mathcal{B}(\mathbf{H}_K(\mathcal{D})))$ such that $S_{\mathbf{x}}(\omega) := \omega(\mathbf{x})$ is Gaussian with mean μ and covariance kernel $\overset{*}{K}$ on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$, where the integral-type kernel $\overset{*}{K}$ of K is given by*

$$\overset{*}{K}(\mathbf{x}, \mathbf{y}) := \int_{\mathcal{D}} K(\mathbf{x}, \mathbf{z})K(\mathbf{y}, \mathbf{z})d\mathbf{z}, \quad \mathbf{x}, \mathbf{y} \in \mathcal{D}.$$

Moreover, the process S has the following expansion

$$S_{\mathbf{x}} = \sum_{k=1}^{\infty} \zeta_k \sqrt{\lambda_k} e_k(\mathbf{x}), \quad \mathbf{x} \in \mathcal{D}, \quad \mathbb{P}^\mu\text{-a.s.},$$

where $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ are the eigenvalues and eigenfunctions of the reproducing kernel K , and ζ_k are independent Gaussian random variables with mean $\hat{\mu}_k := \langle \mu, \sqrt{\lambda_k} e_k \rangle_{K, \mathcal{D}}$ and variance λ_k , $k \in \mathbb{N}$.

Before we prove Lemma 2.1 we remark that we have introduced the integral-type kernel $\overset{*}{K}$ for convenience only. As seen later, in order to “match the spaces”, any other kernel that “dominates” K (in the sense of [16]) could play the role of the integral-type kernel $\overset{*}{K}$.

Proof. We first consider the case when $\mu = 0$. There exist countably many independent standard normal random variables $\{\xi_k\}_{k=1}^{\infty}$ on a probability space $(\Omega_\xi, \mathcal{F}_\xi, \mathbb{P}_\xi)$, i.e., $\xi_k \sim i.i.d. \mathcal{N}(0, 1)$, $k \in \mathbb{N}$. Let $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ be the eigenvalues and eigenfunctions of the reproducing kernel K as in Theorem A.1. We define $S := \sum_{k=1}^{\infty} \xi_k \lambda_k e_k$ \mathbb{P}_ξ -a.s. Note that S is Gaussian with mean 0 and covariance kernel $\overset{*}{K}$. Since $\mathbb{E}(\sum_{k=1}^{\infty} \xi_k^2 \lambda_k) \leq \sum_{k=1}^{\infty} \text{Var}(\xi_k) \lambda_k = \sum_{k=1}^{\infty} \lambda_k < \infty$ indicates that $\sum_{k=1}^{\infty} |\xi_k \sqrt{\lambda_k}|^2 < \infty$ \mathbb{P}_ξ -a.s., Theorem A.1

shows that $S(\cdot, \omega) \in \mathbf{H}_K(\mathcal{D})$ \mathbb{P}_ξ -a.s. Therefore S is a measurable map from $(\Omega_\xi, \mathcal{F}_\xi)$ into $(\Omega_K, \mathcal{F}_K)$ by [3, Chapter 4.3.1] and [16, Lemma 2.1]. On the other hand, the probability measure $\mathbb{P}^0 := \mathbb{P}_\xi \circ S^{-1}$ (also called the *image measure* of \mathbb{P}_ξ under S) is well defined on $(\Omega_K, \mathcal{F}_K)$, i.e., $\mathbb{P}^0(A) := \mathbb{P}_\xi(S^{-1}(A))$ for each $A \in \mathcal{F}_K$. Hence, S is also a Gaussian process with mean 0 and covariance kernel $\overset{*}{K}$ on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^0)$.

Let $S^\mu := S + \mu$ on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^0)$. Then $\mathbb{E}(S^\mu) = \mathbb{E}(S) + \mu(\mathbf{x})$ and $\text{Cov}(S^\mu, S^\mu) = \text{Cov}(S, S)$ with respect to \mathbb{P}^0 . We define a new probability measure \mathbb{P}^μ by $\mathbb{P}^\mu(A) := \mathbb{P}^0(A - \mu)$ for each $A \in \mathcal{F}_K$. It is easy to check that $\Omega_K + \mu = \mathbf{H}_K(\Omega) = \Omega_K$ and $\{\mu + A : A \in \mathcal{F}_K\} = \mathcal{B}(\mathbf{H}_K(\Omega)) = \mathcal{F}_K$. Thus S is Gaussian with mean μ and covariance kernel $\overset{*}{K}$ on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$.

Moreover, since $\mu \in \mathbf{H}_K(\Omega)$, it can be expanded in the form $\mu = \sum_{k=1}^{\infty} \hat{\mu}_k \sqrt{\lambda_k} e_k$, where $\hat{\mu}_k := \langle \mu, \sqrt{\lambda_k} e_k \rangle_{K, \mathcal{D}}$, so that $S^\mu = \sum_{k=1}^{\infty} (\hat{\mu}_k + \sqrt{\lambda_k} \xi_k) \sqrt{\lambda_k} e_k$. But then $\zeta_k \sim \hat{\mu}_k + \sqrt{\lambda_k} \xi_k \sim \mathcal{N}(\hat{\mu}_k, \lambda_k)$ are independent on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$. \square

According to [3, Theorem 4.91], we can also verify that the random variable $V_f(\omega) := \langle \omega, f \rangle_{K, \mathcal{D}}$, $f \in \mathbf{H}_K(\mathcal{D})$, is a scalar normal variable on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$, i.e.,

$$V_f(\omega) \sim \mathcal{N}(m_f, \sigma_f^2), \quad \omega \in \Omega_K = \mathbf{H}_K(\mathcal{D}),$$

where $m_f := \langle \mu, f \rangle_{K, \mathcal{D}}$ and $\sigma_f := \|f\|_{L_2(\mathcal{D})}$. Therefore the probability measure \mathbb{P}^μ defined in Lemma 2.1 is *Gaussian*.

Let $p_X^\mu : \mathbb{R}^N \rightarrow \mathbb{R}$ be the joint probability density function of $S_{\mathbf{x}_1}, \dots, S_{\mathbf{x}_N}$ defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$. Then it is a normal density function with mean $\boldsymbol{\mu} := (\mu(\mathbf{x}_1), \dots, \mu(\mathbf{x}_N))^T$ and covariance matrix $\overset{*}{K} := (\overset{*}{K}(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}$. In analogy to the kriging formulation we can find the optimal mean function $\hat{\mu} \in \mathbf{H}_K(\mathcal{D})$ fitting the data values $\mathbf{y}_0 := (y_1, \dots, y_N)^T$, i.e.,

$$\hat{\mu} := \overset{*}{\mathbf{k}}^T \overset{*}{K}^{-1} \mathbf{y}_0 = \sup_{\mu \in \mathbf{H}_K(\mathcal{D})} p_X^\mu(\mathbf{y}_0) = \sup_{\mu \in \mathbf{H}_K(\mathcal{D})} \mathbb{P}^\mu(\mathbf{S} = \mathbf{y}_0),$$

where $\overset{*}{\mathbf{k}}(\mathbf{x}) := (\overset{*}{K}(\mathbf{x}, \mathbf{x}_1), \dots, \overset{*}{K}(\mathbf{x}, \mathbf{x}_N))^T$.

We now fix any $\mathbf{x} \in \mathcal{D}$. Straightforward calculation shows that the random variable $S_{\mathbf{x}}$, given $S_{\mathbf{x}_1}, \dots, S_{\mathbf{x}_N}$, defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$ has a conditional probability density function

$$p_{\mathbf{x}}^\mu(v|\mathbf{v}) := \frac{1}{\sigma(\mathbf{x})\sqrt{2\pi}} \exp\left(-\frac{(v - m_{\mathbf{x}}^\mu(\mathbf{v}))^2}{2\sigma(\mathbf{x})^2}\right), \quad v \in \mathbb{R}, \quad \mathbf{v} \in \mathbb{R}^N,$$

where $m_{\mathbf{x}}^\mu(\mathbf{v}) := \mu(\mathbf{x}) + \overset{*}{\mathbf{k}}(\mathbf{x})^T \overset{*}{K}^{-1}(\mathbf{v} - \mathbf{m}^\mu)$, $\mathbf{m}^\mu := (\mu(\mathbf{x}_1), \dots, \mu(\mathbf{x}_N))^T$, and $\sigma(\mathbf{x})^2 := \overset{*}{K}(\mathbf{x}, \mathbf{x}) - \overset{*}{\mathbf{k}}(\mathbf{x})^T \overset{*}{K}^{-1} \overset{*}{\mathbf{k}}(\mathbf{x})$. Then the optimal estimator that maximizes the probability

$$\begin{aligned} & \max_{v \in \mathbb{R}} \mathbb{P}^{\hat{\mu}}(\{\omega \in \Omega_K : \omega(\mathbf{x}) = v \text{ s.t. } \omega(\mathbf{x}_1) = y_1, \dots, \omega(\mathbf{x}_N) = y_N\}) \\ &= \max_{v \in \mathbb{R}} \mathbb{P}^{\hat{\mu}}(S_{\mathbf{x}} = v | S_{\mathbf{x}_1} = y_1, \dots, S_{\mathbf{x}_N} = y_N) \end{aligned}$$

is given by

$$\hat{u}(\mathbf{x}) := \hat{\mu}(\mathbf{x}) = \arg \max_{v \in \mathbb{R}} p_{\mathbf{x}}^{\hat{u}}(v | \mathbf{y}_0).$$

Proposition 2.2. *With the above notations, the following equality holds true*

$$p_{\mathbf{x}}^{\hat{u}}(\hat{u}(\mathbf{x}) | \mathbf{y}_0) = \sup_{v \in \mathbb{R}, \mu \in \mathbf{H}_K(\mathcal{D})} p_{\mathbf{x}}^{\mu}(v | \mathbf{y}_0). \quad (2.3)$$

Moreover, for any $\epsilon > 0$,

$$\sup_{\mu \in \mathbf{H}_K(\mathcal{D})} \mathbb{P}^{\mu}(|\hat{u}(\mathbf{x}) - u(\mathbf{x})| \geq \epsilon) \leq \sup_{\mu \in \mathbf{H}_K(\mathcal{D})} \mathbb{P}^{\mu}(\mathcal{E}_{\mathbf{x}}^{\epsilon}) = \operatorname{erfc}\left(\frac{\epsilon}{\sqrt{2}\sigma(\mathbf{x})}\right), \quad (2.4)$$

where

$$\mathcal{E}_{\mathbf{x}}^{\epsilon} := \{\omega \in \Omega_K : |\omega(\mathbf{x}) - \hat{u}(\mathbf{x})| \geq \epsilon \text{ s.t. } \omega(\mathbf{x}_1) = y_1, \dots, \omega(\mathbf{x}_N) = y_N\}.$$

Identity (2.3) follows by direct evaluations. Consequently, taking into account that S is Gaussian, inequality (2.4) follows also immediately.

Remark 2.1. Instead of giving a deterministic (or strong) error bound for the proposed numerical scheme, we provide a *weak type convergence* of the approximated solution \hat{u} to the true solution u , as stated in Proposition 2.2. In fact, inequality (2.4) can be seen as a confidence interval for the estimator \hat{u} with respect to the probability measure \mathbb{P}^{μ} .

In the next section we generalize this stochastic approach to solve elliptic PDEs.

3 Collocation Method for Elliptic PDEs and SPDEs

We begin by setting up Gaussian processes via reproducing kernels with differential and boundary operators.

Suppose that the reproducing-kernel Hilbert space $\mathbf{H}_K(\mathcal{D})$ is embedded into the Sobolev space $\mathcal{H}^m(\mathcal{D})$ where $m > d/2$. Let $n := \lceil m - d/2 \rceil - 1$. By the Sobolev embedding theorem $\mathcal{H}^m(\mathcal{D}) \subset C^n(\overline{\mathcal{D}})$. When the differential operator P and the boundary operator B have the orders $\mathcal{O}(P) < m - d/2$ and $\mathcal{O}(B) < m - d/2$, then the stochastic processes $PS_{\mathbf{x}}(\omega) := (P\omega)(\mathbf{x})$ and $BS_{\mathbf{x}}(\omega) := (B\omega)(\mathbf{x})$ are well-defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^{\mu})$. According to Lemma B.1, we have $PS = \sum_{k=1}^{\infty} \zeta_k \sqrt{\lambda_k} P e_k$ and $BS = \sum_{k=1}^{\infty} \zeta_k \sqrt{\lambda_k} B e_k$. If $\mu \in \mathbf{H}_K(\mathcal{D}) \subseteq \mathcal{H}^m(\mathcal{D})$, then $P\mu \in C(\overline{\mathcal{D}})$ and $B\mu \in C(\partial\mathcal{D})$. Lemma B.2 implies that $P_1 P_2^* K(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{\infty} \lambda_k^2 P e_k(\mathbf{x}) P e_k(\mathbf{y})$ and $B_1 B_2^* K(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{\infty} \lambda_k^2 B e_k(\mathbf{x}) B e_k(\mathbf{y})$ (here we can use the fact that $\operatorname{Cov}(PS_{\mathbf{x}}, PS_{\mathbf{y}}) = P_{\mathbf{x}} P_{\mathbf{y}} \operatorname{Cov}(S_{\mathbf{x}}, S_{\mathbf{y}})$ and $\operatorname{Cov}(BS_{\mathbf{x}}, BS_{\mathbf{y}}) = B_{\mathbf{x}} B_{\mathbf{y}} \operatorname{Cov}(S_{\mathbf{x}}, S_{\mathbf{y}})$). Applying Lemma 2.1, we can obtain the main theorem for the construction of Gaussian processes via reproducing kernels coupled with differential or boundary operators.

Theorem 3.1. *Suppose that the reproducing kernel Hilbert space $\mathbb{H}_K(\mathcal{D})$ is embedded into the Sobolev space $\mathcal{H}^m(\mathcal{D})$ with $m > d/2$. Further assume that the differential operator P and the boundary operator B have the orders $\mathcal{O}(P) < m - d/2$ and $\mathcal{O}(B) < m - d/2$. Given a function $\mu \in \mathbb{H}_K(\mathcal{D})$ there exists a probability measure \mathbb{P}^μ defined on $(\Omega_K, \mathcal{F}_K) = (\mathbb{H}_K(\mathcal{D}), \mathcal{B}(\mathbb{H}_K(\mathcal{D})))$ (as in Lemma 2.1) such that the stochastic processes PS, BS given by*

$$\begin{aligned} PS_{\mathbf{x}}(\omega) &= PS(\mathbf{x}, \omega) := (P\omega)(\mathbf{x}), & \mathbf{x} \in \mathcal{D} \subset \mathbb{R}^d, & \quad \omega \in \Omega_K = \mathbb{H}_K(\mathcal{D}), \\ BS_{\mathbf{x}}(\omega) &= BS(\mathbf{x}, \omega) := (B\omega)(\mathbf{x}), & \mathbf{x} \in \partial\mathcal{D}, & \quad \omega \in \Omega_K = \mathbb{H}_K(\mathcal{D}), \end{aligned}$$

are jointly Gaussian processes with means $P\mu, B\mu$ and covariance kernels $P_1P_2K^*, B_1B_2K^*$ defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$, respectively. In particular, they can be expanded as

$$PS_{\mathbf{x}} = \sum_{k=1}^{\infty} \zeta_k \sqrt{\lambda_k} P e_k(\mathbf{x}), \quad \mathbf{x} \in \mathcal{D}, \quad \text{and} \quad BS_{\mathbf{x}} = \sum_{k=1}^{\infty} \zeta_k \sqrt{\lambda_k} B e_k(\mathbf{x}), \quad \mathbf{x} \in \partial\mathcal{D}, \quad \mathbb{P}^\mu\text{-a.s.},$$

where $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ are the eigenvalues and eigenfunctions of the reproducing kernel K and their related Fourier coefficients are the independent normal variables $\zeta_k \sim \mathcal{N}(\hat{\mu}_k, \lambda_k)$ and $\hat{\mu}_k := \langle \mu, \sqrt{\lambda_k} e_k \rangle_{K, \mathcal{D}}$, $k \in \mathbb{N}$.

Corollary 3.2. *Suppose all notations and conditions are as in Theorem 3.1. Given collocation points $X_{\mathcal{D}} := \{\mathbf{x}_j\}_{j=1}^N \subset \mathcal{D}$ and $X_{\partial\mathcal{D}} := \{\mathbf{x}_{N+j}\}_{j=1}^M \subset \partial\mathcal{D}$, the random vector $\mathbf{S}_{PB} := (PS_{\mathbf{x}_1}, \dots, PS_{\mathbf{x}_N}, BS_{\mathbf{x}_{N+1}}, \dots, BS_{\mathbf{x}_{N+M}})^T$ defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$ has a multivariate normal distribution with mean \mathbf{m}_{PB}^μ and covariance matrix \mathbb{K}_{PB}^* , i.e.,*

$$\mathbf{S}_{PB} \sim \mathcal{N}(\mathbf{m}_{PB}^\mu, \mathbb{K}_{PB}^*),$$

where $\mathbf{m}_{PB}^\mu := (P\mu(\mathbf{x}_1), \dots, P\mu(\mathbf{x}_N), B\mu(\mathbf{x}_{N+1}), \dots, B\mu(\mathbf{x}_{N+M}))^T$ and

$$\mathbb{K}_{PB}^* := \begin{pmatrix} (P_1P_2K^*(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}, & (P_1B_2K^*(\mathbf{x}_j, \mathbf{x}_{N+k}))_{j,k=1}^{N,M} \\ (B_1P_2K^*(\mathbf{x}_{N+j}, \mathbf{x}_k))_{j,k=1}^{M,N}, & (B_1B_2K^*(\mathbf{x}_{N+j}, \mathbf{x}_{N+k}))_{j,k=1}^{M,M} \end{pmatrix}.$$

Remark 3.1. While the covariance matrix \mathbb{K}_{PB}^* may be singular, it is always positive semi-definite and therefore always has a pseudo-inverse $\mathbb{K}_{PB}^{*\dagger}$.

Using Corollary 3.2, we can compute the joint probability density function p_X^μ of \mathbf{S}_{PB} defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$. In the same way, we can also get the joint density function p_J^μ of $(S_{\mathbf{x}}, \mathbf{S}_{PB})$ defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$. By Bayes' rule, we can obtain the conditional probability density function of the random variable $S_{\mathbf{x}}$ given \mathbf{S}_{PB} .

Corollary 3.3. *We follow the notations of Corollary 3.2. For any fixed $\mathbf{x} \in \mathcal{D}$, the random variable $S_{\mathbf{x}}$ given \mathbf{S}_{PB} defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$ has a conditional probability density function*

$$p_{\mathbf{x}}^\mu(v|\mathbf{v}) := \frac{p_J^\mu(v, \mathbf{v})}{p_X^\mu(\mathbf{v})} = \frac{1}{\sigma(\mathbf{x})\sqrt{2\pi}} \exp\left(-\frac{(v - m_{\mathbf{x}}^\mu(\mathbf{v}))^2}{2\sigma(\mathbf{x})^2}\right), \quad v \in \mathbb{R}, \quad \mathbf{v} \in \mathbb{R}^{N+M},$$

where

$$\begin{aligned} m_{\mathbf{x}}^{\mu}(\mathbf{v}) &:= \mu(\mathbf{x}) + \mathbf{k}_{PB}(\mathbf{x})^T \mathbf{K}_{PB}^{\dagger*}(\mathbf{v} - \mathbf{m}_{PB}^{\mu}), \\ \sigma(\mathbf{x})^2 &:= \overset{*}{K}(\mathbf{x}, \mathbf{x}) - \mathbf{k}_{PB}(\mathbf{x})^T \mathbf{K}_{PB}^{\dagger*} \mathbf{k}_{PB}(\mathbf{x}), \\ \mathbf{k}_{PB}(\mathbf{x}) &:= (P_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_1), \dots, P_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_N), B_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_{N+1}), \dots, B_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_{N+M}))^T. \end{aligned}$$

In particular, given the real observation $\mathbf{y} := (y_1, \dots, y_{N+M})^T$, $S_{\mathbf{x}}$ conditioned on $\mathbf{S}_{PB} = \mathbf{y}$ has the probability density $p_{\mathbf{x}}^{\mu}(\cdot|\mathbf{y})$.

This corollary is similar to the features of Gaussian conditional distributions (see [13, Theorem 9.9]).

3.1 Elliptic deterministic PDEs

Suppose that $u \in \mathbf{H}_K(\mathcal{D})$ is the unique solution of the deterministic elliptic PDE

$$\begin{cases} Pu = f, & \text{in } \mathcal{D} \subset \mathbb{R}^d, \\ Bu = g, & \text{on } \partial\mathcal{D}, \end{cases} \quad (3.1)$$

where $f : \mathcal{D} \rightarrow \mathbb{R}$ and $g : \partial\mathcal{D} \rightarrow \mathbb{R}$. Denote by $\{y_j\}_{j=1}^N$ and $\{y_{N+k}\}_{k=1}^M$ the values of f and g at the collocation points $X_{\mathcal{D}}$ and $X_{\partial\mathcal{D}}$, respectively:

$$y_j := f(\mathbf{x}_j), \quad y_{N+k} := g(\mathbf{x}_{N+k}), \quad j = 1, \dots, N, \quad k = 1, \dots, M.$$

From now on we assume that the covariance matrix \mathbf{K}_{PB}^* defined in Corollary 3.2 is nonsingular and we therefore can replace pseudo-inverses with inverses.

Let $\mathbf{y}_0 := (y_1, \dots, y_N, y_{N+1}, \dots, y_{N+M})^T$, and denote by $p_{\mathbf{x}}^{\mu}(\cdot|\cdot)$ the conditional density function defined in Corollary 3.3. We approximate the solution u of (3.1) by the optimal estimator $\hat{u}(\mathbf{x})$ derived in the previous section, i.e., we maximize the conditional probability given the data values \mathbf{y}_0 :

$$u(\mathbf{x}) \approx \hat{u}(\mathbf{x}) = \arg \max_{v \in \mathbb{R}} \sup_{\mu \in \mathbf{H}_K(\mathcal{D})} p_{\mathbf{x}}^{(\mu)}(v|\mathbf{y}_0), \quad \mathbf{x} \in \mathcal{D}.$$

By direct evaluation as in Section 2.2 one finds that

$$\hat{u}(\mathbf{x}) := \mathbf{k}_{PB}(\mathbf{x})^T \mathbf{K}_{PB}^{\dagger*} \mathbf{y}_0, \quad \mathbf{x} \in \mathcal{D},$$

where the basis functions $\mathbf{k}_{PB}(\mathbf{x})$ are defined in Corollary 3.3. Moreover, the estimator $\hat{u} \in \mathbf{H}_K(\mathcal{D})$ fits all the data values: $P\hat{u}(\mathbf{x}_1) = y_1, \dots, P\hat{u}(\mathbf{x}_N) = y_N$ and $B\hat{u}(\mathbf{x}_{N+1}) = y_{N+1}, \dots, B\hat{u}(\mathbf{x}_{N+M}) = y_{N+M}$. This means that we have computed a collocation solution of the PDE (3.1). Also note that \hat{u} can be written as a linear combination of the kernels as in (1.4), i.e.,

$$\hat{u}(\mathbf{x}) = \sum_{k=1}^N c_k P_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_k) + \sum_{k=1}^M c_{N+k} B_2 \overset{*}{K}(\mathbf{x}, \mathbf{x}_{N+k}), \quad \mathbf{x} \in \mathcal{D}, \quad (3.2)$$

where $\mathbf{c} := (c_1, \dots, c_{N+M})^T = \mathbf{K}_{PB}^*^{-1} \mathbf{y}_0 \in \mathbb{R}^{N+M}$.

Finally, we can perform a weak error analysis for $|u(\mathbf{x}) - \hat{u}(\mathbf{x})|$ as in Proposition 2.2, and deduce that

$$\mathbb{P}^\mu(\mathcal{E}_{\mathbf{x}}^\epsilon) = \mathbb{P}^\mu(|S_{\mathbf{x}} - \hat{u}(\mathbf{x})| \geq \epsilon | \mathbf{S}_{PB} = \mathbf{y}_0) = \operatorname{erfc}\left(\frac{\epsilon}{\sqrt{2}\sigma(\mathbf{x})}\right),$$

where $\sigma(\mathbf{x})^2$ is defined in Corollary 3.3, and

$$\mathcal{E}_{\mathbf{x}}^\epsilon := \{\omega \in \Omega_K : |\omega(\mathbf{x}) - \hat{u}(\mathbf{x})| \geq \epsilon \text{ s.t. } P\omega(\mathbf{x}_1) = y_1, \dots, B\omega(\mathbf{x}_{N+M}) = y_{N+M}\}.$$

Because the form of the expression for the variance $\sigma(\mathbf{x})^2$ is analogous to that of the *power function*, we can use the same techniques as in the proofs from [11, 21] to obtain a formula for the order of $\sigma(\mathbf{x})$.

Lemma 3.4.

$$\sigma(\mathbf{x}) = \mathcal{O}(h_X^{m-\rho-d/2}), \quad \mathbf{x} \in \mathcal{D},$$

where $\rho := \max\{\mathcal{O}(P), \mathcal{O}(B)\}$ and h_X is the fill distance of $X_{\mathcal{D}}$ and $X_{\partial\mathcal{D}}$.

Proof. Since there is at least one collocation point $\mathbf{x}_j \in X_{\mathcal{D}} \cup X_{\partial\mathcal{D}}$ such that $\|\mathbf{x} - \mathbf{x}_j\|_2 \leq h_X$ we can use the multivariate Taylor expansion of $\overset{*}{K}(\mathbf{x}, \mathbf{x}_j)$ to introduce the order of $\sigma(\mathbf{x})$, i.e.,

$$\overset{*}{K}(\mathbf{x}, \mathbf{x}_j) = \sum_{|\alpha|, |\beta| < n} \frac{1}{\alpha! \beta!} D_1^\alpha D_2^\beta \overset{*}{K}(\mathbf{x}_j, \mathbf{x}_j) (\mathbf{x} - \mathbf{x}_j)^{\alpha+\beta} + R(\mathbf{x}, \mathbf{x}_j), \quad \alpha, \beta \in \mathbb{N}_0^d,$$

where $R(\mathbf{x}, \mathbf{x}_j) := \sum_{|\alpha|, |\beta| = n} \frac{1}{\alpha! \beta!} D_1^\alpha D_2^\beta \overset{*}{K}(\mathbf{z}_1, \mathbf{z}_2) (\mathbf{x} - \mathbf{x}_j)^{\alpha+\beta}$ for some $\mathbf{z}_1, \mathbf{z}_2 \in \mathcal{D}$ and $n := \lceil m - d/2 \rceil - 1$. The rest of the proof proceeds as in [11, Chapter 14.5] and [21, Chapters 11.3, 16.3]. \square

Using Lemma 3.4 we can deduce the following proposition.

Proposition 3.5. *For any $\epsilon > 0$,*

$$\sup_{\mu \in \mathbf{H}_K(\mathcal{D})} \mathbb{P}^\mu(\mathcal{E}_{\mathbf{x}}^\epsilon) = \mathcal{O}\left(\frac{h_X^{m-\rho-d/2}}{\epsilon}\right), \quad \mathbf{x} \in \mathcal{D},$$

which indicates that

$$\sup_{\mu \in \mathbf{H}_K(\mathcal{D})} \mathbb{P}^\mu(\|u - \hat{u}\|_{L^\infty(\mathcal{D})} \geq \epsilon) \leq \sup_{\mu \in \mathbf{H}_K(\mathcal{D}), \mathbf{x} \in \mathcal{D}} \mathbb{P}^\mu(\mathcal{E}_{\mathbf{x}}^\epsilon) \rightarrow 0, \text{ when } h_X \rightarrow 0.$$

Therefore we say that the estimator \hat{u} converges to the exact solution u of the PDE (3.1) in all probabilities \mathbb{P}^μ when h_X goes to 0.

Sometimes we know only that the solution $u \in \mathcal{H}^m(\mathcal{D})$. In this case, as long as the reproducing kernel Hilbert space is dense in the Sobolev space $\mathcal{H}^m(\mathcal{D})$ with respect to its Sobolev norm, we can still say that \hat{u} converges to u in probability.

3.2 Elliptic stochastic PDEs

Let $\xi : \mathcal{D} \times \Omega_W \rightarrow \mathbb{R}$ be Gaussian with mean 0 and covariance kernel $\Psi : \mathcal{D} \times \mathcal{D} \rightarrow \mathbb{R}$ on the probability space $(\Omega_W, \mathcal{F}_W, \mathbb{P}_W)$. We consider an elliptic PDE driven by a Gaussian additive noise ξ

$$\begin{cases} Pu = f + \xi, & \text{in } \mathcal{D} \subset \mathbb{R}^d, \\ Bu = g, & \text{on } \partial\mathcal{D}, \end{cases} \quad (3.3)$$

and suppose its solution $u \in L_2(\Omega_W; \mathbb{H}_K(\mathcal{D}))$.

Since ξ is a Gaussian process, on some underlying probability space $(\Omega_W, \mathcal{F}_W, \mathbb{P}_W)$ with a known correlation structure, we can simulate the values of ξ at \mathbf{x}_j , $j = 1, \dots, N$. Consequently, we assume that the values $\{y_j\}_{j=1}^N$ and $\{y_{N+k}\}_{k=1}^M$ defined by

$$y_j := f(\mathbf{x}_j) + \xi_{\mathbf{x}_j}, \quad y_{N+k} := g(\mathbf{x}_{N+k}), \quad j = 1, \dots, N, \quad k = 1, \dots, M,$$

are known.

In order to apply the general interpolation framework developed in Section 2.2, we consider the product space

$$\Omega_{KW} := \Omega_K \times \Omega_W, \quad \mathcal{F}_{KW} := \mathcal{F}_K \otimes \mathcal{F}_W, \quad \mathbb{P}_W^\mu := \mathbb{P}^\mu \otimes \mathbb{P}_W.$$

We assume that the random variables defined on the original probability spaces are extended to random variables on the new probability space in the natural way: if random variables $V_1 : \Omega_K \rightarrow \mathbb{R}$ and $V_2 : \Omega_W \rightarrow \mathbb{R}$ are defined on $(\Omega_K, \mathcal{F}_K, \mathbb{P}^\mu)$ and $(\Omega_W, \mathcal{F}_W, \mathbb{P}_W)$, respectively, then

$$V_1(\omega, \tilde{\omega}) := V_1(\omega), \quad V_2(\omega, \tilde{\omega}) := V_2(\tilde{\omega}), \quad \text{for each } \omega \in \Omega_K \text{ and } \tilde{\omega} \in \Omega_W.$$

Note that in this case the random variables have the same probability distributional properties, and they are independent on $(\Omega_{KW}, \mathcal{F}_{KW}, \mathbb{P}_W^\mu)$. This implies that the stochastic processes S , PS , BS and ξ can be extended to the product space $(\Omega_{KW}, \mathcal{F}_{KW}, \mathbb{P}_W^\mu)$ while preserving the original probability distributional properties. In this case, $(y_1, \dots, y_N) \sim \mathcal{N}(\mathbf{f}, \Psi)$, where $\mathbf{f} := (f(\mathbf{x}_1), \dots, f(\mathbf{x}_N))^T$ and $\Psi := (\Psi(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}$ with Ψ being the covariance kernel of ξ . According to Corollary 3.3 we can find the conditional probability density function $p_{\mathbf{x}}^\mu(\cdot|\cdot)$, and the optimal estimator $\hat{u}(\mathbf{x})$

$$\hat{u}(\mathbf{x}) = \arg \max_{v \in \mathbb{R}} \sup_{\mu \in \mathbb{H}_K(\mathcal{D})} p_{\mathbf{x}}^\mu(v|\mathbf{y}_\xi) p_W(\mathbf{y}_\xi), \quad \mathbf{x} \in \mathcal{D},$$

where

$$\begin{aligned} \mathbf{y}_\xi &:= (y_1, \dots, y_{N+M})^T, \quad \mathbf{m}_\xi := \begin{pmatrix} \mathbf{f} \\ 0 \end{pmatrix} \in \mathbb{R}^{N+M}, \quad \Sigma := \begin{pmatrix} \Psi & 0 \\ 0 & 0 \end{pmatrix} \in \mathbb{R}^{(N+M) \times (N+M)}, \\ p_W(\mathbf{v}) &:= \frac{1}{\sqrt{\det(\Sigma)} (2\pi)^{(N+M)/2}} \exp\left(-\frac{1}{2}(\mathbf{v} - \mathbf{m}_\xi)^T \Sigma^\dagger (\mathbf{v} - \mathbf{m}_\xi)\right), \quad \mathbf{v} \in \mathbb{R}^{N+M}. \end{aligned}$$

If $\mathbb{K}_{PB}^* + \Sigma$ is nonsingular, then one optimal solution has the form

$$\hat{u}(\mathbf{x}) := \mathbf{k}_{PB}(\mathbf{x})^T (\mathbb{K}_{PB}^* + \Sigma)^{-1} \mathbf{y}_\xi, \quad \mathbf{x} \in \mathcal{D},$$

where \mathbb{K}_{PB}^* and $\mathbf{k}_{PB}(\mathbf{x})$ are defined in Corollary 3.2 and 3.3.

Similar to the analysis of the error bounds from Section 3.1, we also deduce the following proposition (for more details see [23]).

Proposition 3.6. *Assume $\Psi \in C^{1,1}(\mathcal{D} \times \mathcal{D})$. Then*

$$\lim_{h_X \rightarrow 0} \sup_{\mu \in \mathbb{H}_K(\mathcal{D})} \mathbb{P}_W^\mu (\|u - \hat{u}\|_{L^\infty(\mathcal{D})} \geq \epsilon) = 0, \quad \text{for any } \epsilon > 0.$$

4 Numerical Experiments

We consider the following stochastic heat equation with zero boundary condition

$$\begin{cases} dU_t = \frac{d^2}{dx^2} U_t dt + \sigma dW_{t,i}, & \text{in } \mathcal{D} := (0, 1) \subset \mathbb{R}, \quad 0 < t < T := 1, \\ U_t = 0, & \text{on } \partial\mathcal{D}, \\ U_0(x) = u_0(x) := \sqrt{2} (\sin(\pi x) + \sin(2\pi x) + \sin(3\pi x)), \end{cases} \quad (4.1)$$

driven by two types of space-time white noise (colored in space) W of the form

$$W_{t,i} := \sum_{k=1}^{\infty} W_t^k q_k^i \phi_k, \quad q_k := \frac{1}{k\pi}, \quad \phi_k(x) := \sqrt{2} \sin(k\pi x),$$

where W_t^k , $k \in \mathbb{N}$, is a sequence of independent one-dimensional Brownian motions, and $i = 1, 2$. Note that choosing the larger value of i corresponds to a noise that is smoother in space.

The spatial covariance function $R^i(x, y) = \sum_{k=1}^{\infty} q_k^{2i} \phi_k(x) \phi_k(y)$, $i = 1, 2$, takes the specific forms

$$R^1(x, y) = \min\{x, y\} - xy, \quad 0 < x, y < 1,$$

and

$$R^2(x, y) = \begin{cases} -\frac{1}{6}x^3 + \frac{1}{6}x^3y + \frac{1}{6}xy^3 - \frac{1}{2}xy^2 + \frac{1}{3}xy, & 0 < x < y < 1, \\ -\frac{1}{6}y^3 + \frac{1}{6}xy^3 + \frac{1}{6}x^3y - \frac{1}{2}x^2y + \frac{1}{3}xy, & 0 < y < x < 1. \end{cases}$$

The solution of SPDE (4.1) is given by (for more details see, for instance, [5])

$$U_t(x) = \sum_{k=1}^{\infty} \xi_t^k \phi_k(x), \quad x \in \mathcal{D} := (0, 1), \quad 0 < t < T := 1,$$

where

$$\xi_0^k := \int_{\mathcal{D}} u_0(x) \phi_k(x) dx, \quad \xi_t^k := \xi_0^k e^{-k^2 \pi^2 t} + \frac{\sigma}{q_k^i} \int_0^t e^{k^2 \pi^2 (s-t)} dW_s^k.$$

From this explicit solution we can get that

$$\mathbb{E}(U_t(x)) = \sum_{k=1}^{\infty} \xi_0^k e^{-k^2 \pi^2 t} \phi_k(x), \quad \text{Var}(U_t(x)) = \sum_{k=1}^{\infty} \frac{\sigma^2}{2k^2 \pi^2 q_k^{2i}} (1 - e^{-2k^2 \pi^2 t}) |\phi_k(x)|^2.$$

We discretize the time interval $[0, T]$ with n equal time steps so that $\delta t := T/n$. We also choose the reproducing kernel $K(x, y) := g_{3,2\theta}(x - y)$, where $g_{3,2\theta}$ is the Matérn function with degree $m := 3$ and shape parameter $\theta > 0$ (see Example A.1). As collocation points we select uniform grid points $X_{\mathcal{D}} \subset (0, 1)$ and $X_{\partial\mathcal{D}} := \{0, 1\}$. Let $P := I - \delta t d^2/dx^2$ and $B := I|_{\{0,1\}}$. Using our kernel-based collocation method we can perform the following computations to numerically estimate the sample paths $\hat{u}_j^n \approx U_{t_n}(x_j)$. *Algorithm to solve SPDE (4.1)*:

1. Initialize

- $\hat{\mathbf{u}}^0 := (u_0(x_1), \dots, u_0(x_N))^T$
- $\mathbf{K}_{PB}^* := \begin{pmatrix} (P_1 P_2 K^*(x_j, x_k))_{j,k=1}^{N,N} & (P_1 B_2 K^*(x_j, x_{N+k}))_{j,k=1}^{N,M} \\ (B_1 P_2 K^*(x_{N+j}, x_k))_{j,k=1}^{M,N} & (B_1 B_2 K^*(x_{N+j}, x_{N+k}))_{j,k=1}^{M,M} \end{pmatrix}$
- $\mathbf{B} := \begin{pmatrix} (P_2 K^*(x_j, x_k))_{j,k=1}^{N,N} & (B_2 K^*(x_j, x_{N+k}))_{j,k=1}^{N,M} \end{pmatrix}$
- $\Psi := \sigma^2 \delta t (R(x_j, x_k))_{j,k=1}^{N,N}, \quad \Sigma := \begin{pmatrix} \Psi & 0 \\ 0 & 0 \end{pmatrix}$
- $\mathbf{A} := \mathbf{B}(\mathbf{K}_{PB}^* + \Sigma)^{-1}$

2. Repeat for $j = 1, 2, \dots, n$, i.e., for $t_1, t_2, \dots, t_n = T$

- Simulate $\xi \sim \mathcal{N}(0, \Psi)$
- $\hat{\mathbf{u}}^j := \mathbf{B}(\mathbf{K}_{PB}^* + \Sigma)^{-1} \begin{pmatrix} \hat{\mathbf{u}}^{j-1} + \xi \\ 0 \end{pmatrix} = \mathbf{A} \begin{pmatrix} \hat{\mathbf{u}}^{j-1} + \xi \\ 0 \end{pmatrix}$

Note that in the very last step the matrix \mathbf{A} is pre-computed and can be used for all time steps, and for different sample paths; that makes the proposed algorithm to be quite efficient.

We approximate the mean and variance of $U_t(x)$ by sample mean and sample variance from $s := 10000$ simulated sample paths using the above algorithm, i.e.,

$$\mathbb{E}(U_{t_n}(x_j)) \approx \frac{1}{s} \sum_{k=1}^s \hat{u}_j^n(\omega_k), \quad \text{Var}(U_{t_n}(x_j)) \approx \frac{1}{s} \sum_{i=1}^s \left(\hat{u}_j^n(\omega_i) - \frac{1}{s} \sum_{k=1}^s \hat{u}_j^n(\omega_k) \right)^2.$$

Figure 4.1 shows that the histograms at different values of t and x resemble the theoretical normal distributions. We notice a small shift in the probability distribution function of the solution U , at times closer to zero, and when the noise is equal to W_1 (Figure 4.1, left panel). This shift is due to the fact that W_1 is rougher in space than W_2 .




Figure 4.1: Empirical and theoretical probability distribution of $U_t(x)$ for uniform points $N := 58$ and $M := 2$, equal time steps $n := 800$, $\theta := 26.5$, $\sigma := 1$.

Our use of an implicit time stepping scheme reduces the frequency of the white noise, i.e., $\lim_{\delta t \rightarrow 0} \delta W / \delta t \sim \delta_0$. Consequently, Figure 4.2 shows that the approximate mean is well-behaved but the approximate variance is a little smaller than the exact variance. According to Figure 4.3 we find that this numerical method is convergent as both δt and h_X are refined. Finally, we want to mention that the distribution of collocation points, the shape parameter, and the kernel itself were chosen empirically and based on the authors' experience. As mentioned before, more precise methods are currently not available. A rigorous investigation of these questions, as well as determination of precise rates of convergence is reserved for future work.

5 Final Remarks

This new numerical approach can also be used to approximate systems of elliptic PDEs with vector Gaussian noises ξ_1 and ξ_2 or nonlinear PDEs with Gaussian noise ξ , i.e.,

$$\begin{cases} \mathbf{P}u = \mathbf{f} + \xi_1, & \text{in } \mathcal{D} \subset \mathbb{R}^d, \\ \mathbf{B}u = \mathbf{g} + \xi_2, & \text{on } \partial\mathcal{D}, \end{cases} \quad \text{or} \quad \begin{cases} F(\mathbf{P}u) = \psi(\mathbf{f}, \xi), & \text{in } \mathcal{D} \subset \mathbb{R}^d, \\ G(\mathbf{B}u) = g, & \text{on } \partial\mathcal{D}, \end{cases}$$

where $\mathbf{P} := (P^1, \dots, P^{n_p})^T$ is a vector differential operator and $\mathbf{B} := (B^1, \dots, B^{n_b})^T$ is a vector boundary operator, and $F : \mathbb{R}^{n_p} \rightarrow \mathbb{R}$ and $G : \mathbb{R}^{n_b} \rightarrow \mathbb{R}$ (see [23]).

In addition to the additive noise case discussed here, we can also use the kernel-based collocation method to approximate other well-posed stochastic parabolic equations with




Figure 4.2: Approximate and theoretical mean and standard deviation for uniform points $N := 58$ and $M := 2$, equal time steps $n := 800$, $\theta := 26.5$, $\sigma := 1$.

multiplicative noise, e.g.,

$$\begin{cases} dU_t = \mathcal{A}U_t dt + \psi(U_t) dW_t, & \text{in } \mathcal{D} \subset \mathbb{R}^d, \quad 0 < t < T, \\ BU_t = 0, & \text{on } \partial\mathcal{D}, \\ U_0 = u_0, \end{cases} \quad (5.1)$$

where $\psi : \mathbb{R} \rightarrow \mathbb{R}$. Since $\int_{t_{j-1}}^{t_j} \psi(U_s) dW_s \approx \psi(U_{t_{j-1}}) \delta W_j$, the algorithm for SPDE (5.1) is similar to before:

1. Initialize

- $\hat{u}^0 := (u_0(\mathbf{x}_1), \dots, u_0(\mathbf{x}_N))^T$
- $K_{PB}^* := \begin{pmatrix} (P_1 P_2^* K(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N} & (P_1 B_2^* K(\mathbf{x}_j, \mathbf{x}_{N+k}))_{j,k=1}^{N,M} \\ (B_1 P_2^* K(\mathbf{x}_{N+j}, \mathbf{x}_k))_{j,k=1}^{M,N} & (B_1 B_2^* K(\mathbf{x}_{N+j}, \mathbf{x}_{N+k}))_{j,k=1}^{M,M} \end{pmatrix}$
- $B := \begin{pmatrix} (P_2^* K(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N} & (B_2^* K(\mathbf{x}_j, \mathbf{x}_{N+k}))_{j,k=1}^{N,M} \end{pmatrix}$
- $\Psi_0 := \delta t (R(\mathbf{x}_j, \mathbf{x}_k))_{j,k=1}^{N,N}$, $\Sigma_0 := \begin{pmatrix} \Psi_0 & 0 \\ 0 & 0 \end{pmatrix}$

2. Repeat for $j = 1, 2, \dots, n$, i.e., for $t_1, t_2, \dots, t_n = T$

- $V_1 := \text{diag}(\psi(\hat{u}_1^{j-1}), \dots, \psi(\hat{u}_N^{j-1}))$, $V_2 := \begin{pmatrix} V_1 & 0 \\ 0 & 0 \end{pmatrix}$
- $\Psi := V_1 \Psi_0 V_1$, $\Sigma := V_2 \Sigma_0 V_2$




Figure 4.3: Convergence of mean and variance with respect to refinement of points and time steps for $\sigma := 1$. (The relative RMSE of exact U and approximate \hat{U} is defined by $\text{RMSE}(U, \hat{U}) := \sqrt{\frac{1}{nN} \sum_{j=1}^n \sum_{k=1}^N (U(t_j, x_k) - \hat{U}(t_j, x_k))^2 / \|U(t_k, \cdot)\|_\infty^2}$.)

- Simulate $\xi \sim \mathcal{N}(0, \Psi)$, $A := B(K_{PB}^* + \Sigma)^{-1}$
- $\hat{u}^j := B(K_{PB}^* + \Sigma)^{-1} \begin{pmatrix} \hat{u}^{j-1} + \xi \\ 0 \end{pmatrix} = A \begin{pmatrix} \hat{u}^{j-1} + \xi \\ 0 \end{pmatrix}$

Of course, now the matrix A needs to be updated for each time step and for each sample path so that the algorithm is much costlier.

A Reproducing-Kernel Hilbert Spaces

Definition A.1 ([21, Definition 10.1]). A Hilbert space $H_K(\mathcal{D})$ of functions $f : \mathcal{D} \rightarrow \mathbb{R}$ is called a *reproducing-kernel Hilbert space* with a *reproducing kernel* $K : \mathcal{D} \times \mathcal{D} \rightarrow \mathbb{R}$ if

$$(i) K(\cdot, \mathbf{y}) \in H_K(\mathcal{D}) \quad \text{and} \quad (ii) f(\mathbf{y}) = \langle K(\cdot, \mathbf{y}), f \rangle_{K, \mathcal{D}},$$

for all $f \in H_K(\mathcal{D})$ and all $\mathbf{y} \in \mathcal{D}$. Here $\langle \cdot, \cdot \rangle_{K, \mathcal{D}}$ is the inner product of $H_K(\mathcal{D})$.

According to [21, Theorem 10.4] all reproducing kernels are positive semi-definite. [21, Theorem 10.10] shows that a symmetric positive semi-definite kernel K is always a reproducing kernel of a reproducing-kernel Hilbert space $H_K(\mathcal{D})$.

If \mathcal{D} is open and bounded (pre-compact) and $K \in L_2(\mathcal{D} \times \mathcal{D})$ is symmetric positive definite, then Mercer's theorem [11, Theorem 13.5] guarantees the existence of a countable set of positive values $\lambda_1 \geq \lambda_2 \geq \dots > 0$ with $\sum_{k=1}^{\infty} \lambda_k < \infty$ and an orthonormal base $\{e_k\}_{k=1}^{\infty}$

of $L_2(\mathcal{D})$ such that K possesses the absolutely and uniformly convergent representation

$$K(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{\infty} \lambda_k e_k(\mathbf{x}) e_k(\mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \mathcal{D}.$$

This Mercer series of K implies that

$$\lambda_k e_k(\mathbf{y}) = \int_{\mathcal{D}} K(\mathbf{x}, \mathbf{y}) e_k(\mathbf{x}) d\mathbf{x}, \quad \mathbf{y} \in \mathcal{D}, \quad k \in \mathbb{N}.$$

Definition A.2. The sequences $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ given above are called the *eigenvalues* and *eigenfunctions* of the reproducing kernel K .

Since $\{e_k\}_{k=1}^{\infty}$ is orthonormal in $L_2(\mathcal{D})$ we can compute the series expansion of the integral-type kernel $\overset{*}{K}$ defined in Lemma 2.1, i.e.,

$$\overset{*}{K}(\mathbf{x}, \mathbf{y}) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \int_{\mathcal{D}} \lambda_j e_j(\mathbf{x}) e_j(\mathbf{z}) \lambda_k e_k(\mathbf{z}) e_k(\mathbf{y}) d\mathbf{z} = \sum_{k=1}^{\infty} \lambda_k^2 e_k(\mathbf{x}) e_k(\mathbf{y}).$$

It is easy to check that $\overset{*}{K} \in L_2(\mathcal{D} \times \mathcal{D})$ is symmetric positive definite and $\{\lambda_k^2\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ are the eigenvalues and eigenfunctions of $\overset{*}{K}$.

Theorem A.1 ([21, Theorem 10.29]). *Suppose that $K \in L_2(\mathcal{D} \times \mathcal{D})$ is a symmetric positive definite kernel on a pre-compact $\mathcal{D} \subset \mathbb{R}^d$. Then its reproducing-kernel Hilbert space is given by*

$$\mathbb{H}_K(\mathcal{D}) = \left\{ f \in L_2(\mathcal{D}) : \sum_{k=1}^{\infty} \frac{1}{\lambda_k} \left| \int_{\mathcal{D}} f(\mathbf{x}) e_k(\mathbf{x}) d\mathbf{x} \right|^2 < \infty \right\}$$

and the inner product has the representation

$$\langle f, g \rangle_{K, \mathcal{D}} = \sum_{k=1}^{\infty} \frac{1}{\lambda_k} \int_{\mathcal{D}} f(\mathbf{x}) e_k(\mathbf{x}) d\mathbf{x} \int_{\mathcal{D}} g(\mathbf{x}) e_k(\mathbf{x}) d\mathbf{x}, \quad f, g \in \mathbb{H}_K(\mathcal{D}),$$

where $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ are the eigenvalues and eigenfunctions of K .

We can verify that $\{\sqrt{\lambda_k} e_k\}_{k=1}^{\infty}$ is an orthonormal base of $\mathbb{H}_K(\mathcal{D})$.

Example A.1. The papers [9, 22] show that the *Sobolev spline* (Matérn function) of degree $m > \frac{d}{2}$,

$$g_{m, \theta}(\mathbf{x}) := \frac{2^{1-m-d/2}}{\pi^{d/2} \Gamma(m) \theta^{2m-d}} (\theta \|\mathbf{x}\|_2)^{m-d/2} K_{d/2-m}(\theta \|\mathbf{x}\|_2), \quad \mathbf{x} \in \mathbb{R}^d, \quad \theta > 0,$$

is a full-space Green function of the differential operator $L := (\theta^2 I - \Delta)^m$, i.e., $L g_{m, \theta} = \delta_0$, where $t \mapsto K_{\nu}(t)$ is the modified Bessel function of the second kind of order ν . The kernel function

$$K_{m, \theta}(\mathbf{x}, \mathbf{y}) := g_{m, \theta}(\mathbf{x} - \mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}^d,$$

is a positive definite kernel whose reproducing-kernel Hilbert space is equivalent to the L_2 -based Sobolev space of degree m , i.e., $\mathbf{H}_{K_{m,\theta}}(\mathbb{R}^d) \cong \mathcal{H}^m(\mathbb{R}^d)$. Its inner product has the explicit form

$$\langle f, g \rangle_{K_{m,\theta}, \mathbb{R}^d} = \int_{\mathbb{R}^d} \mathbf{P}_\theta f(\mathbf{x})^T \mathbf{P}_\theta g(\mathbf{x}) d\mathbf{x}, \quad f, g \in \mathbf{H}_{K_{m,\theta}}(\mathbb{R}^d),$$

where $\mathbf{P}_\theta^T := (\mathbf{Q}^0, \mathbf{Q}^1, \dots, \mathbf{Q}^m)$ and

$$\mathbf{Q}^j := \begin{cases} a_j \Delta^k, & j = 2k, \\ a_j \Delta^k \nabla^T, & j = 2k + 1, \end{cases} \quad a_j := \sqrt{\frac{m! \theta^{2m-2j}}{j!(m-j)!}}, \quad k \in \mathbb{N}_0, \quad j = 0, 1, \dots, m.$$

According to [3, Theorem 1.4.6], the reproducing-kernel Hilbert space $\mathbf{H}_{K_{m,\theta}}(\mathcal{D})$ is endowed with the reproducing-kernel norm

$$\|f\|_{K_{m,\theta}, \mathcal{D}} = \inf_{\tilde{f} \in \mathbf{H}_{K_{m,\theta}}(\mathbb{R}^d)} \left\{ \|\tilde{f}\|_{K_{m,\theta}, \mathbb{R}^d} : \tilde{f}|_{\mathcal{D}} = f \right\}, \quad f \in \mathbf{H}_{K_{m,\theta}}(\mathcal{D}).$$

Moreover, if the open bounded domain $\mathcal{D} \subset \mathbb{R}^d$ is regular then $\mathbf{H}_{K_{m,\theta}}(\mathcal{D})$ is equivalent to the L_2 -based Sobolev space of degree m , i.e., $\mathbf{H}_{K_{m,\theta}}(\mathcal{D}) \cong \mathcal{H}^m(\mathcal{D})$.

B Differential and Boundary Operators

In this section we define differential and boundary operators on Sobolev spaces $\mathcal{H}^m(\mathcal{D})$. The differential and boundary operators in this paper are well-defined since we assume that the open and bounded domain \mathcal{D} is *regular*, i.e., it satisfies a strong local Lipschitz condition which implies a uniform cone condition (see [1, Chapter 4.1]). This means that \mathcal{D} has a regular boundary $\partial\mathcal{D}$.

Let the notation for typical derivatives be

$$D^\alpha := \prod_{k=1}^d \frac{\partial^{\alpha_k}}{\partial x_k^{\alpha_k}}, \quad |\alpha| := \sum_{k=1}^d \alpha_k, \quad \alpha := (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d.$$

We extend these derivatives to *weak derivatives* (see [1, Chapter 1.5]) using the same symbol D^α . Using these weak derivatives, the *classical L_2 -based Sobolev space* $\mathcal{H}^m(\mathcal{D})$ is given by

$$\mathcal{H}^m(\mathcal{D}) := \left\{ f \in L_1^{loc}(\mathcal{D}) : D^\alpha f \in L_2(\mathcal{D}), \quad |\alpha| \leq m, \quad \alpha \in \mathbb{N}_0^d \right\}, \quad m \in \mathbb{N}_0,$$

equipped with the natural inner product

$$\langle f, g \rangle_{m, \mathcal{D}} := \sum_{|\alpha| \leq m} \int_{\mathcal{D}} D^\alpha f(\mathbf{x}) D^\alpha g(\mathbf{x}) d\mathbf{x}, \quad f, g \in \mathcal{H}^m(\mathcal{D}).$$

The weak derivative D^α is a bounded linear operator from $\mathcal{H}^m(\mathcal{D})$ into $L_2(\mathcal{D})$ when $|\alpha| \leq m$. Moreover, $D^\alpha f$ is well-posed on the boundary $\partial\mathcal{D}$ when $f \in C^m(\overline{\mathcal{D}})$ and $|\alpha| \leq m - 1$ and we

denote that $D^\alpha|_{\partial\mathcal{D}}f := D^\alpha f|_{\partial\mathcal{D}}$. The book [1] and the paper [10] show that $D^\alpha|_{\partial\mathcal{D}}$ can be extended to a bounded linear operator from $\mathcal{H}^m(\mathcal{D})$ into $L_2(\partial\mathcal{D})$ when $|\alpha| \leq m - 1$ because \mathcal{D} has a regular boundary $\partial\mathcal{D}$. The $L_2(\partial\mathcal{D})$ -inner product is denoted by

$$\langle f, g \rangle_{\partial\mathcal{D}} = \int_{\partial\mathcal{D}} f(\mathbf{x})g(\mathbf{x})dS(\mathbf{x}), \quad \text{when } d \geq 2 \text{ and } \partial\mathcal{D} \text{ is the boundary manifold,}$$

and

$$\langle f, g \rangle_{\partial\mathcal{D}} = f(b)g(b) + f(a)g(a), \quad \text{when } d = 1 \text{ and } \partial\mathcal{D} = \{a, b\}.$$

Definition B.1. A differential operator $P : \mathcal{H}^m(\mathcal{D}) \rightarrow L_2(\mathcal{D})$ is well-defined by

$$P = \sum_{|\alpha| \leq m} c_\alpha D^\alpha, \quad \text{where } c_\alpha \in C^\infty(\overline{\mathcal{D}}) \text{ and } \alpha \in \mathbb{N}_0^d, \quad m \in \mathbb{N}_0,$$

and its *order* is given by $\mathcal{O}(P) := \max \{|\alpha| : c_\alpha \neq 0, |\alpha| \leq m, \alpha \in \mathbb{N}_0^d\}$. A *boundary operator* $B : \mathcal{H}^m(\mathcal{D}) \rightarrow L_2(\partial\mathcal{D})$ is well-defined by

$$B = \sum_{|\alpha| \leq m-1} b_\alpha D^\alpha|_{\partial\mathcal{D}}, \quad \text{where } b_\alpha \in C^\infty(\partial\mathcal{D}) \text{ and } \alpha \in \mathbb{N}_0^d, \quad m \in \mathbb{N},$$

and its *order* is given by $\mathcal{O}(B) := \max \{|\alpha| : b_\alpha \neq 0, |\alpha| \leq m - 1, \alpha \in \mathbb{N}_0^d\}$.

It is obvious that the differential operator P and the boundary operator B are bounded (continuous) linear operators on $\mathcal{H}^m(\mathcal{D})$ with values in L_2 whenever $\mathcal{O}(P) \leq m$ and $\mathcal{O}(B) \leq m - 1$. Much more detail on differential and boundary operators can be found in [1, 10].

If $\mathbb{H}_K(\mathcal{D})$ is embedded¹ into $\mathcal{H}^m(\mathcal{D})$ then the eigenvalues $\{\lambda_k\}_{k=1}^\infty$ and eigenfunctions $\{e_k\}_{k=1}^\infty$ of the reproducing kernel K satisfy

$$\lambda_k \|e_k\|_{m, \mathcal{D}}^2 \leq C^2 \|\sqrt{\lambda_k} e_k\|_{K, \mathcal{D}}^2 = C^2, \quad k \in \mathbb{N},$$

because $\{\sqrt{\lambda_k} e_k\}_{k=1}^\infty$ is an orthonormal base of $\mathbb{H}_K(\mathcal{D})$. When $m > d/2$ then $\mathcal{H}^m(\mathcal{D})$ is embedded into $C(\overline{\mathcal{D}})$ by the Sobolev embedding theorem. This implies that $K \in C(\overline{\mathcal{D}} \times \overline{\mathcal{D}}) \subset L_2(\mathcal{D} \times \mathcal{D})$ because $K(\cdot, \mathbf{y}) \in C(\overline{\mathcal{D}})$ for each $\mathbf{y} \in \mathcal{D}$ and K is symmetric. Based on these properties, we can introduce the following lemma.

Lemma B.1. Consider a differential operator P with order $\mathcal{O}(P) \leq m$ and a boundary operator B with order $\mathcal{O}(B) \leq m - 1$, where $m > d/2$. If the reproducing-kernel Hilbert space $\mathbb{H}_K(\mathcal{D})$ is embedded into the Sobolev space $\mathcal{H}^m(\mathcal{D})$, then

$$Pf = \sum_{k=1}^{\infty} \hat{f}_k \sqrt{\lambda_k} P e_k, \quad Bf = \sum_{k=1}^{\infty} \hat{f}_k \sqrt{\lambda_k} B e_k, \quad f \in \mathbb{H}_K(\mathcal{D}),$$

where $\hat{f}_k = \langle f, e_k \rangle_{K, \mathcal{D}}$ for each $k \in \mathbb{N}$ and $\{\lambda_k\}_{k=1}^\infty$ and $\{e_k\}_{k=1}^\infty$ are the eigenvalues and eigenfunctions of the reproducing kernel K .

¹The recent papers [9, 10, 22] show that there exist kernels K whose reproducing-kernel Hilbert space $\mathbb{H}_K(\mathcal{D})$ is continuously embedded into the Sobolev space $\mathcal{H}^m(\mathcal{D})$.

Proof. According to Theorem A.1 each $f \in \mathbb{H}_K(\mathcal{D})$ can be expanded as $f = \sum_{k=1}^{\infty} \hat{f}_k \sqrt{\lambda_k} e_k$. Since $\{\sqrt{\lambda_k} e_k\}_{k=1}^{\infty}$ is an orthonormal basis we have $\sum_{k=1}^{\infty} |\hat{f}_k|^2 < \infty$. Let $f_n := \sum_{k=1}^n \hat{f}_k \sqrt{\lambda_k} e_k$ for each $n \in \mathbb{N}$. Then

$$\|f_n - f\|_{m, \mathcal{D}}^2 \leq C^2 \|f_n - f\|_{K, \mathcal{D}}^2 \leq C^2 \sum_{k=n+1}^{\infty} |\hat{f}_k|^2 \rightarrow 0, \quad \text{when } n \rightarrow \infty.$$

The proof is completed by remembering that P and B are bounded linear operators on $\mathcal{H}^m(\mathcal{D})$. \square

If $\mathbb{H}_K(\mathcal{D})$ is embedded into $\mathcal{H}^m(\mathcal{D})$, then for each $|\alpha| \leq m$, $|\beta| \leq m$ and $\alpha, \beta \in \mathbb{N}_0^d$, we have

$$\begin{aligned} & \left(\int_{\mathcal{D}} \int_{\mathcal{D}} \left| \sum_{k=1}^{\infty} \lambda_k^2 D^\alpha e_k(\mathbf{x}) D^\beta e_k(\mathbf{y}) \right|^2 d\mathbf{x} d\mathbf{y} \right)^{1/2} \\ & \leq \sum_{k=1}^{\infty} \lambda_k^2 \|D^\alpha e_k\|_{L_2(\mathcal{D})} \|D^\beta e_k\|_{L_2(\mathcal{D})} \leq \sum_{k=1}^{\infty} \lambda_k^2 \|e_k\|_{m, \mathcal{D}}^2 \leq C^2 \sum_{k=1}^{\infty} \lambda_k < \infty, \end{aligned}$$

which implies that $\overset{*}{K} \in \mathcal{H}^{m, m}(\mathcal{D} \times \mathcal{D})$. Let $n := \lceil m - d/2 \rceil - 1$. The Sobolev embedding theorem shows that $\mathcal{H}^{m, m}(\mathcal{D} \times \mathcal{D}) \subset C^{n, n}(\overline{\mathcal{D}} \times \overline{\mathcal{D}})$. Then we can obtain the following lemma.

Lemma B.2. *Consider a differential operator P with order $\mathcal{O}(P) < m - d/2$ and a boundary operator B with order $\mathcal{O}(B) < m - d/2$, where $m > d/2$. If the reproducing-kernel Hilbert space $\mathbb{H}_K(\mathcal{D})$ is embedded into the Sobolev space $\mathcal{H}^m(\mathcal{D})$, then*

$$\begin{aligned} P_1 P_2 \overset{*}{K}(\mathbf{x}, \mathbf{y}) & := P_{z_1} P_{z_2} \overset{*}{K}(z_1, z_2)|_{z_1=\mathbf{x}, z_2=\mathbf{y}} = \sum_{k=1}^{\infty} \lambda_k^2 P e_k(\mathbf{x}) P e_k(\mathbf{y}), \\ B_1 B_2 \overset{*}{K}(\mathbf{x}, \mathbf{y}) & := B_{z_1} B_{z_2} \overset{*}{K}(z_1, z_2)|_{z_1=\mathbf{x}, z_2=\mathbf{y}} = \sum_{k=1}^{\infty} \lambda_k^2 B e_k(\mathbf{x}) B e_k(\mathbf{y}), \\ P_1 B_2 \overset{*}{K}(\mathbf{x}, \mathbf{y}) & := P_{z_1} B_{z_2} \overset{*}{K}(z_1, z_2)|_{z_1=\mathbf{x}, z_2=\mathbf{y}} = \sum_{k=1}^{\infty} \lambda_k^2 P e_k(\mathbf{x}) B e_k(\mathbf{y}), \end{aligned}$$

where $\{\lambda_k\}_{k=1}^{\infty}$ and $\{e_k\}_{k=1}^{\infty}$ are the eigenvalues and eigenfunctions of K . Moreover, $P_1 P_2 \overset{*}{K} \in C(\overline{\mathcal{D}} \times \overline{\mathcal{D}})$, $B_1 B_2 \overset{*}{K} \in C(\partial \mathcal{D} \times \partial \mathcal{D})$ and $P_1 B_2 \overset{*}{K} \in C(\overline{\mathcal{D}} \times \partial \mathcal{D})$.

Acknowledgments

We would like to thank Phaedon-Stelios Koutsourelakis for the inspiration to solve SPDEs with a maximum likelihood-based approach. The work of Igor Cialenco was partially supported by the National Science Foundation (NSF) grant DMS-0908099. Gregory E. Fasshauer and Qi Ye acknowledge support from NSF grants DMS-0713848 and DMS-1115392. The authors would like to thank the anonymous referee and the editors for their helpful comments and suggestions which improved greatly the final manuscript.

References

- [1] Robert A. Adams and John J. F. Fournier, *Sobolev Spaces*, Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [2] Ivo Babuška, Fabio Nobile, and Raúl Tempone, *A stochastic collocation method for elliptic partial differential equations with random input data*, SIAM Rev. **52** (2010), no. 2, 317–355.
- [3] Alain Berlinet and Christine Thomas-Agnan, *Reproducing Kernel Hilbert Spaces in Probability and Statistics*, Kluwer Academic Publishers, Boston, MA, 2004.
- [4] M. D. Buhmann, *Radial Basis Functions: Theory and Implementations*, Cambridge Monographs on Applied and Computational Mathematics, vol. 12, Cambridge University Press, 2003.
- [5] Pao-Liu Chow, *Stochastic Partial Differential Equations*, Chapman & Hall/CRC Applied Mathematics and Nonlinear Science Series, Chapman & Hall/CRC, Boca Raton, FL, 2007.
- [6] G. Da Prato and J. Zabczyk, *Stochastic equations in infinite dimensions*, Encyclopedia of Mathematics and its Applications, vol. 44, Cambridge University Press, Cambridge, 1992.
- [7] Manas K. Deb, Ivo M. Babuška, and J. Tinsley Oden, *Solution of stochastic partial differential equations using Galerkin finite element techniques*, Comput. Methods Appl. Mech. Engrg. **190** (2001), no. 48, 6359–6372.
- [8] G. E. Fasshauer, *Positive definite kernels: Past, present and future*, Dolomite Research Notes on Approximation **4** (2011), 21–63.
- [9] G. E. Fasshauer and Q. Ye, *Reproducing kernels of generalized Sobolev spaces via a Green function approach with distributional operators*, Numer. Math. **119** (2011), no. 3, 585–611.
- [10] G. E. Fasshauer and Q. Ye, *Reproducing kernels of Sobolev spaces via a Green kernel approach with differential operators and boundary operators*, Adv. Comput. Math. (2011), online first DOI: 10.1007/s10444-011-9264-6.
- [11] Gregory E. Fasshauer, *Meshfree Approximation Methods with MATLAB*, Interdisciplinary Mathematical Sciences, vol. 6, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [12] Y. C. Hon and Robert Schaback, *The kernel-based method of lines for the heat equation*, (2010), University of Göttingen, preprint.

- [13] Svante Janson, *Gaussian Hilbert Spaces*, Cambridge Tracts in Mathematics, vol. 129, Cambridge University Press, 1997.
- [14] Arnulf Jentzen and Peter Kloeden, *Taylor expansions of solutions of stochastic partial differential equations with additive noise*, *Ann. Probab.* **38** (2010), no. 2, 532–569.
- [15] Ioannis Karatzas and Steven E. Shreve, *Brownian Motion and Stochastic Calculus*, Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 1991.
- [16] Milan N. Lukić and Jay H. Beder, *Stochastic processes with sample paths in reproducing kernel Hilbert spaces*, *Trans. Amer. Math. Soc.* **353** (2001), no. 10, 3945–3969.
- [17] Thomas Müller-Gronbach, Klaus Ritter, and Tim Wagner, *Optimal pointwise approximation of a linear stochastic heat equation with additive space-time white noise*, *Monte Carlo and Quasi-Monte Carlo Methods 2006*, Springer, Berlin, 2008, pp. 577–589.
- [18] F. Nobile, R. Tempone, and C. G. Webster, *A sparse grid stochastic collocation method for partial differential equations with random input data*, *SIAM J. Numer. Anal.* **46** (2008), no. 5, 2309–2345.
- [19] B. L. Rozovskii, *Stochastic Evolution Systems: Linear Theory and Applications to Nonlinear Filtering*, Mathematics and its Applications (Soviet Series), vol. 35, Kluwer Academic Publishers Group, Dordrecht, 1990, Linear theory and applications to nonlinear filtering.
- [20] Michael Scheuerer, Robert Schaback, and Martin Schlather, *Interpolation of spatial data – a stochastic or a deterministic problem?*, (2010), University of Göttingen, preprint.
- [21] Holger Wendland, *Scattered Data Approximation*, Cambridge Monographs on Applied and Computational Mathematics, vol. 17, Cambridge University Press, 2005.
- [22] Qi Ye, *Reproducing kernels of generalized Sobolev spaces via a Green function approach with differential operators*, Tech. report, Illinois Institute of Technology, 2010.
- [23] ———, *Analyzing reproducing kernel approximation methods via a Green function approach*, Ph.D. thesis, Illinois Institute of Technology, 2012.