

# Aqueous leaf extract of *Moringa oleifera* reduced intracellular ROS production, DNA fragmentation and acrosome reaction in Human spermatozoa in vitro

Faith T. Moichela<sup>1</sup>  | Gbenga A. Adefolaju<sup>2</sup>  | Ralf R. Henkel<sup>3,4</sup>  | Chinyerum S. Opuwari<sup>2</sup> 

<sup>1</sup>Department of Medical Sciences, University of Limpopo, Sovenga, South Africa

<sup>2</sup>Department of Pre-Clinical Sciences, University of Limpopo, Sovenga, South Africa

<sup>3</sup>Department of Medical Biosciences, University of the Western Cape, Bellville, South Africa

<sup>4</sup>American Centre for Reproductive Medicine, Cleveland, OH, USA

## Correspondence

Chinyerum Opuwari, Department of Pre-Clinical Sciences, University of Limpopo, Private Bag X1106, Sovenga 0727, South Africa.  
Email: chinyerum.opuwari@ul.ac.za

## Funding information

National Research Foundation of South Africa, Grant/Award Number: TTK170426228930; University of Limpopo

## Abstract

The effects of aqueous leaf extract of *Moringa oleifera* (MO) on human sperm functions and integrity was studied in vitro. Semen was obtained by masturbation after 3–5 days' abstinence from 34 healthy donors in Western Cape, South Africa. Liquefied semen was washed in human tubular fluid supplemented with 1% bovine serum albumin (HTF-BSA;1:5) with 10 min centrifugation at 300 g. Sperm suspensions were subsequently incubated with MO extract (0.625, 6.25, 62.5 and 625 µg/ml) for 1 hr, where HTF-BSA served as control. Sperm motility, vitality, DNA fragmentation, reactive oxygen species production, mitochondrial membrane potential, capacitation and acrosome reaction were assessed. Sperm motility, vitality, mitochondrial membrane potential and capacitation remained unchanged ( $p > .05$ ). A dose-dependent decrease in sperm reactive oxygen species production ( $p < .0001$ ), DNA fragmentation ( $p < .0001$ ) and acrosome reaction ( $p < .001$ ) was observed. An increase in the percentage of non-capacitated sperm ( $p < .01$ ) was noted at 625 µg/ml. The antioxidant properties of MO actively maintained basic sperm functions, inhibited excess sperm free superoxide production and preserved acrosome reaction and DNA integrity. Further studies are needed to confirm the effect of aqueous MO leaf extract on fertility potential.

## KEYWORDS

antioxidants, infertility, *Moringa oleifera*, reactive oxygen species, spermatozoa

## 1 | INTRODUCTION

Despite the common misconception that men are immune to infertility, 7% of males are diagnosed with some form of infertility in their lifetime (Schwetz et al., 2013), which accounts for about 30 million men worldwide (Agarwal et al., 2015). Males are solely responsible for nearly 20%–30% of the identifiable causes of infertility (Agarwal et al., 2015). Deficiencies in semen quality and quantity have been identified as the most common cause of male infertility, with about 90% of cases attributed to suboptimal spermatogenesis and counts

(Kumar & Singh, 2015). A decline in sperm count has been reported worldwide (Virtanen et al., 2017). About 40% of normozoospermic men still fail to establish a clinical pregnancy with proven fertile females (Cho & Agarwal, 2018). Oxidative stress is mentioned as being central and independent in male infertility (Agarwal et al., 2019). High production of endogenous reactive oxygen species (ROS) is negatively correlated with sperm functional integrity (Tatone et al., 2010). Poor sperm motility and viability decreased mitochondrial membrane potential (Dobrakowski et al., 2017), and high levels of DNA fragmentation (Iommiello et al., 2015), which reduces the

chances of successful fertilisation, embryo integrity and development were found to be secondary to oxidative stress (Nishihara et al., 2018). Despite the advances made with assisted reproductive technology, access to treatments remain unequal due to disparity of income (Adageba et al., 2015; Makuch et al., 2011), this incited the quest for identification and investigation of affordable and indiscriminately accessible therapeutic alternatives of treating infertility such as plant-based medicine.

For many centuries, traditional medicines have been used for the improvement of sexual and reproductive issues through the establishment of healing systems such as Ayurveda, Chinese and African healing systems (Kotta et al., 2013). Several scientific efforts have been concerted towards describing the efficacy of plants and their derivatives in reproductive functions. Some examples of such medicinal plants include *Aloe barbadovera* (Erhabor & Idu, 2017), *Eurycoma longifolia* (Tongkat ali) (Tambi & Imran, 2010), *Kigelia africana* (saw-sage tree) (Azu et al., 2010), *Mondia whitei* (white ginger) (Lampiao et al., 2008) and *Rhoicissus tridentatabushmen grape* (bitter grape) (Rakuambo et al., 2006). The World Health Organisation (WHO) asserts that about 80% of the world population relies exclusively on natural resources for the provision of their health care needs (Yuan et al., 2016).

*Moringa oleifera* (MO) of the family *Moringaceae* is indigenous to South Asian and is now vastly naturalised in other parts of the world (Gandji et al., 2018). *Moringa oleifera*, popularly known as the 'horse-radish' tree, has vast phytochemical and pharmacological attributes in African folklore (Bhattacharya et al., 2018; Farooq et al., 2012), which has led to the widely held conception that MO could be the *panacea* of modern-day herbal therapy (Koul & Chase, 2015). Various parts of MO are used as traditional medicine and possess variable phytochemicals, with the leaves cited as containing high levels of the medicinal fractions (Vergara-Jimenez et al., 2017). Isolated and characterised secondary metabolites include flavonoids, vitamins, alkaloids, glucosinolates (Maldini et al., 2014), tannins and saponins (Kwaghe & Ambali, 2009) phenolic acids (Rodríguez-Pérez et al., 2015), essential amino acids (Prasanth et al., 2011) minerals (Ahmed et al., 2018), simple and complex sugars (Anudeep & Radha, 2018)), proteins (Teixeira et al., 2014), sterols (Leone et al., 2016), carotenoids (Elayaraja et al., 2017) and isothiocyanates (Fahey et al., 2019). Health benefits of MO include anti-hypertensive (Acuram & Chichioco Hernandez, 2019), antimicrobial (Thilza et al., 2010) anti-nociceptive and analgesic (Sulaiman et al., 2008), anti-inflammatory (Waterman et al., 2014), anti-peroxidative and anti-oxidative (Nandave et al., 2009).

Anecdotal evidence points out that traditional healers have been prescribing MO for treatment of sexual inadequacies (Nazim et al., 2017) and fertility-related issues (MacDONalD et al., 2016). For instance, the seeds of MO are used by traditional healers in India as an aphrodisiac (Lalas & Tsaknis, 2002). Several studies have confirmed the anti-oxidative activity of MO leaves in improving reproductive, sexual and seminal functions in rat models (Aprioku & Onyenaturuchi, 2018; Awodele et al., 2012; Cajuday & Pocsidio, 2010; Wafa et al., 2017). This study sought to investigate

the effects of aqueous leaf extracts of *M. oleifera* on human sperm functions in vitro.

## 2 | MATERIALS AND METHODS

### 2.1 | Plant collection

Fresh leaves of MO were collected from the experimental farm of the University of Limpopo, South Africa. The leaves were authenticated by Dr Egan Bronwyn Ann, University of Limpopo. A voucher specimen (121,625) was deposited at the Larry Leach Herbarium of the University of Limpopo.

### 2.2 | Preparation of aqueous leaf extract of *Moringa oleifera*

Fresh leaves of MO were allowed to dry for 7 days at room temperature and pulverised. MO leaf powder was infused in distilled water (100 g/L), allowed to boil for 20 min, filtered using cheesecloth, and centrifuged at 800 g for 10 min at room temperature (Shunmugam, 2016). Thereafter, filtered using Whatman filter papers 4 and 1 respectively, and subsequently freeze-dried. The concentration for this study was calculated based on the assumption that an average healthy man weighs about 80 kg (Shalaweh et al., 2015) together with a previous study where a capsule containing 500 mg MO leaf powder was prescribed to diabetes mellitus type-2 (DM2) patients daily (Taweerutchana et al., 2017) as follows:

$$\text{Therapeutic concentration} = \frac{\text{prescribed dose (mg)}}{\text{Average weight of man (kg)}}$$

where 1 kg = 1,000 ml.

The in vitro therapeutic concentration obtained was 6.25 µg/ml. A stock solution (62.5 mg/ml) was reconstituted in HTF-BSA.

### 2.3 | Source and preparation of semen

The study received ethical approval from the Turfloop Research Ethics Committee (TREC) of the University of Limpopo (TREC/45/PG: 2019) and the Biomedical Research Ethics Committee (BMREC) of the University of Western Cape (BM17/7/14). Following informed consent, semen was collected in sterile vials from 34 healthy donors in the Western Cape, South Africa, by masturbation after a 3–5 days' abstinence and prepared as described by Shalaweh et al. (2015). Semen was classified normozoospermic samples according to WHO (2010) criteria (sperm concentration > 15 million/ml, sperm count > 39 million/ejaculate, sperm motility > 40%). In brief, samples were incubated at 37°C for 30 min for liquefaction. In other to obtain the baseline sperm concentration and motility, liquefied semen (3 µl) was placed on

pre-warmed Leja slides and analysed using sperm class analyser<sup>®</sup> (SCA, version 6.4; Microptic). After that, the semen sample was diluted with HTF-BSA (1:5) and centrifuged at 300 g for 10 min, and the pellet was re-suspended in fresh HTF-BSA. Sperm suspensions (sperm concentration:  $7.5 \times 10^6$ /ml) were mixed with a stock solution of aqueous leaf extract of MO (62.5 mg/ml) to final concentrations of 0.625, 6.25, 62.5 and 625  $\mu$ g/ml and incubated at 37°C for 1 hr to analyse the various sperm parameters. HTF-BSA served as control. Therefore, the total number of samples for each treatment group was 34.

## 2.4 | Determination of sperm motility and kinematic characteristics

Following exposure of sperm to MO aqueous leaf extract, 3  $\mu$ l sperm suspension was loaded on a pre-warmed Leja slide (Leja Products B.V.). Thereafter, sperm motility (total motility (%), progressive motility (%), non-progressive motility (%)) and motion kinematics (hyperactivation (%), linearity index (%), straightness (%), wobble (%), amplitude of lateral head displacement (ALH,  $\mu$ m), average path velocity (VAP,  $\mu$ m/s), beat cross frequency (BCF, Hz), curvilinear velocity (VCL,  $\mu$ m/s), straight line velocity (VSL,  $\mu$ m/s)) were analysed using sperm class analyser. A total of 200 spermatozoa were assessed according to the WHO (2010) criteria.

## 2.5 | Determination of sperm vitality

Sperm vitality was determined with eosin–nigrosin Y stain (Shalaweh et al., 2015; WHO, 2010). After exposure of sperm to the various concentrations of MO aqueous leaf extract, the cells were mixed with eosin–nigrosin dye (1:1). After that, 10  $\mu$ l of the mixture was smeared on frosted slides and left to air-dry. Slides were viewed at 100 $\times$  objective with immersion oil using a light microscope. Live and dead spermatozoa were stained white and red, respectively. Two hundred spermatozoa were counted, and results represented as a percentage of live spermatozoa.

## 2.6 | Determination of sperm mitochondrial membrane potential ( $\Delta\psi_m$ )

Sperm mitochondrial membrane potential was assessed using 5,5',6,6'-tetrachloro-1-1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide (DePsipher<sup>™</sup>, Trevigen), a lipophilic cationic dye. The protocol was applied as previously described (Henkel et al., 2012). In brief, after exposure of spermatozoa to MO leaf extract, they were washed with HTF-BSA. The recovered pellet was re-suspended in 100  $\mu$ l of diluted DePsipher solution, centrifuged for 10 min at 300 g and incubated in 100  $\mu$ l reaction buffer for 20 min at 37°C away from light. After that, the sperm suspension was centrifuged at 300 g for 10 min and the pellet re-suspended in 100  $\mu$ l of pre-warmed

reaction buffer. An aliquot of 10  $\mu$ l from each sample was used to make smears and observed at 1,000 $\times$  magnification using an epifluorescence microscope (Zeiss). Spermatozoa with green fluorescing mid-pieces were regarded as having disturbed mitochondrial membrane potential (MMP), while those with red fluorescence had intact MMP (Erasmus et al., 2012; Shalaweh et al., 2015). A total of 200 spermatozoa were analysed and result expressed as the percentage of sperm with intact MMP.

## 2.7 | Determination of spermatozoa reactive oxygen species production

Reactive oxygen species production in spermatozoa was determined as previously described by Shalaweh et al. (2015). In brief, a stock solution of 20  $\mu$ M dihydroethidium (Molecular Probes) in PBS was prepared (pH of 7.4). Following the treatment of spermatozoa with various concentrations of MO aqueous leaf extract for 1 hr at 37°C, 100  $\mu$ l was diluted with HTF-BSA (1:5) and centrifuged for 10 min at 300 g. The pellet was re-suspended in 200  $\mu$ l of PBS and centrifuged. Then, diluted DHE was mixed with sperm samples (1:5) and incubated at 37°C for 20 min. Subsequently, 10  $\mu$ l of each sample was placed on a slide, covered with a coverslip and viewed under oil immersion using an epifluorescent microscope with 488 nm excitation and 590 emission filters (Zeiss). Two hundred spermatozoa were analysed and result expressed as the percentage of spermatozoa with red/orange fluorescence (indicative of excess ROS production).

## 2.8 | Determination of sperm DNA fragmentation

Sperm DNA fragmentation was determined using the TUNEL assay (terminal deoxynucleotidyl transferase-mediated dUTP-biotin nick-end labelling) (Dead End<sup>™</sup> Fluorometric, Promega) according to Erasmus et al. (2012). Spermatozoa exposed to different concentrations of MO extract for 1 hr were centrifuged for 10 min at 300 g in HTF-BSA. After that, 100  $\mu$ l of the sperm suspension was re-suspended in PBS and centrifuged again. Re-suspended pellets were then smeared on Starfrost slides (Knittel Gläser) and allowed to air-dry. Subsequently, slides were fixed with 4% methanol-free formaldehyde (KIMIX Chemicals) in PBS for 25 min at 4°C, washed with PBS for 5 min and allowed to dry at room temperature. The sperm plasma membrane was permeabilised with 0.2% Triton X-100 (Sigma-Aldrich) in PBS for 5 min, rinsed twice with PBS, and 100  $\mu$ l of equilibration buffer was added for 10 min. After that, 50  $\mu$ l TdT reagent buffer was added, a coverslip placed on slides and incubated for 1 hr in a dark humidified chamber. Then, slides were submerged in 2 $\times$  SSC buffer (Promega) for 15 min, washed three times with PBS for 5 min each and analysed at 1,000 $\times$  using an epifluorescence microscope with excitation and emission filters at 488 and 530 nm, respectively. DNA intact sperm exhibited a bit of a background staining (TUNEL-negative), whereas DNA-fragmented sperm

emitted a bright green fluorescence (TUNEL-positive). Two hundred spermatozoa were counted, and the result expressed as a percentage of TUNEL-positive spermatozoa.

## 2.9 | Determination of sperm capacitation and acrosome reaction

The dual staining method of Hoechst 32258/CTC for assessment of capacitation and acrosome status of human spermatozoa was adopted for this study, according to Green et al. (1996). Chlorotetracycline (CTC; 750  $\mu$ M) solution (containing 130 mM NaCl, 5 mM cysteine and 20 mM Tris-HCl, pH 7.8) was prepared on the day of use. Following the 1 hr incubation of spermatozoa with MO aqueous leaf extracts, Hoechst 32258 was mixed with the sperm suspension in HTF-BSA (1:100), and incubated for 2 min at room temperature and centrifuged with 2% polyvinylpyrrolidone (PVP40; 40 ml) in HTFat 900 g for 5 min. CTC solution (45  $\mu$ l) and 12.5% w/v paraformaldehyde in 0.5 Tris-HCl (pH 7.4) (8  $\mu$ l) was mixed with the Hoechst-stained sperm suspension (45  $\mu$ l). After that, 10  $\mu$ l of the mixture was placed on Starfrost slide and viewed with a 100 $\times$  objective using an epifluorescence microscope. From each slide, 200 live spermatozoa (Hoechst-negative) were identified and assessed for their CTC fluorescence patterns as follows: a uniform fluorescence over the entire sperm head was classified as non-capacitated and acrosome intact spermatozoa (F pattern); fluorescence over the sperm head except the post-acrosomal region was classified as capacitated and acrosome intact spermatozoa (B pattern); and a lack of fluorescent over the entire head with/without a fluorescent post-acrosomal region was regarded as capacitated and acrosome-reacted spermatozoa (AR pattern).

## 2.10 | Data analysis

Data were analysed using GraphPad Prism 6.0.1 for Windows (GraphPad Software). Test for normality was conducted using the D'Agostino-Pearson Omnibus test. One-way analysis of variance (ANOVA) and Tukey's test were used for normally distributed data, while Kruskal-Wallis test and Dunnett's multiple test were employed for non-parametric analysis for comparisons of group means. Results were regarded as statistically significant when  $p < .05$ .

## 3 | RESULTS

Baseline analysis of samples from the donors demonstrated that semen volume (Mean:  $2.8 \pm 0.2$ ; Min: 1; Max: 5 ml), sperm concentration (Mean:  $82.7 \pm 6.5$ ; Min: 18.5; Max: 180.5 million/ml) and percentage of sperm motility (Mean:  $63.9 \pm 2.7$ ; Min: 40.1; Max: 96.3%) were in the range of values defined by WHO (2010).

### 3.1 | Sperm motility parameters

Table 1 shows that aqueous leaf extract of MO did not affect any of the sperm motility parameters or motion kinematics ( $p > .05$ ).

### 3.2 | Sperm vitality and mitochondrial membrane potential

Figures 1a,b demonstrate that the various concentrations of MO aqueous leaf extract had no effect on the human sperm vitality ( $p > .05$ ) and mitochondrial membrane potential ( $p > .05$ ) following a 1-hr incubation.

### 3.3 | Sperm reactive oxygen species production

Aqueous leaf extract of MO caused significant inhibition of sperm intracellular ROS production at 62.5  $\mu$ g/ml ( $p < .01$ ) and 625  $\mu$ g/ml ( $p < .0001$ ) compared to the control (Figure 1c). Repeated measure ANOVA trend analysis revealed a significant dose-dependent decrease ( $p < .0001$ ) in the production of ROS with increasing concentrations of the plant extract. Also, a significant decrease in percentage of intracellular ROS production was observed between treatment groups: 0.625 and 625  $\mu$ g/ml ( $p < .0001$ ), 0.625 and 62.5  $\mu$ g/ml ( $p < .01$ ) and 6.25 and 625  $\mu$ g/ml ( $p < .01$ ).

### 3.4 | Sperm DNA fragmentation

A significant reduction in the percentage of sperm with DNA fragmentation was observed at 625  $\mu$ g/ml ( $p < .0001$ ) of MO leaf extract (Figure 1d) compared to the control. In addition, a significant reduction in DNA fragmentation was observed between treatment groups, that is, 0.625 and 62.5  $\mu$ g/ml ( $p < .0001$ ), 0.625 and 625  $\mu$ g/ml ( $p < .0001$ ) and 6.25 and 625  $\mu$ g/ml ( $p < .0001$ ). Furthermore, repeated measure ANOVA trend analysis revealed a dose-dependent decrease in sperm DNA fragmentation with increasing concentrations of the plant extract ( $p < .0001$ ).

### 3.5 | Sperm capacitation and acrosome reaction

In other to assess capacitation and acrosome reaction, different patterns of CTC staining during the capacitation process were analysed. The highest concentration of MO aqueous leaf extract significantly increased the percentage of uncapacitated and acrosome intact spermatozoa (F pattern;  $p < .01$ ; Figure 2a). Consequently, a significant decrease in the percentage of uncapacitated and acrosome intact spermatozoa was observed between the lowest and highest dose of the extract, that is, between 0.625 and 625  $\mu$ g/ml ( $p < .05$ ), 6.25 and 625  $\mu$ g/ml ( $p < .01$ ), and 62.5 and 625  $\mu$ g/ml ( $p < .05$ ). No significant


**TABLE 1** Sperm motility parameters after 1-hr incubation with various concentrations of aqueous leaf extract of *Moringa oleifera*

Sperm motility parameters	<i>Moringa oleifera</i> aqueous leaf extract ( $\mu\text{g/ml}$ )					p-value
	0	0.625	6.25	62.5	625	
Total motility (%)	24.97 $\pm$ 2.17	23.42 $\pm$ 2.59	24.68 $\pm$ 2.23	24.15 $\pm$ 3.13	24.50 $\pm$ 2.86	0.9188
Progressive motility (%)	9.58 $\pm$ 1.52	8.14 $\pm$ 1.28	7.72 $\pm$ 1.18	6.65 $\pm$ 1.14	7.43 $\pm$ 1.40	0.6594
Non-progressive motility (%)	12.46 $\pm$ 1.24	12.83 $\pm$ 1.51	14.04 $\pm$ 1.37	13.75 $\pm$ 1.92	14.88 $\pm$ 1.90	0.9206
Hyperactivation (%)	4.74 $\pm$ 1.55	6.98 $\pm$ 1.99	4.80 $\pm$ 1.18	3.38 $\pm$ 1.08	5.45 $\pm$ 1.72	0.8197
LIN (%)	39.19 $\pm$ 1.90	37.31 $\pm$ 1.87	38.15 $\pm$ 1.93	36.11 $\pm$ 2.20	37.87 $\pm$ 2.25	0.6833
STR (%)	65.87 $\pm$ 1.76	63.68 $\pm$ 1.73	64.16 $\pm$ 2.01	62.36 $\pm$ 2.26	60.21 $\pm$ 2.34	0.3661
WOB (%)	55.14 $\pm$ 1.49	54.33 $\pm$ 1.52	55.22 $\pm$ 1.55	53.99 $\pm$ 1.63	54.33 $\pm$ 1.91	0.9041
VCL ( $\mu\text{m/s}$ )	71.64 $\pm$ 3.28	71.11 $\pm$ 3.95	67.67 $\pm$ 3.19	67.46 $\pm$ 2.86	66.59 $\pm$ 3.29	0.7873
VAP ( $\mu\text{m/s}$ )	37.95 $\pm$ 1.47	37.00 $\pm$ 1.74	35.79 $\pm$ 1.47	35.17 $\pm$ 1.29	34.82 $\pm$ 1.32	0.5302
VSL ( $\mu\text{m/s}$ )	26.96 $\pm$ 1.30	25.18 $\pm$ 1.32	24.95 $\pm$ 1.26	23.63 $\pm$ 1.38	22.44 $\pm$ 1.29	0.1541
ALH ( $\mu\text{m}$ )	2.44 $\pm$ 0.09	2.44 $\pm$ 0.10	2.35 $\pm$ 0.08	2.36 $\pm$ 0.09	2.31 $\pm$ 0.11	0.9807
BCF (Hz)	15.69 $\pm$ 0.98	14.57 $\pm$ 1.03	14.42 $\pm$ 0.94	15.78 $\pm$ 1.42	13.23 $\pm$ 0.91	0.4147

Note: Data were represented as mean  $\pm$  SEM of 200 spermatozoa from 34 healthy donors.

Abbreviations: ALH, amplitude of lateral head displacement; BCF, beat cross frequency; LIN, linearity; STR, straightness; VAP, average path velocity; VCL, curvilinear velocity; VSL, straight line velocity; WOB, wobble.


**FIGURE 1** Effects of aqueous leaf extracts *Moringa oleifera* on (a) sperm vitality and (b) sperm mitochondrial membrane potential (MMP) (c) intracellular reactive oxygen species (ROS) production in sperm and (d) DNA fragmentation in sperm in vitro. Data were represented as mean  $\pm$  SEM of 200 spermatozoa from 34 healthy donors



changes were observed in the percentage of capacitated spermatozoa (B pattern) in all groups treated with MO leaf extracts when compared to the control ( $p > .05$ ; Figure 2b). Furthermore, no significant change in the percentage of acrosome-reacted spermatozoa was observed by the various concentrations of MO aqueous leaf extract (Figure 2c). However, Mann-Whitney test showed a significant difference at 625  $\mu\text{g/ml}$  of the extract when compared to the control ( $p < .05$ ). Besides, a significant decrease in the percentage of acrosome-reacted spermatozoa was observed between treatment groups, that is 6.25 and 625  $\mu\text{g/ml}$  ( $p < .05$ ), and 62.5 and 625  $\mu\text{g/ml}$  ( $p < .05$ ).

## 4 | DISCUSSION

Pharmacological properties of *M. oleifera* have been studied at length using in vivo and in vitro models for efficacy assessments in nearly all body systems (Igado & Olopade, 2016), and biological activities were ascribed to its antioxidant activities (Vergara-Jimenez et al., 2017). However, most reports on the effects of MO on male reproductive, sexual and seminal functions were conducted using animal models (Prabsattroo et al., 2012). A previous in vivo study demonstrated the fertility properties of MO by the increased FSH and LH levels, the



**FIGURE 2** Effects of aqueous leaf extracts of *Moringa oleifera* on capacitation and acrosome reaction in vitro. (a) F pattern corresponded to non-capacitated and acrosome intact spermatozoa (b) B pattern corresponded to capacitate and acrosome intact spermatozoa and (c) AR pattern corresponded to acrosome-reacted spermatozoa. Data were represented as mean  $\pm$  SEM of 200 spermatozoa from 34 healthy donors

number of Leydig and Sertoli cells, testicular weight, caudal epididymal sperm count and serum testosterone levels, with a resultant decrease in prolactin, in male albino rats (Orji et al., 2016). *Moringa* saponins, flavonoids and alkaloids have been implicated in some of

the observed changes due to their ability to affect androgen synthesis and spermatogenesis (Zade et al., 2013). In other in vivo studies, however, sperm qualitative characteristics in male rats remained unchanged following treatment with MO (Awodele et al., 2012; El-Sheikh et al., 2016). Sperm functional capacity in mammals depends strictly on their morphology, progressive motility, intact plasma and acrosome membranes (Boshoff et al., 2018). To the best of our knowledge, this is the first scientific report to evaluate the effects of aqueous MO leaf extract on the physiological functions of human spermatozoa in vitro.

Our study revealed that MO aqueous leaf extract did not induce any significant changes in percentages of sperm motility parameters, vitality, and capacitation. A possible explanation for this is that the level of sperm ROS produced did not elicit any deleterious effects on sperm membrane integrity, viability and motility (Afolabi et al., 2013). Higher levels of ROS in semen and low levels of antioxidants are observed in infertile men compared to their fertile counterpart (Agarwal et al., 2016; Roychoudhury et al., 2016). Inclusion of semen samples from infertile men (e.g. asthenozoospermia or teratozoospermia) may have demonstrated the anti-oxidative effects of *M. oleifera* leaf on sperm motility and viability.

Furthermore, this study revealed that the level of MMP in sperm remained unchanged following the treatment with MO aqueous leaf extract. MMP is an indicator of the energetic state of mitochondria and determines functional intact mitochondria that have been related mainly to sperm motility (Marchetti et al., 2002). MMP provides insightful information on the sperm fertilising capacity in comparison to the sperm motility, as spermatozoa with high MMP are shown to undergo acrosome reaction (Gallon et al., 2006; Slowińska et al., 2018).

Nutritional antioxidants have been prescribed for pathologies in which oxidative stress was implicated as a cause, including male infertility (Balercia et al., 2005; De Young et al., 2004). *Moringa oleifera* had been shown to possess a high amount of antioxidant with increasing concentration (Luqman et al., 2012), for which quercetin and kaempferol are amongst the abundant flavonoids in MO leaves (Verma et al., 2009). In the present study, the aqueous leaf extract of MO showed a significant decrease in intracellular ROS production in sperm in a dose-dependent manner, specifically at 62.5 and 625 µg/ml, which could be due to scavenging activities of the antioxidants present in the extracts. Addition of quercetin to sperm suspension significantly improved sperm viability, morphology, motility and antioxidants and inhibited lipid peroxidation (Ben Abdallah et al., 2011). Also, alpha-lipoic acid, an organo-sulphur compound derived from octanoic acid, is regarded as a universal antioxidant as it acts as a cofactor for many enzymes and is shown to restore endogenous antioxidants such as oxidised vitamin C and E, thereby protecting organelles, cells and tissue from the damaging effects of excessive ROS production (Taherian et al., 2019). *Moringa oleifera* leaves are also rich in vitamin C and E (Ajantha et al., 2018), which have been indicated as major chain-breaking antioxidants with strong reducing potential that attenuate free radical generating reactions (Brash & Havre, 2002). Seminal intracellular enzyme complexes and

their involvement in male fertility regulation are highly dependent on cofactors such as Zn, Se, Mg, Mn, Fe<sup>3+</sup> and Cu (Dobrakowski et al., 2018) whose presence in MO leaves has been validated (Dilawar et al., 2018). Centrifugation during the processing of semen increases the generation of ROS results in a loss in sperm motility and viability and induces DNA damage in spermatozoa (Henkel et al., 2012; Taherian et al., 2019). Semen samples obtained from normozoospermic men, washed and centrifuged with sperm washing media, in the presence of alpha-lipoic acid (0.02 mM) maintained sperm viability, and motility through the reduction of ROS production as well as prevention DNA damage (Taherian et al., 2019). Therefore, the addition of aqueous leaf extract of MO in the sperm wash media before centrifugation may improve sperm motility and viability. To support this, the addition of crude extracts of MO in bovine semen extender significantly increased Bull's sperm motility (Sokunbi et al., 2015). Hence, the reduced ROS production, as well as the maintained motility and viability seen in this study, may be attributed to the antioxidant activities of the leaf extract of MO.

An inverse relationship exists between oxidative stress and sperm DNA integrity (Dorostghoal et al., 2017). Excessive ROS augments DNA instability through direct modification of nitrogenous bases (Koh et al., 2016), disruption of a maturation stage in spermatogenesis where spermatozoal histones are protaminated (chromatin condensation) (Hamilton et al., 2018), and occurrence of dysregulated abortive apoptosis (Sakkas et al., 2003). Lipid peroxidation and increased membrane fluidity due to polyunsaturated fatty acid (PUFA) oxidation (Tvrdá et al., ), phosphatidyl-serine translocation (Barroso et al., 2000) and caspase and endonuclease activation (Jeng et al., 2015; Wang et al., 2003) are the hallmarks of sperm ROS-induced apoptotic DNA damage (Aitken & Koppers, 2011). Our study revealed a significant reduction in the percentage of TUNEL-positive sperm (DNA fragmentation) following a 1-hr incubation with aqueous leaf extract of MO. Since superoxide availability precedes free radical production in sperm microenvironment (Gosalvez et al., 2017), it is possible that the absence or inhibition of superoxide (as scavenged or inhibited by MO leaf extracts) resulted in subsequent inhibition on the production of peroxidative derivatives such as H<sub>2</sub>O<sub>2</sub> and nitric oxide that initiate and facilitate DNA damage (Verma et al., 2009), hence the significant reduction. Therefore, to the best of our knowledge, this study is first to demonstrate the protective ability of MO leaf extracts on human sperm nuclear DNA and oxidative stress in vitro.

Low levels of ROS are prerequisites for induction of processes crucial for fertilisation such as sperm hyperactivation and capacitation, acrosome reaction, gamete recognition and binding as well as gamete fusion (Aitken et al., 1995; Glenn et al., 2007; de Lamirande & Gagnon, 1993, 1995; Sánchez et al., 2010). Premature acrosome reaction in spermatozoa renders them incapable of fertilisation (Glenn et al., 2007). Aqueous leaf extract of MO significantly increased the percentage of non-capacitated and acrosome intact spermatozoa (F pattern) at 625 µg/ml but had no significant effect on the percentage of acrosome-reacted spermatozoa, which may be associated with the decreased level of ROS observed in this study.

Also, the number of capacitated and acrosome intact spermatozoa (B pattern) remained unchanged at all concentrations of the plant extract. Since MO leaf extracts did not affect sperm hyperactivation, a possible explanation for the unaltered percentage of capacitated spermatozoa could be associated to the fact that the onset of capacitation coincides with hyperactivation.

Moringa, which is a rich source of vitamin E, is accessible, affordable and a readily available source of major essential nutrients and nutraceuticals (Kunyanga et al., 2013). The estimated cost for MO leaf powder is about US\$0.05/g, MO capsule (450 mg dried leaf) is US \$0.07/capsule and compared to vitamin E capsule (1,000 IU, 671 mg) is US\$ 0.21/capsule. Dried MO leaves retain their nutrients and can be stored for up to 1 year provided they are stored in airtight containers, away from light and humidity, and kept below 24°C (Doerr & Cameroon, 2005; Gopalakrishnan et al., 2016). Drying of leaves with an oven at 50°C for 16 hr was shown to maintain most of the nutrients and phytochemicals in the leaves except for vitamin C compared to the freeze-dried leaves. Thus, the mild heating and drying processes of the leaves could be achieved using household appliances such as stove (Yang et al., 2006), making the plant readily available for use in enhancing sperm functions.

To conclude, the aqueous leaf extract of *M. oleifera* maintained human sperm function by the scavenging ability of its antioxidant compounds. Concentration-dependent effects of MO proved that the extract confers significant protection against excess ROS production and DNA fragmentation in sperm population and inhibited premature capacitation and acrosome reaction. The preservation of sperm acrosome integrity and inhibition of spontaneous AR in this study are indicative of the fertility potential of MO extract. MO might be valuable in infertility therapies, and more research is required, particularly in vivo studies, to understand the fertilising capacity of the plant extract and to establish whether moringa products have clinical value. For future work, a study on the antioxidant capacity of MO leaf extract in human sperm should be conducted with standardised compounds such as vitamin E, butylated hydroxytoluene or melatonin for evaluating the efficacy of the plant.

## ACKNOWLEDGEMENTS

This study was supported by National Research Foundation of South Africa (TTK170426228930) and University of Limpopo seed fund. The authors thank the donors. We thank the comparative spermatology laboratory of the University of the Western Cape, under the supervision of Dr. Liana Maree for providing Sperm Class Analyser. Fresh leaves of *M. oleifera* were the gift from Prof Kingsley Ayisi and Ms Paulina Bopape-Mabapa of the University of Limpopo. The authors thank the editor and reviewers for their contribution.

## CONFLICT OF INTEREST

The authors declare no conflict of interests.

## AUTHORS' CONTRIBUTIONS

FTM conducted the experiments, performed the statistical analysis and wrote the first draft of the manuscript. GAA and RRH revised

the manuscript. CSO conceptualised the study and revised the manuscript. All authors read and approved the final manuscript.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

#### ORCID

Faith T. Moichela  <https://orcid.org/0000-0003-0726-4263>

Gbenga A. Adefolaju  <https://orcid.org/0000-0002-4087-1690>

Ralf R. Henkel  <https://orcid.org/0000-0003-1128-2982>

Chinyerum S. Opuwari  <https://orcid.org/0000-0002-0804-2278>

#### REFERENCES

- Acuram, L. K., & Chichico Hernandez, C. L. (2019). Anti-hypertensive effect of *Moringa oleifera* Lam. *Cogent Biology*, 5(1), 1596526. <https://doi.org/10.1080/23312025.2019.1596526>
- Adageba, R. K., Maya, E. T., Annan, J. J., & Damalie, F. J. (2015). Setting up and running a successful IVF program in Africa: Prospects and challenges. *The Journal of Obstetrics and Gynecology of India*, 65(3), 155–157. <https://doi.org/10.1007/s13224-015-0719-4>
- Afolabi, A. O., Aderoju, H. A., & Alagbonsi, I. A. (2013). Effects of methanolic extract of *Moringa oleifera* leaves on semen and biochemical parameters in cryptorchid rats. *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, 10(5), 230–235. <https://doi.org/10.4314/ajtcam.v10i5.3>
- Agarwal, A., Mulgund, A., Hamada, A., & Chyatte, M. R. (2015). A unique view on male infertility around the globe. *Reproductive Biology and Endocrinology: RB&E*, 13, 37. <https://doi.org/10.1186/s12958-015-0032-1>
- Agarwal, A., Parekh, N., Selvam, P., Kumar, M., Henkel, R., Shah, R., Homa, S. T., Ramasamy, R., Ko, E., Tremellen, K., Esteves, S., Majzoub, A., Alvarez, J. G., Gardner, D. K., Jayasena, C. N., Ramsay, J. W., Cho, C.-L., Saleh, R., Sakkas, D., ... Harlev, A. (2019). Male Oxidative Stress Infertility (MOSI): Proposed terminology and clinical practice guidelines for management of idiopathic male infertility. *The World Journal of Men's Health*, 37(3), 296.
- Agarwal, A., Roychoudhury, S., Bjugstad, K. B., & Cho, C.-L. (2016). Oxidation-reduction potential of semen: What is its role in the treatment of male infertility? *Therapeutic Advances in Urology*, 8(5), 302–318. <https://doi.org/10.1177/1756287216652779>
- Ahmed, K. S., Jahan, I. A., Hossain, M. H., Ethane, N. J., & Saha, B. (2018). Mineral and trace element content in different parts of *Moringa oleifera* grown in Bangladesh. *Current Journal of Applied Science and Technology*, 31(5), 1–10. <https://doi.org/10.9734/CJAST/2018/45645>
- Aitken, R. J., & Koppers, A. J. (2011). Apoptosis and DNA damage in human spermatozoa. *Asian Journal of Andrology*, 13(1), 36–42. <https://doi.org/10.1038/aja.2010.68>
- Aitken, R. J., Paterson, M., Fisher, H., Buckingham, D. W., & van Duin, M. (1995). Redox regulation of tyrosine phosphorylation in human spermatozoa and its role in the control of human sperm function. *Journal of Cell Science*, 108(5), 2017–2025.
- Ajantha, A., Kathirvelan, C., Purushothaman, P. M. R., & Visha, P. (2018). Study on nutrients, mineral and vitamin profile of *Moringa oleifera* leaf meal meant for livestock feeding. *International Journal of Current Microbiology and Applied Sciences*, 7, 2478–2481. <https://doi.org/10.20546/ijcmas.2018.705.284>
- Anudeep, S., & Radha, C. (2018). Carbohydrates of *Moringa oleifera* seeds. *International Journal of Research and Analytical Reviews*, 5(4), 103–108.
- Aprioku, J. S., & Onyenaturuchi, P. O. (2018). Comparative phytochemical analysis of *Moringa oleifera* (Moringaceae) extracts and reproductive effect of the aqueous leaf, seed, and bark extracts in rats. *World Journal of Pharmaceutical Research*, 7(9), 42–54.
- Awodele, O., Oreagba, I. A., Odoma, S., da Silva, J. A. T., & Osunkalu, V. O. (2012). Toxicological evaluation of the aqueous leaf extract of *Moringa oleifera* Lam. (Moringaceae). *Journal of Ethnopharmacology*, 139(2), 330–336. <https://doi.org/10.1016/j.jep.2011.10.008>
- Azu, O. O., Duru, F. I. O., Osinubi, A. A., Noronha, C. C., Elesha, S. O., & Okanlawon, A. O. (2010). Preliminary study on the antioxidant effect of *Kigelia africana* fruit extract (Bignoniaceae) in male Sprague-Dawley rats. *African Journal of Biotechnology*, 9(9), 1374–1381. <https://doi.org/10.5897/AJB10.933>
- Balercia, G., Regoli, F., Armeni, T., Kovrech, A., Mantero, F., & Boscaro, M. (2005). Placebo-controlled double-blind randomized trial on the use of L-carnitine, L-acetylcarnitine, or combined L-carnitine and L-acetylcarnitine in men with idiopathic asthenozoospermia. *Fertility and Sterility*, 84(3), 662–671. <https://doi.org/10.1016/j.fertnstert.2005.03.064>
- Barroso, G., Morshedi, M., & Oehninger, S. (2000). Analysis of DNA fragmentation, plasma membrane translocation of phosphatidylserine and oxidative stress in human spermatozoa. *Human Reproduction*, 15(6), 1338–1344. <https://doi.org/10.1093/humrep/15.6.1338>
- Ben Abdallah, F., Zribi, N., & Ammar-Keskes, L. (2011). Antioxidative potential of Quercetin against hydrogen peroxide induced oxidative stress in spermatozoa in vitro. *Andrologia*, 43(4), 261–265. <https://doi.org/10.1111/j.1439-0272.2010.01063.x>
- Bhattacharya, A., Tiwari, P., Sahu, P. K., & Kumar, S. (2018). A review of the phytochemical and pharmacological characteristics of *Moringa oleifera*. *Journal of Pharmacy & Bioallied Sciences*, 10(4), 181. [https://doi.org/10.4103/JPBS.JPBS\\_126\\_18](https://doi.org/10.4103/JPBS.JPBS_126_18)
- Boshoff, N. H., Lambrechts, H., Maree, L., Cloete, S. W. P., & van der Horst, G. (2018). A novel flush technique to simulate natural dispersal of spermatozoa in the female reproductive tract and expedite motility assessment of fresh ejaculated Merino (*Ovis aries*) sperm. *South African Journal of Animal Science*, 48(3), 469–476.
- Brash, D. E., & Havre, P. A. (2002). New careers for antioxidants. *Proceedings of the National Academy of Sciences of the United States of America*, 99(22), 13969–13971. <https://doi.org/10.1073/pnas.232574399>
- Cajuday, L., & Pocsidio, G. (2010). Effects of *Moringa oleifera* Lam. (Moringaceae) on the reproduction of male mice (*Mus musculus*). *Journal of Medicinal Plants Research*, 4, 1115–1121.
- Cho, C.-L., & Agarwal, A. (2018). Role of sperm DNA fragmentation in male factor infertility: A systematic review. *Arab Journal of Urology*, 16(1), 21–34. <https://doi.org/10.1016/j.aju.2017.11.002>
- de Lamirande, E., & Gagnon, C. (1993). Human sperm hyperactivation and capacitation as parts of an oxidative process. *Free Radical Biology & Medicine*, 14(2), 157–166. [https://doi.org/10.1016/0891-5849\(93\)90006-g](https://doi.org/10.1016/0891-5849(93)90006-g)
- de Lamirande, E., & Gagnon, C. (1995). Capacitation-associated production of superoxide anion by human spermatozoa. *Free Radical Biology & Medicine*, 18(3), 487–495. [https://doi.org/10.1016/0891-5849\(94\)00169-k](https://doi.org/10.1016/0891-5849(94)00169-k)
- De Young, L., Yu, D., Bateman, R. M., & Brock, G. B. (2004). Oxidative stress and antioxidant therapy: Their impact in diabetes-associated erectile dysfunction. *Journal of Andrology*, 25(5), 830–836. <https://doi.org/10.1002/j.1939-4640.2004.tb02862.x>
- Dilawar, S., Shah, A., Shahnaz, S., Khan, H. U., Imad, S., & Shirin, K. (2018). Elemental analysis in the indigenous medicinal plant *Moringa oleifera* and their association with ameliorative activity. *Journal of the Chemical Society of Pakistan*, 40(5), 834–841.
- Dobrakowski, M., Kaletka, Z., Machoń-Grecka, A., Kasperczyk, S., Horak, S., Birkner, E., Zalejska-Fiolka, J., & Kasperczyk, A. (2018). The role of



- oxidative stress, selected metals, and parameters of the immune system in male fertility. *Oxidative Medicine and Cellular Longevity*, 2018, 1–8.
- Dobrakowski, M., Kasperczyk, S., Horak, S., Chyra-Jach, D., Birkner, E., & Kasperczyk, A. (2017). Oxidative stress and motility impairment in the semen of fertile males. *Andrologia*, 49(10), e12783. <https://doi.org/10.1111/and.12783>
- Doerr, B., & Cameroon, I. (2005). *Moringa leaf powder*. ECHO Technical Note. Retrieved from <https://www.echonet.org>
- Dorstghoal, M., Kazeminejad, S. R., Shahbazian, N., Pourmehdi, M., & Jabbari, A. (2017). Oxidative stress status and sperm DNA fragmentation in fertile and infertile men. *Andrologia*, 49(10), e12762. <https://doi.org/10.1111/and.12762>
- Elayaraja, M. S., Gopi, R., & Kumarasamy, D. (2017). Growth regulators induced the photosynthetic pigments of drumstick (*Moringa oleifera*). *Advances in Life Science and Technology*, 53, 20–23.
- El-Sheikh, S. M. A., Khairy, M., Fadil, H. A., & Abo-Elmaaty, A. M. A. (2016). Ameliorative effect of *Moringa oleifera* extract on male fertility in paroxetine treated rats. *Zagazig Veterinary Journal*, 44(3), 244–250. <https://doi.org/10.21608/zvjz.2016.7878>
- Erasmus, N., Solomon, M. C., Fortuin, K. A., & Henkel, R. R. (2012). Effect of *Eurycoma longifolia* Jack (Tongkat ali) extract on human spermatozoa in vitro. *Andrologia*, 44(5), 308–314. <https://doi.org/10.1111/j.1439-0272.2012.01282.x>
- Erhabor, J. O., & Idu, M. (2017). Aphrodisiac potentials of the ethanol extract of *Aloe barbadensis* Mill. Root in male Wistar rats. *BMC Complementary and Alternative Medicine*, 17(1), 360. <https://doi.org/10.1186/s12906-017-1866-1>
- Fahey, J. W., Wade, K. L., Stephenson, K. K., Shi, Y., Liu, H., Panjwani, A. A., Warrick, C. R., & Olson, M. E. (2019). A strategy to deliver precise oral doses of the glucosinolates or isothiocyanates from *Moringa oleifera* leaves for use in clinical studies. *Nutrients*, 11(7), 1547. <https://doi.org/10.3390/nu11071547>
- Farooq, F., Rai, M., Tiwari, A., Khan, A. A., & Farooq, S. (2012). Medicinal properties of *Moringa oleifera*: An overview of promising healer. *Journal of Medicinal Plants Research*, 6(27), 4368–4374.
- Gallon, F., Marchetti, C., Jouy, N., & Marchetti, P. (2006). The functionality of mitochondria differentiates human spermatozoa with high and low fertilizing capability. *Fertility and Sterility*, 86(5), 1526–1530. <https://doi.org/10.1016/j.fertnstert.2006.03.055>
- Gandji, K., Chadare, F. J., Idohou, R., Salako, V. K., Assogbadjo, A. E., & Kakaï, R. L. G. (2018). Status and utilisation of *Moringa oleifera* Lam: A review. *African Crop Science Journal*, 26(1), 137–156–156. <https://doi.org/10.4314/acsj.v26i1.10>
- Glenn, D. R. J., McVicar, C. M., McClure, N., & Lewis, S. E. M. (2007). Sildenafil citrate improves sperm motility but causes a premature acrosome reaction in vitro. *Fertility and Sterility*, 87(5), 1064–1070. <https://doi.org/10.1016/j.fertnstert.2006.11.017>
- Gopalakrishnan, L., Doriya, K., & Kumar, D. S. (2016). *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5(2), 49–56. <https://doi.org/10.1016/j.fshw.2016.04.001>
- Gosalvez, J., Tvrdá, E., & Agarwal, A. (2017). Free radical and superoxide reactivity detection in semen quality assessment: Past, present, and future. *Journal of Assisted Reproduction and Genetics*, 34(6), 697–707. <https://doi.org/10.1007/s10815-017-0912-8>
- Green, C. M., Cockle, S. M., Watson, P. F., & Fraser, L. R. (1996). A possible mechanism of action for fertilization promoting peptide, a TRH-related tripeptide that promotes capacitation and fertilizing ability in mammalian spermatozoa. *Molecular Reproduction and Development*, 45(2), 244–252. [https://doi.org/10.1002/\(SICI\)1098-2795\(199610\)45:2<244::AID-MRD19>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1098-2795(199610)45:2<244::AID-MRD19>3.0.CO;2-0)
- Hamilton, T. R. D. S., Siqueira, A. F. P., Castro, L. S. D., Mendes, C. M., Delgado, J. D. C., de Assis, P. M., Mesquita, L. P., Maiorka, P. C., Nichi, M., Goissis, M. D., Visintin, J. A., & Assumpção, M. E. O. D. Á. (2018). Effect of heat stress on sperm DNA: Protamine assessment in ram spermatozoa and testicle [Research article]. *Oxidative Medicine and Cellular Longevity*, 2018, 1–14. <https://doi.org/10.1155/2018/5413056>
- Henkel, R., Fransman, W., Hipler, U.-C., Wiegand, C., Schreiber, G., Menkveld, R., Weitz, F., & Fisher, D. (2012). Typha capensis (Rohrb.) N.E.Br. (bulrush) extract scavenges free radicals, inhibits collagenase activity and affects human sperm motility and mitochondrial membrane potential in vitro: A pilot study. *Andrologia*, 44(s1), 287–294. <https://doi.org/10.1111/j.1439-0272.2011.01179.x>
- Igado, O. O., & Olopade, J. O. (2016). A review on the possible neuro-protective effects of *Moringa oleifera* leaf extract. *Nigerian Journal of Physiological Sciences*, 31(2), 183–187.
- Iommiello, V. M., Albani, E., Di Rosa, A., Marras, A., Menduni, F., Morreale, G., Levi, S. L., Pisano, B., & Levi-Setti, P. E. (2015). Ejaculate oxidative stress is related with sperm DNA fragmentation and round cells. *International Journal of Endocrinology*, 2015, 321901. <https://doi.org/10.1155/2015/321901>
- Jeng, H. A., Pan, C.-H., Chao, M.-R., & Lin, W.-Y. (2015). Sperm DNA oxidative damage and DNA adducts. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 794, 75–82. <https://doi.org/10.1016/j.mrgentox.2015.09.002>
- Koh, S.-A., Sanders, K., & Burton, P. (2016). Effect of male age on oxidative stress markers in human semen. *Journal of Reproductive Biotechnology and Fertility*, 5, 2058915816673242. <https://doi.org/10.1177/2058915816673242>
- Kotta, S., Ansari, S. H., & Ali, J. (2013). Exploring scientifically proven herbal aphrodisiacs. *Pharmacognosy Reviews*, 7(13), 1.
- Koul, B., & Chase, N. (2015). *Moringa oleifera* Lam.: Panacea to several maladies. *Journal of Chemical and Pharmaceutical Research*, 7, 687–707.
- Kumar, N., & Singh, A. K. (2015). Trends of male factor infertility, an important cause of infertility: A review of literature. *Journal of Human Reproductive Sciences*, 8(4), 191–196. <https://doi.org/10.4103/0974-1208.170370>
- Kunyanga, C. N., Imungi, J. K., & Vellingiri, V. (2013). Nutritional evaluation of indigenous foods with potential food-based solution to alleviate hunger and malnutrition in Kenya. *Journal of Applied Biosciences*, 67, 5277–5288. <https://doi.org/10.4314/jab.v67i0.95049>
- Kwaghe, A. V., & Ambali, A. G. (2009). Preliminary phytochemical screening of fresh and dried *Moringa oleifera* leaves and that of chloroform, ethylacetate and n-butanol fractions. *Sahel Journal of Veterinary Sciences*, 8(2), 1–24.
- Lalas, S., & Tsaknis, J. (2002). Extraction and identification of natural antioxidant from the seeds of the *Moringa oleifera* tree variety of Malawi. *Journal of the American Oil Chemists' Society*, 79(7), 677–683.
- Lampiao, F., Krom, D., & du Plessis, S. S. (2008). The in vitro effects of *Mondia whitei* on human sperm motility parameters. *Phytotherapy Research: PTR*, 22(9), 1272–1273. <https://doi.org/10.1002/ptr.2469>
- Leone, A., Spada, A., Battezzati, A., Schiraldi, A., Aristil, J., & Bertoli, S. (2016). *Moringa oleifera* seeds and oil: Characteristics and uses for human health. *International Journal of Molecular Sciences*, 17(12), 2141. <https://doi.org/10.3390/ijms17122141>
- Luqman, S., Srivastava, S., Kumar, R., Maurya, A. K., & Chanda, D. (2012). Experimental assessment of *Moringa oleifera* leaf and fruit for its antioxidant, and scavenging potential using in vitro and in vivo assays. *Evidence-Based Complementary and Alternative Medicine*, 2012, 519084.
- MacDonald, I., Oghale, O.-U., & Hauwa, G. J. (2016). Ethnomedicinal survey of plants used in the treatment of female infertility in Chanchaga Niger State, Nigeria. *Marmara Pharmaceutical Journal*, 20(3), 269–274.
- Makuch, M. Y., Simônia de Padua, K., Petta, C. A., Duarte Osis, M. J., & Bahamondes, L. (2011). Inequitable access to assisted reproductive technology for the low-income Brazilian population: A qualitative study. *Human Reproduction*, 26(8), 2054–2060. <https://doi.org/10.1093/humrep/der158>

- Maldini, M., Maksoud, S. A., Natella, F., Montoro, P., Petretto, G. L., Foddai, M., Nicola, G. R. D., Chessa, M., & Pintore, G. (2014). *Moringa oleifera*: Study of phenolics and glucosinolates by mass spectrometry. *Journal of Mass Spectrometry*, 49(9), 900–910. <https://doi.org/10.1002/jms.3437>
- Marchetti, C., Obert, G., Deffosez, A., Formstecher, P., & Marchetti, P. (2002). Study of mitochondrial membrane potential, reactive oxygen species, DNA fragmentation and cell viability by flow cytometry in human sperm. *Human Reproduction*, 17(5), 1257–1265. <https://doi.org/10.1093/humrep/17.5.1257>
- Nandave, M., Ojha, S. K., Joshi, S., Kumari, S., & Arya, D. S. (2009). *Moringa oleifera* leaf extract prevents isoproterenol-induced myocardial damage in rats: Evidence for an antioxidant, antiperoxidative, and cardioprotective intervention. *Journal of Medicinal Food*, 12(1), 47–55.
- Nazim, K., Sun, M. C., & Neergheen-Bhujun, V. S. (2017). Perceptions and practices on the use of *Moringa oleifera* Lam: A qualitative study amongst elderly Mauritians. *Journal of Complementary and Alternative Medical Research*, 2(2), 1–13. <https://doi.org/10.9734/JOCAMR/2017/31461>
- Nishihara, T., Matsumoto, K., Hosoi, Y., & Morimoto, Y. (2018). Evaluation of antioxidant status and oxidative stress markers in follicular fluid for human in vitro fertilization outcome. *Reproductive Medicine and Biology*, 17(4), 481–486. <https://doi.org/10.1002/rmb2.12229>
- Orji, U. H., Ildigwe, E. E., Akah, P. A., Okonta, M. J., Ajaghaku, D. L., & Ihekwereme, C. P. (2016). *Moringa oleifera* increases serum level of testosterone and gonadotropins in male albino rats. *Journal of International Research in Medical and Pharmaceutical Sciences*, 9(1), 41–48.
- Prabsaturoo, T., Wattanathorn, J., Iamsaard, S., Muchimapura, S., & Thukhammee, W. (2012). *Moringa oleifera* leaves extract attenuates male sexual dysfunction. *American Journal of Neuroscience*, 3(1), 17–24.
- Prasanth, D., Sreedhara, C. S., Deepak, M., Sangli, V. K., & Gururaj, V. M. (2011). Comparative study on estimation of polyphenols in different extracts of *Moringa oleifera* leaves and fruits with respect to tannic acid. *Journal of Pharmacy Research*, 4, 3224–3225.
- Rakuambo, N. C., Meyer, J. J. M., Hussein, A., Huyser, C., Mdlalose, S. P., & Raidani, T. G. (2006). In vitro effect of medicinal plants used to treat erectile dysfunction on smooth muscle relaxation and human sperm. *Journal of Ethnopharmacology*, 105(1–2), 84–88. <https://doi.org/10.1016/j.jep.2005.10.008>
- Rodríguez-Pérez, C., Quirantes-Piné, R., Fernández-Gutiérrez, A., & Segura-Carretero, A. (2015). Optimization of extraction method to obtain a phenolic compounds-rich extract from *Moringa oleifera* Lam leaves. *Industrial Crops and Products*, 66, 246–254. <https://doi.org/10.1016/j.indcrop.2015.01.002>
- Roychoudhury, S., Sharma, R., Sikka, S., & Agarwal, A. (2016). Diagnostic application of total antioxidant capacity in seminal plasma to assess oxidative stress in male factor infertility. *Journal of Assisted Reproduction and Genetics*, 33(5), 627–635. <https://doi.org/10.1007/s10815-016-0677-5>
- Sakkas, D., Seli, E., Bizzaro, D., Tarozzi, N., & Manicardi, G. C. (2003). Abnormal spermatozoa in the ejaculate: Abortive apoptosis and faulty nuclear remodelling during spermatogenesis. *Reproductive BioMedicine Online*, 7(4), 428–432. [https://doi.org/10.1016/S1472-6483\(10\)61886-X](https://doi.org/10.1016/S1472-6483(10)61886-X)
- Sánchez, R., Sepúlveda, C., Risopatrón, J., Villegas, J., & Giojalas, L. C. (2010). Human sperm chemotaxis depends on critical levels of reactive oxygen species. *Fertility and Sterility*, 93(1), 150–153. <https://doi.org/10.1016/j.fertnstert.2008.09.049>
- Schwetz, V., Gumpold, R., Graupp, M., Hacker, N., Schweighofer, N., Trummer, O., Pieber, T. R., Ballon, M., Lerchbaum, E., & Obermayer-Pietsch, B. (2013). Osteocalcin is not a strong determinant of serum testosterone and sperm count in men from infertile couples. *Andrology*, 1(4), 590–594. <https://doi.org/10.1111/j.2047-2927.2013.00095.x>
- Shalaweh, S. M., Erasmus, N., Weitz, F., & Henkel, R. R. (2015). Effect of *Cissampelos capensis* rhizome extract on human spermatozoa in vitro. *Andrologia*, 47(3), 318–327.
- Shunmugam, L. (2016). *Moringa oleifera* crude aqueous leaf extract induces apoptosis in human hepatocellular carcinoma cells via the upregulation of NF- $\kappa$ B and IL-6/STAT3 pathway. [PhD Thesis]. ukzn-dspace.ukzn.ac.za
- Slowińska, M., Liszewska, E., Judycka, S., Konopka, M., & Ciereszko, A. (2018). Mitochondrial membrane potential and reactive oxygen species in liquid stored and cryopreserved turkey (*Meleagris gallopavo*) spermatozoa. *Poultry Science*, 97(10), 3709–3717. <https://doi.org/10.3382/ps/pey209>
- Sokunbi, O., Ajani, O., Lawanson, A., & Amao, E. (2015). Antibiotic potential of *Moringa oleifera* Lam. crude extract in bull semen extender. *European Journal of Medicinal Plants*, 9(2), 1–8. <https://doi.org/10.9734/EJMP/2015/18546>
- Sulaiman, M. R., Zakaria, Z. A., Bujarimin, A. S., Somchit, M. N., Israif, D. A., & Moin, S. (2008). Evaluation of *Moringa oleifera* aqueous extract for antinociceptive and anti-inflammatory activities in animal models. *Pharmaceutical Biology*, 46(12), 838–845.
- Taherian, S., Khayamabed, R., Tavalae, M., & Nasr-Esfahani, M. H. (2019). Alpha-lipoic acid minimises reactive oxygen species-induced damages during sperm processing. *Andrologia*, 51(8), e13314. <https://doi.org/10.1111/and.13314>
- Tambi, M. I. B. M., & Imran, M. K. (2010). *Eurycoma longifolia* Jack in managing idiopathic male infertility. *Asian Journal of Andrology*, 12(3), 376. <https://doi.org/10.1038/aja.2010.7>
- Tatone, C., Di Emidio, G., Vento, M., Ciriminna, R., & Artini, P. G. (2010). Cryopreservation and oxidative stress in reproductive cells. *Gynecological Endocrinology*, 26(8), 563–567. <https://doi.org/10.3109/09513591003686395>
- Taweerutchana, R., Lumlerdkij, N., Vannasaeng, S., Akaraseenont, P., & Sriwijitkamol, A. (2017). Effect of *Moringa oleifera* leaf capsules on glycemic control in therapy-naive type 2 diabetes patients: A randomized placebo controlled study. *Evidence-Based Complementary and Alternative Medicine*, 2017, 6581390.
- Teixeira, E. M. B., Carvalho, M. R. B., Neves, V. A., Silva, M. A., & Arantes-Pereira, L. (2014). Chemical characteristics and fractionation of proteins from *Moringa oleifera* Lam Leaves. *Food Chemistry*, 147, 51–54. <https://doi.org/10.1016/j.foodchem.2013.09.135>
- Thilza, I. B., Sanni, S., Zakari, A. I., Sanni, F. S., Muhammed, T., & Musa, B. J. (2010). In vitro antimicrobial activity of water extract of *Moringa oleifera* leaf stalk on bacteria normally implicated in eye diseases. *Academia Arena*, 2(6), 80–82.
- Tvrđá, E., Massanyi, P., & Lukáč, N. (2017). Physiological and pathological roles of free radicals in male reproduction. In R. Meccariello & R. Chianese (Eds.), *Spermatozoa-facts and perspectives*. IntechOpen. <https://doi.org/10.5772/intechopen.70793>
- Vergara-Jimenez, M., Almatrafi, M., & Fernandez, M. (2017). Bioactive components in *Moringa oleifera* leaves protect against chronic disease. *Antioxidants*, 6(4), 91. <https://doi.org/10.3390/antiox6040091>
- Verma, A. R., Vijayakumar, M., Mathela, C. S., & Rao, C. V. (2009). In vitro and in vivo antioxidant properties of different fractions of *Moringa oleifera* leaves. *Food and Chemical Toxicology*, 47(9), 2196–2201. <https://doi.org/10.1016/j.fct.2009.06.005>
- Virtanen, H. E., Jørgensen, N., & Toppari, J. (2017). Semen quality in the 21st century. *Nature Reviews Urology*, 14(2), 120–130. <https://doi.org/10.1038/nrurol.2016.261>
- Wafa, W., El-Nagar, H., Gabir, A., & Rezk, M. (2017). Impact of dietary *Moringa oleifera* leaves supplementation on semen characteristics, oxidation stress, physiological response and blood parameters of heat stressed buffalo bulls. *Journal of Animal and Poultry Production – Mansoura University*, 8, 367–379. <https://doi.org/10.21608/jappmu.2017.46008>

- Wang, X., Sharma, R. K., Sikka, S. C., Thomas, A. J., Falcone, T., & Agarwal, A. (2003). Oxidative stress is associated with increased apoptosis leading to spermatozoa DNA damage in patients with male factor infertility. *Fertility and Sterility*, 80(3), 531–535. [https://doi.org/10.1016/S0015-0282\(03\)00756-8](https://doi.org/10.1016/S0015-0282(03)00756-8)
- Waterman, C., Cheng, D. M., Rojas-Silva, P., Poulev, A., Dreifus, J., Lila, M. A., & Raskin, I. (2014). Stable, water extractable isothiocyanates from *Moringa oleifera* leaves attenuate inflammation in vitro. *Phytochemistry*, 103, 114–122. <https://doi.org/10.1016/j.phytochem.2014.03.028>
- WHO (2010). WHO | WHO laboratory manual for the examination and processing of human semen. WHO. Retrieved from <http://www.who.int/reproductivehealth/publications/infertility/9789241547789/en/>
- Yang, R.-Y., Chang, L.-C., Hsu, J.-C., Weng, B. B. C., Palada, C., Chadha, M. L., & Levasseur, V. (2006). Nutritional and functional properties of Moringa Leaves – From germplasm, to plant, to food, to health. In Proceedings of the Moringa and other Highly Nutritious Plant Resources: Strategies, Standards and Markets for a Better Impact on Nutrition in Africa, Accra, Ghana, 16–18 November 2006; pp. 1–8.
- Yuan, H., Ma, Q., Ye, L., & Piao, G. (2016). The traditional medicine and modern medicine from natural products. *Molecules*, 21(5), 559. <https://doi.org/10.3390/molecules21050559>
- Zade, V. S., Dabhadkar, D. K., Thakare, V. G., & Pare, S. R. (2013). Effect of aqueous extract of *Moringa oleifera* seed on sexual activity of male albino rats. *Biological Forum – an International Journal*, 5(1), 129–140.

**How to cite this article:** Moichela FT, Adefolaju GA, Henkel RR, Opuwari CS. Aqueous leaf extract of *Moringa oleifera* reduced intracellular ROS production, DNA fragmentation and acrosome reaction in Human spermatozoa in vitro. *Andrologia*. 2021;53:e13903. <https://doi.org/10.1111/and.13903>