

Aromatase activity of human mesenchymal stem cells is stimulated by early differentiation, vitamin D and leptin

Ana María Pino, Juan Manuel Rodríguez, Susana Ríos, Pablo Astudillo, Laura Leiva¹, Germán Seitz², Mireya Fernández and J Pablo Rodríguez

Laboratorio de Biología Celular, INTA, Universidad de Chile, El Líbano 5524, Macul. Casilla 138-11, Santiago, Chile

¹Laboratorio de Envejecimiento y Enfermedades Crónicas Relacionadas con la Nutrición, INTA, Universidad de Chile, Santiago, Chile

²Servicio de Traumatología, Hospital Sótero del Río, Santiago, Chile

(Requests for offprints should be addressed to J P Rodríguez; Email: jprodrig@inta.cl)

Abstract

Human mesenchymal stem cells (hMSCs) are multipotent cells present in bone marrow, which differentiate into osteoblasts and adipocytes, among other lineages. Oestrogens play a critical role in bone metabolism; its action may affect the adipocyte to osteoblast ratio in the bone marrow. In hMSCs, oestrogens are synthesized from C19 steroids by the enzyme aromatase cytochrome P450. In this study, we assessed whether aromatase enzymatic activity varied through early osteogenic (OS) and adipogenic (AD) differentiation. Also, we studied the effect of leptin and 1,25 dihydroxyvitamin D₃ (1,25(OH)₂D₃) on aromatase cell activity. Finally, we analysed whether conditions that modify oestrogen generation by cells affected hMSCs differentiation. For these purposes, hMSCs derived from post-menopausal women (65–86 years old) were cultured under basal, OS or AD conditions, in the presence or the absence of leptin and 1,25(OH)₂D₃. Aromatase activity was

measured by the tritiated water release assay and by direct measurement of steroids synthesized from ³H-labelled androstenedione or testosterone. Our results showed that different OS and AD patterns of aromatase activity developed during the first period of differentiation (up to 7 days). A massive and sharp surge of aromatase activity at 24 h characterized early OS differentiation, while increased but constant aromatase activity was increased through adipogenesis. Both leptin and vitamin D increased aromatase activity during osteogenesis, but not during adipogenesis; finally, we showed that favourable aromatase substrates concentration restrained MSCs adipogenesis but improved osteogenesis. Thus, it could be inferred that a high and early increase of local oestrogen concentration in hMSCs affects their commitment either restraining AD or facilitating OS differentiation, or both.

Journal of Endocrinology (2006) **191**, 715–725

Introduction

Human bone marrow stroma contains mesenchymal stem cells (hMSCs) differentiating along osteogenic, chondrogenic, adipogenic and marrow stromal lineages (Caplan 1991, Bruder *et al.* 1997, Pittenger *et al.* 1999). Changes in the functional characteristics of hMSCs or in the regulation of the differentiation pathways may have consequences in some osteogenic disorders like human postmenopausal osteoporosis (Gimble *et al.* 1996, Nuttall *et al.* 1998, Bianco & Robey 1999, Rodríguez *et al.* 1999).

After the menopause, decreased endogenous oestradiol enhances bone turnover and this is accompanied by a shift in the adipocyte to osteoblast ratio, which favours fat tissue production in the bone marrow (Gambacciani *et al.* 1997, Justesen *et al.* 2001). The directive effect of oestrogen on the skeleton is supported by the developmental failure of bone in males with deficient oestrogen activity as a result of oestrogen receptor dysfunction or aromatase deficiency (Smith *et al.* 1994,

Morishima *et al.* 1995), and the correlation between endogenous oestradiol concentrations and both mineral density and bone loss in men (Amin *et al.* 2000, Khosla *et al.* 2001). In addition, there is evidence that endogenous oestrogen production by CYP 19 aromatase as well as oestrogen receptor signalling play an important role in the development and the distribution of white adipose tissue in the body, as highlighted by analysis of the respective oestrogen receptor- α and ArKO mice (Heine *et al.* 2000, Jones *et al.* 2000).

The biosynthesis of oestrogen from C19 steroids is catalysed by aromatase cytochrome P450 encoded by the *CYP19* gene. In addition to gonads, this enzyme is found in different organs, including adipose tissue, brain, skin, endothelium and bone. Skeletal cells also express a number of other enzymes implicated in sex steroid metabolism (Schweikert *et al.* 1980, 1995, Janssen *et al.* 1999, Compston 2002, Ishida *et al.* 2002, Issa *et al.* 2002), supporting the concept that active androgens and oestrogens can be synthesized within the bone marrow cells from circulating

C19 precursors. Thus, besides contributing to the circulating oestrogen pool, the oestrogen synthesized within bone tissue compartments may be locally active in a paracrine or intracrine way (Labrie *et al.* 1997, Simpson 2000, Simpson & Davis 2001). Therefore, although the total amount of oestrogen synthesized at any given site could be small, local concentrations, could be substantial, giving it functional meaning. The extent, regulation and physiological significance of oestrogen synthesis within the bone remains almost unknown; however, this process could provide mechanisms for tissue-specific responses in the absence of changes in systemic hormone production, and for the preservation of homeostasis in the face of alterations in hormonal status, such as those originated during aging.

Aromatase has been reported to be expressed in hMSCs (Heim *et al.* 2004), in osteoblast or osteoblast-like cells from foetal and adult tissues (Purohit *et al.* 1992, Tanaka *et al.* 1993, Schweikert *et al.* 1995, Sasano *et al.* 1997, Janssen *et al.* 1999), in articular cartilage chondrocytes, in adipocytes adjacent to bone trabeculae, in osteocytes (Sasano *et al.* 1997) and in macrophage/osteoclast-like cells (Shozu *et al.* 1997). The expression of CYP 19 has been shown to be regulated by differential promoter usage, depending on the tissue context. In osteoblasts and adipocytes, aromatase is activated mainly through the I.4 promoter (Shozu & Simpson 1998, Simpson & Davis 2001, Enjuanes *et al.* 2003). In cultures of bone-derived osteoblast or osteoblast-like cells, the regulation of aromatase expression has been studied mainly at the transcriptional level, showing that dexamethasone, vitamin D, testosterone and phytoestrogen genistein, among others, may function as regulatory factors of CYP19 expression (Tanaka *et al.* 1996, Jakob *et al.* 1997, Shozu & Simpson 1998, Shozu *et al.* 2000, Enjuanes *et al.* 2003, Heim *et al.* 2004). Further, transcription of CYP19 has been reported to be induced by physiological or pathological conditions, such as bone differentiation and fractures (Lea *et al.* 1997, Janssen *et al.* 1999, Heim *et al.* 2004), pointing to the importance that local oestrogen generation may have for adequate triggering and ensuing of the differentiation pathway.

Although post-transcriptional modulation of CYP19 has been inferred to account for the differences in cell aromatase enzymatic levels (Tanaka *et al.* 1996, Janssen *et al.* 1999, Heim *et al.* 2004), post-translational modifications, protein stability or cofactor variations have scarcely been studied. These types of mechanisms may be especially relevant during commitment of the common precursor cell to the osteoblastic or adipocytic lineages. The osteogenic (OS) differentiation of cultured hMSCs has been shown to be dependent on the activation of runt-related transcription factor 2 (*runx2*) and extracellular signal-regulated kinase–mitogen-activated protein kinase (ERK–MAPK; Banerjee *et al.* 1997, Ducy *et al.* 1997); while the activation of p38–MAPK and peroxisome proliferator-activated receptor- γ 2 accompanied by suppression of *runx2* expression were shown to induce adipocytic (AD) differentiation (Lecka-Czernik *et al.* 1999). Recently, using selective inhibitors of MEK-1/2 (MAPK/ERK) in bone-derived

osteoblast-like cells, it has been proposed that MAPK could play an important role in aromatase activation at the post-transcriptional level (Shozu *et al.* 2001).

Besides oestradiol, hormones like vitamin D and leptin are recognized as OS agents. Several *in vitro* studies indicate that stromal cells are responsive to leptin, which increases proliferation, differentiation to osteoblastic lineage and the number of mineralized nodules (Takahashi *et al.* 1997, Thomas *et al.* 1999, Reseland *et al.* 2001), but inhibits differentiation to adipocytes (Thomas *et al.* 1999, Hess *et al.* 2005). These observations suggest that leptin may participate in the regulation of bone mass, but the mechanism remains unclear. We have recently demonstrated the presence of high affinity leptin receptors associated with the cell membranes of hMSCs and a direct protective action of leptin on osteogenesis (Hess *et al.* 2005). On the other hand, vitamin D deficiency is an important risk factor for bone mass loss. The more severe deficiencies cause osteomalacia, decreased bone mineralization, bone pain and spontaneous fractures (Bouillo *et al.* 1995). Both 1,25 dihydroxyvitamin D₃ (1,25(OH)₂D₃) and oestradiol may be involved in the regulation of hMSC differentiation (Komm *et al.* 1988, Bouillo *et al.* 1995). Further, 1,25(OH)₂D₃ increases CYP19 transcripts level in bone cells (Tanaka *et al.* 1996, Enjuanes *et al.* 2003), but its effect on hMSCs aromatase activity has been least studied.

Two recent studies on the expression of CYP19 during MSCs differentiation point to the potential importance of distinctive local oestrogen production and action at OS and AD commitment (Janssen *et al.* 1999, Heim *et al.* 2004). Given that aromatase enzymatic activity has not been analysed during hMSCs commitment and early differentiation, and that each differentiation pathway may give rise to specific and exclusive regulation of aromatase activity, we studied in hMSCs: (1) whether early OS and AD differentiation give rise to definite cell aromatase activities; (2) the effect of two hormones involved in bone metabolism, leptin and 1,25(OH)₂D₃, on aromatase cell activity and (3) whether AD differentiation is affected by defined oestrogenic conditions. Our results showed that during the first period of differentiation (up to 7 days), distinctive OS and AD patterns of aromatase activity developed and that a massive and sharp surge of aromatase activity characterized early OS differentiation, while increased but stable aromatase activity was associated with adipogenesis. Both leptin and 1,25(OH)₂D₃ increased aromatase activity during osteogenesis, but not during adipogenesis; finally, we showed that steady oestrogenic conditions restrained MSCs adipogenesis.

Materials and Methods

Subjects

Postmenopausal women aged 65–86 years, patients from the Trauma Section, Hospital Sótero del Río, Santiago, Chile,

were selected as volunteer bone marrow donors. Written informed consent was obtained from all the subjects. Bone marrow was obtained by iliac crest aspiration during surgical procedures (Rodríguez *et al.* 1999); ethical approval was obtained from the Hospital Sôtero del Río and INTA ethics committees. Donors considered themselves healthy, except for fractures and were not using glucocorticoids or oestrogen replacement therapy.

Reagents

Tissue culture reagents were obtained from Gibco/BRL; ICI 182780 (ICI) was purchased from Tocris Cookson Inc., Ellisville, MO, USA. Cell culture dishes were obtained from Nunc, Naperville, IL, USA. Androst-4-ene-3,17 dione, [1- 3 H(N)]-25 \cdot 3 Ci/mmol was purchased from Perkin-Elmer Sciences, Inc., Boston, MA, USA; androst-4-ene-3,17-dione, [1,2,6,7- 3 H(N)] 85 Ci/mmol was from New England Nuclear, Du Pont Co., Wilmington, DE, USA, and [1,2,6,7- 3 H(N)] testosterone 94 Ci/mmol was from Amersham Biosciences Limited, UK. 4-Androsten-4-ol-3,17-dione (Ar-Inh) and all other reagents were supplied by Sigma. Goat polyclonal antibody anti CYP19 (P450 arom) was purchased from Santa Cruz Biotechnology, Santa Cruz, CA, USA; fluorescein isothiocyanate (FITC)-conjugated rabbit-antigoat IgG and peroxidase-conjugated goat anti-rabbit secondary antibodies were from Rockland, Gilbertsville, PA, USA. ECL chemiluminescence reagents were from Amersham Pharmacia Biotech.

Cell preparation and culture methods

hMSCs were isolated from bone marrow as previously described (Jaiswal *et al.* 1997). Briefly, 10 ml bone marrow aspirate were added to 20 ml Dulbecco's modified Eagle's medium high glucose containing 10% foetal bovine serum (basal medium), and it was then centrifuged to pellet the cells, discarding the fat layer. Cells were suspended in basal medium and fractionated on a 70% Percoll density gradient. The hMSCs-enriched low-density fraction was collected, rinsed with culture medium and plated at a density of $1\text{--}2 \times 10^7$ nucleated cells/100 mm dishes. Cells were cultured at 37 °C in a humidified atmosphere of 5% CO₂. After 4 days in culture, non-adherent cells were removed and fresh culture medium was added. Culture medium was replaced by fresh medium twice weekly. When cultures reached near confluence, cells were detached by a mild treatment with trypsin (0.25%, 5 min, 37 °C) and plated at one-third the original density to allow for continued passage. The experiments were performed after the fourth cell passage.

MCF-7 epithelial cell line (Human tumor cell Bank 22, American Type Culture Collection, Rockville, MD, USA), used as control in western blot studies, were cultured as described (Catalano *et al.* 2003).

Osteogenic differentiation

hMSCs ($1\text{--}1.5 \times 10^5$) were maintained in OS culture medium: basal medium supplemented with 0.1 μ M dexamethasone, 10 mM β -glycerophosphate and 50 μ g/ml ascorbic acid (added daily). The medium was changed twice weekly (Rodríguez *et al.* 1999). The ability of hMSCs to differentiate into the osteoblastic lineage *in vitro* was evaluated by measuring alkaline phosphatase activity, as an early osteogenic differentiation marker (Hu *et al.* 2003). At the indicated time (7 days of culture), the culture medium was removed and alkaline phosphatase activity was measured as previously described (Rodríguez *et al.* 2002).

Adipogenic differentiation

hMSCs ($1\text{--}1.5 \times 10^5$ cells/dish, 35 mm) were maintained in AD medium: basal medium supplemented with 1 μ M dexamethasone, 10 μ g/ml insulin, 0.45 mM isobutylmethyl-xanthine and 0.1 mM indomethacin, and this was replaced by fresh medium every 4 days. hMSCs were tested for their lipid content after 14 days of AD treatment by flow cytometry. Cells were placed in freshly diluted Nile Red (1 mg/ml) and analysed by flow cytometry (FACSCalibur, Becton Dickinson, Franklin, NJ, USA; Dennis *et al.* 1999).

Aromatase activity assay

The aromatase activity in hMSCs under specified culture and time conditions was measured by the tritiated water release assay using 8–100 nM [1- 3 H] androst-4-ene-3, 17-dione as substrate, for 2 h at 37 °C (Lephart & Simpson 1991). Cell numbers and protein concentrations were measured in cell monolayers.

Determination of steroid formation from aromatase substrates: $1\text{--}1.2 \times 10^5$ cells/dish were incubated in OS or AD medium in the presence of 30 or 100 nM androstenedione plus 0.3 μ Ci 3 H-androstenedione or 100 nM testosterone plus 0.3 μ Ci 3 H-testosterone, as aromatase substrates. Cells were incubated for 24 h with treatments; the reaction was stopped by placing the plates on ice. The medium was removed, placed in corresponding glass stoppered test tubes. All hormones were extracted four times with a three-fold volume of ethyl ether and the ether phases were pooled. All samples were evaporated to dryness under nitrogen and re-dissolved in 150 μ l ethanol, immediately prior to spotting on TLC plate. Hormones were separated by thin layer chromatography (TLC) using aluminium-backed silica gel-coated plates (60F254, EM Science, Darmstadt, Germany). Each sample included 0.1 μ M standards of oestradiol, oestrone and androstenedione for identification of sample bands. The solvent system (mobile phase) consisted of chloroform/ethyl acetate/(4:1, v/v). Extraction consistencies were controlled using blank incubations without cells that contained known amounts of the radioactive hormones.

Immunofluorescence staining

Cells were seeded on sterile glass coverslips (1.25×10^4 cells/cm²) and placed into 15 mm wells containing basal medium. After 3–5 days, cells were incubated with basal, OS or AD media for 24 h. Cells were washed thrice with PBS, and fixed with ice cold methanol for 20 min at -20°C . The fixed cells were re-hydrated with Tris buffer saline (TBS) and incubated for 1 h in blocking solution (3% BSA in TBS) at room temperature. Cells were incubated with goat polyclonal antibody anti CYP19 (P450 arom), at 1:1000 dilution in 3% BSA–TBS, during 45 min at 37°C and subsequently with the secondary antibody, FITC-conjugated rabbit-antigoat IgG, at a 1:250 dilution in 3% BSA–TBS. Finally, the cells were rinsed in TBS, mounted in DABCO/mowiol and examined with an epifluorescence microscope (100 \times objectives, Nikon, Labophot-2, Tokyo, Japan). In the controls, the first or the second antibodies were omitted.

Western-blot analysis

hMCSs cells were grown in 100 mm dishes to 70–80% confluence. At selected times, cells were lysed in 500 μl of 50 mM HEPES, pH 7.5, 150 mM NaCl, 1.5 mM MgCl, 1 mM EGTA, 10% glycerol, 1% Triton X-100 and a mixture of proteases inhibitors (aprotinin, *p*-methylsulphonyl fluoride and sodium orthovanadate). Further 12 μg protein were separated on 10% SDS-PAGE under reducing conditions. Afterwards, gels were blotted onto polyvinylidene fluoride (PDVF) membrane (Bio-Rad) and aromatase was detected using goat polyclonal antibody anti-CYP19 (P450 arom). Peroxidase-conjugated rabbit anti-goat secondary antibodies

were used. Immunoreactivity was determined using the ECL chemiluminescence reaction. MCF-7 cells were used as a positive control.

Statistical analysis

Statistically significant differences between groups were detected using ANOVA and an *a posteriori* Tukey test. All analyses were performed using STATISTICA 6.1 (StatSoft, Inc., Tulsa, OK, USA; 2004, www.statsoft.com). In all cases, $P < 0.05$ was considered significant.

Results

Aromatase activity in hMSCs under OS and AD differentiation conditions

Aromatase activity of hMSCs showed significant variations after OS and AD differentiation treatments. Very low aromatase activity was observed in cells under basal conditions, its value being slightly higher than blank values (14 ± 5 fmol/mg). However, after OS or AD stimulation, aromatase activity increased significantly. As shown in Fig. 1A, increased aromatase activity was already observed at 12 h of OS differentiation, reaching the maximum level at 24 h of treatment; afterwards the activity decreased sharply and remained low for the rest of the 7 days of treatment. During AD differentiation, a different temporal pattern of aromatase activity was detected, in that increased aromatase activity was maintained up to 3 days, then declining until 7 days of treatment (Fig. 1B).




Figure 1 Aromatase activity during osteogenic (OS) and adipogenic (AD) differentiation of hMSCs. Cells were cultured in OS (A) and AD (B) medium as described in Materials and Methods. At indicated times, the aromatase activity was evaluated by measuring tritiated water released by cells after incubation with 30 nM [$1-\beta$ -³H] androst-4-ene-3, 17-dione as substrate, for 2 h at 37°C . Results were expressed as femtomoles of [^3H]H₂O produced/milligram of protein \times 2 h. Experiments were performed in triplicate from four different samples. Results are the mean \pm s.d. * $P < 0.05$ compared with basal value.

Further characterization of the enzyme was performed in cells after 24 h of OS or AD differentiation treatments. Aromatase activity increased in relation to androstenedione concentration, so that the observed Michaelis–Menten constant (K_m) was 9.22 ± 0.3 nM, although variations in maximal activity were noticed when comparing samples from different donors.

In addition, hMSCs were immunostained for specific aromatase cytochrome P450. Immunostaining occurred in the cytoplasm with a similar distribution in cells cultured under basal OS or AD conditions. No difference in staining intensity of cells was appreciated between basal or differentiation conditions (data not shown).

Results in Table 1 demonstrate that oestradiol and oestrone are found in the incubation medium of hMSCs cultured under OS or AD conditions. Under OS conditions, the amount of oestrogens formed from androstenedione was clearly substrate concentration dependent. As expected, testosterone as substrate (100 nM) allowed the formation of oestradiol as much as that originated from androstenedione (30 nM), but oestrone production was low. No significant difference was appreciated between oestrogens produced under OS and AD conditions at this time point of cell differentiation.

Effect of leptin and 1,25(OH)₂D₃ on aromatase activity

Neither leptin nor 1,25(OH)₂D₃ affected aromatase activity of hMSCs under basal conditions; however, the addition of these compounds to the culture media during OS cell differentiation was associated with increased aromatase activity. At 24 h of OS differentiation the effect was dose dependent (Fig. 2A and B).

After 24 h of OS differentiation 200 nM leptin increased aromatase activity 1.6-fold, compared with the activity in the absence of leptin. The addition of leptin to hMSCs during AD differentiation did not modify aromatase activity compared with the activity in the absence of leptin (Fig. 3A). On the other hand, 1,25(OH)₂D₃ (10 nM) increased 1.8-fold hMSCs aromatase activity after 24 h of OS differentiation, compared with cells cultured in the absence of the seco-steroid hormone; this effect was not observed in hMSCs under AD conditions (Fig. 3B).

Protein aromatase levels were analysed by western blot; no difference in the level of protein immunostained for aromatase cytochrome P450 under basal, OS, AD, OS plus leptin or OS plus 1,25(OH)₂D₃ conditions were detected (Fig. 4A and B). Western blots of proteins from hMSCs demonstrated similar protein migration as the pattern obtained with MCF-7 cells proteins (Fig. 4A).

Figure 5 shows that at 24 h of OS differentiation, increased aromatase activity is dependent, in part, on MAPK-dependent activities, since the inhibitor 2'-amino-3'-methoxyflavone (PD 98059) (25 µM) abolished 60–80% of the aromatase activity. The inhibitor also decreased the stimulatory effect of leptin.

Oestrogens produced by hMSCs affect their differentiation capacity

Table 2 summarizes the extent of AD differentiation of hMSCs under different oestrogenic conditions after 14 days of treatment. Under plain AD condition, the mean number of adipocytes detected by flow cytometry was 1114 ± 384 . This number was not affected by the presence of 50 nM oestradiol in the AD medium. The addition of 0.1 µM androstenedione or 0.5 µM testosterone, substrates for aromatase, during AD differentiation produced a significant inhibition (60%) in the number of adipocytes differentiated from hMSCs. Moreover, this inhibitory effect was blocked by either 0.5 µM 4-androsten-4-ol-3,17-dione or 0.1 µM ICI-182,780, specific inhibitors of the enzyme aromatase (Ar-Inh) and of the oestrogen receptors respectively. On the other hand, the presence of 0.1 µM ICI-182,780 did not have a significant effect on AD differentiation.

A similar experiment was performed to evaluate the effect of the oestrogenic substrates androstenedione and testosterone on OS differentiation of hMSCs. Table 3 shows that under these substrate conditions, OS differentiation is enhanced as evidenced by the alkaline phosphatase activity measurements. This positive effect was decreased by the presence of the inhibitors Ar-Inh and ICI-182,780. Similarly with the AD differentiation, no effect on OS differentiation was observed by the addition of 50 nM oestradiol to the culture medium.

Control studies showed that the presence of inhibitors Ar-Inh and ICI-182,780 did not have a significant effect on OS and AD differentiation (data not shown).

Table 1 Oestrogens produced by hMSCs cultured under OS and AD conditions. Results are the mean \pm s.d.

Culture condition	Substrate (concentration µM)	Oestradiol (pmol/mg protein)	Oestrone (pmol/mg protein)
OS medium	Androstenedione (0.03)	4.90 \pm 0.27	7.96 \pm 1.33
OS medium	Androstenedione (0.1)	11.9 \pm 1.8*	17.9 \pm 2.4*
OS medium	Testosterone (0.1)	5.95 \pm 0.78 [†]	0.93 \pm 0.29 [†]
AD medium	Androstenedione (0.1)	8.8 \pm 1.3	15.1 \pm 4.48

1–1.2 \times 10⁵ hMSCs were cultured under osteogenic or adipogenic conditions during 24 h. Oestradiol and oestrone produced were measured using ³H-labelled substrates as described in Materials and Methods. Experiments were performed in duplicate from two different samples. **P* < 0.05 compared with corresponding value of androstenedione (30 nM). [†]*P* < 0.05 compared with corresponding value of androstenedione (100 nM).




Figure 2 Aromatase activity of hMSCs in response to leptin and 1,25(OH)₂D₃ in different concentrations. hMSCs were cultured in osteogenic conditions for 24 h, supplemented with specified leptin (A) or 1,25(OH)₂D₃ (B) concentrations. The aromatase activity was measured by the tritiated water release assay, as described in legend of Fig. 1. Experiments were performed in triplicate from four different samples. Results are the mean \pm s.d.

Discussion

Differentiation of hMSCs towards osteoblasts or adipocytes requires the sequential expression of genes associated with each of the resulting cell phenotypes (Ren *et al.* 2002, Kobayashi & Kronenberg 2005). The regulation of gene expression or activation during these processes is modulated by several endocrine, paracrine, autocrine and intracrine factors, which determine the phenotype to which the progenitor cells differentiate. Among these factors, oestrogens

play an important role. Moreover, it appears that local aromatization of C19 precursors in bone may contribute significantly to skeletal homeostasis (Labrie *et al.* 1998, Simpson 2000, Simpson & Davis 2001), suggesting that the regulation of aromatase activity by factors present in the local environment play a decisive role in adjusting the levels of bioavailable oestrogenic hormone.

In this study, we demonstrate in hMSCs that OS differentiation promoted an early peak of aromatase CYP 19 activity, followed by a marked decrease of enzyme activity




Figure 3 Effect of leptin and 1,25(OH)₂D₃ on aromatase activity of hMSCs. Cells were cultured for 24 h in osteogenic or adipogenic conditions, supplemented with 200 nM leptin (A) or 10 nM 1,25(OH)₂D₃ (B) and the aromatase activity was measured by the tritiated water release assay as specified in legend of Fig. 1. White bars, no addition; grey bars, leptin; black bars, 1,25(OH)₂D₃. Experiments were performed in triplicate from four different samples. Results are the mean \pm s.d. * P <0.05 compared with osteogenesis values in the presence of hormones. [†] P <0.05 compared with values in the absence of hormone.




Figure 4 Western-blot analysis of aromatase in hMSCs. (A) hMSCs were cultured under basal (lane 1), OS (lane 2), AD (lane 3), OS + leptin (lane 4), and OS + $1,25(\text{OH})_2\text{D}_3$ (lane 5) conditions, positive control (lane 6, MCF-7 cells). Western-blot analysis was performed using an anti CYP19 (P450 arom). (B) Densitometric analysis performed as described in Materials and Methods and normalized with β -actin. Representative data from three independent experiments are shown.

after 48 h; thereafter aromatase activity remained above basal values for up to 7 days. As far as we know, this is the first observation of aromatase activity during commitment and early stages of OS differentiation. A former study, carried out in a human foetal osteoblast cell line (SV-HFO) showed by




Figure 5 Effect of PD98059 on aromatase activity of hMSCs. Cells were cultured for 24 h in osteogenic conditions, supplemented or not with 200 nM leptin, and in the presence or the absence of 25 μM PD98059. The aromatase activity was measured by the tritiated water release assay as specified in legend of Fig. 1. Experiments were performed in triplicate from four different samples. Results are the mean \pm s.d. * $P < 0.05$ compared with values without PD98059.

semi-quantitative analysis that aromatase mRNA expression did not change, while aromatase activity decreased during late stages of OS differentiation (measurements were done from 6 to 21 days of differentiation; Janssen *et al.* 1999). In addition, Heim *et al.* (2004), observed no variation in aromatase transcript levels in the first 7 days of differentiation of hMSCs, however, they did not study aromatase activity. Increased levels of transcripts were observed later during OS differentiation, supporting previous reports of aromatase expression and activity in mature osteoblasts (Sasano *et al.* 1997, Shozu & Simpson 1998). Considering both the results of Heim *et al.* (2004) and ours, it may be concluded that post-transcriptional mechanisms could play an important role in regulating aromatase activity during early OS differentiation.

Although the specific factor responsible for sharply increasing aromatase activity during early OS differentiation is not known, it could result from the dexamethasone present in the OS medium, since a similar increase of aromatase activity has been observed in primary cultured human osteoblasts after 12 h of dexamethasone treatment (Tanaka *et al.* 1996). Moreover, it has been concluded that glucocorticoids regulate transcription of the aromatase gene in bone, adipose and ovarian cells. (Simpson *et al.* 1981, Purohit *et al.* 1992, Shimodaira *et al.* 1996, Tanaka *et al.* 1996, Shozu & Simpson 1998, Enjuanes *et al.* 2003). However, post-transcriptional regulation of the enzyme after dexamethasone has also been suggested by the effects of inhibitors of protein synthesis or MAPK pathway phosphorylation (Tanaka *et al.* 1996, Shozu *et al.* 2001). Thus, the diminished aromatase activity observed after 48 h might result from modulatory changes during osteogenesis.

During early AD differentiation, aromatase activity levels were lower than during OS differentiation, but the enzyme activity remained significantly increased upto 72 h of treatment, suggesting that in addition to the initial effect of dexamethasone, interplay of signals developed during differentiation of MSCs may contribute to define the ensuing level of aromatase activity. Thus, a different pattern of enzyme activity developed after AD treatment of MSCs, despite the fact that the AD medium also contains dexamethasone. There are no previous observations on aromatase activity during early AD differentiation of MSCs, although increased CYP19 transcript levels were observed at this time, compared with control, that decreased considerably by the end of AD maturation (Heim *et al.* 2004). Thus, taking into consideration Heim's report (Heim *et al.* 2004), our observations could indicate that during early adipogenesis, aromatase cell activity could result mainly from transcriptional regulation.

Kinetic parameters of hMSCs aromatase enzyme agree with the values found in human osteoblasts (Tanaka *et al.* 1996), both in the range of apparent maximal velocity and in apparent K_m for androstenedione. Thus, in bone cells, the kinetic properties of both hMSCs and mature osteoblasts (Tanaka *et al.* 1996) indicate a high capacity for conversion of circulating androgens. Direct measurements of oestrogens produced by hMSCs support this conclusion and demonstrate

Table 2 Adipogenic differentiation of hMSCs under different culture conditions. Results are expressed as the relative adipocyte number compared with value obtained in plain adipogenic medium (AD). Experiments were performed in triplicate from four different samples

Culture condition	Adipocytes (relative number)
AD medium	1.00
+50 nM Oestradiol	1.03 ± 0.19
+0.1 µM Androstenedione	0.62 ± 0.36*
+0.5 µM Testosterone	0.53 ± 0.10*
+0.1 µM Androstenedione +0.5 µM Ar-Inh	1.20 ± 0.24
+0.1 µM Androstenedione +0.1 µM ICI-182,780	1.08 ± 0.19
+0.5 µM Testosterone + +0.5 µM Ar-Inh	0.76 ± 0.04*
+0.5 µM Testosterone +0.1 µM ICI-182,780	0.99 ± 0.10

hMSCs were cultured under adipogenic conditions during 14 days. The adipocyte number was determined by flow cytometric analysis. Aromatase inhibitor (Ar-Inh): 4-androsten-4-ol-3,17-dione. Results are the mean ± s.d. * $P < 0.05$ compared with the plain adipogenic condition.

that significant concentration of oestradiol ($1-3 \pm 0.56$ nM) and oestrone ($2-5 \pm 0.8$ nM) is attained, depending on the substrate availability. Our results indicate that during the early stages of differentiation, hMSCs actively biosynthesize oestrogens as described for differentiated osteoblasts. Moreover, the different oestrogen-generating capabilities found among differentiating MSCs suggest that accurate oestrogens signalling may be important for appropriate early bone marrow cell differentiation.

Leptin and $1,25(\text{OH})_2\text{D}_3$, significantly increased aromatase activity only through early OS differentiation. Neither leptin nor $1,25(\text{OH})_2\text{D}_3$ affected aromatase activity of hMSCs under basal conditions, nor during AD differentiation. The dose-response curves support the *in vivo* action of these agents as modulators of aromatase activity, since effective concentrations used in this study are in the physiological range of circulating leptin and $1,25(\text{OH})_2\text{D}_3$.

There are no previous reports on the effect of leptin on bone aromatase cell activity, although there are studies on leptin effects on aromatase gene expression and/or cell activity in luteinized granulosa cells (Kitawaki *et al.* 1999), adipose

stromal cells (Magoffin *et al.* 1999) and MCF-7 cell line (Catalano *et al.* 2003). Leptin activity on immortalized stroma cells from human bone marrow increased their differentiation to osteoblasts, while it inhibited their differentiation to adipocytes, suggesting a role for leptin in bone metabolism (Thomas *et al.* 1999). Previously, we have demonstrated the presence of membrane leptin receptors through early hMSCs differentiation, as well as its direct protective effect on their OS differentiation process (Hess *et al.* 2005). Therefore, from our results it may be suggested that part of the protective influence of leptin on bone tissue may result from its effect on aromatase activity during early differentiation of hMSCs.

The induction of aromatase activity we observed in hMSCs in response to $1,25(\text{OH})_2\text{D}_3$ is consistent with the previous studies in other bone cell types showing that the effect of the hormone on aromatase is dependent on previous or concomitant glucocorticoid treatment (Tanaka *et al.* 1996, Enjuanes *et al.* 2003, Yanase *et al.* 2003). Our results show that the stimulatory effect of $1,25(\text{OH})_2\text{D}_3$ is restricted to early osteogenic hMSCs differentiation and is characterized by a massive increase of aromatase activity that decreases after 48 h

Table 3 Osteogenic differentiation of hMSCs under different culture conditions. Results are the mean ± s.d.

Culture conditions	Alkaline phosphatase activity (µg <i>p</i> -nitrophenol/well)
Basal	0.53 ± 0.18
OS medium	4.84 ± 0.55*
+50 nM Oestradiol	5.24 ± 0.61
+0.1 µM Androstenedione	8.03 ± 0.46 [†]
+0.1 µM Androstenedione +0.5 µM Ar-Inh	6.54 ± 1.85 [‡]
+0.1 µM Androstenedione +0.1 µM ICI-182,780	6.48 ± 0.15 ^{†,‡}
+0.1 µM Testosterone	6.21 ± 1.03 [†]
+0.1 µM Testosterone +0.5 µM Ar-Inh	5.43 ± 1.64
+0.1 µM Testosterone +0.1 µM ICI-182,780	5.32 ± 0.23

hMSCs were cultured under osteogenic conditions during 14 days. Alkaline phosphatase activity was measured at 7 days of culture. Experiments were performed in duplicate from three different samples. Aromatase inhibitor (Ar-Inh): 4-androsten-4-ol-3,17-dione. * $P < 0.05$ compared with basal medium. [†] $P < 0.05$ compared with the plain osteogenic condition. [‡] $P < 0.05$ compared with osteogenic medium + androstenedione.

of differentiation. This effect of $1,25(\text{OH})_2\text{D}_3$ on aromatase cell activity contrasts with the rather modest increase of aromatase mRNA (Enjuanes *et al.* 2003), supporting a post-transcriptional modulation of aromatase by $1,25(\text{OH})_2\text{D}_3$ (Tanaka *et al.* 1996, Yanase *et al.* 2003).

Interestingly, western-blot analysis of hMSCs under basal, OS, OS plus leptin, OS plus $1,25(\text{OH})_2\text{D}_3$ and AD differentiation showed no difference in the expression of protein level associated with immunostained aromatase. Therefore, the amount of protein appeared unrelated to the increase observed in enzyme activity suggesting that aromatase activity increased without changes in the protein concentration.

We observed that much of the increased aromatase activity after 24 h of OS stimulation was abolished by PD 98059, a selective inhibitor of MEK-1/2, supporting the proposition that aromatase activity might be acutely regulated by phosphorylation–dephosphorylation reactions during hMSCs differentiation. This type of post-transcriptional modulation of aromatase activity has been deduced from the inhibitory effects of the selective MEK-1/2 inhibitor on osteoblast-like cells, THP-1 (human peripheral blood) and JEG-3 (human choriocarcinoma) cell lines (Shozu *et al.* 2001). Therefore, specific OS and AD signals could trigger rapid and characteristic changes in aromatase cell activity, avoiding significant variation in aromatase protein and mRNA levels. This may explain the discordance between aromatase immunostained protein and enzyme activity.

We also evaluated whether compounds that modify either oestrogen synthesis or response affected OS and AD capacity of hMSCs. When substrate conditions that favoured aromatase activity existed (addition of androstenedione or testosterone), adipogenesis was significantly inhibited suggesting that high local oestrogen production restrains the process. Further, the effect was abolished by specific inhibitors for either aromatase or the oestrogen receptors, corroborating that increased oestrogen action is required to hold down AD.

During OS differentiation, we noticed that androstenedione added to medium had a positive effect on alkaline phosphatase activity, an early marker of OS differentiation. This favourable effect was diminished by the presence of specific inhibitors of aromatase or oestrogen action.

Osteoblasts (Arts *et al.* 1997, Heim *et al.* 2004) and adipocytes (Mizutani *et al.* 1994, Crandall *et al.* 1998) express oestrogen receptors showing different receptor expression patterns along hMSCs OS and AD differentiation (Heim *et al.* 2004). There is evidence supporting the role of oestrogen as a negative regulator for adipogenesis. *In vivo*, oestrogen receptors knockout mice (Heine *et al.* 2000) and aromatase-deficient mice (Jones *et al.* 2000) have been reported to manifest increased adiposity, although bone marrow adipocytes were not investigated in these reports. Two *in vitro* studies in mouse bone marrow stromal ST2 cell lines (Okazaki *et al.* 2002) and hMSCs (Heim *et al.* 2004) reported reciprocal regulation by oestrogen of osteoblastic and adipocytic differentiation from a common progenitor cell population. Our results confirm these observations and

further underline the effect of oestradiol synthesized from C19 substrates by aromatase activity. In our experiments, these substrates repressed adipogenesis and favoured osteogenesis, while no effect was detected when pharmacological concentration of oestradiol was added to the medium. The lack of a direct effect of oestradiol could result from the presence of both 10% FCS and phenol red in the culture medium. These experimental conditions could provide suboptimal oestrogenic conditions even for cells in basal conditions. We could neither diminish FCS concentration nor use carbon-dextrane-treated serum to reduce oestrogen content in medium, since both treatments diminished viability of hMSCs. Thus, these observations suggest that locally produced oestradiol exerts great impact on hMSCs differentiation. Furthermore, these results support the hypothesis of a threshold oestradiol level for normal skeletal remodelling (Riggs *et al.* 2002, Gennari *et al.* 2004), which could be attained by the activity of endogenous aromatase on appropriate C19 precursors.

Overall, these observations point to critical requirements for the regulation of aromatase activity during the commitment and differentiation of bone hMSCs, suggesting that local production of oestrogen may hold appropriate cell differentiation, its production subjected to subtle adjustments depending on specific local signals. It could be inferred that high and early increases of oestrogen concentration in hMSCs affect their commitment by either restraining AD or facilitating OS differentiation, or both. During aging and some bone disorders, both substrate availability and aromatase regulation might affect the differentiation processes.

Acknowledgements

The authors thank to Dr O Brunser for critical review of the manuscript and valuable comments, Dr L Valladares for TLC analysis and to Dr Marco A Méndez for statistical advice. Also, the authors are grateful to Mrs V Simon for performing the flow cytometric analysis.

Funding

This work was supported by FONDECYT grant number 1050930 and Proyecto Mayor 03/07-2 DI. The authors declare that there is no conflict of interest.

References

- Amin S, Zhang Y, Sawin CT, Evans SR, Hannan MT, Kiel DP, Wilson PW & Felson DT 2000 Association of hypogonadism and estradiol levels with bone mineral density in elderly men from the Framingham study. *Annals of Internal Medicine* **133** 951–963.
- Arts J, Kuiper GG, Janssen JM, Gustafsson JA, Lowik CW, Pols HA & van Leeuwen JP 1997 Differential expression of estrogen receptor α and β mRNA during differentiation of human osteoblast SV-HFO cells. *Endocrinology* **138** 5067–5070.

- Banerjee C, McCabe LR, Choi JY, Hiebert SW, Stein JL, Stein GS & Lian JB 1997 Runt homology domain proteins in osteoblast differentiation: AML3/CBFA1 is a major component of bone-specific complex. *Journal of Cellular Biochemistry* **66** 1–8.
- Bianco P & Robey PG 1999 Diseases of bone and stromal cell lineage. *Journal of Bone and Mineral Research* **14** 336–341.
- Bouillon R, Okamura WH & Norman AW 1995 Structure–function relationships in the vitamin D endocrine system. *Endocrine Reviews* **16** 200–257.
- Bruder SP, Jaiswal N & Haynesworth SE 1997 Growth kinetics, self-renewal, and the osteogenic potential of purified human mesenchymal stem cells during extensive subcultivation and following cryopreservation. *Journal of Cellular Biochemistry* **64** 278–294.
- Caplan AI 1991 Mesenchymal stem cells. *Journal of Orthopaedic Research* **9** 641–650.
- Catalano S, Marsico S, Giordano C, Mauro L, Rizza P, Panno ML & Ando S 2003 Leptin enhances, via AP-1, expression of aromatase in the MCF-7 cell line. *Journal of Biological Chemistry* **278** 28668–28676.
- Compston J 2002 Editorial: Local biosynthesis of sex steroids in bone. *Journal of Clinical Endocrinology and Metabolism* **87** 5398–5400.
- Crandall DL, Busler DE, Novak TJ, Weber RV & Kral JG 1998 Identification of estrogen receptor RNA in human breast and abdominal subcutaneous adipose tissue. *Biochemical and Biophysical Research Communications* **248** 523–526.
- Dennis JE, Merriam A, Awadallah A, Yoo JU, Johnstone B & Caplan AI 1999 A quadripotential mesenchymal progenitor cell isolated from the marrow of an adult mouse. *Journal of Bone and Mineral Research* **14** 700–709.
- Ducy P, Zhang R, Geoffroy V, Riddall AL & Karsenty G 1997 *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. *Cell* **89** 747–754.
- Enjuanes A, Garcia-Giralto N, Supervia A, Nogués X, Mellibovsky L, Carbonell J, Grinberg D, Balcells S & Diez-Perez A 2003 Regulation of *Cyp19* gene expression in primary human osteoblasts: effects of vitamin D and other treatments. *European Journal of Endocrinology* **148** 519–526.
- Gambacciani M, Ciapponi M, Cappagli B, Piaggini L, De Simone L, Orlandi R & Genazzani AR 1997 Body weight, body fat distribution, and hormonal replacement therapy in early postmenopausal women. *Journal of Clinical Endocrinology and Metabolism* **82** 414–417.
- Gennari L, Nuti R & Bilezikian JP 2004 Aromatase activity and bone homeostasis in men. *Journal of Clinical Endocrinology and Metabolism* **89** 5898–5907.
- Gimble JM, Robinson CE, Wu X & Kelly KA 1996 The function of adipocytes in the bone marrow stroma: an update. *Bone* **19** 421–428.
- Heim M, Frank O, Kampmann G, Sochocky N, Pennimpede T, Fuchs P, Hunziker W, Weber P, Martin I & Bendik I 2004 The phytoestrogen genistein enhances osteogenesis and represses adipogenic differentiation of human primary bone marrow stromal cells. *Endocrinology* **145** 848–859.
- Heine PA, Taylor JA, Iwamoto GA, Lubahn DB & Cooke PS 2000 Increased adipose tissue in male and female estrogen receptor- α knockout mice. *PNAS* **97** 12729–12734.
- Hess R, Pino AM, Ríos S, Fernández M & Rodríguez JP 2005 High affinity leptin receptors are present in human mesenchymal stem cells (MSCs) derived from control and osteoporotic donors. *Journal of Cellular Biochemistry* **94** 50–57.
- Hu Y, Chan E, Wang SX & Li B 2003 Activation of p38 mitogen-activated protein kinase is required for osteoblast differentiation. *Endocrinology* **144** 2068–2074.
- Ishida Y, Killinger DW, Khalil MW, Yang K, Strutt B & Heersche JN 2002 Expression of steroid-converting enzymes in osteoblasts derived from rat vertebrae. *Osteoporosis International* **13** 235–240.
- Issa S, Schnabel D, Feix M, Wolf L, Schaefer HE, Russell DW & Schweikert HU 2002 Human osteoblast-like cells express predominantly steroid 5 α -reductase type 1. *Journal of Clinical Endocrinology and Metabolism* **87** 5401–5407.
- Jaiswal N, Haynesworth SE, Caplan AI & Bruder SP 1997 Osteogenic differentiation of purified, culture-expanded human mesenchymal stem cells *in vitro*. *Journal of Cellular Biochemistry* **64** 295–312.
- Jakob E, Siggelkow H, Homann D, Körle J, Adamski J & Schütze N 1997 Local estradiol metabolism in osteoblast- and osteoblast-like-cells. *Journal of Steroid Biochemistry and Molecular Biology* **61** 167–174.
- Janssen JMMF, Bland R, Hewison M, Coughtrie MWH, Sharp S, Arts J, Pols H & van Leeuwen JP 1999 Estradiol formation by human osteoblasts via multiple pathways: relation with osteoblast function. *Journal of Cellular Biochemistry* **75** 528–537.
- Jones ME, Thorburn AW, Britt KN, Hewitt KN, Wreford NG, Proietto J, Oz OK, Leury BJ, Robertson KM, Yao S *et al.* 2000 Aromatase-deficient (ArKO) mice have a phenotype of increased adiposity. *PNAS* **97** 12735–12740.
- Justesen J, Stenderup K, Ebbesen EN, Mosekilde L, Steiniche T & Kassem M 2001 Adipocyte tissue volume in bone marrow is increased with aging and in patients with osteoporosis. *Bogerontology* **2** 165–171.
- Khosla S, Melton LJ III, Atkinson EJ & O’Fallon WM 2001 Relationship of sex steroids to longitudinal changes in bone mineral density and bone resorption in young versus elderly men: effects of estrogen on peak bone mass and on age-related bone loss. *Journal of Clinical Endocrinology and Metabolism* **86** 3555–3561.
- Kitawaki J, Kusuki I, Koshiba H, Tsukamoto K & Honjio H 1999 Leptin directly stimulates aromatase activity in human luteinized granulosa cells. *Molecular Human Reproduction* **8** 708–713.
- Kobayashi T & Kronenberg H 2005 Minireview: transcriptional regulation in development of bone. *Endocrinology* **146** 1012–1017.
- Komm BS, Terpening CM, Benz DJ, Graeme KA, Gallegos A, Kore M, Greene GL, O’Malley BW & Haussler MR 1988 Estrogen binding, receptor mRNA, and biologic response in osteoblast-like osteosarcoma cells. *Science* **241** 81–84.
- Labrie F, Belanger A, Cusan L & Candas B 1997 Physiological changes in dehydroepiandrosterone are not reflected by serum levels of active androgens and oestrogens but of their metabolites: intracrinology. *Journal of Clinical Endocrinology and Metabolism* **82** 2403–2409.
- Labrie F, Belanger A, Luu-The V, Labrie C, Simond J, Cusan L, Gómez JL & Candas B 1998 DHEA and the intracrine formation of androgens and estrogens in peripheral target tissues: its role during aging. *Steroids* **63** 322–328.
- Lea CK, Ebrahim H, Tennant S & Flanagan AM 1997 Aromatase cytochrome P450 transcripts are detected in fractured human bone but not in normal skeletal tissue. *Bone* **21** 433–440.
- Lecka-Czernik B, Gubrij I, Moerman EJ, Kajenova O, Lipschitz DA, Manolagas SC & Jilka RL 1999 Inhibition of *Osf2/Cbfa1* expression and terminal osteoblast differentiation by PPAR γ 2. *Journal of Cellular Biochemistry* **74** 357–371.
- Lephart ED & Simpson ER 1991 Assay of aromatase activity. *Methods in Enzymology* **206** 477–483.
- Magoffin D, Weitsman SR, Aagarwal SK & Jakimiuk AJ 1999 Leptin regulation of aromatase activity in adipose stromal cells from regularly cycling women. *Ginekologia Polska* **70** 1–7.
- Mizutani T, Nishikawa Y, Adachi H, Enomoto T, Ikegami H, Kurachi H, Nomura T & Miyake A 1994 Identification of estrogen receptor in human adipose tissue and adipocytes. *Journal of Clinical Endocrinology and Metabolism* **78** 950–954.
- Morishima A, Grumbach MM, Simpson ER, Fisher C & Qin K 1995 Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *Journal of Clinical Endocrinology and Metabolism* **86** 3689–3698.
- Nuttall ME, Patton AJ, Olivera DL, Nadeau DP & Gowen M 1998 Human trabecular bone cells are able to express both osteoblastic and adipocytic phenotype: implications for osteopenic disorders. *Journal of Bone and Mineral Research* **13** 371–382.
- Okazaki R, Inoue D, Shibata M, Saika M, Kido S, Ooka H, Tomiyama H, Sakamoto Y & Matsumoto T 2002 Estrogen promotes early osteoblast differentiation and inhibits adipocyte differentiation in mouse bone marrow stromal cell lines that express estrogen receptor (ER) α or β . *Endocrinology* **143** 2349–2356.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S & Marshak DR 1999 Multilineage potential of adult human mesenchymal stem cells. *Science* **284** 143–147.
- Purohit A, Flanagan AM & Reed MJ 1992 Estrogen synthesis by osteoblast cell lines. *Endocrinology* **131** 2027–2029.
- Ren D, Collingwood TN, Rebar EJ, Wolfe AP & Camp HS 2002 PPAR γ knockdown by engineered transfection factors: exogenous PPAR γ 2 but not PPAR γ 1 reactivates adipogenesis. *Genes and Development* **16** 27–32.

- Reseland JE, Syversen U, Bakke I, Qvigstad G, Eide LG, Hjertner O, Gordeladze JO & Drevon CA 2001 Leptin is expressed in and secreted from primary cultures of human osteoblasts and promotes bone mineralization. *Journal of Bone and Mineral Research* **16** 1426–1433.
- Riggs BI, Khosla S & Melton LJ III 2002 Sex steroids and the construction and conservation of the adult skeleton. *Endocrine Reviews* **23** 279–302.
- Rodríguez JP, Garat S, Gajardo H, Pino AM & Seitz G 1999 Abnormal osteogenesis in osteoporotic patients is reflected by altered mesenchymal stem cells dynamics. *Journal of Cellular Biochemistry* **75** 414–423.
- Rodríguez JP, Ríos S & González M 2002 Modulation of the proliferation and differentiation of human mesenchymal stem cells by copper. *Journal of Cellular Biochemistry* **85** 92–100.
- Sasano H, Uzuki M, Sawai T, Nagura H, Matsunaga G, Kashimoto O & Harada N 1997 Aromatase in human bone tissue. *Journal of Bone and Mineral Research* **12** 1416–1423.
- Schweikert HU, Rulf W, Niederle N, Schafer HE, Keck E & Krüick F 1980 Testosterone metabolism in human bone. *Acta Endocrinologica* **95** 258–264.
- Schweikert HU, Wolf L & Romalo G 1995 Oestrogen formation from androstenedione in human bone. *Clinical Endocrinology* **43** 37–42.
- Shimodaira K, Fujikawa H, Okura F, Shimizu Y, Saito H & Yanaihara T 1996 Osteoblast cells (MG-63 and HOS) have aromatase and 5 alpha reductase activities. *Biochemistry and Molecular Biology International* **39** 109–116.
- Shozu M & Simpson ER 1998 Aromatase expression of human osteoblast-like cells. *Molecular and Cellular Endocrinology* **139** 117–129.
- Shozu M, Zhao Y & Simpson ER 1997 Estrogen biosynthesis in THP-1 cells is regulated by promoter switching of the aromatase (CYP19) gene. *Endocrinology* **138** 5125–5135.
- Shozu M, Zhao Y & Simpson ER 2000 TGF-beta stimulates expression of the aromatase (CYP19) gene in human osteoblast-like cells and THP-1 cells. *Molecular and Cellular Endocrinology* **160** 123–133.
- Shozu M, Sumitami H, Murakami K, Segawa T, Yang H-J & Inoue M 2001 Regulation of aromatase activity in bone derived cells: possible role of mitogen-activated protein kinase. *Journal of Steroid Biochemistry and Molecular Biology* **79** 61–65.
- Simpson ER 2000 Role of aromatase in sex steroid action. *Journal of Molecular Endocrinology* **25** 149–156.
- Simpson EV & Davis SR 2001 Minireview: Aromatase and the regulation of estrogen biosynthesis: some new perspectives. *Endocrinology* **142** 4589–4594.
- Simpson ER, Ackerman G, Smith M & Mendelson C 1981 Estrogen formation in stromal cells of adipose tissue of women: induction by glucocorticoids. *PNAS* **78** 5690–5694.
- Smith EC, Boyd J, Franck GR, Takahashi H, Cohen RM, Specker B, Williams TC, Lubahn DB & Korach KS 1994 Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *New England Journal of Medicine* **331** 1056–1061.
- Takahashi Y, Okimura Y, Mizuno I, Iida K, Takahashi T, Kaji H, Abe H & Chihara K 1997 Leptin induces mitogen-activated protein kinase-dependent proliferation of C3H10T1/2 cells. *Journal of Biological Chemistry* **272** 12897–12900.
- Tanaka S, Haji M, Nishi Y, Yanase T, Takayanagi R & Nawata H 1993 Aromatase activity in human osteoblast-like osteosarcoma cell. *Calcified Tissue International* **52** 107–109.
- Tanaka S, Haji M, Takayanagi R, Sugioka Y & Nawata H 1996 1,25-Dihydroxy vitamin D₃ enhances the enzymatic activity and expression of the messenger ribonucleic acid for aromatase cytochrome P450 synergistically with dexamethasone depending on the vitamin D receptor level in cultured human osteoblasts. *Endocrinology* **137** 1860–1869.
- Thomas T, Gori F, Khosla S, Jensen MD, Burguera B & Riggs BL 1999 Leptin acts on human marrow stromal cells to enhance differentiation to osteoblasts and to inhibit differentiation to adipocytes. *Endocrinology* **140** 1630–1638.
- Yanase T, Suzuki S, Goto K, Nomura M, Okabe T, Takayanagi R & Nawata H 2003 Aromatase in bone: roles of vitamin D3 and androgens. *Journal of Steroid Biochemistry and Molecular Biology* **86** 393–397.

Received in final form 7 September 2006

Accepted 27 September 2006

Made available online as an Accepted Preprint

12 October 2006