

Artificial Intelligence Techniques for Automated Diagnosis of Neurological Disorders

U. Raghavendra^a U. Rajendra Acharya^{b–d} Hojjat Adeli^{e, f}

^aDepartment of Instrumentation and Control Engineering, Manipal Institute of Technology, Manipal Academy of Higher Education, Manipal, India; ^bDepartment of Electronics and Computer Engineering, Ngee Ann Polytechnic, Clementi, Singapore; ^cDepartment of Biomedical Engineering, School of Science and Technology, SUSS University, Clementi, Singapore; ^dInternational Research Organization for Advanced Science and Technology (IROAST) Kumamoto University, Kumamoto, Japan; ^eDepartment of Neuroscience, The Ohio State University, Columbus, OH, USA; ^fDepartments of Biomedical Informatics and Neuroscience, The Ohio State University, Columbus, OH, USA

Keywords

Neurological disorder · Computer-aided diagnosis · Machine learning · Classification algorithm

Abstract

Background: Authors have been advocating the research ideology that a computer-aided diagnosis (CAD) system trained using lots of patient data and physiological signals and images based on adroit integration of advanced signal processing and artificial intelligence (AI)/machine learning techniques in an automated fashion can assist neurologists, neurosurgeons, radiologists, and other medical providers to make better clinical decisions. **Summary:** This paper presents a state-of-the-art review of research on automated diagnosis of 5 neurological disorders in the past 2 decades using AI techniques: epilepsy, Parkinson's disease, Alzheimer's disease, multiple sclerosis, and ischemic brain stroke using physiological signals and images. Recent research articles on different feature extraction methods, dimensionality reduction techniques, feature selection, and classification techniques are reviewed. **Key Message:** CAD systems using AI

and advanced signal processing techniques can assist clinicians in analyzing and interpreting physiological signals and images more effectively.

© 2019 S. Karger AG, Basel

Introduction

Neurological disorders are the diseases connected with peripheral and central nervous systems. The common symptoms include muscle weakness, paralysis, seizures, pain, poor coordination, and loss of consciousness [1]. There are >600 diseases related to the nervous system such as brain tumor, Parkinson's disease (PD), Alzheimer's disease (AD), multiple sclerosis (MS), epilepsy, dementia, headache disorders, neuroinfections, stroke, and traumatic brain injuries among others. Various viral infections (i.e., HIV, Zika, West Nile Virus, Enteroviruses), bacterial infections (such as *Neisseria meningitides* and *Mycobacterium tuberculosis*), fungal-related infections (such as *Aspergillus* and *Cryptococcus*), and parasitic infections (such as Chagas and malaria) can affect the entire

nervous system [1–6]. The aforementioned neurological symptoms possibly occur due to immune response or infection itself. Hundreds of millions of people worldwide are affected by neurological disorders [7–10]. More than 6 million people die because of stroke each year; majority of these deaths take place in low- and middle-income countries [1, 11, 12]. It is reported that around 50 million people will have epilepsy [1], and 47.5 million people will suffer from dementia [1, 13–17].

The abnormal or the anomalous neurological conditions are commonly identified by a neuropathological examination. Anomalous neurological conditions are found in majority of the population and are not always associated with a neurological disorder [18].

Dementia is usually progressive in nature. The dementia syndromes disturb multiple cortical functions, that is, memory, orientation, thinking, calculation, language, comprehension, judgment, and learning capacity. AD is found to be the most common cause of dementia, which is characterized by neurofibrillary and cortical amyloids accounting for 3 quarters of the cases [16, 19, 20]. Dementia affects mainly older people, above 65 years, but also 2% of the people under 65 years old. The prevalence of dementia doubles with age every 5 years. A genetic polymorphism increases the risk for 25% of entire population [21, 22].

Epilepsy, a chronic neurological disorder, is defined as, “*disorder of brain characterized by enduring predisposition to generate the epileptic seizures.*” [23–25]. The epilepsy definition requires at least occurrence of one epileptic seizure [26, 27]. It affects both male and female sexes and people of all ages. The diagnosis for epileptic seizures is performed by first determining the event of epilepsy and later differentiating between the conditions called provoked or chronic epileptic seizures [28, 29]. The overall incidence of epilepsy is found to be 23–190 per 100,000 of population [30]. The prevalence is lower in early ages and gradually increases with aging [31–34]. Since the pioneering work of Adeli et al. [35], wavelet transform (WT) has been used extensively for electroencephalogram (EEG) analysis, seizure detection, and epilepsy diagnosis [36, 37]. Kugiumtzis et al. [27] investigate the dynamics of epileptiform discharges induced by transcranial magnetic stimulation in epilepsy. Yuan et al. [38] present a method for epileptic seizure prediction using diffusion distance and Bayesian linear discriminate analysis [39] in intracranial EEG.

MS, a disorder caused by a condition called inflammatory demyelinating of the nervous system, is the most common among all neurological disorders. MS causes disabilities in young adults and affects nearly 2.5 million people worldwide. The diagnosis of MS is generally per-

formed by magnetic resonance imaging (MRI). There are no treatments available for this disease [40, 41].

PD is a chronic neurodegenerative disorder often characterized by the presence of predominantly motor symptomatology [42, 43], but it can have nonmotor hyposmia, paresthesia, depression, and pain [44]. PD is a universal disorder with incidence rate of 4.5–19 per 100,000 of population per year [45–47] for both females and males of all ages. The therapy depends on severity, mental status, and age of the patient. Gálvez et al. [48] investigate the short-term effects of Binaural Beats on EEG power, functional connectivity, cognition, gait, and anxiety in PD patients.

Stroke is a clinical syndrome of cerebral deficit that lasts for >24 h with no apparent cause except the vascular one [49]. In the modern developed countries, 75–80% of the strokes are attributed to brain ischemia, and 10–15% are attributed to intracerebral hemorrhage. Stroke diagnosis is made accurately completely based on clinical grounds by a specialist alone.

Traumatic brain injury is one of the foremost causes of disability and death in young adults and children worldwide. More than five million people suffer from the traumatic brain injury disability in the United States alone [50–52].

Authors have been advocating the research ideology that a computer-aided diagnosis (CAD) system trained using lots of patient data and physiological signals and images based on adroit integration of advanced signal processing and artificial intelligence (AI)/machine learning (ML) techniques in an automated fashion can assist neurologists, neurosurgeons, radiologists, and other medical providers to make better clinical decisions. Research in this area has been growing at an accelerating rate in the past decade. In this paper, authors explore and review recent articles on the applications of AI-based CAD systems for the diagnosis of 5 major neurological disorders: epilepsy, AD, PD, MS, and ischemic brain stroke.

Figure 1 shows the functional block diagram of a typical ML-based CAD system consisting of 5 stages: (1) signal transformation, (2) feature extraction, (3) feature dimensionality reduction, (4) optimal feature selection/ranking, and (5) classification.

ML-Based CAD

Input Data Description

Input data for a CAD system are normally signals and/or images. For PD, many CAD systems use speech and




Fig. 1. General block diagram of a typical ML-based CAD system.

EEG for the diagnosis. Image-based approaches normally use MRI and single-photon emission computed tomography scan. For epilepsy, most authors have used Bonn University EEG data set [53]. In MS, T1 and T2 weighted MRI images are commonly used for the diagnosis, where T1 and T2 refer to the time taken between magnetic pulses and the image is taken. Most of the AD systems have used T2-weighted MRI images from Alzheimer Disease Neuro Imaging [54] and Open Access Series of Imaging Studies [55] databases.

Image Transformation

In general, image transformation is performed first where the redundant information is removed and then features are extracted from the transformed images. This step helps in gathering significant information that can be used for feature extraction.

Feature Extraction

Signal-Based Approach

Discrete wavelet transform (DWT) is often used to convert the signal to low- and high-frequency components [56–58]. The curvelet transform that is a higher dimensional DWT represents the images in multiple angles and scales [59]. The higher-order spectra (HOS) features are also used for feature representation and extraction [60, 61]. The extracted features must represent the hidden clues present in the input data.

Image-Based Approach

In image preprocessing methods, intensity normalization, adaptive histogram equalization [62], and background subtractions are performed prior to level set segmentation to detect the region of interest [63]. The gray level co-occurrence matrix features are most commonly used for the images [64]. The entropy and energy features are also used in many articles [65–70]. The wavelet-based energy and entropy features are also employed [71–74]. Various statistical measures such as Hu’s moments [75], Zernike moments [76],

central moments [77], and statistical moments [77] are used as features both in signal- and image- based approaches for developing CAD systems for neurological disorder.

Dimensionality Reduction

Feature extraction techniques often yield a large number of features that may be redundant and result in excessive computational requirements which in turn may make their real-time application impractical or unnecessarily difficult. Hence, various feature dimensionality reduction techniques are commonly used. The most commonly used methods are principal component analysis (PCA) [78], linear discriminant analysis [78], and independent component analysis [79]. Extended versions of PCA such as kernel PCA have also been employed in the literature [80].

Optimal Feature Selection and Ranking

Majority of features exhibit redundant information, which need to be removed to obtain optimum classification performance. The analysis of variance [81] is the most commonly used method when 3 or more classes are involved. Other commonly used optimal feature selection techniques are Student *t* test [82], entropy [83, 84], Wilcoxon rank tests [85–88], Bhattacharyya distance [89], receiver operating characteristic [90], genetic algorithm [91], particle swarm optimization [92, 93], and ant colony optimization [94]. A number of researchers have combined different selection methods in order to obtain the most significant features [95]. The fuzzy logic-based min-redundancy and max-relevance feature selection have been used for diagnosis of PD [96].

Feature Classification

Classification techniques generally have 2 phases: (i) training and (ii) testing. They need to be trained using previously collected data. Once trained, they can be used for classification of new cases. The most commonly used

Table 1. Summary of CAD systems for diagnosis of epilepsy (modality: EEG)

Authors, years	Methodology/features	Classifier	Accuracy, %
Acharya et al. [149], 2009	CD, Hurst exponent (H), ApEn	GMM + SVM	EPacc = 100
Acharya et al. [150], 2011	WPD and HOS	Fuzzy	EPacc = 98.5
Acharya et al. [151], 2011	RQA features	SVM	EPacc = 95.6
Acharya et al. [152], 2012	Entropies + HOS + higuchi FD + hurst/nonlinear features	Fuzzy	EPacc = 99.7
Acharya et al. [153], 2012	Entropy parameters	Fuzzy	EPacc = 98.1
Acharya et al. [154], 2012	Eigen values + WPD coefficients + PCA	GMM	EPacc = 99
Acharya et al. [155], 2013	DWT + ICA	SVM	EPacc = 96
Aslan et al. [156], 2008	–	RBFNN + MLPNN	EPacc = 95.2
Chua et al. [157], 2011	HOS-based features	GMM	EPacc = 93.1
Faust et al. [158], 2010	Frequency parameters + Burg's method	SVM	EPacc = 93.3
Ghosh-Dastidar et al. [159], 2007	Mixed-band feature space	LMBPNN	EPacc = 96.7
Ghosh-Dastidar et al. [160], 2008	Mixed-band feature space + PCA	RBFNN	EPacc = 96.6
Ghosh-Dastidar et al. [161], 2009	Mixed-band feature space	MuSpiNN	EPacc = 94.8
Guler et al. [162], 2005	Lyapunov exponents	RNN	EPacc = 96.79
Guo et al. [163], 2009	Relative wavelet energy	ANN	EPacc = 95.2
Guo et al. [164], 2010	ApEn + wavelet transform	ANN	EPacc = 99.85
Guo et al. [165], 2010	Line length features + wavelet transform	ANN	EPacc = 99.6
Guo et al. [166], 2011	Genetic programming	KNN	EPacc = 99.2
Iscan et al. [167], 2011	Cross correlation and power spectral density	SVM	EPacc = 100
Kannathal et al. [168], 2005	Entropy measures	ANFIS	EPacc = 90
Lima et al. [169], 2010	DWT	SVM	EPacc = 100
Martis et al. [170], 2012	EMD	C4.5	EPacc = 95.33
Nigam and Graupe [171], 2004	Nonlinear filter	LAMSTAR ANN	EPacc = 97.2
Orhan et al. [172], 2011	DWT and K-means clustering	MLPNN	EPacc = 100
Polat and Gunes [173], 2007	FFT	DT	EPacc = 98.72
Polat and Gunes [174], 2008	FFT + PCA	AIRS	EPacc = 100
Polat and Gunes [175], 2008	AR	C4.5	EPacc = 99.32
Sadati et al. [176], 2006	DWT	ANFN	EPacc = 85.9
Srinivasan et al. [177], 2005	Time and frequency domain features	EN	EPacc = 99.6
Srinivasan et al. [178], 2007	ApEn	PNN, EN	EPacc = 100
Subasi [179], 2007	DWT + statistical measures	ME	EPacc = 94.5
Subasi and Gursoy [180], 2010	DWT + PCA	SVM	EPacc = 100
Tzallas et al. [181], 2007	Time–frequency methods	ANN	EPacc = 99.28

Table 1. (continued)

Authors, years	Methodology/features	Classifier	Accuracy, %
Jaiswal and Banka [182], 2017	SpPCA and SubXPCA	SVM	EPacc = 94.60
Ubeyli [183], 2010	AR	LS-SVM	EPacc = 99.56
Wang et al. [184], 2011	Wavelet + entropy	KNN + HKB	EPacc = 100
Swami et al. [185], 2016	DTCWT	GRNN	–
Peker et al. [186], 2016	DTCWT	CVNN	EPacc = 100
Sharma and Pachori [187], 2017	TQWT	LS-SVM + FD	EPacc = 100
Patidar et al. [188], 2017	TQWT and kraskov entropy	LS-SVM	EPacc = 97.75
Gandhi et al. [189], 2011	DWT, energy, SD, entropy	PNN	EPacc = 99.9
Chen [190], 2014	DTCWT, fourier features	NN	EPacc = 100
Swami et al. [191], 2014	Energy, SD, entropy features	SVM	EPacc = 99.53
Pachori and Patidar [192], 2014	EMD and SODP	ANN	EPacc = 100
Sharma and Pachori [193], 2015	EMD and PSR of IMF	LS-SVM	EPacc = 98.67
Bhattacharyya et al. [194], 2017	TQWT and KNN entropies	SVM	EPacc = 100
Bhattacharyya et al. [195], 2016	Empirical wavelet transform	LS-SVM	EPacc = 90
Bhati et al. [196], 2017	Time–frequency localized three-band biorthogonal linear phase wavelet filter bank	MLPNN	EPacc = 99.33
Sharma et al. [197], 2017	Orthogonal wavelet filter banks	LS-SVM	EPacc = 94.25
Kaya et al. [198], 2014	1D – LBP	FT	EPacc = 99.5
Zhu et al. [199], 2014	FWHVA	KNN	EPacc = 100
Samiee et al. [200], 2015	DSTFT	MLPNN	EPacc = 99.8
Riaz et al. [201], 2016	EMD	SVM	EPacc = 96.2
Diykh et al. [202], 2017	Weighted complex network combined with time domain features	LS-SVM	EPacc = 98
Li et al. [203], 2017	MODWT and LND	RFC	EPacc = 100
Acharya et al. [7]	–	CNN	Eac = 88.7
Ghayab et al. [205], 2016	SRS and SFS	LS-SVM	EPacc = 99.9
Sharma et al. [206], 2018	MMSFL-OWFB based KE	SVM	EPacc = 100
Tiwari et al. [207], 2017	Pyramid scheme for key-point localisation and LBP	SVM	EPacc = 99.89
Bajaj and Pachori [208], 2012	EMD based intrinsic mode functions and hilbert transform	LS-SVM	EPacc = 100
Nicolaou and Georgiou [209], 2012	Permutation entropy	SVM	EPacc = 93.55
Xie and Krishnan [210], 2013	Wavelet-based sparse functional linear model	KNN (1NN)	EPacc = 100
Mursalin et al. [211], 2017	ICFS	RFC	EPacc = 100
Upadhyay et al. [212], 2016	DWT based features	LS-SVM	EPacc = 100

Table 1. (continued)

Authors, years	Methodology/features	Classifier	Accuracy, %
Kabir et al. [213], 2016	Optimum allocation technique	LMT	EPacc = 95.33
Murugavel and Ramakrishnan [214], 2016	Wavelet transform based features	H-MSVM with ELM	EPacc = 93.63
Pippa et al. [215], 2016	Time domain and frequency domain features	Bayesian net	EPacc = 95
Kumar et al. [216], 2014	LBP	kNN (1-NN)	EPacc = 98.33
Naser et al. [217], 2019	DWT and approximation and abe entropies	SVM	EPacc = 98.75
Tzimourta et al. [218], 2018	Wavelet transform based features	Random forest classifier	EPacc = 95
Lamhiri and Shmuel [219], 2019	Hurst exponent	k-NN	EPacc = 100
Raghu et al. [220], 2019	Sigmoid entropy	SVM	EPsen = 100
Wang et al. [221], 2019	Symlet wavelet processing, and grid search optimizer	Gradient boosting machine	EPacc = 96.5
Bose et al. [222], 2019	Multifractal detrended fluctuation analysis	SVM	EPacc = 100
Dalal et al. [223], 2019	FAWT and FD	RELS-TSVM	EPacc = 90.2
Sriam et al. [224], 2018	Teager energy feature	Supervised back propagation neural network	EPsen = 96.66
Shaikh et al. [225], 2017	EMD	ANN	EPacc = 96.1
Osman and Alzahrani [226], 2019	SOM	RBFNN	EPacc = 97.47
Sudalaimani et al. [227], 2018	Sub-frequency band features	GRNN	EPacc = 91.6
Raghu and Sriram [228], 2018	NCA	SVM	EPacc = 98.8
Li et al. [37], 2018	GMM and GLCM features, RFE-SVM	SVM	EPacc = 100
Cooman et al. [229], 2018	HRI features	SVM + Adaptive Heuristic classifier	EPsen = 83.3
Li et al. [230], 2018	WPT and KDE	LS-SVM	EPacc = 99.6
Cruz et al. [231], 2018	ACC and EMG	SVM on cloud computing platform	EPsen = 83.3
Kocadagli and Langari [232], 2017	DWT and fuzzy relations	ANN	EPacc = 99.9
Zhang et al. [233], 2018	WPD, fDistIn	kNN	EPsen = 98.33
Feng et al. [234], 2018	WPD	SVM	EPacc = 98.67
Tanveer et al. [235], 2018	FAWT and entropy-based features	RELS-TSVM	EPacc = 100
Choudhury et al. [236], 2018	XHST	kNN	EPacc = 100
Torse et al. [237], 2017	EMD	CSM-SVM	EPacc = 96.4
Tomanik et al. [238], 2019	Complex networks	–	EPacc = 98.2
Wani et al. [239], 2018	DWT	ANN	EPacc = 95

Table 1. (continued)

Authors, years	Methodology/features	Classifier	Accuracy, %
<i>Deep learning based techniques</i>			
Yuan et al. [136], 2017	STFT-Mssda	Softmax	EP _{Pacc} = 93.82
Ullah et al. [137], 2018	P-1D-CNN		EP _{Pacc} = 99.9
Johansen et al. [135], 2016	CNN		AUC = 94.7
Antoniades et al. [134], 2016	CNN		EP _{Pacc} = 87.51
Qi et al. [240], 2014	MCC-based R-SAE model	SVM	EP _{sen} = 100
Lin et al. [241], 2016	SSAE	Softmax	EP _{Pacc} = 96
Gogna et al. [242], 2017	Semi-supervised stacked autoencoder		EP _{Pacc} = 96.9
Acharya et al. [9], 2018	CNN		EP _{Pacc} = 88.67
Tjepkema-Cloostermans et al. [139], 2018	Convolutions (1D and 2D) and/or LSTMs		EP _{spe} = 99.9
Hussein et al. [243], 2019	LSTM + FC		EP _{spe} = 100
Thodoroff et al. [244], 2016	CNN + RNN		EP _{sen} = 85
Emami et al. [245], 2019	CNN		Detection rate = 100
Van Putten et al. [139], 2018	CNN + RNN		EP _{spe} = 100
Jang and Cho [246], 2019	Dual deep neural network	–	EP _{sen} = 100
Zuo et al. [247], 2019	CNN	–	EP _{sen} = 83.23
Wei et al. [248], 2016	Multichannel CNN	–	EP _{Pacc} = 92.40
Achilles et al. [249], 2016	CNN	–	AUC = 78.33
Yuvaraj et al. [250], 2018	CNN	–	EP _{sen} = 86.29
Maria Hugle et al. [251], 2018	CNN	–	EP _{sen} = 96
Thomas et al. [252], 2018	CNN	SVM	EP _{Pacc} = 83.86

CAD, computer aided diagnosis; CD, correlation dimension; EEG, electroencephalogram; ApEn, approximate entropy; SVM, support vector machine; RQA, recurrence quantification analysis; HOS, higher order spectra; PCA, principal component analysis; KNN, k-nearest neighbour; DWT, discrete wavelet transform; ICA, independent component analysis; GMM, gaussian mixer model; PNN, probabilistic neural network; SRS, simple random Sampling; SFS, sequential feature selection; HRI, heart rate increases; XHST, cross hyperbolic S-transform; CNN, convolutional neural network.

classifiers for diagnosis of the neurological disorder are probabilistic neural network classifier [97], support vector machine (SVM) with different kernel functions such as polynomial (Poly) of orders 1, 2, and 3 [58, 98], Naive Bayes [99, 100], k-nearest neighbor [101], linear discriminant analysis, quadratic discriminant analysis [102], decision tree [101], random forest, and Gaussian mixer model [103, 104]. Among these, SVM classifier is one of the most commonly used. More recently, enhanced probabilistic neural network has been used for accurate diagnosis of PD [105].

The consolidated lists of various machine ML-based CAD systems for the diagnosis of neurological disorders are presented in Tables 1–7 for epilepsy, PD, AD, ischemic brain stroke, and MS.

Deep Learning-Based Techniques

To solve the limitations of ML-based techniques, deep learning techniques have recently been advanced. An example of deep learning is the convolutional neural network

Table 2. Summary of CAD systems for diagnosis of PD using measurable indicators (modality: EEG)

Authors, years	Methodology/features	Classifier	Accuracy, %
Yuvaraj et al. [253], 2014	HOS (6 basic emotional states)	KNN, SVM	PDacc: 93.42 (happiness)
Yuvaraj et al. [254], 2014	BNoA (bispectrum feature), ICA	SVM, RBF	PDacc: 76.90±1.08
Yuvaraj et al. [255], 2016	Bispectral functional connectivity index	SVM	PDacc: 51.66±1.02
Nilashi et al. [256], 2017	Data mining technology	CART	–
Tucker et al. [257], 2015 (modality: sensors)	Data mining driven technology	Naïve Bayes	PDacc: 78.00
Prashantha et al. [258], 2018	Machine learning techniques	Logistic regression, random forests, boosted trees, SVM	Acc: >95
<i>Deep learning based techniques</i>			
Ali et al. [259], 2016	DBF (handwriting analysis)	–	Acc: 94.00
Caliskan et al. [260], 2017	Deep neural network (speech dataset)	Softmax classifier	Acc: 86.09, sensitivity: 58.27, specificity: 95.38
Grover et al. [261], 2018	Deep neural network (voice data)	UPDRS	Total UPDRS accuracy: 62.73

CAD, computer-aided diagnosis; PD, Parkinson's disease; EEG, electroencephalogram; HOS, higher-order spectra; ICA, independent component analysis; DBF, deep belief network; SVM, support vector machine.

Table 3. Summary of CAD systems for the detection of PD using brain images (modality: MRI)

Authors, year	Methodology/features	Classifiers	Accuracy, %
Oliveira et al. [262], 2015	Voxel features	SVM	PDacc: 97.86, PDsen: 97.75, PDspe: 98.09
Hirschauer et al. [105], 2015	–	EPNN	PDacc: 92.50
Banerjee et al. [263], 2016	CDT + FA	PGA	PDacc: 98.53, PDsen: 98, PDspe: 100
Cigdem et al. [264], 2018	GM + WM + PCA	SVM	PDacc: 93.75, PDsen: 95, PDspe: 92.50
Segovia et al. [265], 2015 (modality: PET CT)	NB	SVM	PDacc: 78.16
Ahmadlou et al. [266], 2010 (modality: SPECT)	Gaussian Kernel	–	PDacc: 92.5
<i>Deep learning-based techniques</i>			
Choi et al. [267], 2017 (modality: SPECT)	Deep neural network	Softmax classifier	Acc: 98.8, sensitivity: 98.6, Specificity: 100
Sivaranjini et al. [268], 2019	CNN	Alexnet	Acc: 88.90, sensitivity: 84.40, specificity: 88.40

CAD, computer-aided diagnosis; PD, Parkinson's disease; MRI, magnetic resonance imaging; PCA, principal component analysis; NB, Naïve Bayes; CNN, convolutional neural network; SVM, support vector machine; PGA, principal geodesic analysis.

Table 4. Summary of CAD systems for the detection of PD using physiological signals (modality: EEG)

Authors, year	Methodology/features	Classifiers	Accuracy, %
Han et al. [269], 2013	WPE	AR Burg	–
Yuvaraj et al. [270], 2016	HOS, PD diagnosis index	SVM	PDacc: 99.62, PDsen: 100, PDspe: 99.25
Hariharan et al. [271], 2014	Dysphonia + GMM + PCA + LDA + SFS	SVM	PDacc: 100
Zhang et al. [272], 2017	Dysphonia + stacked autoencoders	KNN	PDacc: 94–98
Hlavnicka et al. [273], 2017	Zero-crossing rate + VAC function	GMM	PDacc: 71.30, PDsen: 56.70, PDspe: 80
Joshi et al. [274], 2017	DWT	SVM	PDacc: 90.32
Ornelas-Vences et al. [275], 2017	Turn analysis + biomechanical features	Fuzzy model	–
Sama et al. [276], 2017	Butterworth filter	SVM	PDacc: <90
Oung et al. [277], 2018	EWPT, wavelet energy, entropy	ELM	PDacc: 95.93
Little et al. [278], 2019	Preselection filter, exhaustive Search	SVM	PDacc: 91.4
Shahbaba and Neal [279], 2009	Dirichlet process mixtures	Decision tress, SVM	PDacc: 87.70
Das [280], 2010	Back-propagation learning algorithm	ANN	PDacc: 92.9
Sakar and Kursun [281], 2010	Mutual information-based feature selection	SVM	PDacc: 92.75
Psorakis et al. [282], 2010	Improved mRVMs	SVM	PDacc: 89.47
Guo et al. [283], 2010	Genetic programming and the expectation maximization algorithm	GP-EM	PDacc: 93.10
Ozcift and Gulden [284], 2011	CFS	RF	PDacc: 87.10
Li et al. [285], 2011	Fuzzy-based nonlinear transformation	SVM	PDacc: 93.47
Luukka [286], 2011	Fuzzy entropy measures	Similarity classifier	PDacc: 85.03
Spadoto et al. [287], 2011	PSO, harmony search, gravitational search	Optimum path forest classifier	PDacc: 73.53
Astrom and Koker [288], 2011	Nine layer neural networks	PNN	PDacc: 91.20
Ozcift [289], 2012	RF ensemble of IBk (a KNN variant) algorithm	SVM	PDacc: 97.00
Polat [290], 2012	FCMFW	KNN	PDacc: 97.93
Tsanas et al. [291], 2012	Feature selection: LASSO, mRMR, RELIF, LLBFS	SVM	PDacc: ~90.00
Daliri [292], 2013	STFT, FDR	SVM	PDacc : 91.20
Chen et al. [293], 2013	PCA	FKNN	PDacc: 96.07
Zuo et al. [294], 2013	PSO	FKNN	PDacc: 97.47
Ma et al. [295], 2014	SCFW	KELM	PDacc: 99.49
Drotár et al. [296], 2016	Entropy and energy	SVM, RBF	PDacc: 81.30
Connolly et al. [297], 2015 (modality: sensors)	Feature reduction	SVM	–
Wahid et al. [298], 2015	Neural networks	SVM	PDacc: 86.00

Table 4. (continued)

Authors, year	Methodology/features	Classifiers	Accuracy, %
Smith et al. [299], 2015 (modality: sensors)	Evolutionary algorithms	ROC	PDacc: 78.00
Hirschauer et al. [105], 2015	Neural networks	EPNN	PDacc: 92.5
Shamir et al. [300], 2015	Clinical decision support systems	SVM	PDacc: 71.00
Procházka et al. [301], 2015 (modality: sensors)	Bayesian	Bayesian probability classifiers	PDacc: 94.1
Nilashi et al. [302], 2018	Partial least squares, self organizing map	SVM	UPDRS acc: 46.56
<i>Deep learning based techniques</i>			
Tagaris et al. [140], 2017	Deep neural network	CNN-RNN	Acc: 74
Kim et al. [303], 2018	Neural networks	CNN	PDacc: 85
Oh [304], 2018	CNN	–	Acc: 88.25, sensitivity: 84.71, specificity: 91.77

CAD, computer aided diagnosis; ICA, independent component analysis; EEG, electroencephalogram; HOS, higher order spectra; PD, Parkinson's disease; WPE, wavelet packet entropy; SVM, support vector machine; GMM, gaussian mixture model; KNN, k-nearest neighbour; DWT, discrete wavelet transform; CNN, convolutional neural network; PSO, particle swarm optimization; PNN, probabilistic neural network; PCA, principal component analysis.

(CNN) with a number of advantages over conventional ML-based CAD systems. First and foremost is that it can handle large data sets using its multilayer architecture. It can also handle imbalanced data samples and execute the results without biasing toward majority class. The CNN architecture generally includes a convolution layer, a pooling layer, and fully connected layers [106]. These layers can be extended with few additional layers based on the requirements. Recently, various CNN-based techniques have shown good results in detecting neurological disorders [107–116]. But, detection of such abnormalities in its early stage may require more deeper architecture. Such deeper architecture can pick distinct features from the signals by convolving with kernels defined in each layers [106].

In the past few years, deep neural network learning techniques have been used to solve intractable problems in a variety of disciplines such as image recognition [117–119], pavement crack detection [120], structural damage detection [121, 122], vehicle-type detection [123], transportation systems [124, 125], concrete strength estimation [126], big data time series forecasting [127], and computer-brain interface [128]. Following that trend, a number of researchers have developed CAD systems using

deep learning techniques. These techniques can discover structures or patterns in the biological signals or images. Its effectiveness in signal and image-based CAD systems has been demonstrated recently [117, 127, 129–133].

Deep learning techniques can be broadly classified into 2 types: (1) supervised learning and (2) unsupervised learning. The CNN belongs to the first and auto encoders belong to second type. Antoniadis et al. [134] analyze the epileptic intracranial EEG data using deep learning. Johansen et al. [135] use CNN for epileptiform spike detection. Yuan et al. [136] employ a combination of deep learning and short-time Fourier transform for epileptic seizure detection. Ullah et al. [137] report application of deep learning for epilepsy diagnosis. Martinez-Murcia et al. [138] employ the CNN for analysis of neuroimaging in PD.

The most common architecture of CNN includes a bank of convolution, max-pool, fully connected, and soft-max layers as shown in Figure 2 [129–131]. The 3D CNN and CNN – Recurrent neural network (RNN) are more recent networks used in the diagnosis of epilepsy [139], PD [140] and AD [141]. Other CNN variants are also experimented on PD and ischemic brain stroke detection. The list of approaches used is presented in Tables 1–7. It

Table 5. Summary of CAD systems for the detection of AD (modality: MRI)

Authors, years	Methodology/features	Classifier	Accuracy, %
Al-nuaimi et al. [305], 2015	Tsallis entropy	K-means clustering	ADacc: 77
Silveira and Marques [306], 2010 (modality: PET CT)	Boosting classification method	Boosting classifier	ADacc: 90.97 MCIacc: 79.63
Mahanand et al. [307], 2011	PCA + VBM + ANN	SRAN	Eff: 91.18
Mahmood and Ghimire [308], 2013	PCA + ANN	Feed forward multilayer neural network	ADacc: 90
Yi Ding et al. [309], 2015	VBM + GLCM + gabor filters	SVM	ADacc: 92.86
Herrera et al. [310], 2013	DWT + PCA	SVM	ADacc: 93.90
Sweety and Jiji [311], 2014	PCA + PSO	SVM + decision tree classifier	ADacc: 91.89
Saraswathi et al. [312], 2013	PSO + VBM	GA-ELM-PSO classifier	ADacc: 94.57
Mahanand et al. [313], 2013	VBM + ICGA + ELM	ELM classifier	ADacc: 91.86
Mahanand et al. [314], 2014	VBM + RFE	PBL-McRBFN classifier	ADacc: 89.81
Mathew et al. [315], 2016	DWT + PCA + FDR	SVM	ADacc: 84
Escudero et al. [316], 2013	Machine learning biomarkers	Locally weighted learning	ADacc: 80
Gunawardena et al. [317], 2016	CNN	SVM	Error: 13.13
Zhang et al. [318], 2017	PCA	SVM	ADacc: 84.40
Escudero et al. [319], 2011	K-means clustering + bioprofile analysis	Semi-supervised classifier	–
Zhang et al. [320], 2016	Landmark-based AD diagnosis system	SVM	ADacc: 83.7
Iftikhar et al. [321], 2016	Cortical thickness-based features + volume based features + F-score method	SVM	ADacc: 83.65
Bates et al. [322], 2011	Point-cloud laplacian + spectral signatures + shape analysis	Linear SVM	ADacc: 77.5
Ye et al. [323], 2011	KNN + RAVENS maps + ISOMAP algorithm	LapSVM	ADacc: 94.1
Rabeh et al. [324], 2017	NLMS	Bayes probability	ADacc: 92
Ullah et al. [325], 2018	CNN	SVM	ADacc: 80.25
Donini et al. [326], 2016	MKL RFE	SVM RFE	ADacc: 96.14±3.55
<i>Deep learning based techniques</i>			
Sarraf et al. [327], 2016	CNN	LeNet and GoogleNet binary classifier	ADacc: 99.9 (fMRI) AD acc: 98.84 (MRI)
Cheng and Liu [129], 2017 (Modality: PET CT)	3D-CNN	–	ADacc: 92.2 ADsen: 91.4 ADspe: 92.4
Cullen et al. [130], 2018	CNN	–	ADmean: 0.9969 ADSD: 0.0018

Table 5. (continued)

Authors, years	Methodology/features	Classifier	Accuracy, %
Islam et al. [131], 2018	DCNN	Softmax classifier	ADacc: 93
Liu et al. [141], 2018	CNN-RNN	Softmax classifier	ADacc: 91.2 ADsen: 91.4 ADspe: 91.0
Liu et al. [328], 2018	Cascaded CNN	Softmax classifier	ADacc: 93.26 ADsen: 92.55 ADspe: 93.94
McCrackin et al. [329], 2018	CNN	–	–
Basaia et al. [330], 2019	CNN	–	ADacc: 99.2 ADsen: 98.9 ADspe: 99.5
Hosseini-Asl et al. [331], 2016	CNN	3D-ACNN	ADacc: 97.60
Awate and Bangare [332], 2018	CNN	Softmax classifier	ADacc: 99
Billones et al. [333], 2016	CNN	DemNet	ADacc: 98.33 ADsen: 98.89 ADspe: 97.78
Gunawardena et al. [334], 2017	CNN	Fully connected layer	ADacc: 96.50 ADsen: 96.00 ADspe: 98.00
Farooq et al. [335], 2017	CNN	GoogLeNet	ADacc: 98.8
Luo et al. [336], 2017	CNN	Fully connected layer	ADsen: 100 ADspe: 93.00
Lin et al. [337], 2018	CNN	–	ADacc: 79.9

CAD, computer-aided diagnosis; MRI, magnetic resonance imaging; DWT, discrete wavelet transform; PSO, particle swarm optimization; ICGA, integer coded genetic algorithm; RFE, recursive feature elimination; FDR, fisher discriminant ratio; CNN, convolutional neural networks; VBM, voxel-based morphometry; NLMS, non local means; MKL, multiple kernel learning.

is observed that, in general, CNN-based techniques outperformed conventional handcrafted techniques with the limitation of requiring more training samples to achieve the optimal performance.

Discussion

Research on automated CAD systems is more prevalent first on epilepsy followed by PD and then AD. For seizure detection and epilepsy diagnosis, they are based on EEG signals. For PD, they are developed using different modalities such as EEG, speech, handwriting, and brain images. The automated systems proposed in last 2 decades are summarized in Tables 1–7. Table 1 summarizes the application of AI techniques used for automated seizure detection and

epilepsy diagnosis. It can be seen from Table 1 that WT is the dominant method for multiresolution analysis. The SVM and the back propagation neural network are more commonly used classifiers. Tables 2–4 summarize research on the automated diagnosis of PD using different modalities. WT, HOS coupled with entropy and energy features are found to perform well. SVM is the most commonly used classifier with different kernels for the diagnosis of PD.

For Alzheimer disease, DWT and different dimensionality reduction techniques have been widely used [19, 142–148]. The gray level co-occurrence matrix-based techniques with SVM classifier are the commonly used classifier techniques (Table 5).

Most of the ischemic brain stroke CAD systems are developed using fuzzy clustering and SVM and random forest classifiers (Table 6). It is observed that very little

Table 6. Summary of CAD systems for the detection of ischemic brain stroke (modality: MRI)

Authors, years	Methodology/features	Classifier	Accuracy, %
Li et al. [338], 2004	Manual lesion tracing	MSSC + PVVR	BS _{similarity} : 0.97
Filho et al. [339], 2017 (modality: CT)	ABTD extractor used to extract features	MLP, SVM, kNN, OPF, and Bayesian	BS _{acc} : 95
Lutsep et al. [340], 1997	Radiologic techniques	Acute stroke treatment (TOAST) classification scheme	
Wang et al. [341], 2013	Novel framework	LLC	
Alpert et al. [342], 2012	Probabilistic framework	Clustering algorithm	BS _{fscore} : 0.52
Ji et al. [343], 2017	Asymmetric Distribution	Gaussian Mixture Model	BS _{acc} : 5% more
Sridevi et al. [344], 2019	Image segmentation (ACM and SOM)	Supervised model	–
Agrawal et al. [345], 2014	OBPD method	FCM	–
Monteiro et al. [346], 2018	Novel hybrid GA-BFO-CM algorithm	l1 regularized logistic regression; decision tree; SVM; random forest; Xgboost	BS _{auc} : 0.936
MateSin et al. [347], 2001	Rule-based approach	Expert system	–
Ghosh et al. [348], 2011	HRS	Expert system	–
Mitra et al. [349], 2014	Bayesian-MRF	Random forest	BS _{sensitivity} : 0.53±0.13 BS _{meanpositivevalue} : 0.75±0.18
Maier et al. [350], 2015	FCM approach	Random forest	BS _{dicecoeff} : 0.65
Griffs et al. [351], 2016	Probabilistic tissue segmentation	NB classifier	–
Pustina et al. [352], 2016	Manual lesion tracing	Region wise LSM	BS _{avgcorr} : 0.77
Pennisi et al. [353], 2016	Skin lesion image segmentation	NB, Adaboost, KNN, and random trees	BS _{sensitivity} : 93.5 BS _{specificity} : 87.1
Chen et al. [354], 2017	Novel framework	Convolutional neural networks	BS _{acc} : 0.67
Muda et al. [355], 2015	FCM algorithm	Expert system	BS _{Average dice indices} : 0.73 (acute stroke), 0.68 (tumour) and 0.82 (chronic stroke)
Paul Bentley et al. [356], 2014 (modality: CT)	CT brain machine learning	SVM	BS _{AUC} : 0.744
Lebedev et al. [357], 2018	Artificial intelligence	RetinaNet network detector	–
Subudhi et al. [358], 2018	Delaunay triangulation and fractional order DPSO	Random forest	BS _{sen} : 0.93 BS _{acc} : 0.95
Subudhi et al. [359], 2018	Watershed-based lesion segmentation algorithm	Random forest	BS _{acc} : 0.85 BS _{DSI} : 96
<i>Deep learning based techniques</i> Chin et al. [360], 2017	Deep learning	CNN	BS _{acc} : 0.90

CAD, computer aided diagnosis; MRI, magnetic resonance imaging; NB, Naive Bayes; kNN, k-nearest neighbour; FCM, fuzzy C-means; CNN, convolution neural network; LSM, lesion to symptom mapping; SVM, support vector machine; LLC, locally linear classification; OBPD, optimum boundary point detection; HRS, hierarchical region splitting; MRF, markov random field.




Fig. 2. Conventional architecture of CNN.

Table 7. Summary of CAD systems for the detection of MS (modality: MRI)

Authors, years	Methodology/features	Classifier	Accuracy, %
Elliott et al. [361], 2013	Joint time point bayesian formulation	Bayesian + random-forest	MSsen: 99
Anbeek et al. [362], 2004	MBRASE	KNN	MSsen: 0.791
Ashton et al. [363], 2003	GEORG + DMSS	Bayesian	Coefficient of variation: manual: 8.2, GEORG: 2.4, DMSS: 2.1
Lao et al. [364], 2008	BET	SVM	–
Yamamoto et al. [107], 2010	Rule-based method, level set method	SVM	MSsen: 81.5
Zacharaki et al. [108], 2008	BET	SVM	TVR (% vol increase): 7.58–375, IVR (% vol increase): 11.14–340.32
<i>Deep learning-based techniques</i>			
Zhang et al. [365], 2018	CNN	–	Acc: 98.23
Wang et al. [366], 2018	CNN	–	Acc: 98.77

CAD, computer-aided diagnosis; MRI, magnetic resonance imaging; CNN, convolutional neural network; MS, multiple sclerosis; SVM, support vector machine; kNN, k-nearest neighbour.

research is done on the automated diagnosis of MS. The SVM and k-nearest neighbor classifiers are the most commonly used classifier for the diagnosis of MS.

The transform methods such as DWT and HOS have been widely used with good performances. This can be explained by the fact that after the transformation, these methods can indicate minute changes in the frequency domains. The entropies and energy values of these DWT and HOS coefficients are able to separate the classes.

Conclusion

This study has explored different state-of-the-art AI-based CAD systems developed in last 2 decades for the diagnosis of 5 different neurological disorder. Results are summarized Tables 1–7.

The future research can be in one of 3 directions:

Creation of a Large Public Database for the Validation of All Developed Techniques

As observed in Tables 1–7, the number of subjects used in published research is often small. As such, the conclusions reported should be considered tentative. There is a need to create a large standard public database for all aforementioned neurological disorders.

ML-Based CAD for Early Detection of Neurological Disorders

As more data become available, especially longitudinal data, the research should shift to ML-based CAD systems for early detection of the neurological disorders.

Remote Access by Nonexpert Clinicians

Highly trained and specialized neurologists and associated facilities for diagnosis of neurological disorders may not be available in remote towns and villages. Figure 3




Fig. 3. Prototype of a cloud-based CAD system for diagnosis of neurological disorders using Internet of things. EEG, electroencephalogram; CAD, computer-aided diagnosis.

presents the prototype of a cloud-based CAD system for diagnosis of neurological disorders using Internet of things. The signals or images of patients are transmitted to the cloud where a trained ML-based CAD can provide the diagnosis. The outcome of the model is sent to the cell phone of a nonexpert clinician for a preliminary diagnosis.

Acknowledgments

Authors wish to thank Dr Amir Adeli, Board-Certified Neurologist, Columbus, OH, for his suggestions to improve the quality of the paper.

Statement of Ethics

This paper is a review article with no human/animal participants. Hence, no ethical approval was needed.

References

- 1 World Health Organization. World Health Organization, 27 02 2017. [Accessed December 12, 2018]. Available from: <https://www.who.int/mediacentre/news/releases/2007/pr04/e/>.
- 2 Acharya UR, Hagiwara Y, Adeli H. Automated seizure prediction. *Epilepsy Behav.* 2018; 88:251–61.
- 3 Bairy GM, Lih OS, Hagiwara Y, Subha DP, Faust O, Niranjan UCA, et al. Automated diagnosis of depression electroencephalograph signals using linear prediction coding and higher order spectra features. *J Med Imaging Health Inform.* 2017;7(8): 1857–62.
- 4 Bhat S, Acharya UR, Adeli H, Bairy GM, Adeli A. Autism: cause factors, early diagnosis and therapies. *Rev Neurosci.* 2014;25(6): 841–50.
- 5 Bhat S, Acharya UR, Adeli H, Bairy GM, Adeli A. Automated diagnosis of autism: in search of a mathematical marker. *Rev Neurosci.* 2014;25(6):851–61.
- 6 Sridhar C, Bhat S, Acharya UR, Adeli H, Bairy GM. Diagnosis of attention deficit hyperactivity disorder using imaging and signal processing techniques. *Comput Biol Med.* 2017 Sep;88:93–9.

Disclosure Statement

No conflicts of interest to declare.

Funding Sources

This work has not been supported by any funding agency.

Author Contributions

U.R.A. and H.A. conceived the idea. U.R. and U.R.A. designed the paper. H.A. guided the overall direction and improved the design, coherence, and presentation of the paper.

- 7 Jahmunah V, Lih Oh S, Rajinikanth V, Ciaccio EJ, Hao Cheong K, Arunkumar N, et al. Automated detection of schizophrenia using nonlinear signal processing methods. *Artif Intell Med*. 2019 Sep;100:101698.
- 8 Acharya UR, Oh SL, Hagiwara Y, Tan JH, Adeli H, Subha DP. Automated EEG-based screening of depression using deep convolutional neural network. *Comput Methods Programs Biomed*. 2018 Jul;161:103–13.
- 9 Acharya UR, Oh SL, Hagiwara Y, Tan JH, Adeli H. Deep convolutional neural network for the automated detection and diagnosis of seizure using EEG signals. *Comput Biol Med*. 2018 Sep;100:270–8.
- 10 Ortega-Zamorano F, Jerez JM, Gómez I, Franco L. Layer Multiplexing FPGA Implementation for Deep Back-Propagation Learning. *Integr Comput Aided Eng*. 2017;24(2):171–85.
- 11 Ay B, Yildirim O, Talo M, Baloglu UB, Aydin G, Puthankattil SD, et al. Automated Depression Detection Using Deep Representation and Sequence Learning with EEG Signals. *J Med Syst*. 2019 May;43(7):205.
- 12 Yıldırım Ö, Baloglu UB, Acharya UR. A deep convolutional neural network model for automated identification of abnormal EEG signals. *Neural Comput Appl*. 2018:1–12.
- 13 Khedher L, Illán IA, Górriz JM, Ramírez J, Brahim A, Meyer-Baese A. Independent Component Analysis-Support Vector Machine-Based Computer-Aided Diagnosis System for Alzheimer's with Visual Support. *Int J Neural Syst*. 2017 May;27(3):1650050.
- 14 López-Sanz D, Garcés P, Álvarez B, Delgado-Losada ML, López-Higes R, Maestú F. Network Disruption in the Preclinical Stages of Alzheimer's Disease: From Subjective Cognitive Decline to Mild Cognitive Impairment. *Int J Neural Syst*. 2017 Dec;27(8):1750041.
- 15 Fang C, Li C, Cabrerizo M, Barreto A, Andrian J, Rische N, et al. Gaussian Discriminant Analysis-Based Dual High-Dimensional Decision Spaces for the Diagnosis of Mild Cognitive Impairment in Alzheimer's Disease. *Int J Neural Syst*. 2018;28(8):1850017.
- 16 Valenzuela O, Jiang X, Carrillo A, Rojas I. Multi-Objective Genetic Algorithms to Find Most Relevant Volumes of the Brain Related to Alzheimer's Disease and Mild Cognitive Impairment. *Int J Neural Syst*. 2018 Nov;28(9):1850022.
- 17 Acharya UR, Hagiwara Y, Deshpande SN, Suren S, Wei Koh JE, Lih Oh S, et al. Characterization of focal EEG signals: a review. *Future Gener Comput Syst*. 2019;91:290–9.
- 18 Budka H. Neuropathology of human immunodeficiency virus infection. *Brain Pathol*. 1991 Apr;1(3):163–75.
- 19 Mammone N, Ieracitano C, Adeli H, Bramanti A, Morabito FC. Permutation Jaccard Distance-based Hierarchical Clustering to estimate EEG network density modifications in MCI subjects. *IEEE Trans Neural Netw Learn Syst*. 2018 Feb;29(10):5122–35.
- 20 Acharya UR, Fernandes SJ, WeiKoh JE, Ciaccio EJ, Fabell MK, Tanik U, et al. Automated Detection of Alzheimer's Disease Using Brain MRI Images- A Study with Various Feature Extraction Techniques. *J Med Syst*. 2019 Aug;43(9):302.
- 21 Saunders AM, Strittmatter WJ, Schmechel D, St. George-Hyslop PH, Pericak-Vance MA, Joo SH, et al. Association of apolipoprotein E allele e4 with late-onset familial and sporadic AD. *Neurology*. 1993;43:1467–72.
- 22 Nalbantoglu J, Gilfix BM, Bertrand P, Robitaille Y, Gauthier S, Rosenblatt DS, et al. Predictive value of apolipoprotein E genotyping in Alzheimer's disease: results of an autopsy series and an analysis of several combined studies. *Ann Neurol*. 1994 Dec;36(6):889–95.
- 23 Fisher RS, van Emde Boas W, Blume W, Elger C, Genton P, Lee P, et al. Epileptic seizures and epilepsy. Definitions proposed by the International League against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia*. 2005 Apr;46(4):470–2.
- 24 Guo L, Wang Z, Cabrerizo M, Adjouadi M. A Cross-Related Delay Shift Supervised Learning Method for Spiking Neurons with Application to Interictal Spike Detection in Epilepsy. *Int J Neural Syst*. 2017 May;27(3):1750002.
- 25 Wostyn S, Staljanssens W, De Taeye L, Strobbé G, Gadeyne S, Van Roost D, et al. EEG Derived Brain Activity Reflects Treatment Response from Vagus Nerve Stimulation in Patients with Epilepsy. *Int J Neural Syst*. 2017 Jun;27(4):1650048.
- 26 Martín-López D, Jiménez-Jiménez D, Cabañés-Martínez L, Selway RP, Valentín A, Alarcón G. The role of thalamus versus cortex in epilepsy: evidence from human ictal centro-medial recordings in patients assessed for deep brain stimulation. *Int J Neural Syst*. 2017 Nov;27(7):1750010.
- 27 Kugiumtzis D, Koutlis C, Tsimpiris A, Kimiskidis VK. Dynamics of Epileptiform Discharges Induced by Transcranial Magnetic Stimulation in Genetic Generalized Epilepsy. *Int J Neural Syst*. 2017 Nov;27(7):1750037.
- 28 Varatharajah Y, Iyer RK, Berry BM, Worrell GA, Brinkmann BH. Seizure Forecasting and the Preictal State in Canine Epilepsy. *Int J Neural Syst*. 2017 Feb;27(1):1650046.
- 29 Shanir PP, Khan KA, Khan YU, Farooq O, Adeli H. Automatic Seizure Detection Based on Morphological Features Using One-Dimensional Local Binary Pattern on Long-Term EEG. *Clin EEG Neurosci*. 2018 Sep;49(5):351–62.
- 30 Kotsopoulos IA, van Merode T, Kessels FG, de Krom MC, Knottnerus JA. Systematic review and meta-analysis of incidence studies of epilepsy and unprovoked seizures. *Epilepsia*. 2002 Nov;43(11):1402–9.
- 31 Kotsopoulos I, de Krom M, Kessels F, Lodder J, Troost J, Twellaar M, et al. Incidence of epilepsy and predictive factors of epileptic and non-epileptic seizures. *Seizure*. 2005 Apr;14(3):175–82.
- 32 De Cooman T, Varon C, Hunyadi B, Van Paesschen W, Lagae L, Van Huffel S. Online Automated Seizure Detection in Temporal Lobe Epilepsy Patients Using Single-lead ECG. *Int J Neural Syst*. 2017 Nov;27(7):1750022.
- 33 Li R, Ji GJ, Yu Y, Yu Y, Ding MP, Tang YL, et al. Epileptic discharge related functional connectivity within and between networks in benign epilepsy with centrotemporal spikes. *Int J Neural Syst*. 2017 Nov;27(7):1750018.
- 34 Jiang S, Luo C, Gong J, Peng R, Ma S, Tan S, et al. Aberrant thalamocortical connectivity in juvenile myoclonic epilepsy. *Int J Neural Syst*. 2018 Feb;28(1):1750034.
- 35 Adeli H, Zhou Z, Dadmehr N. Analysis of EEG records in an epileptic patient using wavelet transform. *J Neurosci Methods*. 2003 Feb;123(1):69–87.
- 36 Yuan Q, Zhou W, Xu F, Leng Y, Wei D. Epileptic EEG Identification via LBP Operators on Wavelet Coefficients. *Int J Neural Syst*. 2018 Oct;28(8):1850010.
- 37 Li Y, Cui W, Luo M, Li K, Wang L. Epileptic Seizure Detection Based on Time-Frequency Images of EEG Signals Using Gaussian Mixture Model and Gray Level Co-Occurrence Matrix Features. *Int J Neural Syst*. 2018 Sep;28(7):1850003.
- 38 Yuan S, Zhou W, Chen L. Epileptic Seizure Prediction Using Diffusion Distance and BLDA in Intracranial EEG. *Int J Neural Syst*. 2018;28(1):1750043.
- 39 Schetinin V, Jakaite L, Krzanowski W. Bayesian Learning of Models for Estimating Uncertainty in Alert Systems: Application to Air traffic conflict avoidance. *Integr Comput Aided Eng*. 2018;25(3):1–17.
- 40 Kobelt G, Pugliatti M. Cost of multiple sclerosis in Europe. *Eur J Neurol*. 2005 Jun;12(s1 Suppl 1):63–7.
- 41 Jock Murray T, Allen C. Bowling, Chris Polman, Alan Thompson, John Noseworthy: Multiple sclerosis – The guide to treatment and management. London, 6th Ed. Multiple Sclerosis International Federation. 2006.
- 42 Nutt JG, Wooten GF. Clinical practice. Diagnosis and initial management of Parkinson's disease. *N Engl J Med*. 2005 Sep;353(10):1021–7.
- 43 Bhat S, Acharya UR, Hagiwara Y, Dadmehr N, Adeli H. Parkinson's disease: cause factors, measurable indicators, and early diagnosis. *Comput Biol Med*. 2018 Nov;102:234–41.
- 44 Chaudhuri KR, Yates L, Martinez-Martin P. The non-motor symptom complex of Parkinson's disease: a comprehensive assessment is essential. *Curr Neurol Neurosci Rep*. 2005 Jul;5(4):275–83.
- 45 Marras C, Tanner CM. Epidemiology of Parkinson's disease. In: Watts RL, Koller WC, editors. *Movement disorders, neurologic principles and practice*. 2nd ed. New York: McGraw Hill; 2004. pp. 177–96.
- 46 Shinde S, Prasad S, Saboo Y, Kaushick R, Saini J, Pal PK, et al. Predictive markers for Parkinson's disease using deep neural nets on neuromelanin sensitive MRI. *Neuroimage Clin*. 2019;22:101748.

- 47 Zhang A, San-Segundo R, Panev S, Tabor G, Stebbins K, Whitford A, et al. Automated Tremor Detection in Parkinson's Disease Using Accelerometer Signals. *IEEE/ACM International Conference on Connected Health: Applications, Systems and Engineering Technologies (CHASE)*, Washington, DC, USA, 2018.
- 48 Gálvez G, Recuero M, Canuet L, Del-Pozo F. Short-term effects of Binaural Beats on EEG power, functional connectivity, cognition, gait and anxiety in Parkinson's Disease. *Int J Neural Syst*. 2018 Jun;28(5):1750055.
- 49 Hatano S. Experience from a multicentre stroke register: a preliminary report. *Bull World Health Organ*. 1976;54(5):541–53.
- 50 Kelly DF, Becker DP. Advances in management of neurosurgical trauma: USA and Canada. *World J Surg*. 2001 Sep;25(9):1179–85.
- 51 Fakhry SM, Trask AL, Waller MA, Watts DD; IRTC Neurotrauma Task Force. Management of brain-injured patients by an evidence-based medicine protocol improves outcomes and decreases hospital charges. *J Trauma*. 2004 Mar;56(3):492–9.
- 52 Pain terms: a list with definitions and notes on usage. Recommended by the IASP Subcommittee on Taxonomy. *Pain*. 1979 Jun;6(3):249.
- 53 http://epileptologiebonn.de/cms/front_content.php?idcat=0&idart=0&client=1&lang=3&error=1.
- 54 <http://adni.loni.usc.edu/>.
- 55 <https://www.oasis-brains.org/>.
- 56 Arne Jensen and Anders la Cour-Harbo. Ripples in mathematics: the discrete wavelet transform. Springer Science & Business Media; 2001.
- 57 Abbasi H, Bennet L, Gunn AJ, Unsworth CP. Robust Wavelet Stabilized 'Footprints of Uncertainty' for Fuzzy System Classifiers to Automatically Detect Sharp Waves in the EEG after Hypoxia Ischemia. *Int J Neural Syst*. 2017 May;27(3):1650051.
- 58 Dai H, Cao Z. A wavelet support vector machine-based neural network meta model for structural reliability assessment. *Comput Aided Civ Infrastruct Eng*. 2017;32(4):344–57.
- 59 Candes EJ, Donoho DL. *Curvelets: A surprisingly effective non adaptive representation for objects with edges*. Technical report. DTIC Document; 2000.
- 60 Shao Y, Celenk M. Higher-order spectra (HOS) invariants for shape recognition. *Pattern Recognit*. 2001;34(11):2097–113.
- 61 Chua KC, Chandran V, Acharya UR, Lim CM. Application of higher order statistics/spectra in biomedical signals—a review. *Med Eng Phys*. 2010 Sep;32(7):679–89.
- 62 Pizer SM, Amburn EP, Austin JD, Cromartie R, Geselowitz A, Greer T, et al. Adaptive histogram equalization and its variations. *Comput Vis Graph Image Process*. 1987;39(3):355–68.
- 63 Malladi R, Sethian JA, Vemuri B. Shape modeling with front propagation: A level set approach. *IEEE Trans Pattern Anal Mach Intell*. 1995;17(2):158–75.
- 64 Robert M. Haralick, Karthikeyan Shanmugam, and Its' Hak Dinstein: textural features for image classification. *Systems. IEEE Transactions on Man and Cybernetics*. 1973;6:610–21.
- 65 Renyi A. On measures of entropy and information, in: *Proceedings of the Fourth Berkeley symposium on mathematical statistics and probability*. 1961; 1: 547–561.
- 66 Shannon CE. A mathematical theory of communication. *The Bell System Technical Journal J*. 1948;27(3):379–423.
- 67 Kapur JN. Information of order α and type β . *Proc Indiana Acad Sci*. 1968;68(2):65–75.
- 68 Ghosh M, Chakraborty C, Ray AK. Yager's measure based fuzzy divergence for microscopic color image segmentation. in: *Indian Conference on Medical Informatics and Telemedicine*. Kharagpur, 2013;13–16.
- 69 Chen W, Wang Z, Xie H, Yu W. Characterization of surface EMG signal based on fuzzy entropy. *IEEE Trans Neural Syst Rehabil Eng*. 2007 Jun;15(2):266–72.
- 70 Yin PY. Maximum entropy-based optimal threshold selection using deterministic reinforcement learning with controlled randomization. *Signal Processing*. 2002;82(7):993–1006.
- 71 Sezgin N, Emin Tagluk M. Energy based feature extraction for classification of sleep apnea syndrome. *Comput Biol Med*. 2009 Nov;39(11):1043–50.
- 72 Rosso OA, Blanco S, Yordanova J, Kolev V, Figliola A, Schürmann M, et al. Wavelet entropy: a new tool for analysis of short duration brain electrical signals. *J Neurosci Methods*. 2001 Jan;105(1):65–75.
- 73 Rényi A. On measures of entropy and information. In: *Proceedings of the fourth Berkeley symposium on mathematical statistics and probability*, 1961;547–561.
- 74 Chen J, Li G. Tsallis wavelet entropy and its application in power signal analysis. *Entropy (Basel)*. 2014;16(6):3009–25.
- 75 Hu MK. Visual pattern recognition by moment invariants. *IRE Trans Inf Theory*. 1962; 8(2):179–87.
- 76 Khotanzad A, Hong YH. Invariant image recognition by zernike moments. *IEEE Trans Pattern Anal Mach Intell*. 1990;12(5): 489–97.
- 77 Nixon M, Nixon MS, Aguado AS. *Feature extraction & image processing for computer vision*. Academic Press; 2012.
- 78 Duda RO, Hart PE, Stork DG. *Pattern classification*. 2nd ed. California, USA: Wiley-Interscience; 2000.
- 79 Hyvärinen A, Oja E. Independent component analysis: algorithms and applications. *Neural Netw*. 2000 May-Jun;13(4-5):411–30.
- 80 Scholkopf B, Smola A, Muller KR. Nonlinear component analysis as a kernel eigenvalue problem. *Neural Comput*. 1998;10(5):1299–319.
- 81 Gelman A. Analysis of variance—why it is more important than ever. *Ann Stat*. 2005;33(1):1–33.
- 82 T-test. Student's t-tests. Information. [date accessed on 04.07.17]. Available from: <http://www.physics.csbsju.edu/stats/t-test.html>.
- 83 Dash M, Liu H. Handling large unsupervised data via dimensionality reduction. *ACM SIGMOD Workshop on Research Issues in Data Mining and Knowledge Discovery*. 1999.
- 84 Abe N, Kudo M. Entropy criterion for classifier-independent feature selection. Knowledge-based intelligent information and engineering systems. *Lect Notes Comput Sci*. 2005;3684:689–95.
- 85 Lopes N. Comparing machine learning algorithms with the Wilcoxon Signed Rank Test. Information. [date accessed on 04.07.18]. Available from: <http://www.uc.pt/fctuc/dei/statisticalHypothesis/noel>.
- 86 Hwang T, Sun CH, Yun T, Yi GS. FiGS: a filter-based gene selection workbench for microarray data. *BMC Bioinformatics*. 2010 Jan;11:50.
- 87 Natarajan S, Lipsitz SR, Fitzmaurice GM, Sinha D, Ibrahim JG, Haas J, et al. An extension of the Wilcoxon Rank-Sum test for complex sample survey data. *J R Stat Soc Ser C Appl Stat*. 2012 Aug;61(4):653–64.
- 88 Yuan Y, Van Allen EM, Omberg L, Wagle N, Amin-Mansour A, Sokolov A, et al. Assessing the clinical utility of cancer genomic and proteomic data across tumor types. *Nat Biotechnol*. 2014 Jul;32(7):644–52.
- 89 Kailath T. The divergence and Bhattacharyya distance measures in signal selection. *IEEE Trans Commun Technol*. 1967;15(1): 52–60.
- 90 Obuchowski NA. Receiver operating characteristic curves and their use in radiology. *Radiology*. 2003 Oct;229(1):3–8.
- 91 Mitchell M. *An Introduction to Genetic Algorithms*. USA: MIT Press. Cambridge. 1998.
- 92 Kennedy J, Eberhart R: Particle swarm optimization. *IEEE Int. Conf. Neural Netw*. 1995.
- 93 Shi Y, Eberhart R. A modified particle swarm optimizer. In: *Proceedings of the IEEE World Congress on Computational Intelligence*. Anchorage; 1998.
- 94 Dorigo M, Birattari M, Stutzle T. Ant colony optimization. *IEEE Comput Intell Mag*. 2006; 1(4):28–39.
- 95 Nemati S, Basiri ME, Ghasem-Aghaee N, Aghdam MH. A novel ACO-GA hybrid algorithm for feature selection in protein function prediction. *Expert Syst Appl*. 2009;36(10): 12086–94.
- 96 Peng H, Long F, Ding C. Feature selection based on mutual information: criteria of max-dependency, max-relevance, and min-redundancy. *IEEE Trans Pattern Anal Mach Intell*. 2005 Aug;27(8):1226–38.
- 97 Specht DF. Probabilistic neural networks and the polynomial Adaline as complementary techniques for classification. *IEEE Trans Neural Netw*. 1990;1(1):111–21.
- 98 Kecman V. *Learning and Soft Computing*. Cambridge (MA): MIT Press; 2001.
- 99 Yager RR. An extension of the naive bayesian classifier. *Inf Sci*. 2006;176(5):577–88.

- 100 Heckerman D, Geiger D, Chickering DM. Learning Bayesian networks: the combination of knowledge and statistical data. *Mach Learn*. 1995;20(3):197–243.
- 101 Larose DT. *Discovering Knowledge in Data: An Introduction to Data Mining*. Wiley-Interscience; 2004.
- 102 Cover TM. Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition. *IEEE Trans Electron Comput*. 1965;14(3):326–34.
- 103 Amit Y, Geman D. Shape quantization and recognition with randomized trees. *Neural Comput*. 1997;9(7):1545–88.
- 104 Berge A, Solberg AH. Structured Gaussian components for hyperspectral image classification. *IEEE Trans Geosci Remote Sens*. 2006;44(11):3386–96.
- 105 Hirschauer TJ, Adeli H, Buford JA. Computer-aided diagnosis of Parkinson's disease using an enhanced probabilistic neural network. *J Med Syst*. 2015 Nov;39(11):179.
- 106 Tan JH, Hagiwara Y, Pang W, Lim I, Oh SL, Adam M, et al. Application of stacked convolutional and long short-term memory network for accurate identification of CAD ECG signals. *Comput Biol Med*. 2018 Mar; 94:19–26.
- 107 Yamamoto D, Arimura H, Kakeda S, Magome T, Yamashita Y, Toyofuku F, et al. Computer-aided detection of multiple sclerosis lesions in brain magnetic resonance images: false positive reduction scheme consisted of rule-based, level set method, and support vector machine. *Comput Med Imaging Graph*. 2010 Jul;34(5):404–13.
- 108 Zacharaki EI, Kanterakis S, Bryan RN, Davatzikos C. Measuring brain lesion progression with a supervised tissue classification system. *Proc Int Conf Med Image Comput Comput Assist Interv*. 2008;11:620–27.
- 109 Raghavendra U, Shyamasunder Bhat N, Gudigar A, Acharya UR. Automated system for the detection of thoracolumbar fractures using a CNN architecture. *Future Gener Comput Syst*. 2018;85:184–9.
- 110 Raghavendra U, Fujita H, Bhandary SV, Gudigar A, Tan JH, Acharya UR. Deep convolution neural network for accurate diagnosis of glaucoma using digital fundus images. *Inf Sci*. 2018;441:41–9.
- 111 Yildirim O, Talo M, Ay B, Baloglu UB, Aydin G, Acharya UR. Automated detection of diabetic subject using pre-trained 2D-CNN models with frequency spectrum images extracted from heart rate signals. *Comput Biol Med*. 2019 Aug;113:103387.
- 112 Tan JH, Bhandary SV, Sivaprasad S, Hagiwara Y, Bagchi A, Raghavendra U. Age-related Macular Degeneration detection using deep convolutional neural network. *Future Gener Comput Syst*. 2018;87:127–35.
- 113 Acharya UR, Oh SL, Hagiwara Y, Tan JH, Adam M, Gertych A, et al. A deep convolutional neural network model to classify heartbeats. *Comput Biol Med*. 2017 Oct;89: 389–96.
- 114 Ker J, Wang L, Rao J, Lim T. *Deep Learning Applications in Medical Image Analysis*. *IEEE Access*. 2017;6:9375–89.
- 115 Oh SL, Ng EY, Tan RS, Acharya UR. Automated diagnosis of arrhythmia using combination of CNN and LSTM techniques with variable length heart beats. *Comput Biol Med*. 2018 Nov;102:278–87.
- 116 Yildirim Ö, Plawiak P, Tan RS, Acharya UR. Arrhythmia detection using deep convolutional neural network with long duration ECG signals. *Comput Biol Med*. 2018 Nov; 102:411–20.
- 117 Koziarski M, Cyganek B. Image Recognition with Deep Neural Networks in Presence of Noise – Dealing with and Taking Advantage of Distortions. *Integr Comput Aided Eng*. 2017;24(4):337–49.
- 118 Wang P, Bai X. Regional Parallel Structure Based CNN for Thermal Infrared Face Identification. *Integr Comput Aided Eng*. 2018; 25(3):247–60.
- 119 Chen L, Ye F, Ruan Y, Fan H, Chen Q. An algorithm for highway vehicle detection based on convolutional neural network. *EURASIP J Image Video Process*. 2018; 2018(1):109.
- 120 Zhang A, Wang KC, Li B, Yang E, Dai X, Peng Y, et al. Automated Pixel-Level Pavement Crack Detection on 3D Asphalt Surfaces Using a Deep-Learning Network. *Comput Aided Civ Infrastruct Eng*. 2017; 32(10):805–19.
- 121 Lin YZ, Nie ZH, Ma HW. Structural Damage Detection with Automatic Feature-extraction through Deep Learning. *Comput Aided Civ Infrastruct Eng*. 2017;32(12): 1025–46.
- 122 Gao Y, Mosalam KM. Deep Transfer Learning for Image-based Structural Damage Recognition. *Comput Aided Civ Infrastruct Eng*. 2018;33(9):748–68.
- 123 Molina-Cabello MA, Luque-Baena RM, López-Rubio E, Thurnhofer-Hemsi K. Vehicle Type Detection by Ensembles of Convolutional Neural Networks Operating on Super-resolved Images. *Integr Comput Aided Eng*. 2018;25(4):321–33.
- 124 Nabian MA, Meidani H. Deep Learning for Accelerated Reliability Analysis of Transportation Networks. *Comput Aided Civ Infrastruct Eng*. 2018;33(6):459–80.
- 125 Hashemi H, Abdelghany K. End-to-end deep learning methodology for real-time traffic network management. *Comput Aided Civ Infrastruct Eng*. 2018;33(10):849–63.
- 126 Rafiei MH, Khushefati WH, Demirboga R, Adeli H. Supervised Deep Restricted Boltzmann Machine for Estimation of Concrete Compressive Strength. *ACI Mater J*. 2017;114(2):237–44.
- 127 Torres JF, Galicia A, Troncoso A, Martínez-Álvarez F. A scalable approach based on deep learning for big data time series forecasting. *Integr Comput Aided Eng*. 2018; 25(4):335–48.
- 128 Li W, Li M, Zhou H, Chen G, Jin J, Duan F. A Dual Stimuli Approach Combined with Convolutional Neural Network to Improve Information Transfer Rate of Event-Related Potential-Based Brain-Computer Interface. *Int J Neural Syst*. 2018 Dec;28(10):1850034.
- 129 Cheng D, Liu M. Classification of AD by Cascaded Convolutional Neural Networks Using PET Images. Springer International Publishing AG 2017. Wang Q, et al. (eds). *MLMI 2017*. LNCS 10541. 2017. pp. 106–113.
- 130 Cullen NC, Avants BB. Convolutional Neural Networks for Rapid and Simultaneous Brain Extraction and Tissue Segmentation. *Brain Morphometry*. *Neuroinformatics*. 2018; 136:13–34.
- 131 Islam J, Zhang Y. Brain MRI analysis for Alzheimer's disease diagnosis using an ensemble system of deep convolutional neural networks. *Brain Inform*. 2018 May;5(2):2.
- 132 Hsu WY. A hybrid approach for brain image registration with local constraints. *Integr Comput Aided Eng*. 2017;24(1):73–85.
- 133 Lozano A, Soto-Sánchez C, Garrigós J, Martínez JJ, Ferrández JM, Fernández E. A 3D convolutional neural network to model retinal ganglion cell's responses to light patterns in mice. *Int J Neural Syst*. 2018 Dec; 28(10):1850043.
- 134 Antoniadou A, Spyrou L, Took CC, Sanei S. Deep learning for epileptic intracranial EEG data. Italy: IEEE 26th International Workshop on Machine Learning for Signal Processing (MLSP); 2016.
- 135 Johansen AR, Jin J, Maszczyk T, Dauwels J, Cash SS, Westover MB. Epileptiform spike detection via convolutional neural networks. China: *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*; 2016.
- 136 Yuan Y, Xun G, Jia K, Zhang A. A Multi-view Deep Learning Method for Epileptic Seizure Detection using Short-time Fourier Transform. Proceedings of the 8th ACM International Conference on Bioinformatics, Computational Biology and Health Informatics - ACM-BCB; 2017.
- 137 Ullah I, Hussain M, Qazi EH, Aboalsamh H. An automated system for epilepsy detection using EEG brain signals based on deep learning approach. *Expert Syst Appl*. 2018;107: 61–71.
- 138 Martínez-Murcia FJ, Górriz JM, Ramírez J, Ortiz A. Convolutional Neural Networks for Neuroimaging in Parkinson's Disease: Is Preprocessing Needed? *Int J Neural Syst*. 2018 Dec;28(10):1850035.
- 139 Tjepkema-Cloostermans MC, de Carvalho RC, van Putten MJ. Deep learning for detection of focal epileptiform discharges from scalp EEG recordings. *Clin Neurophysiol*. 2018 Oct;129(10):2191–6.
- 140 Tagaris A, Kollias D, Stafylopatis A. Assessment of Parkinson's Disease Based on Deep Neural Networks. *Commun Comput Inf Sci*. 2017;744:391–403.

- 141 Liu M, Cheng D, Yan W; Alzheimer's Disease Neuroimaging Initiative. Classification of Alzheimer's Disease by Combination of Convolutional and Recurrent Neural Networks Using FDG-PET Images. *Front Neuroinform*. 2018 Jun;12:35.
- 142 Ahmadlou M, Adeli H, Adeli A. New diagnostic EEG markers of the Alzheimer's disease using visibility graph. *J Neural Transm (Vienna)*. 2010 Sep;117(9):1099-109.
- 143 Ahmadlou M, Adeli H, Adeli A. Fractality and a wavelet-chaos-methodology for EEG-based diagnosis of Alzheimer disease. *Alzheimer Dis Assoc Disord*. 2011 Jan-Mar; 25(1):85-92.
- 144 Sankari Z, Adeli H. Probabilistic neural networks for diagnosis of Alzheimer's disease using conventional and wavelet coherence. *J Neurosci Methods*. 2011 Apr;197(1):165-70.
- 145 Sankari Z, Adeli H, Adeli A. Intrahemispheric, interhemispheric, and distal EEG coherence in Alzheimer's disease. *Clin Neurophysiol*. 2011 May;122(5):897-906.
- 146 Sankari Z, Adeli H, Adeli A. Wavelet coherence model for diagnosis of Alzheimer disease. *Clin EEG Neurosci*. 2012 Oct;43(4): 268-78.
- 147 Amezcua-Sanchez JP, Adeli A, Adeli H. A new methodology for automated diagnosis of mild cognitive impairment (MCI) using magnetoencephalography (MEG). *Behav Brain Res*. 2016 May;305:174-80.
- 148 Amezcua-Sanchez JP, Mammone N, Morabito FC, Marino S, Adeli H. A novel methodology for automated differential diagnosis of mild cognitive impairment and the Alzheimer's disease using EEG signals. *J Neurosci Methods*. 2019 Jul;322:88-95.
- 149 Acharya UR, Chua KC, Lim TC, Dorithy JS, Suri JS. Automatic identification of epileptic EEG signals using nonlinear parameters. *J Mech Med Biol*. 2009;9(4):539-53.
- 150 Acharya UR, Sree SV, Suri JS. Automatic detection of epileptic EEG signals using higher order cumulant features. *Int J Neural Syst*. 2011 Oct;21(5):403-14.
- 151 Acharya UR, Sree SV, Chattopadhyay S, Yu W, Ang PC. Application of recurrence quantification analysis for the automated identification of epileptic EEG signals. *Int J Neural Syst*. 2011 Jun;21(3):199-211.
- 152 Acharya UR, Vinitha Sree S, Alvin PC, Yanti R, Suri JS. Application of non-linear and wavelet based features for the automated identification of epileptic EEG signals. *Int J Neural Syst*. 2012;22(2):1250002.
- 153 Acharya UR, Molinari F, Vinitha Sree S, Chattopadhyay S, Kwan-Hoong N, Suri JS. Automated diagnosis of epileptic EEG using entropies. *Biomed Signal Process Control*. 2012;7(4):401-8.
- 154 Acharya UR, Vinitha Sree S, Suri JS. Use of principal component analysis for automatic classification of epileptic EEG activities. *Expert Syst Appl*. 2012;39(10):9072-8.
- 155 Acharya UR, Yanti R, Swapna G, Sree VS, Martis RJ, Suri JS. Automated diagnosis of epileptic electroencephalogram using independent component analysis and discrete wavelet transform for different electroencephalogram durations. *Proc Inst Mech Eng H*. 2013 Mar;227(3):234-44.
- 156 Aslan K, Bozdemir H, Sahin C, Oğulata SN, Erol R. A radial basis function neural network model for classification of epilepsy using EEG signals. *J Med Syst*. 2008 Oct;32(5): 403-8.
- 157 Chua KC, Chandran V, Acharya UR, Lim CM. Application of higher order spectra to identify epileptic EEG. *J Med Syst*. 2011 Dec;35(6):1563-71.
- 158 Faust O, Acharya UR, Min LC, Spath BH. Automatic identification of epileptic and background EEG signals using frequency domain parameters. *Int J Neural Syst*. 2010 Apr;20(2):159-76.
- 159 Ghosh-Dastidar S, Adeli H, Dadmehr N. Mixed-band wavelet-chaos-neural network methodology for epilepsy and epileptic seizure detection. *IEEE Trans Biomed Eng*. 2007 Sep;54(9):1545-51.
- 160 Ghosh-Dastidar S, Adeli H, Dadmehr N. Principal component analysis-enhanced cosine radial basis function neural network for robust epilepsy and seizure detection. *IEEE Trans Biomed Eng*. 2008 Feb;55(2 Pt 1):512-8.
- 161 Ghosh-Dastidar S, Adeli H. A new supervised learning algorithm for multiple spiking neural networks with application in epilepsy and seizure detection. *Neural Netw*. 2009 Dec;22(10):1419-31.
- 162 Guler NF, Ubey ED, Guler I. Recurrent neural network employing Lyapunov exponents for EEG signals classification. *Expert Syst Appl*. 2005;29(3):506-14.
- 163 Guo L, Rivero D, Seoane JA, Pazos A. Classification of EEG signals using relative wavelet energy and artificial neural networks. In: *Conf Proc of the First ACM/SIG-GEVO Summit on Genetic and Evolutionary Computation*. 2009. pp. 177-84.
- 164 Guo L, Rivero D, Pazos A. Epileptic seizure detection using multiwavelet transform based approximate entropy and artificial neural networks. *J Neurosci Methods*. 2010 Oct;193(1):156-63.
- 165 Guo L, Rivero D, Dorado J, Rabuñal JR, Pazos A. Automatic epileptic seizure detection in EEGs based on line length feature and artificial neural networks. *J Neurosci Methods*. 2010 Aug;191(1):101-9.
- 166 Guo L, Rivero D, Dorado J, Munteanu CR, Pazos A. Automatic feature extraction using genetic programming: an application to epileptic EEG classification. *Expert Syst Appl*. 2011;38(8):10425-36.
- 167 Iscan Z, Dokur Z, Demiralp T. Classification of electroencephalogram signals with combined time and frequency features. *Expert Syst Appl*. 2011;38(8):10499-505.
- 168 Kannathal N, Choo ML, Acharya UR, Sadasivan PK. Entropies for detection of epilepsy in EEG. *Comput Methods Programs Biomed*. 2005 Dec;80(3):187-94.
- 169 Lima CA, Coelho AL, Eisenkraft M. Tackling EEG signal classification with least squares support vector machines: a sensitivity analysis study. *Comput Biol Med*. 2010 Aug;40(8):705-14.
- 170 Martis RJ, Acharya UR, Tan JH, Petznick A, Yanti R, Chua CK, et al. Application of empirical mode decomposition (emd) for automated detection of epilepsy using EEG signals. *Int J Neural Syst*. 2012 Dec;22(6): 1250027.
- 171 Nigam VP, Graupe D. A neural-network-based detection of epilepsy. *Neurol Res*. 2004 Jan;26(1):55-60.
- 172 Orhan U, Hekim M, Ozer M. EEG signals classification using the K-means clustering and a multilayer perceptron neural network model. *Expert Syst Appl*. 2011;38(10): 13475-81.
- 173 Polat K, Gunes S. Classification of epileptiform EEG using a hybrid systems based on decision tree classifier and fast Fourier transform. *Appl Math Comput*. 2007;187(2): 1017-26.
- 174 Polat K, Gunes S. Artificial immune recognition system with fuzzy resource allocation mechanism classifier, principal component analysis and FFT method based new hybrid automated identification system for classification of EEG signals. *Expert Syst Appl*. 2008;34(3):2039-48.
- 175 Polat K, Gunes S. A novel data reduction method: distance based data reduction and its application to classification of epileptiform EEG signals. *Appl Math Comput*. 2008;200(1):10-27.
- 176 Sadati N, Mohseni HR, Magshoudi A. Epileptic seizure detection using neural fuzzy networks. In: *Proceedings of the IEEE International Conference on Fuzzy Systems*. 2006. pp. 596-600.
- 177 Srinivasan V, Eswaran C, Sriraam N. Artificial neural network based epileptic detection using time-domain and frequency-domain features. *J Med Syst*. 2005 Dec;29(6):647-60.
- 178 Srinivasan V, Eswaran C, Sriraam N. Approximate entropy-based epileptic EEG detection using artificial neural networks. *IEEE Trans Inf Technol Biomed*. 2007 May; 11(3):288-95.
- 179 Subasi A. EEG Signal classification using wavelet feature extraction and a mixture of expert model. *Expert Syst Appl*. 2007;32(4): 1084-93.
- 180 Subasi A, Gursoy MI. EEG Signal classification using PCA, ICA, LDA and support vector machine. *Expert Syst Appl*. 2010;37(12): 8659-66.
- 181 Tzallas AT, Tsipouras MG, Fotiadis DI. Automatic seizure detection based on time-frequency analysis and artificial neural networks. *Comput Intell Neurosci*. 2007;2007: 80510.

- 182 Jaiswal AK, Banka H. Epileptic seizure detection in EEG signal using machine learning techniques. *Australas Phys Eng Sci Med*. 2018 Mar;41(1):81–94.
- 183 Ubeyli ED. Least squares support vector machine employing model-based methods coefficients for analysis of EEG signals. *Expert Syst Appl*. 2010;37(1):233–9.
- 184 Wang D, Miao D, Xie C. Best basis-based wavelet packet entropy feature extraction and hierarchical EEG classification for epileptic detection. *Expert Syst Appl*. 2011;38(11):14314–20.
- 185 Swami P, Gandhi TK, Panigrahi BK, Tripathi M, Anand S. A novel robust diagnostic model to detect seizures in electroencephalography. *Expert Syst Appl*. 2016;56:116–30.
- 186 Peker M, Sen B, Delen D. A novel method for automated diagnosis of epilepsy using complex-valued classifiers. *IEEE J Biomed Health Inform*. 2016 Jan;20(1):108–18.
- 187 Sharma M, Pachori RB. A novel approach to detect epileptic seizures using a combination of tunable-q wavelet transform and fractal dimension. *J Mech Med Biol*. 2017;17(07):1740003.
- 188 Patidar S, Panigrahi T. Detection of epileptic seizure using Kraskov entropy applied on tunable-q wavelet transform of EEG signals. *Biomed Signal Process Control*. 2017;34:74–80.
- 189 Gandhi T, Panigrahi BK, Anand S. A comparative study of wavelet families for EEG signal classification. *Neurocomputing*. 2011;74(17):3051–7.
- 190 Chen G. Automatic EEG seizure detection using dual-tree complex wavelet-Fourier features. *Expert Syst Appl*. 2014;41(5):2391–4.
- 191 Swami P, Godiyal AK, Santhosh J, Panigrahi BK, Bhatia M, Anand S. Robust expert system design for automated detection of epileptic seizures using SVM classifier. In: *Proceedings of IEEE International Conference on Parallel, Distributed and Grid Computing*; 2014. pp. 219–22.
- 192 Pachori RB, Patidar S. Epileptic seizure classification in EEG signals using second-order difference plot of intrinsic mode functions. *Comput Methods Programs Biomed*. 2014 Feb;113(2):494–502.
- 193 Sharma R, Pachori RB. Classification of epileptic seizures in EEG signals based on phase space representation of intrinsic mode functions. *Expert Syst Appl*. 2015;42(3):1106–17.
- 194 Bhattacharyya A, Pachori RB, Upadhyay A, Acharya UR. Tunable-q wavelet transform based multiscale entropy measure for automated classification of epileptic EEG signals. *Appl Sci (Basel)*. 2017;7(4):385.
- 195 Bhattacharyya A, Sharma M, Pachori RB, Sircar P, Acharya UR. A novel approach for automated detection of focal EEG signals using empirical wavelet transform. *Neural Comput Appl*. 2018;29(8):47–57.
- 196 Bhati D, Sharma M, Pachori RB, Gadre VM. Time-frequency localized threeband biorthogonal wavelet filter bank using semidefinite relaxation and nonlinear least squares with epileptic seizure EEG signal classification. *Digit Signal Process*. 2017;62:259–73.
- 197 Sharma M, Dhere A, Pachori RB, Acharya UR. An automatic detection of focal EEG signals using new class of time–frequency localized orthogonal wavelet filter banks. *Knowl Base Syst*. 2017;118:217–27.
- 198 Kaya Y, Uyar M, Tekin R, Yildirim S. 1D-local binary pattern based feature extraction for classification of epileptic EEG signals. *Appl Math Comput*. 2014;243:209–19.
- 199 Zhu G, Li Y, Wen PP. Epileptic seizure detection in EEGs signals using a fast weighted horizontal visibility algorithm. *Comput Methods Programs Biomed*. 2014 Jul;115(2):64–75.
- 200 Samiee K, Kovács P, Gabbouj M. Epileptic seizure classification of EEG time-series using rational discrete short-time fourier transform. *IEEE Trans Biomed Eng*. 2015 Feb;62(2):541–52.
- 201 Riaz F, Hassan A, Rehman S, Niazi IK, Dremstrup K. Emd-based temporal and spectral features for the classification of EEG signals using supervised learning. *IEEE Trans Neural Syst Rehabil Eng*. 2016 Jan;24(1):28–35.
- 202 Diykh M, Li Y, Wen P. Classify epileptic eeg signals using weighted complex networks based community structure detection. *Expert Syst Appl*. 2017;90:87–100.
- 203 Li M, Chen W, Zhang T. Application of MODWT and log-normal distribution model for automatic epilepsy identification. *Biocybern Biomed Eng*. 2017;37(4):679–89.
- 204 Oh SL, Vicnesh J, Edward JC, Yuvaraj R, Acharya UR. Deep Convolutional Neural Network Model for Automated Diagnosis of Schizophrenia Using EEG Signals. *Appl Sci (Basel)*. 2019;9(14):2870.
- 205 Al Ghayab HR, Li Y, Abdulla S, Diykh M, Wan X. Classification of epileptic EEG signals based on simple random sampling and sequential feature selection. *Brain Inform*. 2016 Jun;3(2):85–91.
- 206 Sharma M, Bhuraneb AA, Acharya UR. MMSFL-OWFB: A novel class of orthogonal wavelet filters for epileptic seizure detection. *Knowl Base Syst*. 2018;160:265–77.
- 207 Tiwari AK, Pachori RB, Kanhangad V, Panigrahi BK, Panigrahi B. Automated diagnosis of epilepsy using key-point based local binary pattern of EEG signals. *IEEE J Biomed Health Inform*. 2017 Jul;21(4):888–96.
- 208 Bajaj V, Pachori RB. Classification of seizure and non-seizure EEG signals using empirical mode decomposition. *IEEE Trans Inf Technol Biomed*. 2012 Nov;16(6):1135–42.
- 209 Nicolau N, Georgiou J. Detection of epileptic electroencephalogram based on Permutation Entropy and Support Vector Machines. *Expert Syst Appl*. 2012;39(1):202–9.
- 210 Xie S, Krishnan S. Wavelet-based sparse functional linear model with applications to EEGs seizure detection and epilepsy diagnosis. *Med Biol Eng Comput*. 2013 Feb;51(1-2):49–60.
- 211 Mursalin M, Zhang Y, Chen Y, Chawla NV. Automated epileptic seizure detection using improved correlation-based feature selection with random forest classifier. *Neurocomputing*. 2017;241:204–14.
- 212 Upadhyay R, Padhy P, Kankar P. A comparative study of feature ranking techniques for epileptic seizure detection using wavelet transform. *Comput Electr Eng*. 2016;53:163–76.
- 213 Kabir E, Siuly, Zhang Y. Epileptic seizure detection from EEG signals using logistic model trees. *Brain Inform*. 2016 Jun;3(2):93–100.
- 214 Murugavel AS, Ramakrishnan S. Hierarchical multi-class SVM with ELM kernel for epileptic EEG signal classification. *Med Biol Eng Comput*. 2016 Jan;54(1):149–61.
- 215 Pippa E, Zacharaki EI, Mporas I, Tsirka V, Richardson MP, Koutroumanidis M, et al. Improving classification of epileptic and non-epileptic EEG events by feature selection. *Neurocomputing*. 2016;171:576–85.
- 216 Kumar Y, Dewal M, Anand R. Epileptic seizures detection in EEG using DWT-based ApEn and artificial neural network. *Signal Image Video Process*. 2014;8(7):1323–34.
- 217 Naser A, Tantawi M, Shedeed H, Tolba M. Detecting Epileptic Seizures Using Abe Entropy, Line Length and SVM Classifier. *International Conference on Advanced Machine Learning Technologies and Applications*. 2019; 169–178.
- 218 Tzimourta K, Tzallas A, Giannakeas N, Astrakas L, Angelidis L, Tsipouras D, et al. A robust methodology for classification of epileptic seizures in EEG signals. *Health Technol*. 2019;9(2):135–42.
- 219 Lahmiri S, Shmuel A. Accurate Classification of Seizure and Seizure-Free Intervals of Intracranial EEG Signals From Epileptic Patients. *IEEE Trans Instrum Meas*. 2019;68(3):791–6.
- 220 Raghu S, Srimaam N, Temel Y, Rao SV, Hegde AS, Kubben PL. Performance evaluation of DWT based sigmoid entropy in time and frequency domains for automated detection of epileptic seizures using SVM classifier. *Comput Biol Med*. 2019 Jul;110:127–43.
- 221 Wang X, Gong G, Li N. Automated recognition of epileptic EEG states using a combination of symlet wavelet processing, gradient boosting machine, and grid search optimizer. *Sensors*. 2019;19(2):219.
- 222 Bose R, Pratiher S, Chatterjee S. Detection of epileptic seizure employing a novel set of features extracted from multifractal spectrum of electroencephalogram signals. *IET Signal Process*. 2019;13(2):157–64.
- 223 Dalal M, Tanveer M, Pachori RB. *Machine Intelligence and Signal Analysis*. Springer Singapore; 2019.

- 224 Sriraam N, Tamanna K, Narayan L, Khanum M, Raghu S, Hegde AS, et al. Multi-channel EEG based inter-ictal seizures detection using Teager energy with backpropagation neural network classifier. *Australas Phys Eng Sci Med*. 2018 Dec;41(4):1047–55.
- 225 Shaikh M, Farooq O, Chandel G. Lecture Notes in Electrical Engineering 509 Advances in System Optimization and Control; 2017.
- 226 Osman AH, Alzahrani AA. New Approach for Automated Epileptic Disease Diagnosis Using an Integrated Self-Organization Map and Radial Basis Function Neural Network Algorithm. *IEEE Access*. 2019;7(7):4741–7.
- 227 Sudalaimani C, Sivakumaran N, Elizabeth TT, Rominus VS. Automated detection of the preseizure state in EEG signal using neural networks. *Biocybern Biomed Eng*. 2019; 39(1):160–75.
- 228 Raghu S, Sriraam N. Classification of focal and non-focal EEG signals using neighborhood component analysis and machine learning algorithms. *Expert Syst Appl*. 2018; 113:18–32.
- 229 De Cooman T, Varon C, Van de Vel A, Jansen K, Ceulemans B, Lagae L, et al. Adaptive nocturnal seizure detection using heart rate and low-complexity novelty detection. *Seizure*. 2018 Jul;59:48–53.
- 230 Li M, Chen W, Zhang T. A novel seizure diagnostic model based on kernel density estimation and least squares support vector machine. *Biomed Signal Process Control*. 2018;41:233–41.
- 231 Cruz NE, Solarte J, Varghas A. *Automated Epileptic Seizure Detection System Based on a Wearable Prototype and Cloud Computing to Assist People with Epilepsy*. *Applied Computer Sciences in Engineering*. Springer; 2018. pp. 204–13.
- 232 Kocadagli O, Langari R. Classification of EEG signals for epileptic seizures using hybrid artificial neural networks based wavelet transforms and fuzzy relations. *Expert Syst Appl*. 2017;88:419–34.
- 233 Zhang T, Chen W, Li M. Fuzzy distribution entropy and its application in automated seizure detection technique. *Biomed Signal Process Control*. 2018;39:360–77.
- 234 Feng B, Zhao J, Fu W. Automated Classification of Epileptic EEG Signals Based on Multi-Feature Extraction. Beijing: IEEE 9th International Conference on Software Engineering and Service Science (ICSESS); 2018. pp. 382–6.
- 235 Tanveer M, Pachori R, Angami N. Entropy based features in FAWT framework for automated detection of epileptic seizure EEG signals. *IEEE Symp Ser Comput Intell*; 2018. pp. 1946–52.
- 236 Choudhury NR, Roy SS, Pal A, Chatterjee S, Bose R. Epileptic Seizure Detection Employing Cross-Hyperbolic Stockwell Transform. Kolkata: *Fourth International Conference on Research in Computational Intelligence and Communication Networks (ICRCICN)*; 2018. pp. 70–4.
- 237 Torse D, Desai V, Khanai R. Classification of EEG Signals in Seizure Detection System Using Ellipse Area Features and Support Vector Machine. Proceedings of the 2nd International Conference on Data Engineering and Communication Technology ICDECT. 2017.
- 238 Tomanik G, Betting L, Luiz A. Campanharo: Automatic Identification of Interictal Epileptiform Discharges with the Use of Complex Networks. Gran Canaria: Advances in Computational Intelligence, Proceedings of 15th International Work-Conference on Artificial Neural Networks, IWANN 2019; 2019. pp. 152–61.
- 239 Wani S, Sabut S, Nalbalwar S. Detection of Epileptic Seizure Using Wavelet Transform and Neural Network Classifier. *Proceedings of ICCASP*; 2018.
- 240 Qi Y, Wang Y, Zhang J, Zhu J, Zheng X. Robust deep network with maximum correntropy criterion for seizure detection. *BioMed Res Int*. 2014;2014:703816.
- 241 Lin Q, Ye S, Huang X, Li S, Zhang M, Xue Y, et al. Classification of Epileptic EEG Signals with Stacked Sparse Autoencoder Based on Deep Learning. In: Huang DS, Han K, Hussain A (eds). *Intelligent Computing Methodologies*. ICIC 2016. Lecture Notes in Computer Science, Springer, Cham; 2016. vol 9773, pp. 802–10.
- 242 Gogna A, Majumdar A, Ward R. Semi-supervised Stacked Label Consistent Autoencoder for Reconstruction and Analysis of Biomedical Signals. *IEEE Trans Biomed Eng*. 2017 Sep;64(9):2196–205.
- 243 Hussein R, Palangi H, Ward RK, Wang ZJ. Optimized deep neural network architecture for robust detection of epileptic seizures using EEG signals. *Clin Neurophysiol*. 2019 Jan;130(1):25–37.
- 244 Thodoroff P, Pineau J, Lim A. Learning robust features using deep learning for automatic seizure detection. In *Proceedings of MLHC*, 2016. pp. 178–90.
- 245 Emami A, Kunii N, Matsuo T, Shinozaki T, Kawai K, Takahashi H. Seizure detection by convolutional neural network-based analysis of scalp electroencephalography plot images. *Neuroimage Clin*. 2019;22:101684.
- 246 Jang HJ, Cho KO. Dual deep neural network-based classifiers to detect experimental seizures. *Korean J Physiol Pharmacol*. 2019 Mar;23(2):131–9.
- 247 Zuo R, Wei J, Li X, Li C, Zhao C, Ren Z, et al. Automated Detection of High-Frequency Oscillations in Epilepsy Based on a Convolutional Neural Network. *Front Comput Neurosci*. 2019 Feb;13:6.
- 248 Wei X, Zhou L, Chen Z, Zhang L, Zhou Y. Automatic seizure detection using three-dimensional CNN based on multi-channel EEG. *BMC Med Inform Decis Mak*. 2018 Dec;18(5 Suppl 5):111.
- 249 Achilles F, Tombari F, Belagiannis V, Loesch A, Noachtar S, Navab N. Convolutional neural networks for real-time epileptic seizure detection. *Comput Methods Biomech Biomed Eng Imaging Vis*. 2016;1163: 264–9.
- 250 Yuvaraj R, Thomas J, Kluge T, Dauwels J. A deep Learning Scheme for Automatic Seizure Detection from Long-Term Scalp EEG. 52nd Asilomar Conf. Signals, Syst. Comput; 2018. pp. 368–72.
- 251 Hüggle M, Heller S, Watter M, Blum M, Manzouri F, Dümpelmann M, et al. Early Seizure Detection with an Energy-Efficient Convolutional Neural Network on an Implantable Microcontroller. *IEEE*; 2018.
- 252 Thomas J, Comoretto L, Jin J, Dauwels J, Cash S, Westover M. EEG Classification Via Convolutional Neural Network-Based Interictal Epileptiform Event Detection. Proceedings of Annual International Conference of the IEEE Engineering in Medicine and Biology Society. 2018. pp. 3148–51.
- 253 Yuvaraj R, Murugappan M, Ibrahim NM, Sundaraj K, Omar MI, et al. Detection of emotions in Parkinson's disease using higher order spectral features from brain's electrical activity. *Biomed. Signal Process Contr*. 2014;14: 108–16.
- 254 Yuvaraj R, Murugappan M, Ibrahim NM, Sundaraj K, Omar MI, Mohamad K, et al. Optimal set of EEG features for emotional state classification and trajectory visualization in Parkinson's disease. *Int J Psychophysiol*. 2014 Dec;94(3):482–95.
- 255 Yuvaraj R, Murugappan M, Acharya UR, Adeli H, Ibrahim NM, Mesquita E. Brain functional connectivity patterns for emotional state classification in Parkinson's disease patients without dementia. *Behav Brain Res*. 2016 Feb;298 Pt B:248–60.
- 256 Nilashi M, Ibrahim O, Ahmadi H, Shahmoradi L. An analytical method for diseases prediction using machine learning techniques. *Comput Chem Eng*. 2017;106:212–23.
- 257 Tucker CS, Behoora I, Nembhard HB, Lewis M, Sterling NW, Huang X. Machine learning classification of medication adherence in patients with movement disorders using non-wearable sensors. *Comput Biol Med*. 2015 Nov;66:120–34.
- 258 Prashantha R., Sumantra Dutta Roy: Early Detection of Parkinson's Disease through Patient Questionnaire and Predictive Modelling. *Int J Med Inform*. 2018;119:75–87.
- 259 Ali H. Al-Fatlawi, Mohammed H. Jabardi, Sai Ho Ling: Efficient Diagnosis System for Parkinson's Disease Using Deep Belief Network. Canada: In 2016 IEEE Congress on Evolutionary Computation (CEC); 2016.
- 260 Caliskan A, Badem H, Baştürk A, Yüksel ME. Diagnosis of The Parkinson Disease By Using Deep Neural Network Classifier. *In Iu-JEEE*. 2017;17(2):3311–8.
- 261 Grover S, Bhartia S, Akshama AY, Seeja KR. Predicting Severity Of Parkinson's Disease Using Deep Learning. *Procedia Comput Sci*. 2018;132:1788–94.

- 262 Oliveira FP, Castelo-Branco M. Computer-aided diagnosis of Parkinson's disease based on [(123)I]FP-CIT SPECT binding potential images, using the voxels-as-features approach and support vector machines. *J Neural Eng*. 2015 Apr;12(2):026008.
- 263 Banerjee M, Okun MS, Vaillancourt DE, Vemuri BC. A method for automated classification of Parkinson's disease diagnosis using an ensemble average propagator template brain map estimated from diffusion MRI. *PLoS One*. 2016 Jun;11(6):e0155764.
- 264 Cigdem O, Beheshti I, Demirel H. Effects of different covariates and contrasts on classification of Parkinson's disease using structural MRI. *Comput Biol Med*. 2018 Aug;99:173–81.
- 265 Segovia F, Illán IA, Górriz JM, Ramírez J, Rominger A, Levin J. Distinguishing Parkinson's disease from atypical parkinsonian syndromes using PET data and a computer system based on support vector machines and Bayesian networks. *Front Comput Neurosci*. 2015 Nov;9(137):137.
- 266 Ahmaddou M, Adeli H. Enhanced probabilistic neural network with local decision circles: a robust classifier. *Integr Comput Aided Eng*. 2010;17(3):197–210.
- 267 Choi H, Ha S, Im HJ, Paek SH, Lee DS. Refining diagnosis of Parkinson's disease with deep learning-based interpretation of dopamine transporter imaging. *Neuroimage Clin*. 2017 Sep;16:586–94.
- 268 Sivaranjini S, Sujatha CM. Deep learning based diagnosis of Parkinson's disease using convolutional neural network. *Multimed Tools Appl*. 2019:1–13.
- 269 Han CX, Wang J, Yi GS, Che YQ. Investigation of EEG abnormalities in the early stage of Parkinson's disease. *Cogn Neurodyn*. 2013 Aug;7(4):351–9.
- 270 Yuvaraj R, Acharya UR, Hagiwara Y. A novel Parkinson's disease diagnosis index using higher-order spectra features in EEG signals. *Neural Comput Appl*. 2018;30(4):1225–35.
- 271 Hariharan M, Polat K, Sindhu R. A new hybrid intelligent system for accurate detection of Parkinson's disease. *Comput Methods Programs Biomed*. 2014 Mar;113(3):904–13.
- 272 Zhang YN. Can a Smartphone Diagnose Parkinson Disease? A Deep Neural Network Method and Telediagnosis System Implementation. *Parkinsons Dis*. 2017; 2017:6209703.
- 273 Hlavnička J, Čmejla R, Tykalová T, Šonka K, Růžička E, Ruzs J. Automated analysis of connected speech reveals early biomarkers of Parkinson's disease in patients with rapid eye movement sleep behaviour disorder. *Sci Rep*. 2017 Feb;7(1):12.
- 274 Joshi D, Khajuria A, Joshi P. An automatic non-invasive method for Parkinson's disease classification. *Comput Methods Programs Biomed*. 2017 Jul;145:135–45.
- 275 Ornelas-Vences C, Sanchez-Fernandez LP, Sanchez-Perez LA, Garza-Rodriguez A, Villegas-Bastida A. Fuzzy inference model evaluating turn for Parkinson's disease patients. *Comput Biol Med*. 2017 Oct;89:379–88.
- 276 Samà A, Pérez-López C, Rodríguez-Martín D, Català A, Moreno-Aróstegui JM, Cabestany J, et al. Estimating bradykinesia severity in Parkinson's disease by analysing gait through a waist-worn sensor. *Comput Biol Med*. 2017 May;84:114–23.
- 277 Oung QW, Muthusamy H, Basah SN, Lee H, Vijean V. Empirical wavelet transform based features for classification of Parkinson's disease severity. *J Med Syst*. 2017 Dec; 42(2):29.
- 278 Little MA, McSharry PE, Hunter EJ, Spielman J, Ramig LO. Suitability of dysphonia measurements for telemonitoring of Parkinson's disease. *IEEE Trans Biomed Eng*. 2009 Apr;56(4):1015–22.
- 279 Shahbaba B, Neal R. Nonlinear models using Dirichlet process mixtures. *J Mach Learn Res*. 2009;10:1829–50.
- 280 Das R. A comparison of multiple classification methods for diagnosis of Parkinson disease. *Expert Syst Appl*. 2010;37(2):1568–72.
- 281 Sakar CO, Kursun O. Telediagnosis of Parkinson's disease using measurements of dysphonia. *J Med Syst*. 2010 Aug;34(4):591–9.
- 282 Psorakis I, Damoulas T, Girolami MA. Multiclass relevance vector machines: sparsity and accuracy. *IEEE Trans Neural Netw*. 2010 Oct;21(10):1588–98.
- 283 Guo PF, Bhattacharya P, Kharma N. Advances in detecting Parkinson's disease. In: Zhang D, Sonka M, (eds). Berlin: Medical biometrics, Lecture Notes in Computer Science; 2010. pp. 306–14.
- 284 Ozcift A, Gulden A. Classifier ensemble construction with rotation forest to improve medical diagnosis performance of machine learning algorithms. *Comput Methods Programs Biomed*. 2011 Dec;104(3):443–51.
- 285 Li DC, Liu CW, Hu SC. A fuzzy-based data transformation for feature extraction to increase classification performance with small medical data sets. *Artif Intell Med*. 2011 May;52(1):45–52.
- 286 Luukka P. Feature selection using fuzzy entropy measures with similarity classifier. *Expert Syst Appl*. 2011;38(4):4600–7.
- 287 Spadoto AA, Guido RC, Carnevali FL, Pagnin AF, Falcao AX, Papa JP. Improving Parkinson's disease identification through evolutionary-based feature selection. In: *Proceedings of the annual international conference of the IEEE engineering in medicine and biology society (EMBC-11)*, Boston; 2011. pp. 7857–60.
- 288 Astrom F, Koker R. A parallel neural network approach to prediction of Parkinson's Disease. *Expert Syst Appl*. 2011;38(10):12470–4.
- 289 Ozcift A. SVM feature selection based rotation forest ensemble classifiers to improve computer-aided diagnosis of Parkinson disease. *J Med Syst*. 2012 Aug;36(4):2141–7.
- 290 Polat K. Classification of Parkinson's disease using feature weighting method on the basis of fuzzy C-means clustering. *Int J Syst Sci*. 2012;43(4):597–609.
- 291 Tsanas A, Little MA, McSharry PE, Spielman J, Ramig LO. Novel speech signal processing algorithms for high-accuracy classification of Parkinson's disease. *IEEE Trans Biomed Eng*. 2012 May;59(5):1264–71.
- 292 Daliri MR. Chi square distance kernel of the gaits for the diagnosis of Parkinson's disease. *Biomed Signal Process Control*. 2013; 8(1):66–70.
- 293 Chen HL, Huang CC, Yu XG, Xuc X, Sund X, Wang G, et al. An efficient diagnosis system for detection of Parkinson's disease using fuzzy k-nearest neighbor approach. *Expert Syst Appl*. 2013;40(1):263–71.
- 294 Zuo WL, Wang ZY, Liu T, Chen HL. Effective detection of Parkinson's disease using an adaptive fuzzy K-nearest neighbour approach. *Biomed Signal Process Control*. 2013;8(4):364–73.
- 295 Ma C, Ouyang J, Chen HL, Zhao XH. An efficient diagnosis system for Parkinson's disease using kernel-based extreme learning machine with subtractive clustering features weighting approach. *Comput Math Methods Med*. 2014;2014:985789.
- 296 Drotár P, Mekyska J, Rektorová I, Masarová L, Smékal Z, Faundez-Zanuy M. Evaluation of handwriting kinematics and pressure for differential diagnosis of Parkinson's disease. *Artif Intell Med*. 2016 Feb;67:39–46.
- 297 Connolly AT, Kaemmerer WF, Dani S, Stanslaski SR, Panken E, Johnson MD, et al. Guiding deep brain stimulation contact selection using local field potentials sensed by a chronically implanted device in Parkinson's disease patients. Montpellier: 7th international conference on neural engineering; 2015. pp. 840–3.
- 298 Wahid F, Begg RK, Hass CJ, Halgamuge S, Ackland DC. Classification of Parkinson's disease gait using spatial-temporal gait features. *IEEE J Biomed Health Inform*. 2015 Nov;19(6):1794–802.
- 299 Smith SL, Lones MA, Bedder M, Alty JE, Cosgrove J, Maguire RJ, et al. Computational approaches for understanding the diagnosis and treatment of Parkinson's disease. *IET Syst Biol*. 2015 Dec;9(6):226–33.
- 300 Shamir RR, Dolber T, Noecker AM, Walter BL, McIntyre CC. Machine learning approach to optimizing combined stimulation and medication therapies for Parkinson's disease. *Brain Stimul*. 2015 Nov-Dec;8(6):1025–32.
- 301 Procházka A, Vysata O, Valis M, Tupa O, Schätz M, Marik V. Bayesian classification and analysis of gait disorders using image and depth sensors of Microsoft Kinect. *Digit Signal Process*. 2015;47:169–77.

- 302 Nilashi M, Ibrahim O, Ahmadi H, Shahmoradi L, Farahmand M. A hybrid intelligent system for the prediction of Parkinson's Disease progression using machine learning techniques. *Biocybern Biomed Eng*. 2018; 38(1):1–15.
- 303 Kim HB, Lee WW, Kim A, Lee HJ, Park HY, Jeon HS, et al. Wrist sensor-based tremor severity quantification in Parkinson's disease using convolutional neural network. *Comput Biol Med*. 2018 Apr;95:140–6.
- 304 Oh SL, Hagiwara Y, Raghavendra U, Yuvraj R, Arunkumar N, Murugappan M, et al. A Deep Learning Approach for Parkinson's Disease Diagnosis from EEG Signals. *Neural Comput Appl*. 2018:1–7.
- 305 Al-nuaimi AH, Jammeh E, Sun L, Ifeakor E. Tsallis Entropy as a Biomarker for Detection of AD. *Conf Proc IEEE Eng Med Biol Soc*. 2015;2015:4166–9.
- 306 Silveira M, Marques J. Boosting Alzheimer Disease Diagnosis using PET images. Turkey: International Conference on Pattern Recognition; 2010.
- 307 Mahanand BS, Suresh S, Sundararajan N, Aswatha Kumar M. Alzheimer's disease detection using a Self-adaptive Resource Allocation Network classifier. San Jose: The 2011 International Joint Conference on Neural Networks; 2011. pp. 1930–4.
- 308 Mahmood R, Ghimire B. Automatic detection and classification of Alzheimer's Disease from MRI scans using principal component analysis and artificial neural networks. Bucharest: 20th International Conference on Systems, Signals and Image Processing (IWSSIP); 2013. pp. 133–7.
- 309 Ding Y, Cong Zhang, Tian Lan, Zhiguang Qin, Xinjie Zhang and Wei Wang: Classification of Alzheimer's disease based on the combination of morphometric feature and texture feature. Washington: IEEE International Conference on Bioinformatics and Biomedicine; 2015. pp. 409–412.
- 310 Herrera LJ, Rojas I, Pomares H, Guillén A, Valenzuela O, Baños O. Classification of MRI Images for Alzheimer's Disease Detection. Alexandria: *International Conference on Social Computing*; 2013. pp. 846–51.
- 311 Sweet ME, Jiji GW. Detection of Alzheimer disease in brain images using PSO and Decision Tree Approach. Ramanathapuram: *IEEE International Conference on Advanced Communications, Control and Computing Technologies*; 2014. pp. 1305–9.
- 312 Saraswathi S, Mahanand BS, Kloczkowski A, Suresh S, Sundararajan N. Detection of onset of Alzheimer's disease from MRI images using a GA-ELM-PSO classifier. Singapore: *Fourth International Workshop on Computational Intelligence in Medical Imaging*; 2013. pp. 42–48.
- 313 Mahanand BS, Suresh S, Sundararajan N, Kumar MA. ICGA-ELM classifier for Alzheimer's disease detection. Kharagpur: *Indian Conference on Medical Informatics and Telemedicine (ICMIT)*; 2013. pp. 48–52.
- 314 Mahanand BS, Babu GS, Suresh S, Sundararajan N. Identification of imaging biomarkers responsible for Alzheimer's Disease using a McRBFN classifier. *International Conference on Cognitive Computing and Information Processing*; 2015.
- 315 Mathew J, Mekkayil L, Ramasangu H, Karthikeyan BR, Manjunath AG. Robust algorithm for early detection of Alzheimer's disease using multiple feature extractions. Bangalore: *IEEE Annual India Conference (INDICON)*; 2016. pp. 1–6.
- 316 Escudero J, Ifeakor E, Zajicek JP, Green C, Shearer J, Pearson S; Alzheimer's Disease Neuroimaging Initiative. Machine learning-based method for personalized and cost-effective detection of Alzheimer's disease. *IEEE Trans Biomed Eng*. 2013 Jan; 60(1):164–8.
- 317 Gunawardena KA, Rajapakse RN, Kodikara ND, Mudalige IU. Moving from detection to pre-detection of Alzheimer's Disease from MRI data. Negombo: Sixteenth International Conference on Advances in ICT for Emerging Regions (ICTer); 2016. p. 324.
- 318 Zhang J, Liu M, Le An Y, Gao Y, Shen D. Alzheimer's Disease Diagnosis Using Landmark-Based Features From Longitudinal Structural MR Images. *IEEE J Biomed Health Inform*. 2017 Nov;21(6):1607–16.
- 319 Escudero J, Zajicek JP, Ifeakor E. Early detection and characterization of Alzheimer's disease in clinical scenarios using Bioprofile concepts and K-means. *Conf Proc IEEE Eng Med Biol Soc*. 2011;2011:6470–3.
- 320 Zhang J, Gao Y, Gao Y, Munsell BC, Shen D. Detecting Anatomical Landmarks for Fast Alzheimer's Disease Diagnosis. *IEEE Trans Med Imaging*. 2016 Dec;35(12):2524–33.
- 321 Iftikhar MA, Idris A. An ensemble classification approach for automated diagnosis of Alzheimer's disease and mild cognitive impairment. International Conference on Open Source Systems & Technologies (ICOSST); 2016. pp. 78–83.
- 322 Bates J, Pafundi D, Kanel P, Liu X, Mio W. Spectral signatures of point clouds and applications to detection of Alzheimer's Disease through Neuroimaging. Chicago: IEEE International Symposium on Biomedical Imaging: From Nano to Macro; 2011. pp. 1851–4.
- 323 Ye DH, Pohl KM, Davatzikos C. Semi-supervised Pattern Classification: Application to Structural MRI of Alzheimer's Disease. Seoul: International Workshop on Pattern Recognition in Neuroimaging; 2011.
- 324 Rabeh AB, Benzarti F, Amiri H. New Method of Classification to Detect Alzheimer Disease. 14th International Conference on Computer Graphics, Imaging and Visualization; 2017.
- 325 Ullah HM, Onik Z, Islam R, Nandi D. Alzheimer's Disease and Dementia Detection from 3D Brain MRI Data Using Deep Convolutional Neural Networks. Pune: 3rd International Conference for Convergence in Technology (I2CT); 2018.
- 326 Donini M, Monteiro JM, Pontil M, Shawe-Taylor J, Mourao-Miranda J. A multimodal multiple kernel learning approach to Alzheimer's disease detection. Vietri sul Mare: IEEE 26th International Workshop on Machine Learning for Signal Processing (MLSP); 2016.
- 327 Sarraf S, Tofighi G, Deep AD. AD Classification via Deep Convolutional Neural Networks using MRI and fMRI. *bioRxiv*. 2016.
- 328 Liu M, Cheng D, Wang K, Wang Y; Alzheimer's Disease Neuroimaging Initiative. Multi-Modality Cascaded Convolutional Neural Networks for AD Diagnosis. *Neuroinformatics*. 2018 Oct;16(3-4):295–308.
- 329 McCrackin L, Bagheri E, Cheung JCK. Early Detection of AD Using Deep Learning. in Springer International Publishing AG, part of Springer Nature 2018 Canadian AI 2018; 2018. pp. 355–359.
- 330 Basaia S, Agosta F, Wagner L, Canu E, Magnani G, Santangelo R, et al.; Alzheimer's Disease Neuroimaging Initiative. Automated classification of Alzheimer's disease and mild cognitive impairment using a single MRI and deep neural networks. *Neuroimage Clin*. 2019;21:101645.
- 331 Hosseini-Asl E, Keynton R, El-Baz A. Ad Diagnostics By Adaptation Of 3d Convolutional Network. In the IEEE 2016 International Conference on Image Processing; 2016.
- 332 Awate GJ, Bangare SL. Detection of AD from MRI using Convolutional Neural Network with Tensorflow. *IEEE Xplore* 2018.
- 333 Billones CD, Demetria OJL, Hostallero DE, Naval PC. DemNet: A Convolutional Neural Network for the Detection of AD and Mild Cognitive Impairment. In IEEE Region 10 Conference. TENCON; 2016.
- 334 Gunawardena KANN, Rajapaksey RN, Kodikara ND. Applying Convolutional Neural Networks for Pre-detection of AD from Structural MRI data. 24th International Conference on Mechatronics and Machine Vision in Practice (M2VIP). IEEE 2017.
- 335 Farooq A, Anwar SM, Awais M, Rehman S. A deep CNN based Multi-Class classification of Alzheimers Disease using MRI. IEEE International Conference on Imaging Systems and Techniques (IST); 2017.
- 336 Luo S, Li X, Li J; for the AD Neuroimaging Initiative. Automatic AD Recognition from MRI Data Using Deep Learning Method. *Z Angew Math Phys*. 2017;5:1892–8.
- 337 Lin W, Tong T, Gao Q, Guo D, Du X, Yang Y, et al.; Alzheimer's Disease Neuroimaging Initiative. Convolutional Neural Networks-Based MRI Image Analysis for the Alzheimer's Disease Prediction From Mild Cognitive Impairment. *Front Neurosci*. 2018 Nov;12:777.

- 338 Li W, Tian J, Li E, Dai J. Robust unsupervised segmentation of infarct lesion from diffusion tensor MR images using multi-scale statistical classification and partial volume voxel reclassification. *Neuroimage*. 2004 Dec;23(4):1507–18.
- 339 Rebouças Filho PP, Sarmento RM, Holanda GB, de Alencar Lima D. New approach to detect and classify stroke in skull CT images via analysis of brain tissue densities. *Comput Methods Programs Biomed*. 2017 Sep;148:27–43.
- 340 Lutsep HL, Albers GW, DeCrespigny A, Kamat GN, Marks MP, Moseley ME. Clinical utility of diffusion-weighted magnetic resonance imaging in the assessment of ischemic stroke. *Ann Neurol*. 1997 May;41(5):574–80.
- 341 Wang Y, Xiang S, Pan C, Wang L, Meng G. Level set evolution with locally linear classification for image segmentation. *Pattern Recognit*. 2013;46(6):1734–46.
- 342 Alpert S, Galun M, Brandt A, Basri R. Image segmentation by probabilistic bottom-up aggregation and cue integration. *IEEE Trans Pattern Anal Mach Intell*. 2012 Feb;34(2):315–26.
- 343 Ji Z, Xia Y, Zheng Y. Robust generative asymmetric GMM for brain MR image segmentation. *Comput Methods Programs Biomed*. 2017 Nov;151:123–38.
- 344 Sridevi M, Mala C. Self-organizing neural networks for image segmentation based on multiphase active contour. *Neural Comput Appl*. 2019;31(2):865–76.
- 345 Agrawal S, Panda R, Dora L. A study on fuzzy clustering for magnetic resonance brain image segmentation using soft computing approaches. *Appl Soft Comput*. 2014;24:522–33.
- 346 Monteiro M, Fonseca AC, Freitas AT, Pinho EMelo T, Francisco AP, Ferro JM, et al. Using machine learning to improve the prediction of functional outcome in ischemic stroke patients. *IEEE/ACM Trans Comput Biol Bioinform*. 2018 Nov–Dec;15(6):1953–9.
- 347 Matesin M, Loncaric S, Petravic D. A rule based approach to stroke lesion analysis from CT brain images. *Proc.of the 2nd Int Symposium on Image and Signal Processing and Analysis*. 2001;219–223.
- 348 Ghosh N, Recker R, Shah A, Bhanu B, Ashwal S, Obenaus A. Automated ischemic lesion detection in a neonatal model of hypoxic ischemic injury. *J Magn Reson Imaging*. 2011 Apr;33(4):772–81.
- 349 Mitra J, Bourgeat P, Frupp J, Ghose S, Rose S, Salvado O, et al. Lesion segmentation from multimodal MRI using random forest following ischemic stroke. *Neuroimage*. 2014 Sep;98:324–35.
- 350 Maier O, Wilms M, von der Gablentz J, Krämer UM, Münte TF, Handels H. Extra tree forests for sub-acute ischemic stroke lesion segmentation in MR sequences. *J Neurosci Methods*. 2015 Jan;240:89–100.
- 351 Griffis JC, Allendorfer JB, Szaflarski JP. Voxel-based Gaussian naïve Bayes classification of ischemic stroke lesions in individual T1-weighted MRI scans. *J Neurosci Methods*. 2016 Jan;257:97–108.
- 352 Pustina D, Coslett HB, Turkeltaub PE, Tustison N, Schwartz MF, Avants B. Automated segmentation of chronic stroke lesions using LINDA: lesion identification with neighborhood data analysis. *Hum Brain Mapp*. 2016 Apr;37(4):1405–21.
- 353 Pennisi A, Bloisi DD, Nardi D, Giampetruzzi AR, Mondino C, Facchiano A. Skin lesion image segmentation using Delaunay Triangulation for melanoma detection. *Comput Med Imaging Graph*. 2016 Sep;52:89–103.
- 354 Chen L, Bentley P, Rueckert D. Fully automatic acute ischemic lesion segmentation in DWI using convolutional neural networks. *Neuroimage Clin*. 2017 Jun;15:633–43.
- 355 Muda AF, Saad NM, Abu-Bakar SAR, Muda S, Abdullah AR. Brain lesion segmentation using fuzzy C-means on diffusion-weighted imaging. *ARPN J Eng Appl Sci*. 2015;10:1138–44.
- 356 Bentley P, Ganesalingam J, Carlton Jones AL, Mahady K, Epton S, Rinne P, et al. Prediction of stroke thrombolysis outcome using CT brain machine learning. *Neuroimage Clin*. 2014 Mar;4(4):635–40.
- 357 Lebedev G, Klimenko H, Kachkovskiy S, Konushin V, Ryabkov I, Gromov A. Application of artificial intelligence methods to recognize pathologies on medical images. *Procedia Comput Sci*. 2018;126:1171–7.
- 358 Subudhi A, Acharya UR, Dash M, Jena S, Sabut S. Automated approach for detection of ischemic stroke using Delaunay Triangulation in brain MRI images. *Comput Biol Med*. 2018;103:116–29.
- 359 Subudhi A, Jena S, Sabut S. Delineation of the ischemic stroke lesion based on watershed and relative fuzzy connectedness in brain MRI. *Med Biol Eng Comput*. 2018 May;56(5):795–807.
- 360 Chin C-L, Lin B-J, Wu G-R, Weng T-C, Yang C-S, Su R-C, Pan Y-J. An Automated Early Ischemic Stroke Detection System using CNN Deep Learning Algorithm. *IEEE 8th International Conference on Awareness Science and Technology (iCAST 2017)*; 2017.
- 361 Elliott C, Arnold DL, Collins DL, Arbel T. Temporally consistent probabilistic detection of new multiple sclerosis lesions in brain MRI. *IEEE Trans Med Imaging*. 2013 Aug;32(8):1490–503.
- 362 Anbeek P, Vincken KL, van Osch MJ, Bisschops RH, van der Grond J. Automatic segmentation of different-sized white matter lesions by voxel probability estimation. *Med Image Anal*. 2004 Sep;8(3):205–15.
- 363 Ashton EA, Takahashi C, Berg MJ, Goodman A, Totterman S, Ekholm S. Accuracy and reproducibility of manual and semiautomated quantification of MS lesions by MRI. *J Magn Reson Imaging*. 2003 Mar;17(3):300–8.
- 364 Lao Z, Shen D, Liu D, Jawad AF, Melhem ER, Launer LJ, et al. Computer-assisted segmentation of white matter lesions in 3D MR images using support vector machine. *Acad Radiol*. 2008 Mar;15(3):300–13.
- 365 Zhang YD, Pan C, Sun J, Tang C. Multiple sclerosis identification by convolutional neural network with dropout and parametric ReLU. *J Comput Sci*. 2018;28:1–10.
- 366 Wang SH, Tang C, Sun J, Yang J, Huang C, Phillips P, et al. Multiple Sclerosis Identification by 14-Layer Convolutional Neural Network With Batch Normalization, Dropout, and Stochastic Pooling. *Front Neurosci*. 2018 Nov;12:818.