

Published in final edited form as:


*J Am Chem Soc.* 2010 March 10; 132(9): 2878–2879. doi:10.1021/ja100955x.

## Aryl-CF<sub>3</sub> Bond-Forming Reductive Elimination from Palladium(IV)

Nicholas D. Ball, Jeff W. Kampf, and Melanie S. Sanford\*

Department of Chemistry, University of Michigan, 930 North University Ave, Ann Arbor, MI 48109

### Abstract



This communication describes oxidatively-induced Ar-CF<sub>3</sub> bond-forming reductive elimination from new Pd<sup>II</sup> complexes of general structure (L~L)Pd<sup>II</sup>(Ar)(CF<sub>3</sub>). The electrophilic fluorinating reagent *N*-fluoro-2,4,6-trimethylpyridinium triflate promotes these reactions in good to excellent yields. The palladium(IV) intermediate (t-Bu-bpy)Pd<sup>IV</sup>(CF<sub>3</sub>)(F)(OTf)(C<sub>6</sub>H<sub>4</sub>F) has been isolated, characterized, and demonstrated to undergo high yielding Ar-CF<sub>3</sub> coupling upon thermolysis. This work provides an attractive conceptual framework for the development of Pd<sup>II/IV</sup>-catalyzed arene trifluoromethylation reactions.

Trifluoromethyl groups feature prominently in a wide variety of medicinal compounds.<sup>1</sup> The substitution of CH<sub>3</sub> for CF<sub>3</sub> can dramatically change the physical properties and biological activity of organic molecules. As a result, tremendous effort has been directed at the introduction of CF<sub>3</sub> groups into organic structures.<sup>2</sup> This has led to numerous methods for the efficient construction of sp<sup>3</sup> carbon-CF<sub>3</sub> bonds using nucleophilic (CF<sub>3</sub><sup>-</sup>), electrophilic (CF<sub>3</sub><sup>+</sup>) and radical (CF<sub>3</sub>•)-based trifluoromethylating reagents.<sup>2</sup>

In contrast, the formation of aryl carbon-CF<sub>3</sub> bonds remains significantly more challenging.<sup>3</sup> This functional group is commonly prepared using the Swarts reaction (which requires high temperatures and reactive SbF<sub>5</sub>)<sup>4</sup> or through the use of stoichiometric quantities of ill-defined and sensitive “Cu-CF<sub>3</sub>” reagents.<sup>5</sup> Transition metal-catalyzed cross-coupling would provide a highly attractive alternative route to aryl-CF<sub>3</sub> linkages. While sporadic reports have suggested the viability of this approach,<sup>6</sup> general, efficient, and robust versions of such transformations have been slow to develop. The major challenge in this area is that CF<sub>3</sub> ligands are typically inert towards C-C bond-forming reductive elimination.<sup>7</sup>


mssanfor@umich.edu.

**Supporting Information Available:** Experimental and crystallographic details and spectroscopic data for new compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

Several groups have made exciting recent progress towards addressing this challenge. For example, Vicic and coworkers have shown that the isolable *N*-heterocyclic carbene (NHC) copper complexes (NHC)Cu–CF<sub>3</sub> react stoichiometrically with aryl iodides to afford Ar–CF<sub>3</sub> products.<sup>8</sup> Vicic's group has also established that (dippe)Ni<sup>II</sup>(Ph)(CF<sub>3</sub>) undergoes H<sub>2</sub>O-promoted Ph–CF<sub>3</sub> coupling in modest (22%) yield.<sup>9</sup> Finally, Grushin has demonstrated stoichiometric Ph–CF<sub>3</sub> coupling from (Xantphos)Pd<sup>II</sup>(Ph)(CF<sub>3</sub>).<sup>10</sup> This latter work is the only reported example of selective Ar–CF<sub>3</sub> bond-forming reductive elimination from a well-defined transition metal aryl/CF<sub>3</sub> complex. However, this reaction was limited to Ar = Ph and was also extremely sensitive to the nature of the ancillary ligand at Pd<sup>II</sup>. A variety of P- and N-donor ligands including dppe,<sup>11</sup> dppb,<sup>11</sup> tmeda,<sup>11</sup> and PPh<sub>3</sub> were examined,<sup>10,12</sup> but only Xantphos was effective at promoting this transformation below 150 °C. Thus, the development of complementary and more general strategies for arene trifluoromethylation from M(Ar)(CF<sub>3</sub>) species remains a topic of great current interest.


Our group<sup>13</sup> and others<sup>14</sup> have shown that Pd<sup>IV</sup> complexes can participate in reductive elimination reactions that are challenging at other metal centers. Since high oxidation state palladium can be accessed following arene C–H activation<sup>15</sup> or transmetalation<sup>16</sup> processes, Ar–CF<sub>3</sub> coupling through this manifold would provide opportunities for the development of diverse new trifluoromethylation reactions. We report herein the first demonstration of Ar–CF<sub>3</sub> bond-forming reductive elimination from a Pd<sup>IV</sup>(Ar)(CF<sub>3</sub>) complex.

Our studies began with the synthesis of a series of new palladium(II) complexes of general structure (L)<sub>2</sub>Pd<sup>II</sup>(Ar)(CF<sub>3</sub>) (**1–3**).<sup>12b</sup> These were prepared by treating the corresponding Pd<sup>II</sup> aryl iodides with CsF followed by TMSCF<sub>3</sub> at 23 °C in THF (eq. 1).<sup>10,21</sup> The products were obtained in 32–76% yield as yellow solids (see Supporting Information for full details).



(1)


Heating complexes **1–3** at 130 °C for 3 d in nitrobenzene-*d*<sub>5</sub> produced <5% of Ar–CF<sub>3</sub> coupling products (eq. 2). This is consistent with literature reports showing that most Pd<sup>II</sup>(Ar)(CF<sub>3</sub>) complexes are poorly reactive towards Ar–CF<sub>3</sub> bond-forming reductive elimination.<sup>10,12</sup> We reasoned that 2e<sup>−</sup> oxidation should yield Pd<sup>IV</sup> species that might undergo more facile Ar–CF<sub>3</sub> coupling.<sup>13–15</sup> However, treatment of **1a** with PhI(OAc)<sub>2</sub>, NCS,<sup>11</sup> or NBS<sup>11</sup> (oxidants known to promote the oxidation of Pd<sup>II</sup> to Pd<sup>IV</sup>)<sup>13</sup> did not afford 1-fluoro-4-trifluoromethylbenzene; instead, the corresponding acetoxyated or halogenated products were obtained (eq. 2). This result suggests that Ar–X (X = OAc, Cl, Br) bond-forming reductive elimination is significantly faster than Ar–CF<sub>3</sub> coupling in this system.



(2)

We next examined the use of *N*-fluoro-2,4,6-trimethylpyridinium triflate (NFTPT) to effect Ar–CF<sub>3</sub> coupling from **1a**. This oxidant was selected based on the hypothesis that fluoride and triflate (the X-type ligands introduced to a putative Pd<sup>IV</sup> intermediate by NFTPT) might undergo slower reductive elimination than CF<sub>3</sub>.<sup>17,18</sup> Gratifyingly, treatment of **1a** with NFTPT at 80 °C for 3 h in nitrobenzene-*d*<sub>5</sub> resulted in clean formation of 1-fluoro-4-trifluoromethylbenzene in 85% yield (as determined by <sup>19</sup>F NMR spectroscopy) (Table 1, entry 1). As predicted, <5% of products derived from C–F or C–OTf coupling were observed. NFTPT also promoted Ar–CF<sub>3</sub> bond-formation from the other Pd<sup>II</sup> complexes **1b–3** (Table 1). These transformations were efficient with strongly electron donating and electron withdrawing arene substituents (entries 2 and 3, respectively). In contrast, reactions of [Cu–CF<sub>3</sub>] are often highest yielding with electron deficient aryl coupling partners.<sup>6</sup> Additionally, unlike Grushin's Ar–CF<sub>3</sub> coupling from Pd<sup>II</sup> (which is very sensitive to the nature of the ancillary ligands),<sup>10</sup> this oxidatively induced transformation proceeded in modest to excellent yield with diverse N- and P-donor ligands, including <sup>t</sup>Bu-bpy, tmeda, and dppe.<sup>11</sup>

In an effort to detect intermediates in the Ar–CF<sub>3</sub> coupling process, we examined the reaction of **1a** with NFTPT at room temperature. In both nitrobenzene and DCE, a single major palladium-containing product (**4**) was observed. This species was isolated from DCE in 53% yield, and the analytically pure yellow solid exhibits four <sup>19</sup>F NMR signals in a 3 : 3 : 1 : 1 ratio. X-ray quality crystals were obtained by vapor diffusion of pentanes into a DCE solution of **4**. The X-ray crystal structure is shown in Figure 1 and confirms that **4** is the octahedral Pd<sup>IV</sup> species (<sup>t</sup>Bu-bpy)Pd<sup>IV</sup>(CF<sub>3</sub>)(F)(OTf)(C<sub>6</sub>H<sub>4</sub>F). To our knowledge this is the first isolated example of a Pd<sup>IV</sup> complex containing a perfluoroalkyl ligand.<sup>19</sup>



The reactivity of **4** towards Ar–CF<sub>3</sub> bond formation was next investigated. Heating a nitrobenzene-*d*<sub>5</sub> solution of **4** at 80 °C for 3 h resulted in smooth reductive elimination to form 1-fluoro-4-trifluoromethylbenzene in 77% yield (eq. 3). Similar to the reactions in Table 1, none of the products derived from Ar–F or Ar–OTf coupling were observed by <sup>19</sup>F NMR spectroscopy. This result demonstrates for the first time that Ar–CF<sub>3</sub> coupling can be kinetically accessible from mono-σ-aryl Pd<sup>IV</sup> complexes; as such, it provides an attractive conceptual framework for the development of Pd<sup>II/IV</sup>-catalyzed arene trifluoromethylation reactions.

In summary, we describe herein a new Ar–CF<sub>3</sub> bond-forming reaction mediated by Pd<sup>IV</sup> centers. These transformations proceed under mild conditions with diverse nitrogen and phosphorus-based ancillary ligands. Efforts to gain further insights into the mechanism as well as to develop related catalytic transformations are currently underway in our laboratory and will be reported in due course.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

We thank the NIH (GM073836 and F31GM089141) and the Research Corporation Cottrell Scholar Program for support of this research. Additional unrestricted support from Dupont is also gratefully acknowledged. Finally, we thank Paul Lennon and Jim Windak for assistance with mass spectroscopy.


## References


1. Kirk KL. *Org. Process Res. Dev* 2008;12:305.
2. (a) Ma J-A, Cahard D. *Chem. Rev* 2004;104:6119. [PubMed: 15584697] (b) Ma J-A, Cahard D. *Chem. Rev* 2008;108 PR1. (c) Prakash GKS, Chacko S. *Curr. Opin. Drug Disc Dev* 2008;11:793. (d) Shibata N, Mizuta S, Kawai H. *Tetrahedron: Asymmetry* 2008;19:2633.
3. Grushin VV. *Acc. Chem. Res* 2010;43:160. [PubMed: 19788304]
4. Swarts F. *Bull. Acad. R. Belg* 1892;24:309.
5. For select examples, see:(a) Kobayashi Y, Kumadaki I. *Tetrahedron Lett* 1969;10:4095. (b) Konderatenko NV, Vechirko EP, Yagupolskii LM. *Synthesis* 1980:932. (c) Matsui K, Tobita E, Ando M, Kondo K. *Chem. Lett* 1981:1719. (d) Suzuki H, Yoshida Y, Osuka A. *Chem. Lett* 1982:135. (e) Burton DJ, Wiemers DM. *J. Am. Chem. Soc* 1985;107:5014. (f) Urata H, Fuchikami T. *Tetrahedron Lett* 1991;32:91.
6. Cu-catalyzed coupling between of ArI and Et<sub>3</sub>SiCF<sub>3</sub>:Oishi M, Kondo H, Amii H. *Chem. Commun* 2009:1909. Pd-catalyzed coupling between ArI and [Zn-CF<sub>3</sub>]:Kitazume T, Ishikawa N. *Chem. Lett* 1982:137.
7. For reviews on [M]-CF<sub>3</sub> and [M]-R<sub>f</sub> (R<sub>f</sub> = perfluoroalkyl) complexes, see:(a) Hughes RP. *Adv. Organomet. Chem* 1990;31:183. (b) Morrison JA. *Adv. Organomet. Chem* 1993;35:211.
8. (a) Dubinina GG, Furutachi H, Vivic DA. *J. Am. Chem. Soc* 2008;130:8600. [PubMed: 18543912] (b) Dubinina GG, Ogikubo J, Vivic DA. *Organometallics* 2008;27:6233.
9. Dubinina GG, Brennessel WW, Miller JL, Vivic DA. *Organometallics* 2008;27:3933.
10. Grushin VV, Marshall WJ. *J. Am. Chem. Soc* 2006;128:12644. [PubMed: 17002347]
11. Xantphos = 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene; <sup>t</sup>Bu-bpy = 4,4'-di-*tert*-butyl-2,2'-bipyridyl; tmeda = *N,N'*-tetramethylethylene diamine; dppe = 1,2-bis(diphenylphosphino)ethane; dippe = 1,2-bis(diisopropylphosphino)ethane; NCS = *N*-chlorosuccinimide; NBS = *N*-bromosuccinimide.
12. (a) Culkin DA, Hartwig JF. *Organometallics* 2004;23:3398. (b) Grushin VV, Marshall WJ. *J. Am. Chem. Soc* 2006;128:4632. [PubMed: 16594700]
13. (a) Dick AR, Kampf JW, Sanford MS. *J. Am. Chem. Soc* 2005;127:12790. [PubMed: 16159259] (b) Whitfield SR, Sanford MS. *J. Am. Chem. Soc* 2007;129:15142. [PubMed: 18004863] (c) Ball ND, Sanford MS. *J. Am. Chem. Soc* 2009;131:3796. [PubMed: 19249867] (d) Racowski JA, Dick AR, Sanford MS. *J. Am. Chem. Soc* 2009;131:10974. [PubMed: 19459631] (e) Arnold PL, Sanford MS, Pearson SM. *J. Am. Chem. Soc* 2009;131:13912. [PubMed: 19788324]
14. (a) Alsters PL, Engel PF, Hogerheide MP, Copijn M, Spek AL, van Koten G. *Organometallics* 1993;12:1831. (b) Kaspi AW, Yahav-Levi A, Goldberg I, Vigalok A. *Inorg. Chem* 2008;47:5. [PubMed: 18052157] (c) Furuya T, Ritter T. *J. Am. Chem. Soc* 2008;130:10060. [PubMed: 18616246] (d) Canty AJ. *Dalton Trans* 2009:10409. [PubMed: 20023859]
15. For recent reviews, see:(b) Lyons TW, Sanford MS. *Chem. Rev.* [Online early access]. DOI: 10.1021/cr900184e. Published online: Jan 15, 2010, <http://pubs.acs.org/doi/full/10.1021/cr900184e>. (b) Muniz K. *Angew. Chem., Int. Ed* 2009;48:9412.
16. Furuya T, Kaiser HM, Ritter T. *Angew. Chem., Int. Ed* 2008;47:5993.
17. For a similar strategy to generate C-N bonds using NFTPT, see:Mei TS, Wang X, Yu JQ. *J. Am. Chem. Soc* 2009;131:10806. [PubMed: 19606861]
18. For use of similar oxidants in Ar-F coupling, see:(a) Hull KL, Anani WQ, Sanford MS. *J. Am. Chem. Soc* 2006;128:7134. [PubMed: 16734446] (b) Wang X, Mei TS, Yu JQ. *J. Am. Chem. Soc* 2009;131:7520. [PubMed: 19435367]

19. Notably, related structures have been proposed as transient intermediates in the reactions of perfluoroalkyl iodides with Pd<sup>II</sup>. See: Hughes RP, Overby JS, Williamson A, Lam K-C, Concolino TE, Rheingold AL. *Organometallics* 2000;19:5190.



**Figure 1.**  
ORTEP Diagram of 4.



**Table 1**Oxidatively Induced Ar–CF<sub>3</sub> Coupling from Complexes **1–3**


Entry	Compound	L <sub>2</sub>	Ar	Yield Ar–CF <sub>3</sub> <sup>a</sup>
1	<b>1a</b>	<sup>t</sup> Bu-bpy	<i>p</i> -FC <sub>6</sub> H <sub>4</sub>	85%
2	<b>1b</b>	<sup>t</sup> Bu-bpy	<i>p</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	61%
3	<b>1c</b>	<sup>t</sup> Bu-bpy	<i>p</i> -CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	86%
4	<b>1d</b>	<sup>t</sup> Bu-bpy	<i>p</i> -PhC <sub>6</sub> H <sub>4</sub>	83%
5	<b>1e</b>	<sup>t</sup> Bu-bpy	<i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	85%
6	<b>2</b>	tmeda	<i>p</i> -FC <sub>6</sub> H <sub>4</sub>	89%
7	<b>3</b>	dppe	Ph	29%

<sup>a</sup>Determined by <sup>19</sup>F NMR spectroscopy