


## Assessing groundwater contamination risk using the DASTI/IDRISI GIS method: coastal system of western Mamora, Morocco

Brahim Ben Kabbour · Lahcen Zouhri ·  
Jacky Mania · Jean-Pierre Colbeaux

Published online: 29 August 2007  
© Springer-Verlag 2007

**Erratum to: Bull Eng Geol Environ (2006) 65:463–470**  
DOI 10.1007/s10064-004-0265-2

This article contains two errors in Figs. 7 and 8. The correct figures are given below:



**Fig. 7** Vulnerability map according to the DRASTIC method [I. Vulnerability ranges (Aller et al. 1987), II. intervals of Id values]


The online version of the original article can be found under doi:10.1007/s10064-004-0265-2.

B. B. Kabbour  
FST Beni Mellal, Département de Géologie,  
Laboratoire d'Aménagement du Territoire,  
Université Cadi Ayyad, Marrakech, Morocco

L. Zouhri (✉)  
UMR CNRS 8110, Processus et Bilans des Domaines  
Sédimentaires, UFR des Sciences de la Terre,  
Université des Sciences et Technologies de Lille (USTL),  
59655 Villeneuve d'Ascq Cedex, France  
e-mail: lahcen.zouhri@laposte.net

J. Mania  
UMR CNRS 8107, LML, Ecole Polytechnique de Lille,  
Avenue Paul Langevin, 59655 Villeneuve d'Ascq Cedex, France

J.-P. Colbeaux  
UFR de Géologie, Bât SN5, Université de Lille I,  
59655 Villeneuve d'Ascq Cedex, France



**Fig. 8** Sensibility map according to the DRASTIC method and land use analysis