

ASSOCIATED AND PERSPECTIVE SIMPLEXES

BY

LEON GERBER

ABSTRACT. A set of $n + 1$ lines in n -space such that any $(n - 2)$ -dimensional flat which meets n of the lines also meets the remaining line is said to be an associated set of lines. Two simplexes are associated if the joins of corresponding vertices are associated. A simple criterion is given for simplexes to be associated and an analogous one for simplexes to be perspective. These are used to give a brief proof of the following generalization of the theorem of Pappus.

Let A and B be n -simplexes and let p be a permutation on the vertices of B . If A and B are associated (respectively perspective) and A and Bp are associated (perspective) then A and Bp^k are associated (perspective) for any integer k . Very short proofs are given of extensions to n -dimensions of many theorems from Neuberg's famous *Memoir sur le Tétraèdre*, such as: the altitudes of a simplex are associated.

1. Introduction. A set of $n + 1$ lines in n -space having the property that any $(n - 2)$ -dimensional flat which meets n of the lines meets the remaining line is said to be an associated set of lines (see [7], [2]). In 1884, in a famous Memoir [13], J. Neuberg established many of the properties of associated tetrahedra, that is, tetrahedra the joins of whose corresponding vertices are associated lines. In 1905 Berzolari [2] gave a necessary and sufficient condition for two n -simplexes to be associated, proved that the altitudes of an n -simplex are associated (which result was considered new when proved more than fifty years later [10]), and extended to n -dimensions both Neuberg's theorem on orthological tetrahedra and, in a very special way, Pappus' theorem on perspective triangles. In 1907 Neuberg [14] gave a much shorter proof of Berzolari's criterion, valid only in 3-space, and a number of applications.

In this paper we give a proof of Berzolari's criterion for associativity as well as a criterion for perspectivity which is very similar. In the sequel we give a large number of very short proofs of extensions of Neuberg's theorems to n -dimensions as well as the n -dimensional version of Pappus' theorem in its full generality.

Received by the editors August 13, 1973.

AMS (MOS) subject classifications (1970). Primary 50B99; Secondary 50D05.

Key words and phrases. Associated lines, hyperbolic group of lines, perspective, simplex, orthological simplexes, polar simplex, Pappus theorem.

Copyright © 1975, American Mathematical Society

Since the Neuberg Memoir is not easily accessible, we have included more accessible references as well.

A few remarks about associated lines may help orient the reader. In 2-space, given two distinct lines there is a single point which lies on both lines and any line through this point is associated with the given lines. In higher dimensions, then, associated lines may be thought of as a generalization of concurrent lines and many theorems of plane geometry can be extended to higher dimension in this way.

For $n = 3$ there is a 1-parameter family of lines which meet three given skew lines. These form one set of rulings of a hyperboloid of one sheet if there is no plane parallel to the three given lines, and the rulings of a hyperbolic paraboloid if there is such a plane [9, pp. 14–15]. Any line, distinct from the given lines, of the 1-parameter family of conjugate rulings is associated with the given lines. This explains why some authors call a set of associated lines a hyperbolic set [2], [11]. In the plane, every set of three concurrent lines is the set of altitudes of some triangle, but in 3-space a set of four associated lines need not be the altitudes of some tetrahedron [3, p. 75].

For $n = 4$, there is a 2-parameter family of planes which meet four given skew lines and in general there is a unique fifth line which meets these planes, i.e. is associated with the given lines [1, pp. 115–116]. For $n > 4$ there is in general no line which meets the $(n - 2)$ -parameter family of $(n - 2)$ -flats each of which meets n given lines, i.e. no line associated with n given lines.

2. Notation and preliminaries. The setting for our theorems is Euclidean n -dimensional space augmented by an **improper point** (point at infinity) on each line which is defined to be the 1-dimensional vector space determined by pairs of points on the line. Any nonzero vector in this space is a **representative** of the improper point. The duality principle holds: in any true incidence statement we may replace “point” by “**prime**” ($(n - 1)$ -flat) and “line” by “**secundum**” ($(n - 2)$ -flat), providing we reverse any inclusions. Thus any $n + 1$ secunda are said to be **associated** if any line which meets n of the secunda also meets the remaining secundum, and associated secunda are a generalization of collinear points. It is clear that any **correlation** (a transformation that sends k -flats into $(n - 1 - k)$ -flats and preserves incidence) sends a set of associated lines into a set of associated secunda and vice versa.

Let I represent that set $\{0, 1, \dots, n\}$, $I(i) = I - \{i\}$, and $I' = I(0)$, and let the ranges of Σ , $\Sigma^{(i)}$, Σ' be I , $I(i)$, and I' respectively. Throughout this paper A will be a proper n -simplex with corresponding vertices and faces given by $A_i, A_p, i \in I$. In general, an upper case letter represents a point and a lower case boldface letter its position vector if it is proper or a representative if

it is improper. Any point B has an expression in the form $\mathbf{b} = \sum b_i \mathbf{a}_i$ where $s(B) = \sum b_i = 1$ if B is proper and $s(B) = 0$ if B is improper. If B is proper the expression is unique, while if B is improper there is a unique expression for each representative of B . The b_i are called the **weights** (barycentric coordinates) of B (with respect to A) and we write $B = (b_0, \dots, b_n) = (b_i)$. The b_i are sometimes called areal coordinates: If c_n is the n -dimensional content (signed volume) function and B_i is the simplex obtained from A by replacing the vertex A_i by B , then $b_i = c_n(B_i)/c_n(A)$, for $i \in I$. Dually, the weights (x_i) of the points X of any prime F satisfy a unique (to within a multiplicative constant) homogeneous equation $\sum u_i x_i = 0$. The u_i are called the **weights** of F and we write $F = (u_i)$. Thus $A_i = (\delta_{ij})$ where δ_{ij} is 1 if $j = i$ and 0 otherwise, and $(1, 1, \dots, 1)$ is (the weights of) the prime which consists of all improper points. The u_i are sometimes called tangential coordinates: they are proportional to the lengths of parallel cevians from the A_i to F . (A cevian of a simplex is a line passing through a vertex.)

If B is a proper point, let d_i be the distance of B from A_i , positive if B is on the same side of A_i as A_i and negative if B is on the other side, $i \in I$. The d_i are called the **heights** (absolute normal coordinates) of B . Let $\overrightarrow{A_i H_i} = \mathbf{h}_i$ be the altitude from A_i and h_i its length. If $B = (b_i)$, then $d_i = b_i h_i$ in magnitude and sign. Let $v_n = |c_n|$ be the n -dimensional volume function, $v = v_n(A)$ and $f_i = v_{n-1}(A_i)$, $i \in I$. Then $f_i h_i = nv$ and $f_i d_i = nv b_i$.

Finally, B will denote an n -simplex, which may degenerate or have improper vertices, in the space of A with corresponding vertices and faces given by $B_i, B_p, i \in I$. The matrix (b_{ij}) will have for its i th row the weights of $B_p, i \in I$. The equation of B_i is $D_i = 0$ where D_i is the determinant of the matrix obtained from (b_{ij}) by replacing row i by (x_0, \dots, x_n) . Thus the matrix of weights of the B_i has as its elements the cofactors of (b_{ij}) .

3. Fundamental result. We now prove a simple theorem about concurrent cevians whose generalization to associated cevians will be our basic tool. Let us call the matrix (b_{ij}) **composite** if there exist numbers $y_p, z_p, i \in I$, such that $b_{ij} = y_i z_j$ for $i \neq j$.

THEOREM 3.1. *The cevians $A_i B_p, i \in I$, are concurrent if and only if the matrix (b_{ij}) is composite.*

PROOF. The lines $A_i B_i$ concur at $P = (p_j)$ if and only if, for $i \in I$, $\mathbf{b}_i = u_i \mathbf{a}_i + w_i \mathbf{p}$ for some u_p, w_i where $u_i = 1 - w_i, -w_i, 1$, or 0 according as both P and B_i, P only, B_i only, or neither is proper. If the lines concur at P , then it is clear that $b_{ij} = w_i p_j$ for $i \neq j$. Conversely, suppose $b_{ij} = y_i z_p, i \neq j$.

Suppose $\sum z_i = z \neq 0$. Then $b_{ij} = w_i p_j, i \neq j$, where $w_i = y_i z, p_j = z_j/z$ and $\sum p_j = 1$. Let $s_i = s(B_i)$. Then

$$b_{ii} = s_i - \sum_j^{(i)} b_{ij} = s_i - \sum_j^{(i)} w_i p_j = (s_i - w_i) + w_i p_i$$

so $b_i = (s_i - w_i)a_i + w_i p$ where $P = (p_i)$. If $\sum z_i = 0$, then

$$b_{ii} = s_i - \sum_j^{(i)} y_i z_j = s_i + y_i z_i$$

so $b_i = s_i a_i + y_i p$ where $P = (z_i)$, which completes the proof.

We observe that if (b_{ij}) is composite with constants $y_p, z_i, i \in I$, it follows that $(z_i/y_i)b_{ij} = (z_j/y_j)b_{ji}$. We shall call the matrix (b_{ij}) **semisymmetric** if we can find numbers $x_p, i \in I$, not all zero, such that $x_i b_{ij} = x_j b_{ji}$ for $i, j \in I$. Let M be the class of semisymmetric matrices. It follows that if a matrix $A \in M$, then its transpose A^T , the matrix obtained by multiplying a row or column by a nonzero constant, and the matrix of cofactors are all semisymmetric. The class of composite matrices is also closed under these operations. We are now ready for our basic tool ([2], [14, p. 299]).

THEOREM 3.2. *The cevians $A_i B_p, i \in I$, are associated if and only if the matrix (b_{ij}) is semisymmetric.*

PROOF. Suppose $(b_{ij}) \in M$ with multipliers $x_p, i \in I$. For $i = 2, \dots, n$, let P_i lie on $A_i B_i$ so $p_i = w_i a_i + u_i b_i$ where $u_i = s(P_i) - w_i$. The secundum determined by the P_i meets $A_0 B_0$ if and only if A_0, B_0 , and the P_i lie in some prime, i.e.

$$0 = \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ b_{00} & b_{01} & b_{02} & \cdots & b_{0n} \\ u_2 b_{20} & u_2 b_{21} & u_2 b_{22} + w_2 & \cdots & u_2 b_{2n} \\ & & \cdots & & \\ u_n b_{n0} & u_n b_{n1} & u_n b_{n2} & \cdots & u_n b_{nn} + w_n \end{vmatrix}$$

$$= \begin{vmatrix} b_{01} & b_{02} & \cdots & b_{0n} \\ u_2 b_{21} & u_2 b_{22} + w_2 & \cdots & u_2 b_{2n} \\ u_n b_{n1} & u_n b_{n2} & \cdots & u_n b_{nn} + w_n \end{vmatrix} \cdot$$

Multiplying row zero by x_0 and row i by x_i/u_p , $i = 2, \dots, n$, results in a determinant which is symmetric in the subscripts 0 and 1, i.e. the secundum also meets A_1B_1 . (If some u_i is zero, first expand in terms of the corresponding row.)

To prove the converse, assume the cevians A_iB_p , $i \in I$, are associated. Dividing the rows of the last determinant above by the u_p , $i = 2, \dots, n$, and setting $r_i = b_{ii} + w_i/u_i$ we see that

$$\begin{vmatrix} b_{01} & b_{02} & b_{03} & \cdots & b_{0n} \\ b_{21} & r_2 & b_{23} & \cdots & b_{2n} \\ b_{31} & b_{32} & r_3 & \cdots & b_{3n} \\ & & \cdots & & \\ b_{n1} & b_{n2} & b_{n3} & \cdots & r_n \end{vmatrix} = 0$$

implies

$$\begin{vmatrix} b_{10} & b_{12} & b_{13} & \cdots & b_{1n} \\ b_{20} & r_2 & b_{23} & \cdots & b_{2n} \\ b_{30} & b_{32} & r_3 & \cdots & b_{3n} \\ & & \cdots & & \\ b_{n0} & b_{n2} & b_{n3} & \cdots & r_n \end{vmatrix} = 0$$

for all values of the r_i . As P_i varies, r_i takes on all real values. Thus as a polynomial in the r_i the second is b_{01}/b_{10} times the first. The respective coefficients of $r_3 \cdots r_n$ are $-b_{21}b_{02}$ and $-b_{12}b_{20}$ so $(b_{02}/b_{20})b_{21} = (b_{01}/b_{10})b_{12}$. Working with the coefficients of the other terms of degree $n - 2$ we get

$$(b_{0i}/b_{i0})b_{i1} = (b_{01}/b_{10})b_{1i} \text{ for } i = 2, 3, \dots, n.$$

Next, form the pair of determinant equations in which the special subscripts are 1 and 2 (instead of 0 and 1) and compare coefficients of $r_0r_4r_5 \cdots r_n$ to get $(b_{12}/b_{21})b_{23} = (b_{13}/b_{31})b_{32}$. Since $(b_{02}/b_{20})b_{21} = (b_{01}/b_{10})b_{12}$. and $(b_{03}/b_{30})b_{31} = (b_{01}/b_{10})b_{13}$ we obtain

$$(b_{02}/b_{20})b_{23} = (b_{03}/b_{30})b_{32}.$$

In a similar fashion we get

$$(b_{0i}/b_{i0})b_{ij} = (b_{0j}/b_{j0})b_{ji} \text{ for all distinct } i, j \in I'.$$

If we take $x_0 = 1, x_i = b_{0i}/b_{i0}$ for $i \in I'$, the proof is complete.

Referring to the determinant of the direct proof we see that for each choice of parameters w_3, \dots, w_n we can choose w_2 so the determinant vanishes. This proves the assertion that there is an $(n-2)$ -parameter family of secunda which meet n of the lines.

Although we do not wish to discuss quadrics in this paper, the reader familiar with the subject will observe that an immediate consequence of Theorem 3.2 is the following ([2], [5], [14], [15]).

COROLLARY. *The lines $A_i B_p$, $i \in I$, are associated if and only if A and B are polar with respect to some quadric.*

This generalizes the known result for $n=2$ (Chasles' theorem and its converse [18, p. 125, Exercise 14]): The lines $A_i B_p$, $i=0, 1, 2$, are concurrent if and only if the triangles A and B are polar with respect to some conic. An independent proof of the "if" part of the corollary for the case of the sphere is given later as Theorem 6.3.

Following Neuberg [14, p. 304] (cf. [10, p. 409]) consider the polar of a simplex A with respect to the absolute. In this case the polar of a face of A is the improper point on a perpendicular to the face so the associated lines $A_i B_i$ of the Corollary are the altitudes of A . A metric proof that the altitudes of a simplex are associated is given later as Theorem 5.1.

The dual of Theorem 3.2 is ([2], [14]):

THEOREM 3.3. *Let B_i be a prime in the space of A with weights (b_{ij}) , $i \in I$. Then the secunda $A_i \cap B_i$ are associated if and only if the matrix (b_{ij}) is semisymmetric.*

In the sequel we use these theorems to give short proofs of many theorems, some of which are new.

4. Desargues', Pappus', and other theorems. Each of the next two theorems reduces to Desargues' theorem for $n=2$. (For a different extension of Desargues' theorem see [11].)

THEOREM 4.1 [18, p. 54, Exercise 26]. *Let B be a simplex in the space of A . Then the lines $A_i B_p$, $i \in I$, are concurrent at a point P if and only if the secunda $A_i \cap B_p$, $i \in I$, lie in a prime F . In either case we say A and B are perspective with P and F as center and axis of perspectivity.*

This theorem is its own dual as is the following analogue ([2], cf. [3, pp. 119–120], [14]).

THEOREM 4.2. *Let B be a simplex in the space of A . Then the lines $A_i B_p$, $i \in I$, are associated if and only if the secunda $A_i \cap B_p$, $i \in I$, are associated. In either case we say the simplexes A and B are associated.*

PROOF OF THEOREMS 4.1 AND 4.2. The matrix of the weights of the B_i is the matrix of cofactors of the weights of the B_i . Thus the hypothesis of each theorem is equivalent to the conclusion, via the remark preceding Theorem 3.2.

Our next result concerns the theorem of Pappus which is usually stated as:

If the six vertices of a hexagon lie alternately on two lines, then the three points of intersection of pairs of opposite sides are collinear.

However, the theorem can also be stated as follows [18, p. 100]:

If triangle $A_0A_1A_2$ is perspective with $B_0B_1B_2$ and with $B_1B_2B_0$, then it is also perspective with $B_2B_0B_1$.

In this form we can extend Pappus' theorem to n -space in its full generality (cf. [2]) in two ways. In each, p is a permutation on I , ip denotes the image of i under p , and Bp is the simplex with vertices $B_{0p}, B_{1p}, \dots, B_{np}$.

THEOREM 4.3. *If A and B are perspective and A and Bp are perspective, then A and Bp^k are perspective for every integer k*

THEOREM 4.4. *If A and B are associated and A and Bp are associated, then A and Bp^k are associated for every integer k*

To prove Theorem 4.4 we prove the equivalent proposition: Let q be the inverse of p . If A and Bq are associated and A and B are associated, then A and Bp are associated.

Let the matrix (of the weights of the vertices) of simplex B be (b_{ij}) . By hypothesis, we can find nonzero numbers $x_i, i \in I$, such that $x_i b_{ij} = x_j b_{ji}$, and the matrix of $Bq = (b_{iq,j}) \in M$, the class of semisymmetric matrices. Applying p to the row and column subscripts of the latter (equivalent to relabelling the corresponding vertices of A and Bq) we get $(b_{i,ip}) \in M$. Its transpose $(b_{j,ip}) \in M$ and so $(x_j b_{j,ip}) \in M$. The last matrix equals $(x_{ip} b_{ip,j})$ so $(b_{ip,j}) \in M$ and this is the matrix of Bp . To prove Theorem 4.3 replace "associated" and "semisymmetric" by "perspective" and "composite" in the above proof.

The next three theorems are easy corollaries of Theorem 3.2. P' is called the isotomic (respectively isogonal) conjugate of the point P with respect to A if the weights (heights) of P' are inversely proportional to those of P . Thus [14, p. 301]:

THEOREM 4.5 (cf. [3, p. 139] and [13, p. 28]). *Let B'_i denote the isotomic conjugate of B_i with respect to A . Then the cevians $A_i B'_i, i \in I$, are associated (concurrent) if and only if the cevians $A_i B''_i, i \in I$, are associated (concurrent).*

THEOREM 4.6 [13, p. 28]. *Let B'_i denote the isogonal conjugate of B_i*

with respect to A . Then the cevians $A_i B_i$, $i \in I$, are associated (concurrent) if and only if the cevians $A_i B'_i$, $i \in I$, are associated (concurrent).

Dually, we say that F' is the reciprocal transversal of the prime F if its weights are inversely proportional to those of F . Thus we have

THEOREM 4.7. *Let B'_i be the reciprocal transversal of B_i with respect to A . Then the secunda $A_i \cap B_i$, $i \in I$, are associated (coprime) if and only if the secunda $A_i \cap B'_i$ are associated (coprime).*

The point P and prime F are said to be pole and polar with respect to A if the weights of P are inversely proportional to those of F . (In this case F is the axis of perspectivity for A and the pedal simplex of P with respect to A .) Thus

THEOREM 4.8. *Let B_i and F_i be pole and polar with respect to A , $i \in I$. Then the cevians $A_i B_i$ are associated (concurrent) if and only if the secunda $A_i \cap F_i$ are associated (coprime).*

5. Perpendiculars. Our first theorem is the recently rediscovered result on altitudes [2], [7], [10], [13, p. 23], [14, p. 304].

THEOREM 5.1. *The altitudes of a simplex are associated.*

PROOF. H_i , the foot of the altitude from A_i , is the orthogonal projection of A_i on A_j . Thus its j th weight with respect to A_j , and hence with respect to A , is the $(n-1)$ -content of the projection of A_j on A_i divided by the $(n-1)$ -content of A_j , $j \in I(i)$, i.e. $h_{ii} = 0$ and $h_{ij} = f_j f_i^{-1} \cos(A_i, A_j)$ for $j \in I(i)$. Thus $f_i^2 h_{ij} = f_i f_j \cos(A_i, A_j) = f_i^2 h_{ji}$.

COROLLARY (cf. [13, p. 29]). *The altitudes concur if and only if the matrix $(\cos(A_i, A_j))$ is composite.*

COROLLARY [3, p. 76, Exercise 5]. *The perpendiculars to the prime faces of a simplex at their centroids are associated.*

PROOF. The dilatation with center at the centroid and constant $-1/n$ sends the altitudes into these perpendiculars.

The case of concurrency (for $n = 3$) in the next theorem is due to Steiner [3, pp. 173–174, p. 342] in which case A and B are said to be mutually **orthological**. The associated case (for $n = 3$) is due to Neuberg [13, p. 27], in which case A and B are called **skew orthological**.

THEOREM 5.2. *Let B be a proper simplex in the space of A . If the perpendiculars from A_i to B_j , $i, j \in I$, are associated (concurrent), then the perpen-*

diculars from B_i to $A_i, i \in I$, are associated (concurrent).

PROOF. Let P_i be the foot of the perpendicular from A_i to B_i and p_i its length, and c_i^j the cosine of the angle between A_j and B_i . It is easy to see that for $j \in I(i)$ the signed distance of P_i from A_j is $p_i c_i^j$ so the j th weight of P_i with respect to A is $(1/n)f_j p_i c_i^j$. It follows that (c_i^j) is semisymmetric (composite) and hence (c_j^i) is semisymmetric (composite). The conclusion follows by expressing the weights of the feet of the perpendiculars from B_i to A_i with respect to B .

6. Spheres. Let $S(S, r)$ be the sphere with center S and radius r . Two proper points X and P are conjugate with respect to S if, with S as origin, $\mathbf{p} \cdot \mathbf{x} = r^2$; if either X or P is improper, the condition is $\mathbf{p} \cdot \mathbf{x} = 0$. Thus the locus of points conjugate to a k -flat F is an $(n - 1 - k)$ -flat F' , completely perpendicular to F , called the polar of F and their unique common perpendicular passes through S . In particular, the polar of S is the improper prime and if F contains S then F' is entirely improper. The transformation that sends a flat into its polar is a correlation, called a polarity, hence

THEOREM 6.1. *If $F_i, i \in I$, is an associated set of lines, then $F'_i, i \in I$, is an associated set of secunda.*

The simplex A'' which has for its vertices A'_i and for its faces $A''_i, i \in I$, is called the polar simplex of A .

THEOREM 6.2. *If A and B are associated simplexes, so are A'' and B'' .*

PROOF. The hypothesis is that the lines $A_i B_i, i \in I$, are associated. By Theorem 6.1, the secunda $A'_i \cap B'_i = A''_i \cap B''_i, i \in I$, are associated. The conclusion follows from Theorem 4.2.

The preceding theorem states that a polarity preserves an existing associative relation. The next theorem shows that a polarity generates an associative relation [2], [5], [14], [15].

THEOREM 6.3. *A simplex and its polar simplex are associated.*

PROOF. Let A be the given simplex and let $B_i = (b_{ij})$ be the polar of A_i with respect to $S(S, r)$. Let $s_i = s(A_i)$ and let $X = (x_i)$ be a proper point of B_i . The equation of B_i is

$$0 = \mathbf{a}_i \cdot \mathbf{x} - s_i r^2 = \mathbf{a}_i \cdot \sum x_j \mathbf{a}_j - s_i r^2 \sum s_j x_j = \sum (\mathbf{a}_i \cdot \mathbf{a}_j - s_i s_j r^2) x_j$$

so $b_{ij} = \mathbf{a}_i \cdot \mathbf{a}_j - s_i s_j r^2 = b_{ji}$. The result follows from Theorems 3.3 and 4.2.

Applying the result to the special case $r = 0$ we get

THEOREM 6.4 [3, p. 244, EXERCISE 36], [16]. *Let S be any point in the space of A and let F_i be a prime through S perpendicular to SA_i , $i \in I$. Then the secunda $F_i \cap A_i$ are associated, $i \in I$.*

The tangent primes to the circumsphere of a simplex at its vertices are the polars of the vertices with respect to the circumsphere and form the **tangential simplex**. Thus [3, pp. 117–118], [13, p. 24]

THEOREM 6.5. *A simplex and its tangential simplex are associated.*

For $n = 2$ we recognize this theorem as a special case of Brianchon's theorem. Reversing the roles of the two simplexes in the previous theorem we get [3, p. 118], [13, p. 24]:

THEOREM 6.6. *Let B_i be the point of contact of the inscribed sphere of A with A_i , $i \in I$. Then A and B are associated.*

THEOREM 6.7 [13, p. 23], [17, p. 61]. *The lines joining the vertices of a simplex to the incenters of the opposite faces are associated.*

PROOF. Let $I_i = (w_{ij})$ be the incenter of A_i . Let $A_{ij} = A_i \cap A_j$, let $f_{ij} = v_{n-2}(A_{ij})$ and let $k_i = \sum_j^{(i)} f_{ij}$. Then $w_{ij} = f_{ij}/k_i$ [6, p. 453] so $k_i w_{ij} = f_{ij} = k_j w_{ji}$.

We remark in passing that the lines joining the vertices of a simplex to the circumcenters of the opposite faces are not in general associated.

We next prove a generalization of the theorem that the bisectors of the angles of a triangle concur. A **corner** with vertex A_i , the analogue of an angle, is what remains of the simplex A when the face A_i is deleted. An **isoclinal prime** of a corner is a prime which cuts equal segments on the edges of the corner; all such primes are related by a dilatation with center A_i . A **centroidal line** of a corner is a line which joins the vertex to the centroid of the $(n-1)$ -simplex determined by an isoclinal prime. The centroidal lines of a triangle are the angle bisectors. Let $|AB|$ denote the length of segment AB . An n -simplex A is said to be **isodynamic** if there exist positive numbers t_i , $i \in I$, such that $|A_i A_j| = t_i t_j$, $i \neq j$. (A full discussion of isodynamic n -simplexes for $n = 3$ will be found in [3, pp. 314–330] and [17].) We have the following result. (The associated case is [3, p. 140, Example 2].)

THEOREM 6.8. *The centroidal lines of a simplex are associated; they concur if and only if the simplex is isodynamic.*

PROOF. For the vertex A_i choose D_j on $A_i A_j$ so that $|A_i D_j| = c_i$, $j \in I(i)$, so $\overrightarrow{A_i D_j} = c_i |A_i A_j|^{-1} \overrightarrow{A_i A_j}$. The centroid B_i of the isoclinal prime determined by the D_j is given by

$$\overrightarrow{A_i B_i} = c_i n^{-1} \sum_j^{(i)} |A_i A_j|^{-1} \overrightarrow{A_i A_j}.$$

If we choose c_i so that $c_i^{-1} = n^{-1} \sum_j^{(i)} |A_i A_j|^{-1}$, the expression on the right is independent of the origin and defines B_i as a point of A_i with weights $b_{ij} = c_i n^{-1} |A_i A_j|^{-1}$, $j \neq i$. Thus $c_i^{-1} b_{ij} = n^{-1} |A_i A_j|^{-1} = c_j^{-1} b_{ji}$ and the lines are associated.

The lines will concur if and only if there exist numbers $x_i, y_i, i \in I$, such that $x_i y_j = c_i n^{-1} |A_i A_j|^{-1}$ for $i \neq j$. Then

$$|A_i A_j| = |A_i A_j|^{1/2} |A_j A_i|^{1/2} = (n c_i^{-1} x_i^{-1} y_i^{-1})^{1/2} (n c_j^{-1} x_j^{-1} y_j^{-1})^{1/2} = t_i t_j,$$

so the simplex is isodynamic and conversely.

Let a prime meet the edges $A_0 A_i$ of A in $B_i, i \in I'$. If the $2n$ points $A_i, B_i, i \in I'$ are cospherical, then A_0 and B_0 are said to be antiparallel sections of the corner at A_0 . In this case B_0 is parallel to the prime tangent to the circumsphere at A_0 and conversely.

THEOREM 6.9 [3, p. 288, Exercise 6], [13, p. 30]. *The lines joining the vertices of a simplex to the centroids of sections antiparallel to the corresponding faces are associated, and concur if and only if the simplex is isodynamic.*

PROOF. Let p_0 be the power of A_0 with respect to the sphere through A_j and $B_j, j \in I'$. Then $\overrightarrow{A_0 B_j} = p_0 |A_0 A_j|^{-2} \overrightarrow{A_0 A_j}, j \in I'$, so the weights of the centroid M_0 of this section are given by $m_{0j} = n^{-1} p_0 |A_0 A_j|^{-2}, j \in I'$. Similarly, $M_i = (m_{ij})$ with $m_{ij} = n^{-1} p_i |A_i A_j|^{-2}, j \in I(i)$. Since $p_i \neq 0, p_i^{-1} m_{ij} = n^{-1} |A_i A_j|^{-2} = p_j^{-1} m_{ji}$ and the lines are associated. The case of concurrency is handled as in the preceding theorem.

7. The first Lemoine point of a simplex. Since there does not seem to be any discussion of this point in English (cf. [13], [17, pp. 147–149] for $n = 3$, and [8] for the general case) we prove some preliminary results. We recall that $f_i = v_{n-1}(A_i)$.

THEOREM 7.1. *If P is a proper point in the space of A with heights (d_i) then $\sum d_i^2 \geq n^2 v^2 / \sum f_i^2$. The minimum is attained at a unique point $K = (w_i)$ called the first Lemoine point of A where $w_i = f_i^2 / \sum f_i^2$.*

PROOF. An application of the Schwarz inequality yields:

$$\sum d_i^2 = n^2 v^2 \sum (w_i / f_i)^2 \geq n^2 v^2 \left[\sum (w_i / f_i) f_i \right]^2 / \sum f_i^2 = n^2 v^2 / \sum f_i^2.$$

Equality is attained if and only if (w_i / f_i) and (f_i) are proportional. Since

$\sum w_i = 1$, we have the result.

THEOREM 7.2 (cf. [17, pp. 27–28]). *There are, in general, n primes which divide $n + 1$ given ordered line segments in the same ratio.*

PROOF. Let the segments be $A_i B_i$, $i \in I$. Let $B_i = (b_{ij})$ with respect to A and let $p_i = (1 - r)a_i + rb_i$. The P_i lie in a prime if and only if $\det(p_{ij}) = 0$. Add each column to the first to get a column of 1's. Expand by the first column to get a polynomial of degree n in r .

THEOREM 7.3 (cf. [17, pp. 148–149]). *Any prime which divides the altitudes of a simplex in the same ratio contains the first Lemoine point.*

PROOF. Let the point B_i lie on the altitude $A_i H_i$ so that $\overrightarrow{A_i B_i} = r \overrightarrow{A_i H_i}$, $i \in I$. Define the point P by $(\sum f_i^2)p = \sum f_i^2 b_i$ so that if the B_i are coprime then P lies in this prime. Let $'$ denote projection on A_0 . Then

$$\begin{aligned} \left(\sum f_i^2\right) |PP'| &= \sum f_i^2 |B_i B_i'| = (1 - r)f_0^2 |A_0 H_0| + r \sum f_i^2 |H_i H_i'| \\ &= (1 - r)f_0^2 |A_0 H_0| + r \sum f_i^2 |A_i H_i| \cos(A_i, A_0) \\ &= nu \left[(1 - r)f_0 + r \sum f_i \cos(A_i, A_0) \right] = rnvf_0 \end{aligned}$$

and similarly for the other normal coordinates of P . Thus $P = K$.

THEOREM 7.4 (cf. [13, p. 23]). *The lines joining the vertices of a simplex to the first Lemoine points of the opposite faces are associated.*

PROOF. Replace " f_{ij} " by " f_{ij}^2 " in the proof of Theorem 6.7.

Thanks are due to H. S. M. Coxeter for encouragement and for references [2] and [15].

REFERENCES

1. H. F. Baker, *Principles of geometry*. Vol. IV, Cambridge Univ. Press, New York, 1940.
2. L. Berzolari, *Sui sistemi di $n + 1$ rette dello spazio ad n dimensioni situati in posizione di Schläfli*, Rend. Circ. Mat. Palermo (2) 20 (1905), 229–247.
3. N. A. Court, *Modern pure solid geometry*, 2nd ed., Chelsea, New York, 1964.
4. H. S. M. Coxeter, *Introduction to geometry*, Wiley, New York, 1961. MR 23 #A1251.
5. H. S. M. Coxeter and J. A. Todd, *Solution of advanced problem 4079*, Amer. Math. Monthly 51 (1944), 599–600.
6. L. Gerber, *Spheres tangent to all the faces of a simplex*, J. Combinatorial Theory 12 (1972), 453–456. MR 45 #7607.
7. ———, *The altitudes of a simplex are associated*, Math. Mag. 46 (1973), 155–157.
8. H. Hanani, *On a point of minimum sum of distances-squares from the faces of a simplex*, Riveon Lematematika 7 (1954), 10–12. MR 15, 451. (Hebrew)

9. D. Hilbert and S. Cohn-Vossen, *Geometry and the imagination*, Springer-Berlin, 1932; English transl., Chelsea, New York, 1952. MR 13, 766.
10. S. R. Mandan, *Attitudes of a general n -simplex*, J. Austral. Math. Soc. 5 (1965), 409–415. MR 32 #6277.
11. ———, *Desargues' theorem in n -space*, J. Austral. Math. Soc. 1 (1959/60), 311–318. MR 23 #A4045.
12. ———, *Isodynamic & isogonic simplexes*, Ann. Mat. Pura Appl. (4) 53 (1961), 45–55. MR 23 #A1253.
13. J. Neuberg, *Memoir sur le tétraèdre*, Académie Royale des Sciences des lettres et des beaux-arts de Belgique, Mémoires couronnés 37 (1884), 1–72.
14. ———, *Über hyperboloidische Würfe*, Arch. Math. Phys. (3) 12 (1907), 297–305.
15. L. Schläfli, *Erweiterung des Satzes, das zwei polare Dreiecke perspectivisch liegen, auf eine beliebige Zahl von Dimensionen*, J. Reine Angew. Math. 65 (1866), 189–197 (= *Gesammelte mathematische Abhandlungen*. Bd. III, Verlag Birkhäuser, Basel, 1956, pp. 9–20. MR 17, 814).
16. H. M. Taylor, *Solution of question 15886*, Educational Times Reprints (2) 10 (1906), 48.
17. V. Thébault, *Parmi les belles figures de la géométrie dans l'espace*, Librairie Vuibert, Paris, 1955. MR 16, 737.
18. O. Veblen and W. J. Young, *Projective geometry*, Blaisdell, New York, 1938.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S UNIVERSITY, JAMAICA, NEW YORK 11439