

# UCSF

## UC San Francisco Previously Published Works

### Title

Association Between Comorbid Depression and Rates of Postoperative Complications, Readmissions, and Revision Arthroscopic Procedures After Elective Hip Arthroscopy.

### Permalink

<https://escholarship.org/uc/item/8gz7f2gh>

### Journal

Orthopaedic journal of sports medicine, 9(9)

### ISSN

2325-9671

### Authors

Freshman, Ryan D  
Salesky, Madeleine  
Cogan, Charles J  
et al.

### Publication Date

2021-09-01

### DOI

10.1177/232596712111036493

Peer reviewed

# Association Between Comorbid Depression and Rates of Postoperative Complications, Readmissions, and Revision Arthroscopic Procedures After Elective Hip Arthroscopy

Ryan D. Freshman,<sup>\*†</sup> MD, Madeleine Salesky,<sup>†</sup> AB, Charles J. Cogan,<sup>†</sup> MD, Drew A. Lansdown,<sup>†</sup> MD, and Alan L. Zhang,<sup>†</sup> MD

*Investigation performed at University of California–San Francisco, San Francisco, California, USA*

**Background:** Depression and related psychiatric diagnoses are common in patients undergoing hip arthroscopy for femoroacetabular impingement syndrome (FAIS). The effects of depression on postoperative complications, readmissions, and additional ipsilateral hip surgery are not well studied.

**Hypothesis:** Patients with preoperative depression who undergo hip arthroscopy for FAIS would experience higher rates of 90-day postoperative complications and readmissions, with an increased risk of additional ipsilateral hip procedures, as compared with patients without depression.

**Study Design:** Cohort study; Level of evidence, 3.

**Methods:** A retrospective cohort study between 2010 and 2019 was performed using the Mariner/PearlDiver database. Current Procedural Terminology and International Classification of Diseases codes were used to compare patients with and without preexisting depression who underwent hip arthroscopy for FAIS. Patients were matched at a 1:1 ratio based on age, sex, Charlson Comorbidity Index, body mass index, and tobacco use. Patients undergoing shoulder or knee arthroscopy were also identified to compare lifetime preoperative depression prevalence amongst groups.

**Results:** The lifetime preoperative depression prevalence was significantly higher in patients undergoing hip arthroscopy as compared with patients undergoing shoulder or knee arthroscopy (25.4% vs 22.2% vs 19.8%;  $P < .001$ ). When compared with the patients without depression, patients with preoperative depression had higher rates of 90-day readmissions (2.4% vs 1.5%) and complications, including urinary tract infection (36.2% vs 28.9%), pneumonia (12.9% vs 9.1%), hematoma formation (3.1% vs 1.9%), acute kidney injury (4.0% vs 2.6%), deep venous thrombosis/pulmonary embolism (2.6% vs 1.7%), and superficial infection (4.9% vs 2.8%;  $P < .01$  for all comparisons). Preoperative depression was associated with significantly higher odds of undergoing revision hip arthroscopy within 2 years (6.3% vs 2.4%;  $P < .001$ ).

**Conclusion:** Patients with preexisting depression experienced higher rates of 90-day postoperative complications and hospital readmissions after elective hip arthroscopy for FAIS and were more likely to undergo revision hip arthroscopy within 2 years of the index procedure.

**Keywords:** depression; hip; arthroscopy; femoroacetabular impingement; postoperative complications

Hip arthroscopy is a minimally invasive procedure most commonly utilized to treat femoroacetabular impingement syndrome (FAIS), which has become increasingly recognized as a source of hip pain and disability.<sup>3,28</sup> While FAIS reduces patients' ability to perform activities of daily living or participate in sports,<sup>28</sup> arthroscopic management results in improved functional patient-reported outcomes (PROs) and return to sports at midterm follow-up.<sup>4,5,22,26,35</sup> Recent studies have reported superior

outcomes after hip arthroscopy as compared with nonoperative management, with patients who undergo arthroscopy being more likely to achieve the minimal clinically important difference for several validated PRO scores.<sup>10,11</sup>

A significant proportion of patients with FAIS have psychiatric disorders, with up to 35% of patients reported to have a diagnosis of depression or anxiety before elective hip arthroscopy.<sup>16</sup> These conditions are associated with inferior PROs after hip arthroscopy,<sup>8,21,23,24</sup> as well as an increased risk of chronic opioid use and higher health care costs.<sup>15</sup> Although several studies have demonstrated that patients with depression and anxiety experience less improvement in PROs after arthroscopic management of FAIS,<sup>23,34</sup> data

The Orthopaedic Journal of Sports Medicine, 9(9), 23259671211036493  
DOI: 10.1177/23259671211036493  
© The Author(s) 2021

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

are lacking on the effect of mental health conditions on postoperative complication rates, revision hip arthroscopy rates, and conversion to total hip arthroplasty (THA).

The purpose of this study was to determine the prevalence of depression and related psychiatric disorders among patients undergoing hip arthroscopy for FAIS and to assess the association between preoperative depression and postoperative complications, readmissions, and revision hip arthroscopy. We hypothesized that patients with these psychiatric disorders undergoing hip arthroscopy would experience increased rates of 90-day postoperative complications and readmissions as well as increased risk of undergoing early revision hip arthroscopy (within 2 years of the index procedure) as compared with patients without these psychiatric comorbidities.

## METHODS

Using PearlDiver (PearlDiver Technologies), we searched for deidentified data from the Mariner database, which comprises 122 million patient lives from multiple insurance payer types, including private insurance, Medicare, Medicaid, government, and cash. A retrospective cohort study was performed using patient data between 2010 and 2019. Complete patient cohort allocation is outlined in Figure 1.

### Identification of Arthroscopic Hip, Knee, and Shoulder Procedures

Arthroscopic hip procedures for FAIS were identified using Current Procedural Terminology (CPT) codes 29914, 29915, 29916, 29860, 29861, 29862, and 29863. Patients who underwent hip arthroscopy with an associated encounter code for septic arthritis or hip trauma, including acetabular fracture, femoral head fracture, intra-articular foreign/loose bodies, and hip dislocation, were excluded (Appendix Table A1). Patients with a preexisting ipsilateral artificial hip joint before hip arthroscopy were also excluded. Arthroscopic knee and shoulder procedures were also identified using CPT codes in order to compare lifetime prevalence of preoperative depression (Appendix Table A2).

### Identification of Depression and Related Psychiatric Disorders

A diagnosis of depression or related psychiatric disorders was identified using International Classification of Diseases, Ninth Revision (ICD-9) and Tenth Revision (ICD-10), codes (Appendix Table A3). Patients with a reported diagnosis code within the year preceding hip arthroscopy were included in

the “depression” group (experimental group), while patients with no such diagnosis preceding hip arthroscopy were included in the “no depression” group (control group). Patients without at least 1 year of records before an arthroscopic hip procedure were excluded to reduce selection bias.

Baseline characteristics were collected for the experimental and control groups. To control for baseline differences, the groups were matched by age, sex, body mass index, tobacco use, and Charlson Comorbidity Index (CCI). The CCI is a widely used and validated adjustment index that accounts for multiple comorbidities to provide an overall assessment of a patient’s health.<sup>6</sup>

### Identification of 90-Day Readmissions, Postoperative Complications, and Revision Hip Procedures

Patient outcomes, including readmission and medical complications, were tracked for 90 days after hip arthroscopy. Appendix Table A4 provides a comprehensive list of tracked complications with their corresponding ICD-9 and ICD-10 codes.

Revision hip arthroscopy was defined as any hip arthroscopy performed on the ipsilateral hip after the index procedure; laterality of the index and revision procedures was defined via ICD-10 codes for hip pain and dysfunction (Appendix Table A5). Conversion to THA was defined as an ipsilateral THA after the index procedure; laterality was defined as in revision hip arthroscopy. Given that these laterality codes were implemented in 2015 with the advent of ICD-10, analysis of revision hip arthroscopy and conversion THA was limited to the period between 2015 and 2018.

Patients with a diagnosis of depression or related psychiatric disorder who underwent revision hip arthroscopy or conversion to THA within 2 years were included in the “depression” group (experimental group), while patients without such diagnoses who underwent these procedures within the defined times were included in the “no depression” group (control group). Patients without at least 2 years of records after an arthroscopic hip procedure were excluded to reduce selection bias.

### Statistical Analysis

Rates of postoperative readmission, complications, and revision hip arthroscopy were compared using chi-square analysis; odds ratios (ORs) and 95% confidence intervals (CIs) were calculated. Statistical significance was set at  $P < .5$  for all tests. Statistical tests were performed with the R Statistical Package.


\*Address correspondence to Ryan D. Freshman, MD, Department of Orthopedic Surgery, University of California–San Francisco, 500 Parnassus Avenue, MU W320, San Francisco, CA 94143, USA (email: ryan.freshman@ucsf.edu).

<sup>†</sup>Department of Orthopedic Surgery, University of California–San Francisco, San Francisco, California, USA.

Final revision submitted May 1, 2021; accepted May 13, 2021.

One or more of the authors has declared the following potential conflict of interest or source of funding: D.A.L. has received grants from Arthrex, Smith & Nephew, and Evolution Surgical. A.L.Z. has received education payments from Arthrex, grants from Zimmer, and consulting fees from DePuy Mitek and Stryker. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was waived by the University of California–San Francisco (21-33336).



**Figure 1.** CONSORT diagram outlining study cohort allocations. CPT, Current Procedural Terminology; FAIS, femoroacetabular impingement syndrome; ICD-10, International Classification of Diseases, Tenth Revision.

**RESULTS**


Among the entire Mariner population, 52,352 patients underwent elective hip arthroscopy for FAIS between 2010 and 2019. The lifetime and 1-year preprocedure prevalence of depression in patients undergoing hip arthroscopy was 25.4% and 14.5%, respectively, as compared with 19.8% and 11.6% in patients undergoing elective knee arthroscopy and 22.2% and 12.9% in patients undergoing elective shoulder arthroscopy ( $P < .001$  for all intergroup comparisons). Between 2011 and 2018, the prevalence of depression in patients undergoing elective hip arthroscopy increased from 11.6% to 18.9% (Figure 2).

Characteristics of the study groups are listed in Table 1. Before matching, patients with depression were more likely to be older, female, and obese and more likely to use tobacco and have a higher CCI.

**90-Day Readmissions and Complications**

The matched analysis for 90-day readmissions and complications consisted of 5814 patients with a diagnosis of depression or related psychiatric disorder within 1 year of hip arthroscopy and 5634 patients without such diagnoses (Table 2).

When comparing these matched groups, we found a significantly increased odds of 90-day readmission in the depression group (OR, 1.59 [95% CI, 1.21-2.10];  $P < .001$ ). The depression group also had increased odds of multiple 90-day postoperative medical complications, such as urinary tract infection (OR, 1.39 [95% CI, 1.29-1.51];  $P < .001$ ), pneumonia (OR, 1.49 [95% CI, 1.32-1.67];  $P < .001$ ), hematoma formation (OR, 1.64 [95% CI, 1.29-2.08];  $P < .001$ ), acute kidney injury (OR, 1.58 [95% CI, 1.28-1.95];  $P < .001$ ), deep venous thrombosis/pulmonary



**Figure 2.** Year-by-year percentage of patients undergoing elective hip arthroscopy with a recorded diagnosis of depression within 1 year before surgery.

**TABLE 1**  
Descriptive Data of Unmatched Study Cohorts Undergoing Hip Arthroscopy for FAIS<sup>a</sup>

	Depression (n = 6054)	No Depression (n = 31,526)
Age, y	41.1 ± 13.5	39.8 ± 15.2
Female sex	4838 (79.9)	21,389 (67.8)
Obese	2165 (35.8)	6407 (20.3)
Tobacco use	2053 (33.9)	5087 (16.1)
CCI	0.64 ± 1.27	0.38 ± 0.92

<sup>a</sup>Data are reported as mean ± SD or n (%). Difference between cohorts for each variable:  $P < .001$ . CCI, Charlson Comorbidity Index; FAIS, femoroacetabular impingement syndrome.

embolism (OR, 1.51 [95% CI, 1.16-1.95];  $P = .002$ ), and superficial wound infection (OR, 1.76 [95% CI, 1.44-2.14];  $P < .001$ ) (Table 3).

### Revision Hip Arthroscopy and Conversion to THA

The matched analysis for early revision hip arthroscopy and conversion to THA (as defined by ipsilateral hip arthroscopy or THA within 2 years of the index procedure) consisted of 1513 patients with a diagnosis of depression or related psychiatric disorder within 1 year of hip arthroscopy and 1513 patients without such diagnoses. The overall rates of early revision hip arthroscopy and conversion to THA were 4.4% and 3.1%, respectively. The depression group had increased odds of early revision hip arthroscopy (OR, 2.78 [95% CI, 1.88-4.11];  $P < .001$ ) but not early conversion to THA ( $P = .40$ ) (Table 4).

## DISCUSSION

This study demonstrated that a diagnosis of depression or related psychiatric disorders is correlated with increased

**TABLE 2**  
Descriptive Data of Matched Study Cohorts Undergoing Hip Arthroscopy for FAIS<sup>a</sup>

	Depression (n = 5814)	No Depression (n = 5634)	P Value
Age, y	41.0 ± 13.4	41.0 ± 13.4	.83
Female sex	4728 (80.1)	4728 (80.1)	≥.99
Obese	2055 (34.8)	2055 (34.8)	≥.99
Tobacco use	1954 (33.1)	1954 (33.1)	≥.99
CCI	0.53 ± 0.97	0.53 ± 0.97	≥.99

<sup>a</sup>Data are reported as mean ± SD or n (%). CCI, Charlson Comorbidity Index; FAIS, femoroacetabular impingement syndrome.

risk for multiple short-term postoperative complications after hip arthroscopy for FAIS. In addition, depression was associated with increased risk of undergoing ipsilateral revision hip arthroscopy within 2 years of primary arthroscopy.

In our study, the lifetime prevalence of preoperative depression in patients undergoing elective hip arthroscopy was 25.4%. It has been reported that up to 35% of patients undergoing hip arthroscopy carry a diagnosis of anxiety or depression,<sup>10</sup> as compared with 20% of patients undergoing total knee arthroplasty<sup>9</sup> and 12% of patients undergoing shoulder arthroplasty.<sup>27</sup> In a large database study, Iglinski-Benjamin et al<sup>14</sup> found that depression was the most common comorbid psychiatric diagnosis in patients undergoing hip arthroscopy and that patients in this cohort had a higher incidence of psychiatric conditions than did patients undergoing anterior cruciate ligament reconstruction or shoulder stabilization surgery.

While the prevalence of depression in patients with FAIS has been well documented, the relationship between FAIS and depression is not well understood but could stem from a variety of patient-related factors, such as the high costs and significant wait time between onset of symptoms and diagnosis of a labral tear.<sup>17</sup> It is also possible that patients with depression have increased sensitivity to somatic injuries,<sup>7</sup> which can lead to amplified FAI symptoms necessitating surgical intervention. Despite these gaps in the literature, prior studies have shown that depression and anxiety are associated with higher rates of opioid use and increased costs of care for patients undergoing hip arthroscopy.<sup>2,15</sup> Given our findings that patients with preoperative depression experience higher rates of postoperative complications, the prevalence of depression in patients undergoing hip arthroscopy has the possibility to represent a significant burden to our health care system.

While a connection between mental health and PROs after hip arthroscopy for FAIS has been demonstrated,<sup>8,21,23,24</sup> our study is the first to our knowledge to show how mental health may affect postoperative complication and readmission rates in a large population of patients undergoing hip arthroscopy. Although this concept is novel to the field of hip arthroscopy, it has been

TABLE 3  
Comparison of 90-Day Postoperative Complications After Hip Arthroscopy

	Complications, No. (%)		Odds Ratio (95% CI)	P Value <sup>a</sup>
	Depression (n = 5814)	No Depression (n = 5634)		
Readmission within 90 d	137 (2.4)	84 (1.5)	1.59 (1.21-2.10)	<b>&lt;.001</b>
Urinary tract infection	2103 (36.2)	1630 (28.9)	1.39 (1.29-1.51)	<b>&lt;.001</b>
Pneumonia	751 (12.9)	511 (9.1)	1.49 (1.32-1.67)	<b>&lt;.001</b>
Hematoma formation	182 (3.1)	109 (1.9)	1.64 (1.29-2.08)	<b>&lt;.001</b>
Superficial wound infection	284 (4.9)	160 (2.8)	1.76 (1.44-2.14)	<b>&lt;.001</b>
Acute kidney injury	235 (4.0)	146 (2.6)	1.58 (1.28-1.95)	<b>&lt;.001</b>
Deep venous thrombosis/pulmonary embolism	151 (2.6)	98 (1.7)	1.51 (1.16-1.95)	<b>.002</b>
Cardiac arrest	29 (0.5)	25 (0.4)	1.12 (0.66-1.92)	.77
Lower extremity nerve injury	31 (0.5)	36 (0.6)	0.83 (0.51-1.35)	.54
Heterotopic ossification	34 (0.6)	23 (0.4)	1.44 (0.84-2.44)	.23
Septic arthritis	19 (0.3)	17 (0.3)	1.08 (0.56-2.09)	.94
Need for transfusion	111 (1.9)	88 (1.6)	1.23 (0.93-1.63)	.18

<sup>a</sup>Bold *P* values indicate statistically significant difference between cohorts (*P* < .05).

TABLE 4  
Comparison of Additional Ipsilateral Hip Procedures Within 2 Years

	Additional Procedures, No. (%)		Odds Ratio (95% CI)	P Value <sup>a</sup>
	Depression (n = 1513)	No Depression (n = 1513)		
Revision hip arthroscopy	96 (6.3)	36 (2.4)	2.78 (1.88-4.11)	<b>&lt;.001</b>
Conversion to total hip arthroplasty	43 (2.8)	52 (3.4)	0.82 (0.55-1.24)	.40

<sup>a</sup>Bold *P* value indicates statistically significant difference between cohorts (*P* < .05).

demonstrated in patients with depression undergoing hip, knee, and shoulder arthroplasty,<sup>27,31</sup> who have been shown to be at increased risk of postoperative anemia, renal failure, and prosthetic joint infection. The mechanisms underlying this interaction are likely multifactorial but may include the upregulation of inflammatory cytokines (eg, tumor necrosis factor  $\alpha$ , interleukin 1 $\beta$ , and the cyclo-oxygenase 2 pathway), which has been observed in depression and other psychiatric diagnoses.<sup>25,29,30</sup> High levels of these cytokines may lead to an inflammatory condition in the perioperative period and thus negatively affect patient outcomes. A diagnosis of depression itself may also indicate worse patient health and higher rates of chronic medical illness, which in turn could contribute to an increased risk of postoperative complications in patients with depression.<sup>18</sup>

Despite our knowledge of the relationship between depression and patient outcomes, it is unknown how exactly preoperative treatment for depression affects postoperative outcomes after hip arthroscopy. While there is mixed evidence surrounding this concept within the arthroplasty literature,<sup>13,20</sup> no studies to date have examined this topic in a population of patients undergoing hip arthroscopy; thus, further investigation is warranted to optimize patient outcomes and reduce postoperative complications.

In our study, patients with depression were nearly 3 times more likely to undergo revision hip arthroscopy within 2 years of the index procedure. These patients underwent

similar rates of conversion to THA as compared with controls likely because we included age (a good surrogate for Tönnis grade) in our matching criteria. The overall rate of early revision hip arthroscopy in our study (4.4%) is consistent with rates reported in the literature.<sup>19,36</sup> While radiographic and intraoperative findings such as residual FAI are cited as common reasons for revision hip arthroscopy, few studies have examined patient characteristic factors associated with revision hip arthroscopy. Kester et al<sup>19</sup> and West et al<sup>36</sup> identified patient age, female sex, and index procedure performed by a low-volume surgeon as risk factors for undergoing revision hip arthroscopy within 2 years. Although studies have associated these and other numerous risk factors with early conversion from hip arthroscopy to THA,<sup>1,32</sup> previous studies to our knowledge have not reported an association between depression and revision hip arthroscopy.

One potential explanation for our findings is lower pre- and postoperative PROs disclosed by patients with depression and other mental health diagnoses. Although these patients can achieve the minimal clinically important difference for the Hip Outcome Score and modified Harris Hip Score,<sup>23</sup> Shapira et al<sup>33</sup> demonstrated a threshold effect for the modified Harris Hip Score at 1 year after hip arthroscopy that predicted need for a secondary hip procedure. Depression also has a synergistic relationship with pain,<sup>12</sup> and unimproved or worsening postoperative pain may influence a surgeon's decision to perform revision hip arthroscopy. Future investigation into the risk factors and decision-

making processes surrounding revision hip arthroscopy in the setting of depression is necessary to help surgeons counsel patients considering revision hip arthroscopy and to prevent unnecessary revision procedures in patients with continued symptoms despite unclear surgical indications.

The findings of our study should be interpreted in the context of its limitations. The validity of our results is reliant on accurate ICD-10 and CPT coding, although specific inclusion and exclusion criteria were applied to minimize potential misclassification. Given the limitations of the Mariner database, we were not able to directly control for multiple variables, such as the severity of depression symptoms, pharmacologic treatment of depression, radiographic measurements of FAI, labral tear morphology, degree of preexisting osteoarthritis, and completeness of impingement correction. The Mariner database does not provide granular data on preoperative function, intraoperative findings, or surgical technique, which limited us to perform a matched-cohort analysis instead of a multivariate logistic regression analysis. Additionally, we were unable to evaluate pre- and postoperative PROs to investigate possible correlation between outcome scores and revision hip arthroscopy. Finally, given the retrospective nature of this database study, it is important to note that we were able to report only on correlations between depression and incidence of postoperative complications; determination of causation lies outside the scope of this study.

## CONCLUSION

Patients with a diagnosis of depression or similarly related psychiatric disorders experience higher rates of 90-day complications and hospital readmissions after elective hip arthroscopy for FAIS and are more likely to undergo revision hip arthroscopy within 2 years of the index procedure. Given the prevalence of depression in this population, mental health screening should be considered part of a standard preoperative patient assessment.

## ACKNOWLEDGMENT

The authors thank Scott Ellison for his assistance with database coding and data collection.

## REFERENCES

- Allahabadi S, Hinman AD, Horton BH, Avins AL, Coughlan MJ, Ding DY. Risk factors for conversion of hip arthroscopy to total hip arthroplasty: a large closed-cohort study. *Arthrosc Sports Med Rehabil*. 2020;2(5):e599-e605.
- Baron JE, Khazi ZM, Duchman KR, Wolf BR, Westermann RW. Increased prevalence and associated costs of psychiatric comorbidities in patients undergoing sports medicine operative procedures. *Arthroscopy*. 2021;37(2):686-693.
- Bozic KJ, Chan V, Valone FH III, Feeley BT, Vail TP. Trends in hip arthroscopy utilization in the United States. *J Arthroplasty*. 2013; 28(8 suppl):140-143.
- Byrd JW, Jones KS. Arthroscopic management of femoroacetabular impingement: minimum 2-year follow-up. *Arthroscopy*. 2011;27(10): 1379-1388.
- Byrd JW, Jones KS, Gwathmey FW. Femoroacetabular impingement in adolescent athletes: outcomes of arthroscopic management. *Am J Sports Med*. 2016;44(8):2106-2111.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383.
- Chiu YH, Silman AJ, Macfarlane GJ, et al. Poor sleep and depression are independently associated with a reduced pain threshold: results of a population based study. *Pain*. 2005;115(3):316-321.
- Dick AG, Smith C, Bankes MJK, George M. The impact of mental health disorders on outcomes following hip arthroscopy for femoroacetabular impingement syndrome: a systematic review. *J Hip Preserv Surg*. 2020;7(2):195-204.
- Duivenvoorden T, Vissers MM, Verhaar JA, et al. Anxiety and depressive symptoms before and after total hip and knee arthroplasty: a prospective multicentre study. *Osteoarthritis Cartilage*. 2013;21(12): 1834-1840.
- Dwyer T, Whelan D, Shah PS, Ajrawat P, Hoit G, Chahal J. Operative versus nonoperative treatment of femoroacetabular impingement syndrome: a meta-analysis of short-term outcomes. *Arthroscopy*. 2020;36(1):263-273.
- Griffin DR, Dickenson EJ, Wall PDH, et al. Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHION): a multicentre randomised controlled trial. *Lancet*. 2018;391(10136):2225-2235.
- Gureje O. Treating chronic pain in the context of comorbid depression. *Pain*. 2008;134(1-2):3-4.
- Halawi MJ, Gronbeck C, Savoy L, Cote MP, Lieberman JR. Depression treatment is not associated with improved patient-reported outcomes following total joint arthroplasty. *J Arthroplasty*. 2020;35(1): 28-31.
- Iglinski-Benjamin KC, Xiao M, Safran MR, Abrams GD. Increased prevalence of concomitant psychiatric diagnoses among patients undergoing hip arthroscopic surgery. *Orthop J Sports Med*. 2019; 7(1):2325967118822451.
- Jacobs CA, Hawk GS, Jochimsen KN, et al. Depression and anxiety are associated with increased health care costs and opioid use for patients with femoroacetabular impingement undergoing hip arthroscopy: analysis of a claims database. *Arthroscopy*. 2020;36(3): 745-750.
- Jochimsen KN, Magnuson JA, Kocan KR, et al. Anxiety and depression are associated with lower preoperative quality of life and function but not duration of symptoms in patients with femoroacetabular impingement syndrome. *J Hip Preserv Surg*. 2019;6(3):207-213.
- Kahlenberg CA, Han B, Patel RM, Deshmene PP, Terry MA. Time and cost of diagnosis for symptomatic femoroacetabular impingement. *Orthop J Sports Med*. 2014;2(3):2325967114523916.
- Katon WJ. Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues Clin Neurosci*. 2011;13(1):7-23.
- Kester BS, Capogna B, Mahure SA, Ryan MK, Mollon B, Youm T. Independent risk factors for revision surgery or conversion to total hip arthroplasty after hip arthroscopy: a review of a large statewide database from 2011 to 2012. *Arthroscopy*. 2018;34(2):464-470.
- Kohring JM, Erickson JA, Anderson MB, Gililand JM, Peters CL, Pelt CE. Treated versus untreated depression in total joint arthroplasty impacts outcomes. *J Arthroplasty*. 2018;33(7):S81-S85.
- Kuroda Y, Saito M, Cinar EN, Norrish A, Khanduja V. Patient-related risk factors associated with less favourable outcomes following hip arthroscopy. *Bone Joint J*. 2020;102(7):822-831.
- Kyin C, Maldonado DR, Go CC, Shapira J, Lall AC, Domb BG. Mid to long-term outcomes of hip arthroscopy: a systematic review. *Arthroscopy*. 2021;37(3):1011-1025.
- Lansdown DA, Ukwuani G, Kuhns B, Harris JD, Nho SJ. Self-reported mental disorders negatively influence surgical outcomes after arthroscopic treatment of femoroacetabular impingement. *Orthop J Sports Med*. 2018;6(5):2325967118773312.
- Martin RL, Christoforetti JJ, McGovern R, et al. The impact of depression on patient outcomes in hip arthroscopic surgery. *Orthop J Sports Med*. 2018;6(11):2325967118806490.

25. Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol Psychiatry*. 2009;65(9):732-741.

26. Minkara AA, Westermann RW, Rosneck J, Lynch TS. Systematic review and meta-analysis of outcomes after hip arthroscopy in femoroacetabular impingement. *Am J Sports Med*. 2019;47(2):488-500.

27. Mollon B, Mahure SA, Ding DY, Zuckerman JD, Kwon YW. The influence of a history of clinical depression on peri-operative outcomes in elective total shoulder arthroplasty: a ten-year national analysis. *Bone Joint J*. 2016;98B(6):818-824.

28. Philippon MJ, Maxwell RB, Johnston TL, Schenker M, Briggs KK. Clinical presentation of femoroacetabular impingement. *Knee Surg Sports Traumatol Arthrosc*. 2007;15(8):1041-1047.

29. Pollak Y, Yirmiya R. Cytokine-induced changes in mood and behaviour: implications for "depression due to a general medical condition," immunotherapy and antidepressive treatment. *Int J Neuropsychopharmacol*. 2002;5(4):389-399.

30. Postal M, Appenzeller S. The importance of cytokines and autoantibodies in depression. *Autoimmun Rev*. 2015;14(1):30-35.

31. Rasouli MR, Menendez ME, Sayadipour A, Purtill JJ, Parvizi J. Direct cost and complications associated with total joint arthroplasty in patients with preoperative anxiety and depression. *J Arthroplasty*. 2016;31(2):533-536.

32. Redmond JM, Gupta A, Dunne K, Humayun A, Yuen LC, Domb BG. What factors predict conversion to THA after arthroscopy? *Clin Orthop Relat Res*. 2017;475(10):2538-2545.

33. Shapira J, Bheem R, Kyin C, et al. Can patient-reported outcomes predict the need for secondary surgeries after hip arthroscopy? *Am J Sports Med*. 2021;49(1):97-103.

34. Sochacki KR, Brown L, Cenkus K, Di Stasi S, Harris JD, Ellis TJ. Preoperative depression is negatively associated with function and predicts poorer outcomes after hip arthroscopy for femoroacetabular impingement. *Arthroscopy*. 2018;34(8):2368-2374.

35. Weber AE, Nakata H, Mayer EN, et al. Return to sport after hip arthroscopy for femoroacetabular impingement syndrome in NCAA Division I athletes: experience at a single institution. *Orthop J Sports Med*. 2020;8(5):2325967120918383.

36. West CR, Bedard NA, Duchman KR, Westermann RW, Callaghan JJ. Rates and risk factors for revision hip arthroscopy. *Iowa Orthop J*. 2019;39(1):95-99.

APPENDIX

TABLE A1  
ICD-9 and ICD-10 Codes for Exclusion Criteria<sup>a</sup>

Hip trauma

ICD-10-D-M24051, ICD-10-D-M24351, ICD-10-D-M24352, ICD-10-D-M24451, ICD-10-D-M24452, ICD-10-D-M247, ICD-10-D-S32401A, ICD-10-D-S32401B, ICD-10-D-S32401D, ICD-10-D-S32401G, ICD-10-D-S32401 K, ICD-10-D-S32401 S, ICD-10-D-S32402A, ICD-10-D-S32402B, ICD-10-D-S32402D, ICD-10-D-S32402G, ICD-10-D-S32402 K, ICD-10-D-S32402 S, ICD-10-D-S32409A, ICD-10-D-S32409B, ICD-10-D-S32409D, ICD-10-D-S32409G, ICD-10-D-S32409 K, ICD-10-D-S32409 S, ICD-10-D-S32411A, ICD-10-D-S32411B, ICD-10-D-S32411D, ICD-10-D-S32411 K, ICD-10-D-S32411 S, ICD-10-D-S32412A, ICD-10-D-S32412D, ICD-10-D-S32412 S, ICD-10-D-S32413A, ICD-10-D-S32413D, ICD-10-D-S32413 S, ICD-10-D-S32414A, ICD-10-D-S32414D, ICD-10-D-S32414 S, ICD-10-D-S32415A, ICD-10-D-S32415D, ICD-10-D-S32415 S, ICD-10-D-S32416A, ICD-10-D-S32416D, ICD-10-D-S32421A, ICD-10-D-S32421B, ICD-10-D-S32421D, ICD-10-D-S32421 K, ICD-10-D-S32421 S, ICD-10-D-S32422A, ICD-10-D-S32422B, ICD-10-D-S32422D, ICD-10-D-S32422G, ICD-10-D-S32422 K, ICD-10-D-S32422 S, ICD-10-D-S32423A, ICD-10-D-S32423D, ICD-10-D-S32424A, ICD-10-D-S32424D, ICD-10-D-S32424G, ICD-10-D-S32424 S, ICD-10-D-S32425A, ICD-10-D-S32425D, ICD-10-D-S32425 S, ICD-10-D-S32426A, ICD-10-D-S32431A, ICD-10-D-S32431B, ICD-10-D-S32431D, ICD-10-D-S32431G, ICD-10-D-S32431 K, ICD-10-D-S32432A, ICD-10-D-S32432B, ICD-10-D-S32432D, ICD-10-D-S32432G, ICD-10-D-S32432 K, ICD-10-D-S32432 S, ICD-10-D-S32433A, ICD-10-D-S32433D, ICD-10-D-S32433 S, ICD-10-D-S32434A, ICD-10-D-S32434D, ICD-10-D-S32434G, ICD-10-D-S32434 S, ICD-10-D-S32435A, ICD-10-D-S32435B, ICD-10-D-S32435D, ICD-10-D-S32435 K, ICD-10-D-S32435 S, ICD-10-D-S32436A, ICD-10-D-S32492A, ICD-10-D-S32492B, ICD-10-D-S32492D, ICD-10-D-S32492G, ICD-10-D-S32492 K, ICD-10-D-S32492 S, ICD-10-D-S32499A, ICD-10-D-S32499D, ICD-10-D-S32499 S, ICD-10-D-S72051A, ICD-10-D-S72051B, ICD-10-D-S72051D, ICD-10-D-S72051G, ICD-10-D-S72051 K, ICD-10-D-S72051 S, ICD-10-D-S72052A, ICD-10-D-S72052D, ICD-10-D-S72052G, ICD-10-D-S72052 H, ICD-10-D-S72052 J, ICD-10-D-S72052 K, ICD-10-D-S72052P, ICD-10-D-S72052 S, ICD-10-D-S72059A, ICD-10-D-S72059C, ICD-10-D-S72059D, ICD-10-D-S72059 S, ICD-10-D-S72061A, ICD-10-D-S72061B, ICD-10-D-S72061D, ICD-10-D-S72061E, ICD-10-D-S72061 H, ICD-10-D-S72061P, ICD-10-D-S72061 S, ICD-10-D-S72062A, ICD-10-D-S72062D, ICD-10-D-S72062E, ICD-10-D-S72062G, ICD-10-D-S72062 S, ICD-10-D-S72063A, ICD-10-D-S72064A, ICD-10-D-S72064C, ICD-10-D-S72064D, ICD-10-D-S72064F, ICD-10-D-S72064 S, ICD-10-D-S72065A, ICD-10-D-S72065D, ICD-10-D-S72065 S, ICD-10-D-S72091A, ICD-10-D-S72091B, ICD-10-D-S72091C, ICD-10-D-S72091D, ICD-10-D-S72091G, ICD-10-D-S72091 K, ICD-10-D-S72091 M, ICD-10-D-S72091P, ICD-10-D-S72091 S, ICD-10-D-S72092A, ICD-10-D-S72092B, ICD-10-D-S72092C, ICD-10-D-S72092D, ICD-10-D-S72092E, ICD-10-D-S72092G, ICD-10-D-S72092 H, ICD-10-D-S72092 K, ICD-10-D-S72092 M, ICD-10-D-S72092 N, ICD-10-D-S72092P, ICD-10-D-S72092 S, ICD-10-D-S72099A, ICD-10-D-S72099B, ICD-10-D-S72099D, ICD-10-D-S72099F, ICD-10-D-S72099 K, ICD-10-D-S72099 S, ICD-10-D-S73004A, ICD-10-D-S73004D, ICD-10-D-S73004 S, ICD-10-D-S73005A, ICD-10-D-S73005D, ICD-10-D-S73005 S, ICD-10-D-S73014A, ICD-10-D-S73014D, ICD-10-D-S73014 S, ICD-10-D-S73015A, ICD-10-D-S73015D, ICD-10-D-S73024D, ICD-10-D-S73025A, ICD-10-D-S73034A, ICD-10-D-S73034D, ICD-10-D-S73034 S, ICD-10-D-S73035A, ICD-10-D-S73035D, ICD-10-D-S73035 S, ICD-10-D-S73044A, ICD-10-D-S73044D, ICD-10-D-S73044 S, ICD-10-D-S73045A, ICD-10-D-S73045D, ICD-9-D-71865, ICD-9-D-8080, ICD-9-D-8081, ICD-9-D-83500, ICD-9-D-83501, ICD-9-D-83502, ICD-9-D-83503

Preexisting hip arthroplasty

ICD-10-D-Z96641, ICD-10-D-Z96642, ICD-10-D-Z96643, ICD-10-D-Z96649, ICD-9-D-V4364

Septic arthritis

ICD-10-D-M00051, ICD-10-D-M00052, ICD-10-D-M00251, ICD-10-D-M00252, ICD-10-D-M00851, ICD-10-D-M00852, ICD-10-D-M00859, ICD-10-D-M0089, ICD-10-D-M009, ICD-10-D-M01X51, ICD-10-D-M01X52, ICD-10-D-M01X59, ICD-9-D-71100, ICD-9-D-71105, ICD-9-D-71108, ICD-9-D-71109

<sup>a</sup>ICD-9, International Classification of Diseases, Ninth Revision; ICD-10, International Classification of Diseases, Tenth Revision.



TABLE A2  
Codes for arthroscopic shoulder and knee procedures

---

Arthroscopic Shoulder procedures

CPT-29805, CPT-29806, CPT-29807, CPT-29819, CPT-29820, CPT-29821, CPT-29822, CPT-29823, CPT-29824, CPT-29825, CPT-29826, CPT-29827, CPT-29828, CPT-S2300

Arthroscopic Knee Procedures

CPT-29850, CPT-29851, CPT-29866, CPT-29867, CPT-29868, CPT-29870, CPT-29871, CPT-29873, CPT-29874, CPT-29875, CPT-29876, CPT-29877, CPT-29879, CPT-29880, CPT-29881, CPT-29882, CPT-29883, CPT-29884, CPT-29885, CPT-29886, CPT-29887, CPT-G0289, CPT-S2112

---

TABLE A3  
ICD-9 and ICD-10 Codes for Depression and Related Psychiatric Diagnoses<sup>a</sup>

---

Depression and related diagnoses

ICD-10-D-F3130, ICD-10-D-F3131, ICD-10-D-F3132, ICD-10-D-F314, ICD-10-D-F315, ICD-10-D-F320, ICD-10-D-F321, ICD-10-D-F322, ICD-10-D-F323, ICD-10-D-F324, ICD-10-D-F325, ICD-10-D-F328, ICD-10-D-F3281, ICD-10-D-F3289, ICD-10-D-F329, ICD-10-D-F330, ICD-10-D-F331, ICD-10-D-F332, ICD-10-D-F333, ICD-10-D-F3340, ICD-10-D-F3341, ICD-10-D-F3342, ICD-10-D-F338, ICD-10-D-F339, ICD-10-D-F341, ICD-10-D-F4320, ICD-10-D-F4321, ICD-10-D-F4322, ICD-10-D-F4323, ICD-10-D-F4324, ICD-10-D-F4325, ICD-10-D-F4329, ICD-9-D-2962, ICD-9-D-29620, ICD-9-D-29621, ICD-9-D-29622, ICD-9-D-29623, ICD-9-D-29624, ICD-9-D-29625, ICD-9-D-29626, ICD-9-D-2963, ICD-9-D-29630, ICD-9-D-29631, ICD-9-D-29632, ICD-9-D-29633, ICD-9-D-29634, ICD-9-D-29635, ICD-9-D-29636, ICD-9-D-29651, ICD-9-D-29652, ICD-9-D-29653, ICD-9-D-29654, ICD-9-D-29655, ICD-9-D-29656, ICD-9-D-3004, ICD-9-D-3090, ICD-9-D-3091, ICD-9-D-30921, ICD-9-D-30922, ICD-9-D-30923, ICD-9-D-30924, ICD-9-D-30928, ICD-9-D-30929, ICD-9-D-3093, ICD-9-D-3094, ICD-9-D-3098, ICD-9-D-30981, ICD-9-D-30982, ICD-9-D-30983, ICD-9-D-30989, ICD-9-D-3099, ICD-9-D-311

---

<sup>a</sup>ICD-9, International Classification of Diseases, Ninth Revision; ICD-10, International Classification of Diseases, Tenth Revision.

TABLE A4  
ICD-9 and ICD-10 Codes for Tracked Complications<sup>a</sup>

---

Acute kidney injury

ICD-10-D-N170, ICD-10-D-N171, ICD-10-D-N172, ICD-10-D-N178, ICD-10-D-N179, ICD-9-D-5845, ICD-9-D-5846, ICD-9-D-5847, ICD-9-D-5848, ICD-9-D-5849

Cardiac arrest

ICD-10-D-I462, ICD-10-D-I468, ICD-10-D-I469, ICD-9-D-42741, ICD-9-D-4275

Deep venous thrombosis/pulmonary embolism

ICD-10-D-I2601, ICD-10-D-I2602, ICD-10-D-I2609, ICD-9-D-4151, ICD-9-D-41511, ICD-9-D-41512, ICD-9-D-41513, ICD-9-D-41519, ICD-10-D-I2690, ICD-10-D-I2692, ICD-10-D-I2694, ICD-10-D-I2699, ICD-9-D-4532, ICD-9-D-4533, ICD-9-D-4534, ICD-9-D-45382, ICD-9-D-45384, ICD-9-D-45385, ICD-9-D-45386

Hematoma

ICD-10-D-E89811, ICD-10-D-G9751, ICD-10-D-G9752, ICD-10-D-I97418, ICD-10-D-I97610, ICD-10-D-I97618, ICD-10-D-I97620, ICD-10-D-J9561, ICD-10-D-J95830, ICD-10-D-J95831, ICD-10-D-K9161, ICD-10-D-K91840, ICD-10-D-K91841, ICD-10-D-L7602, ICD-10-D-L7621, ICD-10-D-L7622, ICD-10-D-M96810, ICD-10-D-M96830, ICD-10-D-M96831, ICD-10-D-N99820, ICD-10-D-N99821, ICD-10-D-T888XXA, ICD-9-D-99811, ICD-9-D-99812, ICD-9-D-99813

---

(continued)

TABLE A4 (continued)

## Nerve injury

ICD-10-D-S7400XA, ICD-10-D-S7400XD, ICD-10-D-S7400XS, ICD-10-D-S7401XA, ICD-10-D-S7401XD, ICD-10-D-S7401XS, ICD-10-D-S7402XA, ICD-10-D-S7402XD, ICD-10-D-S7402XS, ICD-10-D-S7410XA, ICD-10-D-S7410XD, ICD-10-D-S7410XS, ICD-10-D-S7411XA, ICD-10-D-S7411XD, ICD-10-D-S7411XS, ICD-10-D-S7412XA, ICD-10-D-S7412XD, ICD-10-D-S7412XS, ICD-10-D-S7420XA, ICD-10-D-S7420XS, ICD-10-D-S7421XA, ICD-10-D-S7421XD, ICD-10-D-S7421XS, ICD-10-D-S7422XA, ICD-10-D-S7422XD, ICD-10-D-S7422XS, ICD-10-D-S748X1A, ICD-10-D-S748X1D, ICD-10-D-S748X2A, ICD-10-D-S748X2D, ICD-10-D-S748X9A, ICD-10-D-S748X9D, ICD-10-D-S7490XA, ICD-10-D-S7490XD, ICD-10-D-S7491XA, ICD-10-D-S7491XD, ICD-10-D-S7491XS, ICD-10-D-S7492XA, ICD-10-D-S7492XD, ICD-10-D-S7492XS, ICD-10-D-S8400XA, ICD-10-D-S8400XD, ICD-10-D-S8400XS, ICD-10-D-S8401XA, ICD-10-D-S8401XD, ICD-10-D-S8401XS, ICD-10-D-S8402XA, ICD-10-D-S8402XD, ICD-10-D-S8402XS, ICD-10-D-S8410XA, ICD-10-D-S8410XD, ICD-10-D-S8410XS, ICD-10-D-S8411XA, ICD-10-D-S8411XD, ICD-10-D-S8411XS, ICD-10-D-S8412XA, ICD-10-D-S8412XD, ICD-10-D-S8412XS, ICD-10-D-S8420XA, ICD-10-D-S8420XD, ICD-10-D-S8421XA, ICD-10-D-S8421XD, ICD-10-D-S8421XS, ICD-10-D-S8422XA, ICD-10-D-S8422XD, ICD-10-D-S8422XS, ICD-10-D-S84801A, ICD-10-D-S84801D, ICD-10-D-S84801 S, ICD-10-D-S84802A, ICD-10-D-S84802D, ICD-10-D-S84802 S, ICD-10-D-S84809A, ICD-10-D-S8490XA, ICD-10-D-S8490XD, ICD-10-D-S8490XS, ICD-10-D-S8491XA, ICD-10-D-S8491XD, ICD-10-D-S8491XS, ICD-10-D-S8492XA, ICD-10-D-S8492XD, ICD-10-D-S8492XS, ICD-10-D-S9400XA, ICD-10-D-S9400XD, ICD-10-D-S9400XS, ICD-10-D-S9401XA, ICD-10-D-S9401XD, ICD-10-D-S9401XS, ICD-10-D-S9402XA, ICD-10-D-S9402XD, ICD-10-D-S9402XS, ICD-10-D-S9410XA, ICD-10-D-S9410XD, ICD-10-D-S9410XS, ICD-10-D-S9411XA, ICD-10-D-S9411XD, ICD-10-D-S9411XS, ICD-10-D-S9412XA, ICD-10-D-S9412XD, ICD-10-D-S9412XS, ICD-10-D-S9420XA, ICD-10-D-S9420XD, ICD-10-D-S9420XS, ICD-10-D-S9421XA, ICD-10-D-S9421XD, ICD-10-D-S9421XS, ICD-10-D-S9422XA, ICD-10-D-S9422XD, ICD-10-D-S9422XS, ICD-10-D-S9430XA, ICD-10-D-S9430XD, ICD-10-D-S9431XA, ICD-10-D-S9431XD, ICD-10-D-S9431XS, ICD-10-D-S9432XA, ICD-10-D-S9432XD, ICD-10-D-S9432XS, ICD-10-D-S948X1A, ICD-10-D-S948X1D, ICD-10-D-S948X1 S, ICD-10-D-S948X2A, ICD-10-D-S948X2D, ICD-10-D-S948X2 S, ICD-10-D-S948X9A, ICD-10-D-S948X9D, ICD-10-D-S948X9 S, ICD-10-D-S9490XA, ICD-10-D-S9490XD, ICD-10-D-S9490XS, ICD-10-D-S9491XA, ICD-10-D-S9491XD, ICD-10-D-S9491XS, ICD-10-D-S9492XA, ICD-10-D-S9492XD, ICD-10-D-S9492XS, ICD-9-D-9560, ICD-9-D-9561, ICD-9-D-9562, ICD-9-D-9563, ICD-9-D-9564, ICD-9-D-9565, ICD-9-D-9566, ICD-9-D-9567, ICD-9-D-9568, ICD-9-D-9569

## Pneumonia

ICD-10-D-J120, ICD-10-D-J121, ICD-10-D-J122, ICD-10-D-J123, ICD-10-D-J1281, ICD-10-D-J1289, ICD-10-D-J129, ICD-10-D-J13, ICD-10-D-J14, ICD-10-D-J150, ICD-10-D-J151, ICD-10-D-J1520, ICD-10-D-J15211, ICD-10-D-J15212, ICD-10-D-J1529, ICD-10-D-J153, ICD-10-D-J154, ICD-10-D-J155, ICD-10-D-J156, ICD-10-D-J157, ICD-10-D-J158, ICD-10-D-J159, ICD-10-D-J160, ICD-10-D-J168, ICD-10-D-J17, ICD-10-D-J180, ICD-10-D-J181, ICD-10-D-J182, ICD-10-D-J188, ICD-10-D-J189, ICD-9-D-4800, ICD-9-D-4801, ICD-9-D-4802, ICD-9-D-4808, ICD-9-D-4809, ICD-9-D-481, ICD-9-D-4820, ICD-9-D-4821, ICD-9-D-48230, ICD-9-D-48232, ICD-9-D-48239, ICD-9-D-48240, ICD-9-D-48241, ICD-9-D-48242, ICD-9-D-48249, ICD-9-D-48281, ICD-9-D-48282, ICD-9-D-48283, ICD-9-D-48284, ICD-9-D-48289, ICD-9-D-4829, ICD-9-D-4830, ICD-9-D-4831, ICD-9-D-4838, ICD-9-D-4843, ICD-9-D-4846, ICD-9-D-4848, ICD-9-D-485, ICD-9-D-486

## Septic arthritis

ICD-10-D-M00051, ICD-10-D-M00052, ICD-10-D-M00251, ICD-10-D-M00252, ICD-10-D-M00851, ICD-10-D-M00852, ICD-10-D-M00859, ICD-10-D-M0089, ICD-10-D-M009, ICD-10-D-M01X51, ICD-10-D-M01X52, ICD-10-D-M01X59, ICD-9-D-71100, ICD-9-D-71105, ICD-9-D-71108, ICD-9-D-71109

## Superficial wound infection

ICD-10-D-T8131XA, ICD-10-D-T8131XD, ICD-10-D-T8131XS, ICD-10-D-T8141XA, ICD-10-D-T8141XD, ICD-10-D-T8141XS, ICD-9-D-99832, ICD-9-D-99851, ICD-9-D-99859, ICD-9-D-99883

## Transfusion

ICD-10-P-30230N0, ICD-10-P-30233H0, ICD-10-P-30233H1, ICD-10-P-30233J1, ICD-10-P-30233K1, ICD-10-P-30233L1, ICD-10-P-30233N0, ICD-10-P-30233N1, ICD-10-P-30233P1, ICD-10-P-30233R1, ICD-10-P-30233S1, ICD-10-P-30233V1, ICD-10-P-30243H0, ICD-10-P-30243K1, ICD-10-P-30243N1, ICD-10-P-30243R1, ICD-10-P-30243S1, ICD-10-P-30243Y0, ICD-10-P-30253N0, ICD-10-P-30253N1, ICD-10-P-30263G1, ICD-10-P-30263N1, ICD-10-P-30273N1, ICD-10-P-30283B1, ICD-9-P-9904

## Urinary tract infection

ICD-10-D-N390, ICD-9-D-5990

<sup>a</sup>ICD-9, International Classification of Diseases, Ninth Revision; ICD-10, International Classification of Diseases, Tenth Revision.

TABLE A5

ICD-10 Codes for Revision Procedures: Laterality Modifiers and Total Hip Arthroplasty Procedures<sup>a</sup>

## Laterality modifiers

ICD-10-D-M24152, ICD-10-D-M24852, ICD-10-D-M25852, ICD-10-D-S73192A, ICD-10-D-S73192D, ICD-10-D-S73192 S, ICD-10-D-M24151, ICD-10-D-M24851, ICD-10-D-M25851, ICD-10-D-S73191A, ICD-10-D-S73191D, ICD-10-D-S73191S

## Total hip arthroplasty procedures

ICD-10-P-0SRB019, ICD-10-P-0SRB01A, ICD-10-P-0SRB01Z, ICD-10-P-0SRB029, ICD-10-P-0SRB02A, ICD-10-P-0SRB02Z, ICD-10-P-0SRB039, ICD-10-P-0SRB03A, ICD-10-P-0SRB03Z, ICD-10-P-0SRB049, ICD-10-P-0SRB04A, ICD-10-P-0SRB04Z, ICD-10-P-0SRB069, ICD-10-P-0SRB06A, ICD-10-P-0SRB06Z, ICD-10-P-0SRB07Z, ICD-10-P-0SRB0J9, ICD-10-P-0SRB0JA, ICD-10-P-0SRB0JZ, ICD-10-P-0SRB0KZ, ICD-10-P-0SR9019, ICD-10-P-0SR901A, ICD-10-P-0SR901Z, ICD-10-P-0SR9029, ICD-10-P-0SR902A, ICD-10-P-0SR902Z, ICD-10-P-0SR9039, ICD-10-P-0SR903A, ICD-10-P-0SR903Z, ICD-10-P-0SR9049, ICD-10-P-0SR904A, ICD-10-P-0SR904Z, ICD-10-P-0SR9069, ICD-10-P-0SR906A, ICD-10-P-0SR906Z, ICD-10-P-0SR90J9, ICD-10-P-0SR90JA, ICD-10-P-0SR90JZ

<sup>a</sup>ICD-10, International Classification of Diseases, Tenth Revision.