

RESEARCH ARTICLE

Association between Personality Traits and Sleep Quality in Young Korean Women

Han-Na Kim¹, Juhee Cho^{2,3,4,5}, Yoosoo Chang^{2,6}, Seungho Ryu^{2,6}, Hocheol Shin⁷*, Hyung-Lae Kim¹✉*

1 Department of Biochemistry, School of Medicine, Ewha Womans University, Seoul, South Korea, **2** Center for Cohort Studies, Total Healthcare Center, Kangbuk Samsung Hospital, School of Medicine, Sungkyunkwan University, Seoul, South Korea, **3** Department of Health Sciences and Technology, SAHIST, Sungkyunkwan University, Seoul, South Korea, **4** Biostatistics and Clinical Epidemiology Center, Research Institute for Future Medicine, Samsung Medical Center, School of Medicine, Sungkyunkwan University, Seoul, South Korea, **5** Department of Health, Behavior and Society and Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States of America, **6** Department of Occupational Medicine, Kangbuk Samsung Hospital, School of Medicine, Sungkyunkwan University, Seoul, South Korea, **7** Department of Family Medicine and Health Screening Center, Kangbuk Samsung Hospital, School of Medicine, Sungkyunkwan University, Seoul, South Korea

✉ These authors contributed equally to this work.

* hcfm.shin@samsung.com (HS); hyung@ewha.ac.kr (HLK)



CrossMark
click for updates

OPEN ACCESS

Citation: Kim H-N, Cho J, Chang Y, Ryu S, Shin H, Kim H-L (2015) Association between Personality Traits and Sleep Quality in Young Korean Women. PLoS ONE 10(6): e0129599. doi:10.1371/journal.pone.0129599

Academic Editor: Martin Voracek, University of Vienna, AUSTRIA

Received: November 25, 2014

Accepted: May 11, 2015

Published: June 1, 2015

Copyright: © 2015 Kim et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This research was supported by the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (NRF-2014R1A2A2A04006291) and the Ministry of Education (NRF-2013R1A1A2062702).

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Personality is a trait that affects behavior and lifestyle, and sleep quality is an important component of a healthy life. We analyzed the association between personality traits and sleep quality in a cross-section of 1,406 young women (from 18 to 40 years of age) who were not reporting clinically meaningful depression symptoms. Surveys were carried out from December 2011 to February 2012, using the Revised NEO Personality Inventory and the Pittsburgh Sleep Quality Index (PSQI). All analyses were adjusted for demographic and behavioral variables. We considered beta weights, structure coefficients, unique effects, and common effects when evaluating the importance of sleep quality predictors in multiple linear regression models. Neuroticism was the most important contributor to PSQI global scores in the multiple regression models. By contrast, despite being strongly correlated with sleep quality, conscientiousness had a near-zero beta weight in linear regression models, because most variance was shared with other personality traits. However, conscientiousness was the most noteworthy predictor of poor sleep quality status (PSQI \geq 6) in logistic regression models and individuals high in conscientiousness were least likely to have poor sleep quality, which is consistent with an OR of 0.813, with conscientiousness being protective against poor sleep quality. Personality may be a factor in poor sleep quality and should be considered in sleep interventions targeting young women.

Introduction

Sleep is a universal part of life and one of the important behavioral components of sustaining health state. Sleep disturbance is linked to the development of physical and psychological

problems, including cardiovascular disease, diabetes, obesity, anxiety, depression, and all-cause mortality [1–7]. An estimated 50–70 million US adults have a sleep or wakefulness disorder [8], and the prevalence of insomnia in the adult Korean population is greater than 20% [9]. Currently, most assessments of sleep quality depend upon individual perception, although it can be presented as objective sleep parameter through itemizing and measuring. Such perception is likely to depend upon the personality of the individual. Personality has been reported as a predictor of the objective sleep parameters as well [10, 11]. This suggests that the link between personality and sleep may be important in the assessment of sleep quality and in intervention. An increasing number of behavioral therapy intervention trials seek to improve sleep quality by tailoring interventions to individuals' needs and stages of change [12–14]. Thus, it is important to identify individual-level variables associated with poor sleep quality, particularly personality traits.

The five-factor model (FFM) of personality has emerged as a promising predictor of health behaviors [15]. The five-factor model (FFM) is a comprehensive yet manageable taxonomy of traits that is generalizable across cultures, including that of Korea [16, 17]. According to the FFM, personality traits can be described using five broad dimensions, also known as the Big Five personality domains: Neuroticism (N), Extraversion (E), Openness to Experience (O), Agreeableness (A), and Conscientiousness (C). Each factor is defined by lower-order traits, known as facets. Of these five factors, neuroticism and conscientiousness have been frequently linked to a variety of health-risk behaviors and outcomes [18]. For example, high neuroticism or low conscientiousness was associated with lower subjective [19] and physician-rated health [20], chronic illnesses [21], and mortality [22].

Recently, Duggan reported that low conscientiousness and high neuroticism were the best predictors of poor sleep (i.e., poor sleep hygiene, low sleep quality, and increased sleepiness) [23]. They created new variables using principal component analysis (PCA) to remove multicollinearity between the original personality factor scores. When they re-ran analyses using men-centered personality traits, the conscientiousness parameter was in the same direction but became non-significant, whereas the neuroticism parameter remained significant in the multiple linear models. Hintsanen et al., also reported that the conscientiousness was significantly associated with sleep deficiency when it was examined separately as a predictor, but when all other personality traits were additionally added, the associations attenuated to non-significant [24]. In the presence of correlation among predictors, focusing solely on regression weights yields at best limited information and, in some cases, erroneous interpretation in the multiple regression model [25]. Some researchers have suggested that structure coefficients must be interpreted in conjunction with the standardized beta weights when predictors are correlated [26, 27]. Williams et al. examined the independent and interactive contributions of neuroticism and conscientiousness to sleep quality, but they also interpreted only beta weights in the regression model [28]. Thus, it seems important to understand both the unique and common (i.e. shared) effects of personality traits for the sleep quality.

Several other studies have explored the relationship between sleep quality and neuroticism or conscientiousness, but most of them have been conducted in university students, clinical samples, or small samples [23, 28–31]. Recently, sleep quality has emerged as an important issue in young adult, particularly workers. Doi et al. found that being younger was a risk factor for poor sleep quality, arguing that younger workers may be more vulnerable to stressors because they are less able to cope [32, 33]. In young adults, good sleep quality might make them healthy and sustain productive activity. However, there little research of sizable samples to identify personality traits related with sleep quality in young adults. Thus, there is a need to assess the relationship between sleep quality and personality for effective health-care and preventive health interventions in young adults.

Many studies have reported sex differences in personality traits, and the prevalence of insomnia and depression [34, 35]. Several studies have reported a relationship between personality traits and sleep health in Caucasians [23, 29]. Lichstein et al. have documented racial differences in sleep [36]. Although there are several studies in Asians, most of them have been conducted in adolescents, university students, or small samples [30, 37, 38]. Besides, they focused on only the neuroticism. It needs to be examined whether the sleep quality would be related with the five factors of personality in broad range of age.

Our cross-sectional, survey-based study, which analyzes the five broad dimensions of personality as well as specific facets, was carried out in a sample of young Korean women not reporting clinically depression symptoms. This study was performed in a cohort that based on health screening for middle-aged workers and their spouses. Our aim was to determine whether an individual's personality traits significantly are associated with her sleep quality in young women.

Materials and Methods

Participants

Participants were recruited from the Kangbuk Samsung Health Study, which is a cohort study of Korean men and women who undergo a comprehensive annual or biennial examination at Kangbuk Samsung Hospital Screening Centers in South Korea [39]. The sample for this analysis included all 2,213 female participants between the ages of 20 and 40 who underwent a comprehensive health checkup and completed a sleep quality questionnaire. The Revised NEO Personality Inventory (NEO PI-R) manual provides a protocol for validity checking based on acquiescence, naysaying, and randomness of response. Based on item response patterns in the personality assessment, 93 subject surveys were classified as invalid and eliminated from this analysis. These subjects provided repetitive answers or had a pattern of acquiescing or naysaying that would have invalidated formal scoring and interpretation of the NEO PI-R. In addition, 106 subjects with one or more missing values, or 21 subjects who showed the abnormal habitual sleep efficiency according to the sleep assessment were excluded. We also excluded participants with a history of sleep-related problems ($N = 18$). Additionally, we excluded participants who worked night shifts ($N = 110$) and those diagnosed with mental disorders such as depression and panic disorder ($N = 78$). Depression symptoms were measured using the Korean version [40] of the Center for Epidemiologic Studies Depression Scale (CES-D) [41]. Participants who had symptoms of depression ($CES-D \geq 16$) [42] and missing values were excluded from the analysis ($N = 426$). In the end, the total number of subjects included in the study was 1,406 (Fig 1).

Statement of ethics

The Institutional Review Board of Kangbuk Samsung Hospital approved this study, and written informed consent was obtained from all participants. All applicable institutional and governmental regulations concerning the ethical use of human volunteers were followed during our research.

Sleep Variables

Sleep quality was measured using the Korean version of the Pittsburgh Sleep Quality Index (PSQI) [43]. The PSQI is a validated, self-administered questionnaire used to generate seven component scores. It contains 19 items, which measure subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication, and




Fig 1. Sample selection procedure. Some participants were excluded for multiple criteria.

doi:10.1371/journal.pone.0129599.g001

daytime function. Each component score is weighed equally on a scale from 0 to 3, which are summed together to yield a global PSQI score. This score can range from 0 to 21, with higher scores (≥ 6) indicating poor sleep quality.

Personality Assessment

Personality traits were assessed online using the Korean short version of the original NEO PI-R [44], a 90-item measure of the five factors of personality (PSI Consulting Corp., Seoul, Korea). The Korean version of the NEO PI-R has been used in the Korean population with good reliability and validity [45]. This instrument has a robust factor structure that has been replicated in more than 50 other cultures [16] in addition to that of Korea [45, 46]. It measures 30 facets, six for each of the five major dimensions of personality. Items were answered on a 5-point Likert-type scale with responses ranging from “strongly disagree” to “strongly agree.” There were no missing responses in our dataset, because the subjects could complete the questionnaire if there are no missing data in every page online. In the present study, the Cronbach’s alpha coefficients for neuroticism, extraversion, openness, agreeableness, and conscientiousness were 0.83, 0.82, 0.76, 0.69, and 0.78, respectively. Raw scores were converted to *T*-scores (mean = 50, standard deviation = 1) using Korean combined-sex norms (N = 7,418) to confirm similarities with the Korean normative sample data (PSI Consulting Corp., Seoul, Korea).

Demographic and behavioral characteristics

Age, marital status, working status, education, caffeine intake, alcohol use, smoking, and physical activity were considered covariates in this analysis. Marital status was categorized as “married” or “single/separated/divorced.” Working status was categorized as “employed” or “unemployed.” Education was categorized as “high school,” “college,” “university,” or “graduated university.” Participants were asked how many alcoholic drinks and caffeinated beverages (coffee, tea, other) they consumed on average per day. Participants were also categorized as

“never smokers,” “former smokers,” or “current smokers.” Physical activity was categorized as “low,” “moderate,” or “high” following the categorical scoring of the International Physical Activity Questionnaire (IPAQ) [47].

Statistical Analysis

All statistical analyses were conducted using SAS version 9.3 for Windows and RStudio version 0.98.983 for Mac. Descriptive statistics were calculated to characterize the sample. Sleep quality was categorized as good (PSQI <6) or poor (PSQI \geq 6). The two-sample t-test was used to compare the means of continuous variables of interest for these two groups, while the chi-square test was used to examine differences in categorical variables.

Pearson (for continuous variables) or Spearman (for categorical variables) correlation coefficients (zero-order r) were used in preliminary investigations of the relationships among all variables. Sleep quality residuals were normally distributed, and multicollinearity diagnostics were performed on the five personality factors. Next, we performed multivariate linear regression analysis using two models. In Model I, five linear regression analyses were conducted on each personality factor to examine its unique effect on the global PSQI score. In Model II, one multiple linear regression analysis including all five personality factors was conducted, to identify independent predictors of the global PSQI score. The covariates were only used in the regression model if the bivariate correlation indicated they were significantly correlated ($p < .05$) with the global PSQI score. All statistical tests were two-sided ($\alpha = .05$).

To further analyze the relationships among the personality traits in the multiple regression models, we examined structure coefficients as well as beta weights (β) and performed commonality analyses. A structure coefficient (r_s) is a correlation between an observed predictor variable and the predicted criterion score (\hat{Y}) [48]. It is calculated by dividing the Pearson correlation (r) between an independent variable and the dependent variable by the multiple correlation (R). A squared structure coefficient (r_s^2) reveals how much variance the predictor can explain of the observed R^2 effect (not of the total dependent variable), and therefore provides a sense of how much each predictor contributes to the explanatory power of the entire model [48]. Commonality analysis provides separate measures of unique variance explained for each predictor in addition to measures of shared variance for all combinations of predictors [49].

Next, multivariate logistic regression analysis was conducted to identify independent determinants of poor sleep (global PSQI score \geq 6) using two models. In Model I, five different logistic regression analyses were conducted on each personality factor to examine its unique effect on poor sleep. In Model II, one multiple logistic regression analysis including all five personality factors was conducted, to identify independent predictors of poor sleep. All demographic variables were included in both models. In addition, we performed a stepwise discriminant function analysis (DFA) to determine which of five personality factors could best differentiate good sleepers and poor sleepers. The stepwise method of discriminant analyses was preferred over the standard method in order to eliminate discriminating variables that do not add to the power of the discriminant functions.

Results

Sample characteristics

Descriptive statistics for the participants are shown in [Table 1](#). Their average age was 33.1 ± 3.9 , and roughly 80% of participants were married. The number of employed and unemployed women was roughly equal. Most participants had an educational background of university or higher (71%) and were never smokers (90%). In a univariate analyses, we found no significant

Table 1. Descriptive statistics for study participants.

Variable	Overall	Sleep quality		p
		Good (PSQI <6)	Poor (PSQI ≥6)	
Age ^a	33.34 (3.79)	33.40 (3.84)	33.14 (36.2)	0.28
Sex	1,406 (Female)	1,093	313	-
Marital status ^b				0.33
Married	1,138 (81)	888 (81)	250 (80)	
Single/Separated/Divorced	268 (19)	205 (19)	63 (20)	
Working status ^b				0.08
Employed	736 (52)	586 (54)	150 (48)	
Unemployed	670 (48)	507 (46)	163 (52)	
Education ^b				0.77
High school	149 (11)	118 (11)	31 (10)	
College	235 (17)	177 (16)	58 (19)	
University	751 (53)	588 (54)	163 (52)	
≥Graduate school	271 (19)	210 (19)	61 (19)	
Caffeine per Day ^a	1.22 (1.37)	1.24 (1.38)	1.13 (1.34)	0.22
Alcohol per Day ^a	2.62 (2.81)	2.61 (2.81)	2.70 (2.85)	0.62
Smoking ^b				0.01
Never smoker	1,262 (90)	993 (91)	269 (86)	
Former/Current smoker	144 (10)	100 (9)	44 (14)	
Physical Activity ^b				0.98
Low	813 (58)	632 (58)	181 (58)	
Moderate	486 (34)	377 (34)	109 (35)	
High	107 (8)	84 (8)	23 (7)	
PSQI ^a				
Subjective sleep quality	1.12 (0.29)	0.95 (0.46)	1.73 (0.58)	<0.001
Sleep latency	0.77 (0.82)	0.56 (0.64)	1.51 (0.93)	<0.001
Sleep duration	0.53 (0.74)	0.34 (0.58)	1.19 (0.86)	<0.001
Habitual sleep efficiency	0.24 (0.62)	0.07 (0.28)	0.85 (0.99)	<0.001
Sleep disturbances range	0.7 (0.48)	0.63 (0.49)	0.95 (0.38)	<0.001
Use of sleeping medication	0.01 (0.11)	0.00 (0.06)	0.03 (0.21)	<0.001
Daytime dysfunction	0.65 (0.68)	0.52 (0.59)	1.12 (0.76)	<0.001
Global score (scale range 0–21)	4.03 (2.29)	3.07 (1.31)	7.39 (1.73)	<0.001
Personality ^a (T-score, mean = 50, SD = 10)				
Neuroticism	55.18 (9.16)	54.74 (9.12)	56.74 (9.12)	<0.001
Extraversion	44.85 (9.75)	44.93 (9.78)	44.57 (9.67)	0.56
Openness	54.06 (12.52)	53.85 (12.55)	54.81 (12.41)	0.23
Agreeableness	48.41 (11.77)	48.65 (11.72)	47.58 (11.95)	0.16
Conscientiousness	41.98 (8.90)	42.46 (8.88)	40.31 (8.77)	<0.001

Note. PSQI: Pittsburgh Sleep Quality Index; SD: Standard Deviation

^a Quantitative variable. Mean (standard deviation) and p-value from t-test are shown.

^b Nominal scale. Frequency (percentage) and p-value from χ^2 test are shown.

doi:10.1371/journal.pone.0129599.t001

associations between demographic and behavioral variables and sleep quality, except for smoking status. The mean sleep quality (PSQI) score was 4.03 (SD = 2.29), and poor sleepers represented 22% of the total. The mean of our subjects was lower than those reported in university students and older women in Caucasian [23, 29, 50]. Overall, the T-scores of personality traits

for the entire sample, which never deviate more than 10 points from the standardized men score of 50, indicated that they don't differ substantially from the normative sample data.

Association between personality traits and sleep quality

[Table 2](#) provides the correlation coefficients among all variables. All dimensions except openness were significantly associated with sleep quality. Neuroticism and conscientiousness had a significant correlation with PSQI scores ($r = 0.170$ and -0.104 , respectively). Big five personality traits showed non-zero correlations with each other. Among them, neuroticism and conscientiousness had the highest Pearson's correlation coefficient ($r = -0.448$, $p < .001$). The correlation matrix among the five personality traits indicated that the absolute values of the correlation coefficients were lower than the acceptable cut-off point of 0.8 for inclusion in multiple regression analysis [51]. The correlation matrix between the facets of five personality traits and sleep quality was showed in [S1 Table](#).

Although the absolute values of the correlation coefficients between the five dimensions of personality were lower than the acceptable cut-off point of 0.8, we investigated potential multicollinearity among the variables by calculating variance inflation factors (VIFs) and tolerance values for the multiple regression analysis. VIF scores and tolerance values ranged from 1.120 to 1.400 and from 0.715 to 0.893, respectively, indicating that multicollinearity was not a problem in our model.

Associations between personality traits and sleep quality (continuous PSQI global scores), controlling for demographic and behavioral variables, are shown in [Table 3](#). The results of a series of linear regression analyses for each individual personality domain (Model I) showed that all dimensions except openness were significantly associated with sleep quality. With a standardized regression (β) weight of 0.157 ($P < .001$), neuroticism showed the strongest positive association with PSQI scores. Extraversion, agreeableness, and conscientiousness showed significant negative associations with PSQI scores. In a multiple linear regression model (Model II), the combination of age, marital status, smoking, and working status explained 1.9% of the variance on sleep quality. The five personality dimensions explained an additional 3.1% of variance in sleep quality. The multiple regression equation including all five personality domains was significant at the 0.01 level ($F = 8.08$, $p < .001$, $R^2 = 0.050$, Adjusted $R^2 = 0.043$). Of the five personality dimensions, neuroticism was found to contribute most significantly to the prediction of sleep quality ($\beta = -0.123$, $p < .001$). Although extraversion, agreeableness, and conscientiousness were significantly correlated with sleep quality in zero-order bivariate analyses, in combination with the other personality domains they were not significant predictors. In contrast, openness was shown significant association with sleep quality in the multiple linear regressions, although it was not correlated with sleep quality in zero-order bivariate analyses.

The associations between the facets of five personality traits and sleep quality were showed in [S1 Table](#). At facet level, anxiety (N1) depression (N3), and impulsiveness (N5), facets of neuroticism, were positively associated with PSQI global scores. Gregariousness (E2), activity (E4), and positive emotions (E6) of extraversion, feeling (O3) and values (O6) of openness, trust (A1) and straightforwardness (A2) of agreeableness, and self-discipline (C5) of conscientiousness were negatively associated with PSQI global scores.

We considered structure coefficients (r_s) to interpret the multiple linear regression as well as beta weights. If two or more predictors explain some of the same part of the criterion, the sum of squared structure coefficients for all predictors will be greater than 1.00 [52]. In our model, the sum of five dimensions of personality was 1.090, suggesting a small amount of multicollinearity was present. After computing the structure coefficients for this multiple regression model, neuroticism was still the most noteworthy predictor of sleep quality ($r_s = 0.764$) and

Table 2. Correlation coefficients for sleep quality, covariates, and personality traits.

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. PSQI global score ^a	1													
2. Age ^a	-0.075**	1												
3. Marital status ^b	0.056*	-0.377***	1											
4. Education ^b	-0.006	-0.055*	0.044*	1										
5. Caffeine ^a	-0.019	0.111***	0.045	-0.012	1									
6. Alcohol ^a	0.043	-0.167***	0.200***	-0.109***	0.080**	1								
7. Smoking ^b	0.064*	-0.028	0.129***	0.010	0.028	0.109***	1							
8. Physical activity ^b	-0.009	-0.018	0.102***	0.035	0.005	0.043	0.031	1						
9. Working status ^b	-0.074**	-0.171***	0.274***	0.145***	0.128***	0.169***	0.055*	0.031	1					
10. Neuroticism ^a	0.170***	-0.084**	0.015	-0.074**	-0.015	-0.014	0.013	-0.084**	-0.089**	1				
11. Extraversion ^a	-0.063*	-0.058*	-0.002	0.013	0.006	0.120***	0.040	0.104***	0.066*	-0.262***	1			
12. Openness ^a	0.035	-0.127***	0.138***	0.167***	-0.022	0.069**	0.072*	0.115***	0.043	-0.085**	0.399***	1		
13. Agreeableness ^a	-0.095***	0.171***	-0.116***	-0.009	-0.061*	-0.107***	-0.102***	-0.017	-0.088**	-0.231***	0.023	-0.001	1	
14. Conscientiousness ^a	-0.104***	0.054*	0.014	0.150***	0.009	-0.046	-0.057*	0.144***	0.109***	-0.448***	0.204***	0.201***	0.065*	1

N = 1,406

^a Pearson's correlation coefficients

^b Spearman correlation coefficients

**p*<.05

***p*<.01

****p*<.001

doi:10.1371/journal.pone.0129599.t002

Table 3. Linear regression analysis investigating the association between personality traits and sleep quality, as measured by PSQI global score.

Factors	Model I ^a	Model II ^b		r_s^2	Unique ^d	Common ^e
	β^c	β^c	Structure Coefficient (r_s)			
Neuroticism (N)	0.157***	0.123***	0.764	0.584	0.011	0.018
Extraversion (E)	-0.065*	-0.047	-0.285	0.081	0.002	0.002
Openness (O)	0.021	0.061*	0.156	0.024	0.003	-0.002
Agreeableness (A)	-0.084**	-0.053	-0.428	0.183	0.003	0.007
Conscientiousness (C)	-0.087***	-0.033	-0.467	0.218	0.001	0.010

N = 1,406

Note. All models were adjusted for age, marital status, smoking, and working status.

^a Model I: linear regression model including a single domain of personality as an independent variable ($R^2 = 0.043, 0.023, 0.020, 0.026, \text{ and } 0.027$; adjusted $R^2 = 0.040, 0.020, 0.016, 0.022, \text{ and } 0.023$, $F = 12.67, 6.70, 5.61, 7.45, \text{ and } 7.68$; $P < .001, < .001, < 0.001, < .001, \text{ and } < 0.001$ in N, E, O, A, and C, respectively).

^b Model II: multiple linear regression model including all five domains of personality as independent variables ($R^2 = 0.0524$, adjusted $R^2 = 0.041$, $F = 4.80$, $P < .001$).

^c β : standardized coefficient in linear regression analyses.

^d Unique = proportion of criterion variance explained uniquely by the predictor.

^e Common = proportion of criterion variance explained by the predictor that is also explained by one or more other predictors. Unique + Common = r^2 , r = zero-order correlation coefficient

* $p < .05$

** $p < .01$

*** $p < .001$

doi:10.1371/journal.pone.0129599.t003

could account for 58.4% ($r_s^2 = 0.584$, partial $R^2 = 0.029$) of the explained variance ($R^2 = 0.050$). The rank of structure coefficient for conscientiousness was second ($r_s = -0.467$, $r_s^2 = 0.218$) among the five personality factors, not fifth as the beta weights suggested. Openness showed the smallest structure coefficient and could account for 2% of the obtained effect by itself ($r_s = 0.156$, $r_s^2 = 0.024$), although the beta weight was significant and the rank of that was second. Yet openness has insignificant zero-order correlation and small structure coefficients with sleep quality but a sizeable non-zero beta weight. In this case openness is serving as a suppressor variable and helping the other predictor variables do a better job of predicting the sleep quality even though openness itself is unrelated to the outcome. This affect R^2 by allowing the beta weights for other predictors to deviate further from zero than the boundary of their respective zero-order correlations with sleep quality, thus making some of the other product terms within the equation larger, and thus making R^2 larger [27]. Inspection of the structure coefficients suggests that neuroticism was a very strong indicator of PSQI scores, conscientiousness was a moderate indicator of PSQI scores, and openness was a suppressor variable in the multiple regression models.

We considered commonality coefficients to find why the conscientiousness vanished the significance of a beta weight in multiple linear regression models although its contribution for PSQI scores was identified by correlation coefficients and structure coefficients. Commonality analysis decomposes the squared multiple correlation (R^2) to determine both the unique and common explained variance of the sleep quality by the predictor (Table 3 and S2 Table). In total, the five personality domains individually accounted for 38% (sum of the unique/ $R^2 = 0.019/0.050$) of the regression effect. The remaining 62% was due to variance in PSQI global scores that the predictors shared. The unique coefficient for neuroticism (0.011) indicated that neuroticism uniquely explained 1.1% of the variance in PSQI scores. This amount of variance

is more than any other partition, representing 22.03% of the R^2 effect (0.050). The unique coefficient for conscientiousness (0.001) was the smallest of the unique effects and indicated that the regression model only improved slightly with the addition of conscientiousness. The common effects represented the proportion of PSQI scores variance that could be jointly explained by two or more predictors together. Neuroticism and conscientiousness together explained 0.5% of the outcome, which represented 10.92% of the total effect size (S2 Table). The common effects also showed that openness served as a suppressor variable, yielding a unique effect greater than its total contribution to the regression effect and negative commonality coefficients [25]. Overall, commonality analysis could help to quantify the location and amount of variance explained by suppression and multicollinearity. We also investigated the potential interaction between neuroticism and conscientiousness in predicting sleep quality. In a regression analysis including neuroticism, conscientiousness and their interaction, the interaction term was not significant ($\beta = -0.118, p = 0.380$).

Additionally, S3 Table was shown the associations between five personality traits and seven components of sleep quality as measured by PSQI. The subjective sleep quality scores were associated with extraversion and openness. The sleep latency scores were showed the positive association with neuroticism but the negative association with extraversion. The sleep duration scores and use of sleeping medication scores were not significantly associated with any personality trait. The habitual sleep efficiency scores were negatively associated with conscientiousness. The daytime dysfunction scores were shown the positive association with neuroticism.

Personality factors and facets predict poor sleepers

Results from the t-test indicated significant personality differences between sleep quality groups (Table 1). Neuroticism score was higher in poor sleepers than good sleeper ($t = -3.42, p < 0.001$), while conscientiousness was lower in poor sleepers than good sleepers ($t = 3.79, p < 0.001$). The other personality domains showed no difference between groups.

As noted earlier, a logistic regression analysis was used to examine whether personality traits could predict poor quality status identified by PSQI scores (global PSQI ≥ 6) (Table 4). In the logistic regression models that considered the personality traits individually (Model I), both neuroticism and conscientiousness were significant, separate predictors of poor sleep quality status. The significant association between neuroticism and poor sleep in Model I vanished when investigated in a multiple regression model adding the other personality domains (Model II). In contrast to the results of the linear regression, conscientiousness was the most noteworthy predictor of sleep quality in logistic regression models (Model II). Individuals high in conscientiousness were least likely to have poor sleep quality, which is consistent with an OR of 0.813, with conscientiousness being protective against poor sleep quality, and specifies that the logistic regressions are predicting poor sleep quality status rather than good sleep quality status. The facets of neuroticism and conscientiousness were analyzed using the same models (Model I and II). Among the facets of neuroticism, depression (N3) were the strongest predictors of poor sleep quality status (OR = 1.175, 95% CI = 1.004 to 1.375). The score of deliberation (C6) was significantly lower in poor sleepers than in good sleepers (OR = 0.878, 95% CI = 0.781 to 0.988).

Because of slightly inter-correlation among some of the five personality domains, a stepwise DFA was computed with covariates to evaluate which measures uniquely maximized the separation between good sleepers and poor sleepers (Table 5). This analysis also allowed us to determine a more parsimonious set of personality traits. The stepwise DFA indicated conscientiousness and neuroticism were statistically significant and independent contributors to group separation. The classification accuracy of the DFA model was 60.4% for good sleepers

Table 4. Logistic regression models predicting poor sleep quality based on personality characteristics.

Factors	Poor sleepers, n = 313 (reference: good sleepers, n = 1,093)	
	OR ^a (95% CI)	
	Model I ^b	Model II ^c
Neuroticism (N)	1.246 (1.084, 1.433)**	1.137 (0.966, 1.338)
Extraversion (E)	0.960 (0.843, 1.093)	0.982 (0.848, 1.138)
Openness to Experience (O)	1.050 (0.947, 1.163)	1.101 (0.983, 1.233)
Agreeableness (A)	0.939 (0.842, 1.047)	0.966 (0.863, 1.083)
Conscientiousness (C)	0.784 (0.680, 0.904)***	0.813 (0.692, 0.956)*
Facets ^d		
N1: Anxiety	1.150 (1.015, 1.303)*	1.075 (0.923, 1.253)
N2: Angry Hostility	1.127 (1.007, 1.261)*	1.012 (0.885, 1.157)
N3: Depression	1.211 (1.076, 1.381)**	1.175 (1.004, 1.375)*
N4: Self-consciousness	1.043 (0.936, 1.161)	0.914 (0.800, 1.044)
N5: Impulsiveness	1.178(1.047, 1.326)**	1.137 (0.999, 1.294)
N6: Vulnerability	1.127 (1.006, 1.263)*	1.012 (0.875, 1.171)
C1: Competence	0.896 (0.805, 0.999)*	0.945 (0.839, 1.065)
C2: Order	0.949 (0.832, 1.082)	1.110 (0.947, 1.302)
C3: Dutifulness	0.879 (0.782, 0.988)*	0.961 (0.837, 1.104)
C4: Achievement Striving	0.873 (0.784, 0.971)*	0.943 (0.821, 1.084)
C5: Self-Discipline	0.844 (0.749, 0.952)**	0.905 (0.782, 1.047)
C6: Deliberation	0.856 (0.770, 0.951)**	0.878 (0.781, 0.988)*

N = 1,406

Note. CI: Wald Confidence Interval

^a Odds Ratios (ORs) per 10 T-score increase in a given personality trait, controlling for age, marital status, working status, education, caffeine intake, alcohol use, smoking status, and physical activity.

^b Model I: logistic regression model including a single domain or facet of personality as an independent variable.

^c Model II: multiple logistic regression model including all five domains of personality as independent variables. At a facet level, six facets of each domain were included in the model.

^d Logistic regression analyses of the facet level were performed in neuroticism and conscientiousness.

**p* < .05

***p* < .01

****p* < .001

doi:10.1371/journal.pone.0129599.t004

and 53.4% for poor sleepers. At facet levels, depression (N3), impulsiveness (N5), self-discipline (C5), and deliberation (C6) were statistically significant and independent contributors to group separation. The classification accuracies for the specific facets of neuroticism and conscientiousness were 59.6% and 60.8% for good sleepers, respectively, and 54.3% and 49.5% for poor sleepers, respectively.

Supplementary Analyses

To determine the robustness of our findings, we re-ran analyses using PCA-derived personality factors, as described by Duggan et al. [23]. PCA-derived personality factors obtained using the orthogonal varimax rotation. Thus, PCA removed the substantial multicollinearity between the personality traits. The results were consistent with the findings of bivariate analyses and Model

Table 5. Discriminant function analysis (DFA) by the stepwise method for the separation between good sleepers and poor sleepers.

Variable	Standardized coefficients	Wilk's λ	F	p	DFA entry order
Factors					
Neuroticism	0.452	0.983	7.916	<0.0001	2
Conscientiousness	-0.553	0.990	14.396	<0.0001	1
Smoking (covariate)	0.488	0.986	9.933	<0.0001	3
Facets: Neuroticism					
N3: Depression	0.605	0.992	11.572	0.001	1
N5: Impulsiveness	0.470	0.984	7.554	<0.0001	2
Smoking (covariate)	0.494	0.987	8.991	<0.0001	3
Facets: Conscientiousness					
C5: Self-Discipline	0.550	0.993	9.519	0.002	1
C6: Deliberation	0.497	0.986	6.753	<0.0001	2
Smoking (covariate)	-0.521	0.989	7.797	<0.001	3

N = 1,406

doi:10.1371/journal.pone.0129599.t005

I of logistic analyses (S4 Table). In the simultaneous multiple linear regression model, PCA-derived neuroticism showed a positive association with PSQI global scores ($\beta = 0.136, p < .001$), while PCA-derived extraversion, agreeableness, and conscientiousness were negatively associated with the scores (extraversion: $\beta = -0.052, p < .05$, agreeableness: $\beta = -0.069, p < .01$, and conscientiousness: $\beta = -0.059, p < .05$). In logistic regression models, PCA-derived conscientiousness was the strongest predictor of sleep quality (OR = 0.822, 95% CI = 0.724 to 0.933) while PCA-derived neuroticism was the second-best predictor (OR = 1.171, 95% CI = 1.031 to 1.330). At the facet level, PCA-derived depression (N3), impulsiveness (N5), self-discipline (C5), and deliberation (C6) showed significant associations with sleep quality.

Discussion

The main findings of this study were that neuroticism and conscientiousness were associated with PSQI global scores, even after controlling for demographic and behavioral characteristics. Among young women not reporting clinically meaningful depression symptoms, poor sleepers scored significantly lower than good sleepers on conscientiousness. We examined not only the factors of personality traits, but also facet-level analysis, which permits the examination of multifaceted constructs, which could be relevant to understanding the relationship between personality and sleep quality. This study examined how both unique and common effects of each personality trait explain variance in sleep quality. In particular, the current study provided a rich interpretation of our sleep quality regression models, by identifying personality trait suppressor effects and patterns of shared variance that would have gone undiscovered through reliance on beta weights alone. In Table 3, we simultaneously considered beta weights, structure coefficients, unique effects, and common effects when evaluating the importance of PSQI global score predictors in the multiple linear regression models. The each coefficient has distinct advantages and limitations, and that multiple statistical measures complement each other in the perspective they provide regarding regression findings [53]. We found the best predictors of sleep quality were different domains of five personality traits in accordance with the approach for the PSQI global score, either continuous variables or discrete variables.

In our multiple linear regression models, we found that neuroticism was the best contributor to PSQI global scores, though all dimensions except openness were significantly associated with PSQI global scores in the bivariate analyses. Conscientiousness was highly correlated with sleep quality in the zero-order bivariate linear relationship, but it was not significant in a multiple linear regression model. Moreover, conscientiousness did not moderate the association between neuroticism and sleep quality. These results were consistent with prior findings [28]. Many researchers have recommended that both beta weights and structure coefficients be considered when interpreting the results of regression models containing correlated predictor variables [25, 27]. Interpreting both beta weights and structure coefficients could help to not only find existence of a small amount of multicollinearity, but also identify the best predictor and suppressor for sleep quality in the multiple linear regressions. We found a reason why conscientiousness was not a significant predictor in multiple regression models. In the commonality analysis, conscientiousness explained more shared than unique variance in PSQI global scores. Williams et al. also examined the unique and interactive effects of neuroticism and conscientiousness for sleep quality using moderation model of the regression analysis, but they interpreted only beta weights in the regression model [28]. We quantified the amount of variance explained by multicollinearity using the commonality analysis. Another interesting finding was openness's role as a suppressor in predicting PSQI global scores. It was confirmed in both structure coefficients and commonality coefficients.

In the logistic analyses, conscientiousness remained a significant predictor of poor sleep quality status even after inclusion of several lifestyle factors in our models. Several demographic risk factors for poor sleep, such as sex [50], being elderly [50], marriage status [54], education [55], and income [56], have been reported. According to the results of the multiple linear and logistic regressions, we suggest that neuroticism might be the best predictor for sleep quality using PSQI global scores as continuous variables, while conscientiousness might be the best predictor at risk for poor sleep quality status using PSQI global scores as discrete variables, even though neuroticism was the second-best predictor in DFA and PCA.

Our findings replicate and extend previous research on the association between personality traits and sleep quality [28, 29]. At the seven component of sleep quality, neuroticism was associated with relatively subjective component such as sleep disturbances and daytime dysfunction, while it was not related to more objective aspects such as sleep duration. Conscientiousness was the only personality factor associated with the habitual sleep efficiency. The more objective component, such as sleep duration and use of sleeping medication were not shown association with any personality factor. Perhaps people low in conscientiousness and high in neuroticism have poor sleep quality because both factors are associated with difficulty regulating emotions (e.g., anxiety, depression) and behavior (e.g., sleep hygiene) [23]. The propensity for highly neurotic individuals to be sensitive to signs of threat and dysfunction may place them at greater risk for dysfunction in response to perceived sleep disruption [28]. Individuals who score low in conscientiousness tend to engage in risky behaviors that contribute to poor physical health [18].

To the best of our knowledge, this is also the sizeable study to evaluate the effects of personality on sleep quality in Asians. Despite our study's contributions, it does have some limitations. First, because this study was based on cross-sectional data, we could not control for the confounding influence of time-invariant common causes or infer causal relationships between variables. In addition, this study was confined to women.

Identifying the role of personality traits in sleep quality not only provides insight into the etiology of poor sleep but can suggest points of intervention and help tailor prevention and treatment strategies to individual poor sleepers. For example, when faced with a poor sleeper high in conscientiousness, a physician's or nurse's advice to change his or her sleep hygiene or

control bed/wake time would likely be met by a high degree of self-directed effort on the part of the patient. However, for poor sleepers low in conscientiousness, this advice may need to be accompanied by short-term incentives, regular monitoring, and behavior modification reminders, either by a healthcare provider or some other expert [57]. In addition, certain personality styles, such as those who are high in neuroticism and low in conscientiousness, could be flagged as being at much higher risk for sleep quality. Such reports could be developed with the help of physician and epidemiologists [57]. A few studies have reported that interventions such as cognitive behavioral therapy (CBT), interpersonal psychotherapy, or generic therapy were effective in the reduction of neuroticism levels [58, 59], anxiety, and depression [60] or in the increasing conscientiousness [59, 61]. Thus, knowledge of patients' personality profiles could aid in differential treatment planning [62].

In conclusion, this study demonstrated that neuroticism is the strongest personality factor influencing PSQI global scores and conscientiousness is the best predictor of poor quality status in young Korean women. Personality may account for a relatively small but significant amount (~4%) of the variance in sleep quality. However, given the prevalence of sleep disturbance worldwide, even small improvements could have considerable public health impact. Early intervention and prevention efforts targeting sleep quality are important for improving the health of young women.

Supporting Information

S1 Table. Correlations between PSQI global score and facets of five personality domains.
(DOC)

S2 Table. Commonality coefficients.
(XLS)

S3 Table. Associations between personality traits and the basic components of sleep quality as measured by PSQI.
(DOC)

S4 Table. Multiple regressions of sleep quality and PCA-derived personality components.
(DOC)

Author Contributions

Conceived and designed the experiments: HNK JC HS HLK. Performed the experiments: YC SR. Analyzed the data: HNK JC. Contributed reagents/materials/analysis tools: YC SR. Wrote the paper: HNK HS HLK.

References

1. Mallon L, Broman JE, Hetta J. Sleep complaints predict coronary artery disease mortality in males: a 12-year follow-up study of a middle-aged Swedish population. *Journal of internal medicine*. 2002; 251(3):207–16. PMID: [11886479](#).
2. Schwartz S, McDowell Anderson W, Cole SR, Cornoni-Huntley J, Hays JC, Blazer D. Insomnia and heart disease: a review of epidemiologic studies. *Journal of psychosomatic research*. 1999; 47(4):313–33. PMID: [10616226](#).
3. Knutson KL, Ryden AM, Mander BA, Van Cauter E. Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus. *Archives of internal medicine*. 2006; 166(16):1768–74. doi: [10.1001/archinte.166.16.1768](#) PMID: [16983057](#).
4. Bass J, Turek FW. Sleepless in America: a pathway to obesity and the metabolic syndrome? *Archives of internal medicine*. 2005; 165(1):15–6. doi: [10.1001/archinte.165.1.15](#) PMID: [15642868](#).

5. Breslau N, Roth T, Rosenthal L, Andreski P. Sleep disturbance and psychiatric disorders: a longitudinal epidemiological study of young adults. *Biological psychiatry*. 1996; 39(6):411–8. PMID: [8679786](#).
6. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes care*. 2010; 33(2):414–20. doi: [10.2337/dc09-1124](#) PMID: [19910503](#); PubMed Central PMCID: PMC2809295.
7. Castro-Costa E, Dewey M, Ferri C, Uchoa E, Firmo J, Rocha F, et al. Association between Sleep Duration and All-Cause Mortality in Old Age: 9-Year Follow-up of the Bambui Cohort Study, Brazil. *J Epidemiol Commun H*. 2011; 65:A229–A30. doi: [10.1136/Jech.2011.142976h.73](#) PMID: [WOS:000293901801137](#).
8. Colten HR, Altevogt BM. In: Colten HR, Altevogt BM, editors. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. The National Academies Collection: Reports funded by National Institutes of Health. Washington (DC), US: National Academies Press; 2006.
9. Cho YW, Shin WC, Yun CH, Hong SB, Kim J, Earley CJ. Epidemiology of insomnia in Korean adults: prevalence and associated factors. *Journal of clinical neurology*. 2009; 5(1):20–3. doi: [10.3988/jcn.2009.5.1.20](#) PMID: [19513329](#); PubMed Central PMCID: PMC2686894.
10. van de Laar M, Verbeek I, Pevernagie D, Aldenkamp A, Overeem S. The role of personality traits in insomnia. *Sleep medicine reviews*. 2010; 14(1):61–8. doi: [10.1016/j.smrv.2009.07.007](#) PMID: [19897388](#).
11. Kajimura N, Kato M, Sekimoto M, Watanabe T, Takahashi K, Okuma T, et al. A polysomnographic study of sleep patterns in normal humans with low- or high-anxiety personality traits. *Psychiatry and clinical neurosciences*. 1998; 52(3):317–20. doi: [10.1046/j.1440-1819.1998.00401.x](#) PMID: [9681584](#).
12. Tan E, Healey D, Gray AR, Galland BC. Sleep hygiene intervention for youth aged 10 to 18 years with problematic sleep: a before-after pilot study. *BMC pediatrics*. 2012; 12:189. doi: [10.1186/1471-2431-12-189](#) PMID: [23216856](#); PubMed Central PMCID: PMC3538572.
13. Kamdar BB, King LM, Collop NA, Sakamuri S, Colantuoni E, Neufeld KJ, et al. The effect of a quality improvement intervention on perceived sleep quality and cognition in a medical ICU. *Critical care medicine*. 2013; 41(3):800–9. doi: [10.1097/CCM.0b013e3182746442](#) PMID: [23314584](#); PubMed Central PMCID: PMC3579000.
14. Berger AM, Kuhn BR, Farr LA, Lynch JC, Agrawal S, Chamberlain J, et al. Behavioral therapy intervention trial to improve sleep quality and cancer-related fatigue. *Psycho-oncology*. 2009; 18(6):634–46. doi: [10.1002/pon.1438](#) PMID: [19090531](#).
15. Raynor DA, Levine H. Associations Between the Five-Factor Model of Personality and Health Behaviors Among College Students. *J Am Coll Health*. 2009; 58(1):73–81. PMID: [ISI:000268121400010](#). doi: [10.3200/JACH.58.1.73-82](#)
16. McCrae RR, Terracciano A, Personality Profiles of Cultures P. Universal features of personality traits from the observer's perspective: data from 50 cultures. *J Pers Soc Psychol*. 2005; 88(3):547–61. doi: [10.1037/0022-3514.88.3.547](#) PMID: [15740445](#).
17. McCrae RR, John OP. An introduction to the five-factor model and its applications. *Journal of personality*. 1992; 60(2):175–215. PMID: [1635040](#).
18. Sutin AR, Terracciano A, Deiana B, Uda M, Schlessinger D, Lakatta EG, et al. Cholesterol, triglycerides, and the Five-Factor Model of personality. *Biological psychology*. 2010; 84(2):186–91. doi: [10.1016/j.biopsycho.2010.01.012](#) PMID: [20109519](#); PubMed Central PMCID: PMC2933037.
19. Lockenhoff CE, Sutin AR, Ferrucci L, Costa PT. Personality traits and subjective health in the later years: The association between NEO-PI-R and SF-36 in advanced age is influenced by health status. *Journal of Research in Personality*. 2008; 42(5):1334–46. doi: [10.1016/J.Jrp.2008.05.006](#) PMID: [WOS:000260301600015](#).
20. Chapman BP, Lyness JM, Duberstein P. Personality and medical illness burden among older adults in primary care. *Psychosomatic medicine*. 2007; 69(3):277–82. doi: [10.1097/PSY.0b013e3180313975](#) PMID: [17401059](#); PubMed Central PMCID: PMC3807736.
21. Goodwin RD, Friedman HS. Health status and the five-factor personality traits in a nationally representative sample. *Journal of health psychology*. 2006; 11(5):643–54. doi: [10.1177/1359105306066610](#) PMID: [16908463](#).
22. Terracciano A, Lockenhoff CE, Zonderman AB, Ferrucci L, Costa PT Jr. Personality predictors of longevity: activity, emotional stability, and conscientiousness. *Psychosomatic medicine*. 2008; 70(6):621–7. doi: [10.1097/PSY.0b013e31817b9371](#) PMID: [18596250](#); PubMed Central PMCID: PMC2505356.
23. Duggan KA, Friedman HS, McDevitt EA, Mednick SC. Personality and healthy sleep: the importance of conscientiousness and neuroticism. *PloS one*. 2014; 9(3):e90628. doi: [10.1371/journal.pone.0090628](#) PMID: [24651274](#); PubMed Central PMCID: PMC3961248.
24. Hintsanen M, Puttonen S, Smith K, Tornroos M, Jokela M, Pulkki-Raback L, et al. Five-factor personality traits and sleep: evidence from two population-based cohort studies. *Health psychology: official*

- journal of the Division of Health Psychology, American Psychological Association. 2014; 33(10):1214–23. doi: [10.1037/hea0000105](https://doi.org/10.1037/hea0000105) PMID: [25133841](https://pubmed.ncbi.nlm.nih.gov/25133841/).
25. Kraha A, Turner H, Nimon K, Zientek LR, Henson RK. Tools to support interpreting multiple regression in the face of multicollinearity. *Frontiers in psychology*. 2012; 3:44. doi: [10.3389/fpsyg.2012.00044](https://doi.org/10.3389/fpsyg.2012.00044) PMID: [22457655](https://pubmed.ncbi.nlm.nih.gov/22457655/); PubMed Central PMCID: PMC3303138.
 26. Thompson B, Borrello GM. The importance of structure coefficients in regression research. *Educ Psychol Meas*. 1985; 45:203–9.
 27. Courville T, Thompson B. Use of structure coefficients in published multiple regression articles: beta is not enough. *Educ Psychol Meas*. 2001; 61(2):229–48. doi: [10.1177/0013164401612006](https://doi.org/10.1177/0013164401612006) PMID: [WOS:000167722100006](https://pubmed.ncbi.nlm.nih.gov/WOS:000167722100006/).
 28. Williams PG, Moroz TL. Personality vulnerability to stress-related sleep disruption: Pathways to adverse mental and physical health outcomes. *Pers Indiv Differ*. 2009; 46(5–6):598–603. doi: [10.1016/j.paid.2008.12.017](https://doi.org/10.1016/j.paid.2008.12.017) PMID: [WOS:000264255400006](https://pubmed.ncbi.nlm.nih.gov/WOS:000264255400006/).
 29. Gray EK, Watson D. General and specific traits of personality and their relation to sleep and academic performance. *Journal of personality*. 2002; 70(2):177–206. PMID: [11908845](https://pubmed.ncbi.nlm.nih.gov/11908845/).
 30. Yoshiike T, Kuriyama K, Honma M, Ikeda H, Kim Y. Neuroticism relates to daytime wakefulness and sleep devaluation via high neurophysiological efficiency in the bilateral prefrontal cortex: a preliminary study. *Psychophysiology*. 2014; 51(4):396–406. doi: [10.1111/psyp.12180](https://doi.org/10.1111/psyp.12180) PMID: [24660887](https://pubmed.ncbi.nlm.nih.gov/24660887/).
 31. Glozier N, O'Dea B, McGorry PD, Pantelis C, Amminger GP, Hermens DF, et al. Delayed sleep onset in depressed young people. *BMC psychiatry*. 2014; 14:33. doi: [10.1186/1471-244X-14-33](https://doi.org/10.1186/1471-244X-14-33) PMID: [24506941](https://pubmed.ncbi.nlm.nih.gov/24506941/); PubMed Central PMCID: PMC3938136.
 32. Doi Y, Minowa M, Tango T. Impact and correlates of poor sleep quality in Japanese white-collar employees. *Sleep*. 2003; 26(4):467–71. PMID: [12841374](https://pubmed.ncbi.nlm.nih.gov/12841374/).
 33. Knudsen HK, Ducharme LJ, Roman PM. Job stress and poor sleep quality: data from an American sample of full-time workers. *Social science & medicine*. 2007; 64(10):1997–2007. doi: [10.1016/j.socscimed.2007.02.020](https://doi.org/10.1016/j.socscimed.2007.02.020) PMID: [17363123](https://pubmed.ncbi.nlm.nih.gov/17363123/); PubMed Central PMCID: PMC1933584.
 34. Zhang BWYKM. Sex differences in insomnia: A meta-analysis *Sleep*. 2006; 29(1):85–93. PMID: [16453985](https://pubmed.ncbi.nlm.nih.gov/16453985/)
 35. Klein DN, Kotov R, Bufferd SJ. Personality and depression: explanatory models and review of the evidence. *Annual review of clinical psychology*. 2011; 7:269–95. doi: [10.1146/annurev-clinpsy-032210-104540](https://doi.org/10.1146/annurev-clinpsy-032210-104540) PMID: [21166535](https://pubmed.ncbi.nlm.nih.gov/21166535/); PubMed Central PMCID: PMC3518491.
 36. Lichstein KL, Durrence HH, Reidel BW, Taylor DJ, Bush AJ. *Epidemiology of Sleep: Age, Gender, and Ethnicity*. Mahwah, NJ: Lawrence Erlbaum Associates, Inc; 2004.
 37. Cheng SH, Shih CC, Lee IH, Hou YW, Chen KC, Chen KT, et al. A study on the sleep quality of incoming university students. *Psychiatry research*. 2012; 197(3):270–4. doi: [10.1016/j.psychres.2011.08.011](https://doi.org/10.1016/j.psychres.2011.08.011) PMID: [22342120](https://pubmed.ncbi.nlm.nih.gov/22342120/).
 38. Gau SF. Neuroticism and sleep-related problems in adolescence. *Sleep*. 2000; 23(4):495–502. PMID: [10875556](https://pubmed.ncbi.nlm.nih.gov/10875556/).
 39. Kim CW, Yun KE, Jung HS, Chang Y, Choi ES, Kwon MJ, et al. Sleep duration and quality in relation to non-alcoholic fatty liver disease in middle-aged workers and their spouses. *Journal of hepatology*. 2013; 59(2):351–7. doi: [10.1016/j.jhep.2013.03.035](https://doi.org/10.1016/j.jhep.2013.03.035) PMID: [23578884](https://pubmed.ncbi.nlm.nih.gov/23578884/).
 40. Cho MJ, Kim KH. Use of the Center for Epidemiologic Studies Depression (CES-D) Scale in Korea. *The Journal of nervous and mental disease*. 1998; 186(5):304–10. PMID: [9612448](https://pubmed.ncbi.nlm.nih.gov/9612448/).
 41. Berkman LF, Berkman CS, Kasl S, Freeman DH Jr., Leo L, Ostfeld AM, et al. Depressive symptoms in relation to physical health and functioning in the elderly. *American journal of epidemiology*. 1986; 124(3):372–88. PMID: [3740038](https://pubmed.ncbi.nlm.nih.gov/3740038/).
 42. Radloff L. The CES-D scale: a self-report depression scale for research in the general population. *Applied psychological measurement*. 1977; 1:385–401.
 43. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry research*. 1989; 28(2):193–213. PMID: [2748771](https://pubmed.ncbi.nlm.nih.gov/2748771/).
 44. Costa PT Jr., McCrae RR. Professional manual: Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) Professional Manual. Odessa, FL: Psychological Assessment Resources; 1992.
 45. Ahn CK, Chae JH. Standardization of the Korean version of the Revised NEO Personality Inventory. *Korean Journal of Counseling & Psychotherapy*. 1997; 9:443–73.

46. Piedmont RL, Chae JH. Cross-cultural generalizability of the Five-Factor Model of personality: Development and validation of the NEO-PI-R for Koreans. *Journal of Cross-Cultural Psychology*. 1997; 28:131–55.
47. Craig CL, Marshall AL, Sjoström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Medicine and science in sports and exercise*. 2003; 35(8):1381–95. doi: [10.1249/01.MSS.0000078924.61453.FB](https://doi.org/10.1249/01.MSS.0000078924.61453.FB) PMID: [12900694](https://pubmed.ncbi.nlm.nih.gov/12900694/).
48. Thompson B. *Foundation of Behavioral Statistics: An Insight-Based Approach*. New York: Guilford Press.; 2006.
49. Pedhazur EJ. *Multiple regression in behavioral research: explanation and prediction*. 3rd ed. Forth Worth: Harcourt Brace College Publishers; 1997. xiv, 1058 p. p.
50. Vitiello MV, Larsen LH, Moe KE. Age-related sleep change: Gender and estrogen effects on the subjective-objective sleep quality relationships of healthy, noncomplaining older men and women. *Journal of psychosomatic research*. 2004; 56(5):503–10. doi: [10.1016/S0022-3999\(04\)00023-6](https://doi.org/10.1016/S0022-3999(04)00023-6) PMID: [15172206](https://pubmed.ncbi.nlm.nih.gov/15172206/).
51. Benny WDF S. *Multiple Regression in Practice*. Thousand Oaks, CA: Sage Publications; 1985.
52. Henson RK, Hull DM, Williams CS. Methodology in Our Education Research Culture: Toward a Stronger Collective Quantitative Proficiency. *Educ Researcher*. 2010; 39(3):229–40. doi: [10.3102/0013189x10365102](https://doi.org/10.3102/0013189x10365102) PMID: [WOS:000283009800004](https://pubmed.ncbi.nlm.nih.gov/WOS:000283009800004/).
53. Nathans LL, Oswald FL, Nimón K. Interpreting Multiple Linear Regression: A Guidebook of Variable Importance. *Practical Assessment, Research & Evaluation*. 2012; 17(9):1–19.
54. Troxel WM, Buysse DJ, Matthews KA, Kravitz HM, Bromberger JT, Sowers M, et al. Marital/cohabitation status and history in relation to sleep in midlife women. *Sleep*. 2010; 33(7):973–81. PMID: [20614858](https://pubmed.ncbi.nlm.nih.gov/20614858/); PubMed Central PMCID: PMC2894440.
55. Assari S, Moghani Lankarani M, Kazemi Saleh D, Ahmadi K. Gender modifies the effects of education and income on sleep quality of the patients with coronary artery disease. *International cardiovascular research journal*. 2013; 7(4):141–6. PMID: [24757639](https://pubmed.ncbi.nlm.nih.gov/24757639/); PubMed Central PMCID: PMC3987449.
56. Friedman EM, Love GD, Rosenkranz MA, Urry HL, Davidson RJ, Singer BH, et al. Socioeconomic status predicts objective and subjective sleep quality in aging women. *Psychosomatic medicine*. 2007; 69(7):682–91. doi: [10.1097/PSY.0b013e31814ceada](https://doi.org/10.1097/PSY.0b013e31814ceada) PMID: [17766692](https://pubmed.ncbi.nlm.nih.gov/17766692/).
57. Deary IJ, Weiss A, Batty GD. Intelligence and personality as predictors of illness and death: How researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychological Science in the Public Interest*. 2010; 11(2):53–79.
58. Hemalatha A, Gayathri A. Behavioral Interventions and Neuroticism in Young Adulthood. *Global Journal for Research Analysis*. 2013; 2(2):188–9.
59. Krasner MS, Epstein RM, Beckman H, Suchman AL, Chapman B, Mooney CJ, et al. Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *Jama*. 2009; 302(12):1284–93. doi: [10.1001/jama.2009.1384](https://doi.org/10.1001/jama.2009.1384) PMID: [19773563](https://pubmed.ncbi.nlm.nih.gov/19773563/).
60. Hooke GR, Page AC. Predicting outcomes of group cognitive behavior therapy for patients with affective and neurotic disorders. *Behavior modification*. 2002; 26(5):648–59. PMID: [12375379](https://pubmed.ncbi.nlm.nih.gov/12375379/).
61. Chapman BP, Hampson S, Clarkin J. Personality-informed interventions for healthy aging: conclusions from a National Institute on Aging work group. *Dev Psychol*. 2014; 50(5):1426–41. doi: [10.1037/a0034135](https://doi.org/10.1037/a0034135) PMID: [23978300](https://pubmed.ncbi.nlm.nih.gov/23978300/); PubMed Central PMCID: PMC3940665.
62. Sanderson C, Clarkin J. Further use of the NEO-PI-R personality dimensions in differential treatment planning. Costa PT Jr., Widiger TA, editors. Washington (DC), US: American Psychological Association; 2002. 351–75 p.