


RESEARCH ARTICLE

# Association between social support and post-traumatic stress disorder symptoms among Chinese patients with ovarian cancer: A multiple mediation model

Chunli Liu<sup>1,2</sup>, Yi Zhang<sup>3</sup>, Hong Jiang<sup>4</sup>, Hui Wu<sup>2\*</sup>

**1** Library of China Medical University, Shenyang, Liaoning, China, **2** Department of Social Medicine, College of Public Health, China Medical University, Shenyang, Liaoning, China, **3** The First Affiliated Hospital of China Medical University, Heping District, Shenyang, Liaoning, China, **4** School of Public Health, China Medical University, Shenyang, Liaoning, China

\* [hwu@cmu.edu.cn](mailto:hwu@cmu.edu.cn)



## Abstract

Post-traumatic stress disorder (PTSD) symptoms can develop after person experiences one or more traumatic events. Little research, however, has been done on PTSD symptoms of patients with ovarian cancer. The present study aimed to estimate the prevalence of PTSD symptoms in patients with ovarian cancer in China; the effects of demographic and clinical variables on PTSD symptoms; multiple mediation roles in the association between social support and PTSD symptoms in patients with ovarian cancer in China. We collected demographic and clinical information of patients with ovarian cancer in the first and second hospitals of China Medical University between January 1, 2014 and December 31, 2015. Qualified patients were asked to complete the Posttraumatic Stress Disorder Checklist-Civilian Version (PCL-C), Duke-UNC Functional Social Support Questionnaire, Herth Hope Index (HHI), and Resilience Scale-14 (RS-14). 201 patients provided responses. We performed hierarchical linear regression to assess the correlation between social support and PTSD symptoms and bootstrapping to test the mediating role of hope and resilience as potential mediators. After controlling demographic and clinical characteristics, social support negatively correlated with PTSD symptoms ( $\beta = -0.406$ ,  $P < 0.01$ ). Social support explained 14.7% of the variance in PTSD symptoms. Hope and resilience explained 17.0% of the variance in PTSD symptoms. The proportion of the hope mediating effect was 43.37% for social support and the proportion of the resilience mediating effect was 10.64% for social support. Hope and resilience partly mediated the correlation between social support and PTSD symptoms despite accounting for different proportions of the mediating effect. Future intervention plans should pay more attention to social support as well as hope and resilience to prevent, relieve and treat PTSD symptoms.

## OPEN ACCESS

**Citation:** Liu C, Zhang Y, Jiang H, Wu H (2017) Association between social support and post-traumatic stress disorder symptoms among Chinese patients with ovarian cancer: A multiple mediation model. PLoS ONE 12(5): e0177055. <https://doi.org/10.1371/journal.pone.0177055>

**Editor:** Yutaka J. Matsuoka, National Cancer Center, JAPAN

**Received:** August 18, 2016

**Accepted:** April 23, 2017

**Published:** May 5, 2017

**Copyright:** © 2017 Liu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The authors received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Ovarian cancer is a gynecological malignancy disease that causes more deaths than other female reproductive cancers. Most patients are diagnosed ovarian cancer at a late stage, and the prognosis is poor [1–3]. The incidence rate and survival rate vary by geographical region and population. According to epidemiology statistics in UK, ovarian cancer is the sixth most common cancer in women [4], while it is the greatest cause of gynecological cancer death, with a 46% 5-year survival rate. In the US [5], while it is the eighth most common cancer among women, ovarian cancer is the fifth leading cause of cancer-related death among women, and is the deadliest of gynecologic cancers, with a 45.6% 5-year survival rate. In China, ovarian cancer is the second most common cancer in women behind cervical cancer, and the mortality rate is 21.6%, ranking as the deadliest cancer among women [6].

Cancer and its treatment can bring significant physical and psychological trauma. Patients and even their partners are continuously bearing the fear of death as well as pain, stress, and the cost [7–10]. Besides the economic burden and mental pressure, patients with ovarian cancer have to suffer terrible pain from surgery and side effects of chemical and radiation therapy. Approximately one fifth of women self-reported different degrees of distress (more than half with high psychological stress or mental disorder) due to the diagnosis of ovarian cancer and corresponding treatment [11].

Post-traumatic stress disorder (PTSD) is the appearance of delayed issues and continued mental disturbance after an individual experiences, witnesses, or encounters one or more deaths of someone related to themselves or others, or a threat of death or serious injury, according to the Fifth Edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5). The diagnostic criteria of PTSD emphasize direct exposure, witnessing in person, learning that a close relative or close friend experienced trauma, or repeated indirect exposure to adverse events. PTSD includes three types of symptoms: hyper vigilance (always being on alert, trouble sleeping, irritability, difficulty concentrating), avoidance of stimuli related to the traumatic event (such as persons, places, or situations) and intrusive thoughts (re-experiencing the trauma, such as nightmares, extreme distress, frequent and distressing memories). PTSD symptoms can also result in job loss, substance abuse, and other stressful problems [12].

Some researchers have found a high prevalence of PTSD-related symptoms in patients who have suffered from newly diagnosed cancer or cancer resection [13–17]. Among women, a few studies have focused on prevalence and risk factors of PTSD symptoms in patients with breast cancer [18–21] and ovarian cancer [22–23]. For example, there is evidence of a decline in PTSD symptoms rates over time in a sample of 121 UK outpatients with ovarian cancer [22]. Another study surveyed 108 Australian women with ovarian cancer, and found that 9.25% suffered from PTSD symptoms, 5.6% from depression, and 13.9% from anxiety [23]. However, there is little study concerning the prevalence of PTSD symptoms among ovarian cancer patients in China. In addition, PTSD symptoms vary according to a series of variables: time since diagnosis or treatment, type of treatment, income, marital status, cancer stage, and treatment type [24–25]. Similarly, little evidence concerning which demographic and clinical characteristic of patients significantly show impact on the PTSD symptoms among ovarian cancer patients in China has been demonstrated.

Social support refers not only to material support but also to spiritual help that actually or apparently comes from the community, social network, and other people who can be trusted. Social support primarily covers emotional support, social integration or network support, support to improve self-esteem, financial support, information support, and accompanying support, and so on. Social support [26–27] has been found to promote mental adjustment in conditions with chronic high stress like cancer. People with higher levels of social support may

adjust to crises more smoothly [28]. On the contrary, people with lower social support may have more difficulty recovering from trauma.

Does social support or other positive psychological factors show effect on PTSD symptoms after cancer? Recently, social support has shown negative correlations with PTSD symptoms [29–30]. Having a good social support system can prevent you from PTSD symptoms. Dinenberg RE[31] concluded that social support was strongly associated with future PTSD symptoms in patients with cardiovascular disease. They indicated that to help patients from risk of developing PTSD symptoms, optimize the social support level might be a good intervention method. In turn, PTSD symptoms may affect social support. A study on United States veterans from Afghanistan/Iraq era suggested that veterans with PTSD symptoms might have difficulty in accepting or benefitting from social support[32]. Moreover, some studies indicated that social support correlates with recovery from PTSD symptoms [33–34]. Given the collectivist culture in China, social support may be relevant to the outcomes in that social support is particularly important to Chinese women. However, little work has investigated the impact of social support on PTSD symptoms in Chinese patients with ovarian cancer.

Furthermore, if social support and PTSD symptoms are related, hope and resilience may explain or mediate this relationship. Resilience is an individual characteristic of power or quality. For example, psychological resilience, which serves as an ability to adapt to the changing environment flexibly, can accelerate recovery from negative experiences [35–36]. Resilience can give individuals psychological strength to solve a series of problems after disaster or trauma. Resilient individuals are able to utilize their skills and strengths to recover from stress or challenges. However, little evidence has documented the complex relationship between resilience and PTSD symptoms in patients with cancer, especially in ovarian survivors [37]. Hope is also an important impact factor for the lives of cancer patients. Miller defined hope as anticipation of a continued good state or an improved state [38]. To the ill populations, hope is anticipation for a good future, based on the ability of coping with illness, the relationships with others, psychological well-being, and others [38]. What is more, hope means not only the sense of hope, but also the desire for interpersonal relationships, even engagement in the relationships, and the controlling of emotional responses[39]. He developed a model of hope with three dimensions: inner sense of temporality and future (such as presence of goals; scared about the future); positive readiness and expectancy (for example, a sense of direction; life has value and worth); interconnectedness (for instance, give and receive caring/love) [40]. To the cancer patients, continuous negative mood will affect physical and mental state. Many cancer patients have to struggle with the PTSD symptoms including insomnia, nightmares, anxiety and hyper-vigilance and even enormous despair [41–43]. In this case, hope is very essential to the adaption to illness and the recovery from the PTSD symptoms.

Then, is there any evidence that social support correlates with hope or resilience? Quite a few years ago, some researchers have studied this question and provided evidence that social support promote the resilience and hope. Jennifer L. Mattioli[44] has conducted semi-structured interviews to explore and describe what hope and social support mean to the patients who are receiving chemotherapy. Social support from family, friends, and healthcare providers is helpful for patients to deal with their disease treatment. Hope aids in people overall health and well-being. Therefore, it is understandable that people with good social support keep more optimistic toward their life, and then become more active and hopeful to conquer or coexist with cancer. In a word, there may be direct relationship between social support and hope. When social support increases, hope will elevate accordingly[45–46].

Lauren M. Sippel [47] has studied the correlation between social support and individual resilience. The findings suggest that high social support can increase self-confidence, foster more active problem solving, and then elevate individual resilience in the capacity to adapt

well in the face of adversity. Also, findings[48] from laboratory-based studies in which participants undergo cardiovascular monitoring have provided evidence that social support can nourish further emotional resilience, or in other word, social support may be effective in elevating resilience. Phoenix Mo [49] examined the relationship between social supports, resilience, hopelessness, posttraumatic growth among children of HIV-infected parents in China. Results suggested that social support had a significant positive relationship with resilience and Posttraumatic growth. Wilks SE found that social support from family positively influenced resilience in Alzheimer's disease caregivers[50].

In this perspective, we posit that in our sample of ovarian cancer patients, social support show negative effect on PTSD symptoms and positive effect on both hope and resilience. Besides that, hope and resilience may mediate the relation between social support and PTSD symptoms. The aims of the present study were to estimate: (1) the prevalence of PTSD in patients with ovarian cancer in China; and (2) the effects of demographic and clinical variables on PTSD symptoms; and (3) a multiple mediation in the association between social support and PTSD symptoms in patients with ovarian cancer in China.

## Methods

### Ethics statement

The Committee for Human Trials of the China Medical University revised and approved the research and study procedures were in accordance with the ethical standards. All patients gave their permission to participate after being verbally informed of the study protocol. Participation was completely voluntary and anonymous. We protected the privacy of personal data processing and maintained confidentiality of individual records and accounts. Participation in this study did not affect future free medical examination and treatment that is standard in China.

### Study design and sample

From January 1, 2014 to December 31, 2015, we used convenience sampling to recruit patients with ovarian cancer in the first and second hospitals of China Medical University, which are important providers of gynecologic surgery in the northeastern region of China. Patients who satisfied the following inclusion criteria were enrolled as potential subjects: (1) at least 18 years old when diagnosed with ovarian cancer, (2) histological evidence of ovarian cancer, (3) able to speak and read Chinese and fill in the questionnaire, and (4) clear consciousness and cognition (can correctly answer place and characters within 30 seconds). Exclusion criteria were as follows: (1) those with psychiatric problems such as depression, anxiety, or other psychiatric disorders before being diagnosed with cancer, (2) those with intellectual abnormalities, and (3) patients who had other types of cancers. After obtaining written informed consent from patients, clinical data were collected from their medical records and a structured questionnaire was distributed. A total of 220 patients with ovarian cancer were enrolled. Five patients refused to join the survey. Of the 215 eligible patients, 14 were excluded from analysis (more than 30% missing data). Finally, we received effective responses from 201 patients (effective response rate of 93.5%).

### Demographic and clinical characteristics

We obtained four demographic characteristics including age, education and marital status. Educational level was divided into primary school, secondary school, and junior college or above. Income was divided into  $\leq 2000$ , 2001–3000 and  $> 3000$ . Three clinical characteristics

obtained were cancer stage, treatment type, and metastasis. Cancer stage was divided into I, II, III, and IV. Treatment type was divided into no treatment, chemotherapy, surgery, and combined treatment. Metastasis was divided into no and yes.

### Post-traumatic stress disorder symptoms

We used the PTSD Checklist-Civilian Version (PCL-C) [51], a 17-item questionnaire to assess the degree of PTSD symptoms. Each item was rated on a 5-point Likert scale (1 = not at all; 5 = extremely) based on how much the symptom was bothersome in the last month. Total scores range from 17 to 85. Scores equal to or greater than 44 are considered to indicate PTSD symptoms [52]. In this study, patients were asked to comment on items of the PCL-C in their experience with ovarian cancer and its treatment. The Chinese version of the PCL-C has been used in the Chinese population, demonstrating sufficient reliability [53–54]. In our study, Cronbach's alpha for the total scale was 0.940.

### Social support

We chose the Duke-UNC Functional Social Support Questionnaire [55] to assess social support. This is an 8-item questionnaire, where each item is rated on a 5-point Likert scale (1 = much less than I would like; 5 = as much as I would like) based on satisfaction with perceived social support. Studies using the Duke-UNC Functional Social Support in Chinese populations have shown that it has good reliability [56]. In our study, Cronbach's alpha for the total scale was 0.886.

### Hope

We selected the Herth Hope Index (HHI) [57], which is an adaptation of Herth Hope Scale (HHS) to assess hope of patients. It is a questionnaire including 12 items, and each item is rated on a 4-point Likert scale (1 = strongly disagree; 4 = strongly agree). The total score ranges from 12 to 48. The Chinese version of the HHI has been used in Chinese patients with cancer, and demonstrated satisfactory reliability and internal consistency [58]. In this study, Cronbach's alpha for the total scale was 0.840.


### Resilience

We used the Resilience Scale-14 (RS-14) [59] to test the resilience level of patients. The RS-14 is a 14-item questionnaire, where each item is rated on a 4-point Likert scale (1 = entirely not match; 4 = completely match). The total score ranges from 14 to 56. Some studies have verified the reliability of the Chinese version [60]. In our study, the Cronbach's alpha value for the total scale was 0.906.

### Statistical analysis

We used SPSS 17.0 for Windows to perform all analyses. All statistical tests were two-sided ( $\alpha = 0.05$ ). Descriptive statistics for demographic and clinical variables are shown with mean, standard deviation (SD), number (N), and percentage (%), as appropriate. Variation in PTSD symptoms was examined with regard to demographic and clinical variables using independent samples t-tests and one-way analyses of variance (ANOVAs). Correlations among PTSD symptoms, social support, hope, and resilience were preliminarily examined by Pearson's correlation analysis.

We performed hierarchical multiple regression analysis to explore the effects of groups of independent variables on PTSD symptoms. There were three blocks of independent



**Fig 1. Theoretical model of the mediating role of hope and resilience on the relationship between social support and PTSD symptoms.** (c) The association between social support and PTSD symptoms; (a<sub>1</sub>) The relationship between social support and hope; (b<sub>1</sub>) the relationship of hope with post-traumatic stress disorder (PTSD) symptoms after controlling the independent variables; (a<sub>2</sub>) The relationship between social support and resilience; (b<sub>2</sub>) the relationship of resilience with post-traumatic stress disorder (PTSD) symptoms after controlling the independent variables; (c') the association between social support and PTSD symptoms after adding hope and resilience as mediators.

<https://doi.org/10.1371/journal.pone.0177055.g001>

variables. In block 1, all demographic and clinical variables were entered as control variables. Because marital status, educational level, income, cancer stage, treatment type, and metastasis are categorical variables without a linear trend, we set dummy variables for them [61]. In block 2, social support was added. Hope and resilience were added in block 3.

We used asymptotic and resampling strategies to examine the mediating roles of hope and resilience on the association between social supports on PTSD symptoms [62]. Social support was modeled as an independent variable, with PTSD symptoms as the dependent variable, and hope and resilience as mediators (as shown in Fig 1). Age, income and cancer stage were considered as covariates. In the first step, the aim is to identify the correlation between social support and PTSD symptoms (the c path). In the second step, the aim is to examine the mediation of hope and resilience (the a × b path). Standards we used to judge the mediating role are the same as we have used in previous studies [63]. We used bias-corrected and accelerated 95% confidence intervals (BCa 95% CIs). If zero is not included in the BCa 95% CI, then the mediation role (c') is statistically significant. In the present study, we used 5000 bootstrap samples.

Fig 1 depicts a multiple mediation model with two mediators. Step 1 represents the total effect of social support on PTSD symptoms (path c). Step 2 represents both the direct effect of social support on PTSD symptoms (path c') and the indirect effects of social support on PTSD symptoms via mediator hope and resilience. The specific indirect effect is the product of a and b. Hence, the total indirect effect can be calculated by the equation:  $c - c' = a_1 \times b_1 + a_2 \times b_2$

**Table 1. Demographic and clinical characteristics (N = 201).**

Variable	N (%)	PTSD symptoms(Mean ± SD)	P-value
Demographic variables			
Age (years)			0.474
≤45	22 (10.9)	32.82 ± 10.82	
46–55	82 (40.8)	35.23 ± 11.09	
≥ 56	97 (48.3)	35.99 ± 10.99	
Marital status			0.263
Married/living with a partner	183 (91.0)	35.06 ± 10.85	
Single/widowed/divorced	18 (9.0)	38.11 ± 12.47	
Educational level			0.137
Primary school	54 (26.8)	37.28 ± 9.33	
Secondary school	120 (59.7)	35.18 ± 11.87	
Junior college or above	27 (13.4)	32.15 ± 9.51	
Income (yuan per month)			0.024
≤ 2000	98 (48.7)	37.27 ± 10.80	
2001–3000	61 (30.3)	34.59 ± 11.99	
> 3000	42 (20.8)	31.90 ± 9.05	
Clinical variables			
Cancer stage			0.010
I	36 (17.9)	30.69 ± 8.96	
II	19 (9.5)	33.58 ± 7.38	
III and IV	146 (72.6)	36.71 ± 11.54	
Treatment type			0.864
No treatment	13 (6.5)	36.62 ± 13.96	
Chemotherapy	60 (29.8)	35.55 ± 13.11	
Surgery	13 (6.5)	33.08 ± 5.12	
Combined treatment	115 (57.2)	35.33 ± 9.96	
Metastasis			0.299
No	187 (93.0)	35.11 ± 11.24	
Yes	14 (7.0)	38.29 ± 6.72	

<https://doi.org/10.1371/journal.pone.0177055.t001>

## Results

### Demographic and clinical statistics and prevalence of PTSD symptoms

Demographic and clinical characteristics of the participants are shown in [Table 1](#). Of the 201 respondents, age ranged from 24 to 79, and the average age was  $55.28 \pm 9.65$  years. Approximately 91.0% of respondents were married or living with a partner, and 59.7% received secondary school education. Nearly one half of respondents had a household income equal to or less than 2000. Of all respondents, 72.6% were diagnosed at III and IV cancer stage, and 57.2% received combined treatment. Metastasis was not present in 93.0% of respondents. Based on the cut-off values recommended by Blanchard, E. B. ( $\geq 44$ ), the prevalence of PTSD symptoms among ovarian cancer patients in China was 21.9%.

### Effects of demographic and clinical variables on PTSD symptoms

As shown in [Table 1](#), respondents whose income was equal to or less than 2000 had a high level of PTSD symptoms ( $37.27 \pm 10.80$ ), as compared to respondents with income in the 2001–3000 range ( $34.59 \pm 11.99$ ) or higher than 3000 ( $31.90 \pm 9.05$ ). In addition, respondents

**Table 2. Means, SDs, and correlations among study variables.**

Variables	Mean ± SD	Social support	Hope	Resilience	PTSD symptoms
Social Support	28.98 ± 6.88	1	0.46**	0.25**	-0.41**
Hope	35.40 ± 4.11		1	0.39**	-0.54**
Resilience	37.29 ± 7.28			1	-0.36**
PTSD symptoms	35.33 ± 11.00				1

\*\*P < 0.01.

<https://doi.org/10.1371/journal.pone.0177055.t002>

whose cancer stage was III and IV had a high level of PTSD symptoms (36.71 ± 11.54), as compared to stages I (30.69 ± 8.96) and II (33.58 ± 7.38).

### Pearson’s correlations among PTSD symptoms, social support, hope, and resilience

Pearson’s correlation coefficients were calculated between social support, hope, resilience, and PTSD symptoms. As shown in Table 2, PTSD symptoms were negatively associated with the three positive psychological variables (social support:  $r = -0.41$ ,  $P < 0.01$ ; hope:  $r = -0.54$ ,  $P < 0.01$ ; resilience:  $r = -0.36$ ,  $P < 0.01$ ). Social support was positively related with hope ( $r = 0.46$ ,  $P < 0.01$ ) and resilience ( $r = 0.25$ ,  $P < 0.01$ ). Moreover, hope was positively correlated with resilience ( $r = 0.39$ ,  $P < 0.01$ ).

### Association between social support and PTSD symptoms by hierarchical linear regression

Table 3 shows the results of the hierarchical regression analysis. After controlling for age, income, and cancer stage, social support was negatively associated with PTSD symptoms ( $\beta = -0.406$ ,  $P < 0.01$ ) and explained 14.7% of the variance in PTSD symptoms. While hope and resilience were negatively associated with PTSD symptoms ( $\beta = -0.459$ ,  $P < 0.01$ ;  $\beta = -0.135$ ,  $P < 0.05$ ) and explained 17.0% of the variance in PTSD symptoms. In addition, after adding hope and resilience in the regression model of PTSD symptoms, the regression coefficient (absolute value of regression coefficient when it is negative) for social support reduced (from  $\beta = -0.406$  to  $\beta = -0.136$ ). Hence, we preliminarily consider that hope and resilience might play a mediating role between social support and PTSD symptoms. However, this point still needs further test by bootstrapping.

### Bootstrapping test of the mediating role of hope and resilience

Path coefficients a (between social support and mediators) and b (between mediators and PTSD symptoms), a×b products are presented in Table 4. Social support was significantly and positively associated with hope and resilience. Consistent with the results from hierarchical multiple regression analysis, hope and resilience were significantly and negatively associated with PTSD symptoms after controlling for age, income, cancer stage and social support. Each BCa 95% CI for a × b of hope and resilience excluding 0 indicated their significant mediation when they are added in the model. Thus, significant mediating roles of hope (a×b = -0.274, BCa 95% CI: -0.405, -0.158), and resilience (a×b = -0.067, BCa 95% CI: -0.145, -0.008) on the association between social support and PTSD symptoms were revealed among patients with ovarian cancer in China.

We used the formula (a × b / c) to calculate the proportion of mediation role. The proportion of the mediating effect accounted for by hope was 43.37% for social support, and that of



**Table 3. Hierarchical linear regression for exploring the associated variables of PTSD symptoms.**

Variables	PTSD symptoms		
	Step 1 ( $\beta$ )	Step 2 ( $\beta$ )	Step 3 ( $\beta$ )
<b>Block 1</b>			
Age	0.052	0.088	0.015
Income 1	0.196	0.130	0.046
Income 2	0.253*	0.241*	0.175
Cancer stage 1	0.063	0.075	0.024
Cancer stage 2	0.223*	0.162*	0.158*
<b>Block 2</b>			
Social support		-0.406**	-0.136
<b>Block 3</b>			
Hope			-0.459**
Resilience			-0.135*
F	1.779	4.769**	8.891**
Adjusted R <sup>2</sup>	0.045	0.197	0.372
$\Delta R^2$	0.102	0.147**	0.170**

\*P< 0.05,

\*\*P< 0.01.

Income 1 means 2001–3000 vs. > 3000; Income 2 means ≤2000 vs.>3000;

Cancer stage 1 means II vs. I; Cancer stage 2 means III and IV vs. I.

<https://doi.org/10.1371/journal.pone.0177055.t003>

resilience was 10.64%. Hence, the total effect of social support on PTSD symptoms by two mediators' role (from hope and resilience together) was 54.00%.

## Discussion

We used self-report checklist to measure the prevalence of PTSD symptoms among patients with ovarian cancer in China. The prevalence of PTSD symptoms was 21.89%, which is higher than that among Chinese individuals with hematological malignancies(10.7%) [64], Australian women with ovarian cancer (9.25%) [23], and lower than that among Chinese women with bladder and kidney cancer (25.2%) [65]. The prevalence of PTSD symptoms correlates with the PTSD checklist version[66] and the cut-off score that is used [67–68]. In addition, the comparison of the prevalence of PTSD would be challenging without considering the different study procedures and populations across studies [69].

**Table 4. Bootstrapping test of the indirect effect of hope and resilience on the relation between social support and PTSD symptoms.**

(N = 201)	Mediators	c	a	b	c'	a × b (BCa 95% CI)
PTSD symptoms	Hope	-0.631**	0.263**	-1.034**	-0.292**	-0.274 (-0.405, -0.158)
	Resilience	-0.631**	0.283**	-0.238*	-0.292**	-0.067 (-0.145, -0.008)

Notes: c: correlation of social support with PTSD symptoms; a<sub>1</sub>: correlation of social support with hope; b<sub>1</sub>: correlation of hope with PTSD symptoms after controlling for the predictor variables; a<sub>2</sub>: correlation of social support with resilience; b<sub>2</sub>: correlation of resilience with PTSD symptoms after controlling for the predictor variables; c': the association between social support and PTSD symptoms after adding hope and resilience as two mediators; a × b: the product of a and b; BCa 95% CI: the bias-corrected and accelerated 95% confidence interval; age, income and cancer stage were covariates;

\* p<0.05,

\*\* p<0.01.

<https://doi.org/10.1371/journal.pone.0177055.t004>

Demographic and clinical characteristics also affect the level of PTSD symptoms. Income and cancer stage are two significant factors that affect PTSD symptoms. Given the significant economic pressure in the treatment of ovarian cancer, the psychological burden of patients with a lower income (less than 2000) was heavier than it was for those with a higher income (more than 3000). It is indicated from our results that the level of PTSD symptoms in stages III and IV was higher than those in stage I. Thus, in psychological counseling and clinical and family care, we should be more concerned with two important factors to PTSD: income and cancer stage.

The present results indicate that social support significantly and negatively correlates with PTSD symptoms in patients with ovarian cancer in China ( $P < 0.01$ ), in accordance with prior studies. Jacobsen [70] found that lower social support predicts greater PTSD symptoms severity. In other words, higher levels of PTSD symptoms are associated with less social support [71]. In recent studies, Yang [65] showed that social support from one's family is significantly associated with PTSD symptoms in Chinese individuals with bladder and kidney cancers. Liu [64] demonstrated that PTSD symptoms are negatively associated with social support in patients with hematological malignancies in China. Wang [72] also confirmed that social support is negatively associated with PTSD symptoms in Chinese patients with central system tumors.

In addition, our results show that hope and resilience correlated negatively with PTSD symptoms, concordant with prior results [73] despite different samples. However, among Chinese patients with hematological malignancies, hope and resilience have not shown significant negative associations with PTSD symptoms [64]. In contrast, Besser [74] showed that resilience negatively associated with acute PTSD symptoms. Resilience is inversely correlated with self-reported PTSD symptoms in most survivors of critical illness [75]. Another study found that nurses with PTSD symptoms had significantly lower total resilience scores compared to those without PTSD symptoms [76]. Wright [77] reported that kidnapping victims without PTSD symptoms reported higher individual resilience. Joscelyne [78] found that responders with high levels of resilience reported minimal symptoms of PTSD symptoms. In general, to our knowledge, relevant studies remain limited. Therefore, we provided reliable evidence of the association between hope, resilience, and PTSD symptoms in Chinese patients with ovarian cancer.

Moreover, the results of the bootstrapping test indicate that hope and resilience act as mediators in the association between social support and PTSD symptoms in Chinese patients with ovarian cancer. The mediator tested in our study differs from most previous studies [72, 79–81], in which authors have mostly chosen self-efficacy [72], emotional abuse [79], sleep disturbances [80], and negative perception of social support [81]. Although Li [73] also suggested that resilience partially mediated the relation between psychological stress with depressive and anxiety symptoms, their research differed from ours in terms of population, dependent and independent variables, and even the resilience scale to some extent. Although we all used the Resilience Scale-14, the original scale of the Chinese revision we used is from Block J [59]. We used a 4-point Likert scale, while Li [73] used a 7-point Likert scale. Both studies, however, showed that resilience is a significant negative mediator.

It is noteworthy that the proportion of the mediating effect of hope for social support (43.37%) was higher than the proportion of the mediating effect of resilience (10.64%). This might be because of potentially different features of hope and resilience. Hope is a multidimensional dynamic life force. Hope is characterized with confident and uncertain expectation of achieving a good future. If the individuals are hopeful, they will be confident that things will turn out to be fine. Then they are more likely to dedicate themselves to meaningful work, for example, participating in the social network; committing to family relationships; relieving stress and tension; and keeping optimistic attitude toward life. Hence, hope probably plays a relatively more important role in tapering off the PTSD symptoms. Resilience represents a homeostatic rebounding to a prior level of functioning after a stressor or trauma. If a patient is

high in resilience, she will handle or rebound from adversity successfully. Although the mediating effect of resilience is not as high as hope for social support, we should not ignore it. Inter-correction between hope and resilience may occur. An individual who is more resilient or, who has gone through a resilient recovery, probably may be more hopeful. In addition, an individual, who has seen the hope of the future, may be more likely to resume their normal pre-stressor level of functioning. A strong correlation between resilience and hope has been indicated among advanced cancer patients (0.63;  $p < 0.05$ ) [82] and middle aged adults (0.59;  $p < 0.05$ ) [83].

In the present study, social support explained 14.7% of the variance in PTSD symptoms, while hope and resilience together explained 17.0% of the variance in PTSD symptoms. It seems that intrapersonal psychological resources (hope and resilience) are more important than perceived interpersonal resources (social support), or at least that they are about equal. Future studies should examine the effect of these different kinds of psychological resources on PTSD symptoms.

Finally, it is necessary to pay more attention to and invest more resources into not only social support, but also hope and resilience concerning the prevention of PTSD symptoms among patients with ovarian cancer. Since social support correlates with PTSD symptoms negatively, we should pay enough emphasis on the assessment for deficits in social support in ovarian cancer patients. In addition, clinical doctors and researchers should be aware that hope and resilience might play a role in the outcomes among cancer patients. For example, Fava GA developed psychotherapeutic strategy to increase psychological well-being and resilience in mood and anxiety disorders [84].

Some limitations should be mentioned. First, the present study was a cross-sectional study. Accordingly, we could not draw important conclusions regarding the trends of the relation among social support, hope, resilience, and PTSD symptoms by time. Thus, causality of each variable could not be determined. However, there might be reverse causality. For example, high resilience can possibly promote the social support [85]. An evidence has confirmed that greater resilience enables individuals maintain better social networks and seek new social support resources, and own higher perceived social support; Moreover, high-hope individuals tend to be social beings, they are good at interpersonal interactions and prefer to explore social support [86]. Furthermore, to individuals with PTSD symptoms, it is possible that they will feel inferior, autistic, and refuse to mention their misfortune story to others and even just explore social support. For individuals with PTSD, the acceptance of social support may be particularly difficult because of core symptoms such as avoidance, alienation, detachment and emotional numbing [32].

Second, we used convenience sampling, not stratified sampling, and we only recruited patients with ovarian cancer from two hospitals of China Medical University. Third, we only focused on some psychological resources; other factors such as self-efficacy, optimism, and self-esteem needed further consideration. Despite these limitations, we assessed the mediation with multiple mediators (hope and resilience) in the association between social support and PTSD symptoms in patients with ovarian cancer in China.

In the future, we will make further research: to use longitudinal designing methods to infer causality; to consider the relation between self-efficacy, self-esteem and optimism with PTSD symptoms; to recruit ovarian patients from the south and west regions in China.

## Conclusions

This study provides a self-report prevalence of PTSD symptoms among Chinese patients with ovarian cancer. After controlling demographic and clinical characteristics, we demonstrated

multiple mediation roles of hope and resilience in the association between social support and PTSD symptoms. Social support correlated negatively with PTSD symptoms. Social support explained 14.7% of the variance in PTSD symptoms. When hope and resilience were added, we found they together explained 17.0% of the variance in PTSD symptoms. The findings indicated that hope and resilience partly mediated the correlation between social support and PTSD symptoms. Patients with higher levels of hope and resilience might show a strengthened effect of social support on PTSD symptoms. Interventions to relieve PTSD symptoms should be focused on not only social support, but also hope and resilience.

## Supporting information

**S1 Data. Supporting dataset.** The supporting dataset includes the data underlying our findings in this study.  
(XLSX)

## Author Contributions

**Conceptualization:** HW.

**Data curation:** CLL HW.

**Formal analysis:** CLL HW.

**Investigation:** YZ HJ HW.

**Methodology:** HW CLL.

**Resources:** CLL.

**Software:** HW.

**Supervision:** HW.

**Writing – original draft:** CLL HW.

**Writing – review & editing:** CLL HW.

## References

1. Suryawanshi S, Vlad AM, Lin HM, Mantia-Smaldone G, Laskey R, Lee M, et al. Plasma microRNAs as novel biomarkers for endometriosis and endometriosis-associated ovarian cancer. *Clin Cancer Res.* 2013; 19(5):1213–24. <https://doi.org/10.1158/1078-0432.CCR-12-2726> PMID: 23362326
2. Bjørge T, Engeland A, Hansen S, Tropé CG. Prognosis of patients with ovarian cancer and borderline tumours diagnosed in Norway between 1954 and 1993. *Int J Cancer.* 1998; 75(5):663–70. PMID: 9495231
3. Mackay HJ, Brady MF, Oza AM, Reuss A, Pujade-Lauraine E, Swart AM, et al. Prognostic relevance of uncommon ovarian histology in women with stage III/IV epithelial ovarian cancer. *Int J Gynecol Cancer.* 2010; 20(6):945–52. <https://doi.org/10.1111/IGC.0b013e3181dd0110> PMID: 20683400
4. Office for National Statistics. Cancer Registration Statistics. England Statistical bulletins. July 2015. <http://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/cancerregistrationstatisticsengland/2014>.
5. National Cancer Institute. SEER Stat Fact Sheets: Ovarian Cancer. 2016. <http://seer.cancer.gov/statfacts/html/ovary.html>
6. Chen W, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, et al. Cancer Statistics in China, 2015. *CA Cancer J Clin.* 2016; 66(2): 115–132.
7. Eskander R., Chang J, Ziogas A, Anton-Culver H, Bristow RE. Hospital readmission (30-day) following surgery for advanced-stage ovarian cancer: Analysis of risk factors and cost using the SEER-Medicare database. *Gynecologic Oncology.* 2014; 133:44. <http://dx.doi.org/10.1016/j.ygyno.2014.03.127>

8. Plamadeala V, Kelley JL, Chan JK, Krivak TC, Gabrin MJ, Brower SL, et al. A cost-effectiveness analysis of a chemoresponse assay for treatment of patients with recurrent epithelial ovarian cancer. *Gynecologic Oncology*. 2015; 136(1):94–8. <https://doi.org/10.1016/j.ygyno.2014.11.019> PMID: [25462203](https://pubmed.ncbi.nlm.nih.gov/25462203/)
9. Poonawalla IB, Lairson DR, Chan W, Piller LB, Du XL. Cost-Effectiveness of Neoadjuvant Chemotherapy versus Primary Surgery in Elderly Patients with Advanced Ovarian Cancer. *Value in Health*. 2015; 18(4):387–395. <https://doi.org/10.1016/j.jval.2015.01.005> PMID: [26091592](https://pubmed.ncbi.nlm.nih.gov/26091592/)
10. Hamilton AB. Psychological aspects of ovarian cancer. *Cancer Invest*, 1999; 17(5):335–41. PMID: [10370361](https://pubmed.ncbi.nlm.nih.gov/10370361/)
11. Norton TR, Manne SL, Rubin S, Carlson J, Hernandez E, Edelson MI, et al. Prevalence and predictors of psychological distress among women with ovarian cancer. *J Clin Oncol*. 2004; 22(5):919–926 <https://doi.org/10.1200/JCO.2004.07.028> PMID: [14990648](https://pubmed.ncbi.nlm.nih.gov/14990648/)
12. Kangas M, Henry JL, Bryant RA. Posttraumatic stress disorder following cancer: A conceptual and empirical review. *Clinical psychology review*. 2002; 22(4): 499–524. [http://dx.doi.org/10.1016/S0272-7358\(01\)00118-0](http://dx.doi.org/10.1016/S0272-7358(01)00118-0). PMID: [12094509](https://pubmed.ncbi.nlm.nih.gov/12094509/)
13. Jeantieu M, Gaillat F, Antonini F, Azoulay E, Martin C, Thomas P, et al. Postoperative pain and subsequent PTSD-related symptoms in patients undergoing lung resection for suspected cancer. *J Thorac Oncol*. 2014; 9(3):362–9. <https://doi.org/10.1097/JTO.000000000000084> PMID: [24496000](https://pubmed.ncbi.nlm.nih.gov/24496000/)
14. Posluszny DM, Dougall AL, Johnson JT, Argiris A, Ferris RL, Baum A, et al. Posttraumatic Stress Disorder (PTSD) Symptoms in Newly Diagnosed Head and Neck Cancer Patients and their Partners. *Head Neck*. 2015; 37(9): 1282–1289. <https://doi.org/10.1002/hed.23760> PMID: [24817018](https://pubmed.ncbi.nlm.nih.gov/24817018/)
15. Palgi Y, Shrira A, Haber Y, Wolf JJ, Goldray O, Shacham-Shmueli E, et al. Comorbidity of posttraumatic stress symptoms and depressive symptoms among gastric cancer patients. *Eur J Oncol Nurs*. 2011; 15(5):454–8. <https://doi.org/10.1016/j.ejon.2010.11.011> PMID: [21220210](https://pubmed.ncbi.nlm.nih.gov/21220210/)
16. Thomas KS, Bower JE, Williamson TJ, Hoyt MA, Wellisch D, Stanton AL, et al. Post-traumatic disorder symptoms and blunted diurnal cortisol production in partners of prostate cancer patients. *Psychoneuroendocrinology*. 2012; 37(8):1181–90. <https://doi.org/10.1016/j.psyneuen.2011.12.008> PMID: [22222119](https://pubmed.ncbi.nlm.nih.gov/22222119/)
17. Dahl AA, Østby-Deglum M, Oldenburg J, Bremnes R, Dahl O, Klepp O, et al. Aspects of posttraumatic stress disorder in long-term testicular cancer survivors: cross-sectional and longitudinal findings. *J Cancer Surviv*. 2016; 10(5):842–9 <https://doi.org/10.1007/s11764-016-0529-4> PMID: [26920871](https://pubmed.ncbi.nlm.nih.gov/26920871/)
18. Vin-Raviv N, Hillyer GC, Hershman DL, Galea S, Leoce N, Bovbjerg DH, et al. Racial disparities in post-traumatic stress after diagnosis of localized breast cancer: the BQUAL study. *J Natl Cancer Inst*. 2013; 105(8):563–72. <https://doi.org/10.1093/jnci/djt024> PMID: [23434900](https://pubmed.ncbi.nlm.nih.gov/23434900/)
19. Palesh O, Koopman C. Breast cancer: Post-traumatic stress disorder—prevalent and persistent. *Nat Rev Clin Oncol*. 2013; 10(5):252–4. <https://doi.org/10.1038/nrclinonc.2013.49> PMID: [23568418](https://pubmed.ncbi.nlm.nih.gov/23568418/)
20. Kwakkenbos L, Coyne JC, Thombs BD. Prevalence of posttraumatic stress disorder (PTSD) in women with breast cancer. *J Psychosom Res*. 2014; 76(6):485–6. <https://doi.org/10.1016/j.jpsychores.2014.03.103> PMID: [24840144](https://pubmed.ncbi.nlm.nih.gov/24840144/)
21. O'Connor M, Christensen S, Jensen AB, Møller S, Zachariae R. How traumatic is breast cancer? Post-traumatic stress symptoms (PTSS) and risk factors for severe PTSS at 3 and 15 months after surgery in a nationwide cohort of Danish women treated for primary breast cancer. *Britain Journal of Cancer*. 2011; 104(3):419–26.
22. Gonçalves V, Jayson G, Tarrier N. A longitudinal investigation of posttraumatic stress disorder in patients with ovarian cancer. *J Psychosom Res*. 2011; 70(5):422–31. <https://doi.org/10.1016/j.jpsychores.2010.09.017> PMID: [21511072](https://pubmed.ncbi.nlm.nih.gov/21511072/)
23. Shand LK, Brooker JE, Burney S, Fletcher J, Ricciardelli LA. Symptoms of posttraumatic stress in Australian women with ovarian cancer. *Psycho-oncology*. 2015; 24(2):190–196 <https://doi.org/10.1002/pon.3627> PMID: [25044162](https://pubmed.ncbi.nlm.nih.gov/25044162/)
24. Kangas M, Henry JL, Bryant RA. Posttraumatic stress disorder following cancer: A conceptual and empirical review. *Clinical Psychology Review*. 2002; 22(4):499–524. PMID: [12094509](https://pubmed.ncbi.nlm.nih.gov/12094509/)
25. Andrykowski MA, Cordova MJ. Factors associated with PTSD symptoms following treatment for breast cancer: test of the Andersen model. *Journal of Traumatic Stress*. 1998; 11(2):189–203. <https://doi.org/10.1023/A:1024490718043> PMID: [9565911](https://pubmed.ncbi.nlm.nih.gov/9565911/)
26. Dunkel SC. Social Support and Cancer: Findings Based on Patient Interviews and Their Implications. *Journal of Social Issues*. 1984; 40(4):77–98.
27. Holland KD, Holahan CK. The relation of social support and coping to positive adaptation to breast cancer. *Psychology & Health*. 2003; 18(1):15–29. <http://dx.doi.org/10.1080/0887044031000080656>

28. Pinar G, Okdem S, Buyukgonenc L, Ayhan A. The relationship between social support and the level of anxiety, depression, and quality of life of Turkish women with gynecologic cancer. *Cancer Nursing*. 2012; 35(35):229–35.
29. McCanlies EC, Mnatsakanova A, Andrew ME, Burchfiel CM, Violanti JM. Positive psychological factors are associated with lower PTSD symptoms among police officers: post Hurricane Katrina. *Stress Health*. 2014; 30(5):405–15. <https://doi.org/10.1002/smi.2615> PMID: 25476965
30. Sripada RK, Pfeiffer PN, Rauch SA, Bohnert KM. Social support and mental health treatment among persons with PTSD: results of a nationally representative survey. *Psychiatr Serv*. 2015; 66(1):65–71. <https://doi.org/10.1176/appi.ps.201400029> PMID: 25269889
31. Dinenberg RE, McCaslin SE, Bates MN, Cohen BE. Social support may protect against development of posttraumatic stress disorder: findings from the Heart and Soul Study. *Am J Health Promot*. 2014; 28(5):294–7. <https://doi.org/10.4278/ajhp.121023-QUAN-511> PMID: 23941102
32. Brancu M, Thompson NL, Beckham JC, Green KT, Calhoun PS, Elbogen EB, et al. The impact of social support on psychological distress for U.S. Afghanistan/Iraq era veterans with PTSD and other psychiatric diagnoses. *Psychiatry Res*. 2014; 217(1–2):86–92. <https://doi.org/10.1016/j.psychres.2014.02.025> PMID: 24679515
33. Auxéméry Y, L'Encephale. Posttraumatic stress disorder (PTSD) as a consequence of the interaction between an individual genetic susceptibility, a traumatogenic event and a social context. *Encephale*. 2012; 38(5):373–80 <https://doi.org/10.1016/j.encep.2011.12.003> PMID: 23062450
34. Dai W, Chen L, Tan H, Wang J, Lai Z, Kaminga AC, et al. Association between social support and recovery from post-traumatic stress disorder after flood: a 13–14 year follow-up study in Hunan, China. *BMC Public Health*. 2016; 16:194. <https://doi.org/10.1186/s12889-016-2871-x> PMID: 26924178
35. Luthar S, Cicchetti D, Becker B. The Construct of Resilience: A Critical Evaluation and Guidelines for Future Work. *Child Development*. 2000; 71(3): 543–562. PMID: 10953923
36. Kathleen T, Janyce D. Resilience: A historical review of the construct. *Holistic Nursing Practice*. 2004; 10: 45–48.
37. Wenzel LB, Donnelly JP, Fowler JM, Habbal R, Taylor TH, Aziz N, et al. Resilience, reflection, and residual stress in ovarian cancer survivorship: a gynecologic oncology group study. *Psychooncology*. 2002; 11(2):142–53. PMID: 11921330
38. Miller Fitzgerald J. Hope: a construct central to nursing. *Nurs Forum*. 2007; 1:12–9.
39. Herth K. Development and refinement of an instrument to measure hope. *Research & Theory for Nursing Practice*, 1991; 5(5):39–51.
40. Phillips-Salimi CR, Haase JE, Kintner EK, Monahan PO, Azzouz F. Psychometric properties of the Herth Hope Index in adolescents and young adults with cancer. *Journal of nursing measurement*, 2007, 15(1): 3–23. PMID: 17665817
41. Tanimukai H, Murai T, Okazaki N, Matsuda Y, Okamoto Y, Kabeshita Y, et al. An observational study of insomnia and nightmare treated with trazodone in patients with advanced cancer. *American Journal of Hospice and Palliative Medicine*, 2013, 30(4): 359–362. <https://doi.org/10.1177/1049909112452334> PMID: 22777411
42. Mehnert A, Berg P, Henrich G, Herschbach P. Fear of cancer progression and cancer—related intrusive cognitions in breast cancer survivors. *Psycho—Oncology*, 2009, 18(12): 1273–1280. <https://doi.org/10.1002/pon.1481> PMID: 19267364
43. Miller K, Massie MJ. Depression and anxiety. *The Cancer Journal*, 2006, 12(5): 388–397. PMID: 17034675
44. Mattioli JL, Repinski R, Chappy SL. The meaning of hope and social support in patients receiving chemotherapy. *Oncology Nursing Forum*. 2008; 35(5):822–9. <https://doi.org/10.1188/08.ONF.822-829> PMID: 18765329
45. Bastani F, Sayahi S, Haghani H. Social support and relationship with hope among mothers of a child with leukemia. *Journal of Clinical Nursing and Midwifery*. 2012; 1(1): 23–32. Available from: <http://en.journals.sid.ir/ViewPaper.aspx?ID=377094>
46. Naoko M, Hongjun Y, Donald H. Functional declines, socialsupport and mental health in the elderly: Does living in a state supportive of home andcommunity-based services make a difference? *Social Science & Medicine*. 2010; 70(7):1050–1058.
47. Sippel LM, Pietrzak RH, Charney DS, Mayes LC, Southwick SM. How does social support enhance resilience in the trauma-exposed individual? *Ecology & Society*. 2015; 20(4):10. Available from: [https://www.researchgate.net/publication/283080749\\_How\\_does\\_social\\_support\\_enhance\\_resilience\\_in\\_the\\_trauma-exposed\\_individual](https://www.researchgate.net/publication/283080749_How_does_social_support_enhance_resilience_in_the_trauma-exposed_individual)
48. Howard S, Hughes BM. Benefit of social support for resilience-building is contingent on social context: examining cardiovascular adaptation to recurrent stress in women. *Anxiety Research*. 2012; 25(4):411–23.

49. Mo PKH, Lau JTF, Yu X, Gu J. The role of social support on resilience, posttraumatic growth, hopelessness, and depression among children of HIV-infected parents in mainland China. *AIDS Care*. 2014; 26(12):1526–33. <https://doi.org/10.1080/09540121.2014.923810> PMID: [24922584](#)
50. Wilks SE, Croom B. Perceived stress and resilience in Alzheimer's disease caregivers: testing moderation and mediation models of social support. *Aging & Mental Health*. 2008; 12(3):357–65.
51. Weathers FW, Litz BT, Herman DS, Huska JA, Keane TM. The PTSD Checklist (PCL): Reliability, Validity, and Diagnostic Utility. Paper presented at the Annual Convention of the International Society for Traumatic Stress Studies: San Antonio, TX, 1993.
52. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA. Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy*. 1996; 34(8), 669–673. PMID: [8870294](#)
53. Wu KK, Chan SK, Yiu VF. Psychometric properties and confirmatory factor analysis of the Posttraumatic Stress Disorder Checklist (PCL) for Chinese survivors of motor vehicle accident. *Hong Kong J Psychiatry*. 2008; 18:144–151. Available from: [http://easap.asia/journal\\_file/0804\\_V18N4\\_p144.pdf](http://easap.asia/journal_file/0804_V18N4_p144.pdf)
54. Li RH, Fan M, Hu MS, Ran MS, Fang DZ. Reduced severity of posttraumatic stress disorder associated with Val allele of Val66Met polymorphism at brain-derived neurotrophic factor gene among Chinese adolescents after Wenchuan earthquake. *Psychophysiology*. 2016; 53(5):705–11. <https://doi.org/10.1111/psyp.12603> PMID: [26751724](#)
55. Broadhead WE, Gehlbach SH, Gruy FV, Kaplan BH. The Duke-UNC Functional Social Support Questionnaire: measurement of social support in family medicine patients. *Med Care*. 1988; 26:709–723. PMID: [3393031](#)
56. Wu M, Yao WQ, Wang L, Jiang FX, Song W, Zhang Q, et al. Effect of psychological capital on social support and quality of life among patients with HIV/AIDS. *Chinese Journal of Public Health*. 2015; 31(1):101–103 [Article in Chinese]
57. Herth K. Abbreviated instrument to measure hope: development and psychometric evaluation. *J Adv Nurs*. 1992; 17:1251–1259. PMID: [1430629](#)
58. Wang YH. Study on feasibility of Chinese version of Herth Hope Index for cancer patients. *Chinese Nursing Research*. 2010; 24: 20–21. [Article in Chinese]
59. Block J, Kremen. IQ and Ego—Resiliency: Conceptual and empirical connections and Separateness. *Journal of Personality and Social Psychology*. 1996; 70(2):349–361 PMID: [8636887](#)
60. Li YX, Luo PC, Tan YM. Research of rural left-behind children's mental elastic. *Journal of Henan University (Social Science)*. 2008; 48(1):13–18 [Article in Chinese]
61. Cohen J, Cohen P, West SG, Aiken LS. *Journal of the Royal Statistical Society*. 2003; 52(4): 227–229. Available from: <http://tocs.ulb.tu-darmstadt.de/218422970.pdf>
62. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods*. 2008; 40(3):879–91. PMID: [18697684](#)
63. Hao J, Wu D, Liu L, Li X, Wu H. Association between Work-Family Conflict and Depressive Symptoms among Chinese Female Nurses: The Mediating and Moderating Role of Psychological Capital. *Int J Environ Res Public Health*. 2015; 12(6):6682–99. <https://doi.org/10.3390/ijerph120606682> PMID: [26075725](#)
64. Liu L, Yang YL, Wang ZY, Wu H, Wang Y, Wang L. Prevalence and Positive Correlates of Posttraumatic Stress Disorder Symptoms among Chinese Patients with Hematological Malignancies: A Cross-Sectional Study. *PLoS One*. 2015; 10(12):e0145103 <https://doi.org/10.1371/journal.pone.0145103> PMID: [26669841](#)
65. Yang YL, Liu L, Li MY, Shi M, Wang L. Psychological Disorders and Psychosocial Resources of Patients with Newly Diagnosed Bladder and Kidney Cancer: A Cross-Sectional Study. *PLoS One*. 2016; 11(5):e0155607. <https://doi.org/10.1371/journal.pone.0155607> PMID: [27191964](#)
66. Mehnert A, Koch U. Prevalence of acute and post-traumatic stress disorder and comorbid mental disorders in breast cancer patients during primary cancer care: A prospective study. *Psycho-Oncology*, 2007; 16, 181–188. <https://doi.org/10.1002/pon.1057> PMID: [16856147](#)
67. McDonald SD, Calhoun PS. The diagnostic accuracy of the PTSD checklist: A critical review. *Clinical Psychology Review*. 2010; 30: 976–987. <https://doi.org/10.1016/j.cpr.2010.06.012> PMID: [20705376](#)
68. Terhakopian A, Sinai N, Engel CC, Schnurr PP, Hoge CW. Estimating population prevalence of post-traumatic stress disorder: an example using the PTSD checklist. *Journal of Traumatic Stress*. 2008; 21(3):290–300. <https://doi.org/10.1002/jts.20341> PMID: [18553416](#)
69. Andrykowski MA, Cordova M. Factors associated with PTSD symptoms following treatment for breast cancer: test of the Andersen model. *Journal of Traumatic Stress*. 1998; 11: 189–203. <https://doi.org/10.1023/A:1024490718043> PMID: [9565911](#)
70. Jacobsen PB, Sadler IJ, Booth-Jones M, Soety E, Weitzner MA, Fields KK. Predictors of posttraumatic stress disorder symptomatology following bone marrow transplantation for cancer. *J Consult Clin Psychol*. 2002; 70(1):235–40. PMID: [11860050](#)

71. Andrykowski MA, Cordova MJ. Factors associated with PTSD symptoms following treatment for breast cancer: test of the Andersen model. *J Trauma Stress*. 1998; 11(2):189–203. <https://doi.org/10.1023/A:1024490718043> PMID: 9565911
72. Wang Y, Bao Y, Liu L, Ramos A, Wang Y, Wang L. The mediating effect of self-efficacy in the relationship between social support and post-traumatic stress disorder symptoms among patients with central system tumors in China: a cross-sectional study. *Psychooncology*. 2015; 24(12):1701–7. <https://doi.org/10.1002/pon.3838> PMID: 25958841
73. Li M, Wang L. The Associations of Psychological Stress with Depressive and Anxiety Symptoms among Chinese Bladder and Renal Cancer Patients: The Mediating Role of Resilience. *PLoS One*. 2016; 11(4):e0154729. <https://doi.org/10.1371/journal.pone.0154729> PMID: 27128438
74. Besser A, Weinberg M, Zeigler-Hill V, Neria Y. Acute symptoms of posttraumatic stress and dissociative experiences among female israeli civilians exposed to war: the roles of intrapersonal and interpersonal sources of resilience. *J Clin Psychol*. 2014; 70(12):1227–39. <https://doi.org/10.1002/jclp.22083> PMID: 24619400
75. Maley JH, Brewster I, Mayoral I, Siruckova R, Adams S, McGraw KA, et al. Resilience in Survivors of Critical Illness in the Context of the Survivors' Experience and Recovery. *Ann Am Thorac Soc*. 2016; 13(8): 1351–60. <https://doi.org/10.1513/AnnalsATS.201511-782OC> PMID: 27159794
76. Mealer M, Schmiede SJ, Meek P. The Connor-Davidson Resilience Scale in Critical Care Nurses: A Psychometric Analysis. *J Nurs Meas*. 2016; 24(1):28–39. <https://doi.org/10.1891/1061-3749.24.1.28> PMID: 27103241
77. Wright AM, Talia YR, Aldhalimi A, Broadbridge CL, Jamil H, Lumley MA, et al. Kidnapping and Mental Health in Iraqi Refugees: The Role of Resilience. *J Immigr Minor Health*. 2016 Jan 19. [Epub ahead of print]
78. Joscelyne A, Knuckey S, Satterthwaite ML, Bryant RA, Li M, Qian M, et al. Mental Health Functioning in the Human Rights Field: Findings from an International Internet-Based Survey. *PLoS One*. 2015; 10(12):e0145188. <https://doi.org/10.1371/journal.pone.0145188> PMID: 26700305
79. Evren C, Umut G, Bozkurt M, Evren B, Agachanli R. Mediating role of childhood emotional abuse on the relationship between severity of ADHD and PTSD symptoms in a sample of male inpatients with alcohol use disorder. *Psychiatry Res*. 2016; 239:320–4. <https://doi.org/10.1016/j.psychres.2016.03.049> PMID: 27058158
80. Short NA, Ennis CR, Oglesby ME, Boffa JW, Joiner TE, Schmidt NB. The mediating role of sleep disturbances in the relationship between posttraumatic stress disorder and self-injurious behavior. *J Anxiety Disord*. 2015; 35:68–74. <https://doi.org/10.1016/j.janxdis.2015.09.004> PMID: 26398304
81. Ardino V, Milani L, Di Blasio P. PTSD and re-offending risk: the mediating role of worry and a negative perception of other people's support. *Eur J Psychotraumatol*. 2013; 4.
82. Solano JPC, Silva AG, Soares IA, Ashmawi HA, Vieira JE. Resilience and hope during advanced disease: a pilot study with metastatic colorectal cancer patients. *BMC Palliative Care*. 2016; 15(1): 70.
83. Shetty V. Resiliency, hope, and life satisfaction in midlife. *IOSR journal of humanities and social science*. 2015; 20(6):29–32. <https://doi.org/10.9790/0837-20632932>
84. Fava GA, Tomba E. Increasing psychological well-being and resilience by psychotherapeutic methods. *Journal of Personality*. 2009; 77(6), 1903–34. <https://doi.org/10.1111/j.1467-6494.2009.00604.x> PMID: 19807860
85. Ruiz-Robledillo N, Andrés-García SD, Pérez-Blasco J, González-Bono E, Moya-Albiol L. Highly resilient coping entails better perceived health, high social support and low morning cortisol levels in parents of children with autism spectrum disorder. *Research in Developmental Disabilities*. 2014; 35(3):686–95. <https://doi.org/10.1016/j.ridd.2013.12.007> PMID: 24405793
86. Snyder CR. *Handbook of hope: Theory, measures, and applications*. Academic press, 2000.