

Association of autophagy related gene polymorphisms with neutrophilic airway inflammation in adult asthma

Duy Le Pham^{1,2}, Seung-Hyun Kim², Purevsuren Losol³, Eun-Mi Yang², Yoo Seob Shin², Young-Min Ye², and Hae-Sim Park^{1,2}

Departments of ¹Biomedical Sciences and ²Allergy and Clinical Immunology, Ajou University School of Medicine, Suwon, Korea; ³Department of Molecular Biology and Genetics, Health Sciences University of Mongolia, Ulaanbaatar, Mongolia

Received: December 26, 2014

Revised: June 2, 2015

Accepted: August 17, 2015

Correspondence to
Hae-Sim Park, M.D.

Department of Allergy and Clinical Immunology, Ajou University School of Medicine, 206 World cup-ro, Yeongtong-gu, Suwon 16499, Korea
 Tel: +82-31-219-5196
 Fax: +82-31-219-5154
 E-mail: hspark@ajou.ac.kr

Background/Aims: Role of autophagy in neutrophil function and the association of autophagy and autophagy related (ATG) gene polymorphisms with asthma susceptibility were suggested. In this study, we investigated the genetic association of *ATG5* and *ATG7* polymorphisms with asthma risk, severity and neutrophilic airway inflammation.

Methods: We recruited 408 asthma patients and 201 healthy controls. Sputum neutrophil counts were determined by H&E staining. Serum interleukin 8 (IL-8) levels were measured by enzyme-linked immunosorbent assay (ELISA). Genetic polymorphisms of *ATG5* (-769T>C, -335G>A, and 8830C>T) and *ATG7* (-100A>G and 25108G>C) were genotyped. The functional activities of *ATG5* -769T>C and -335G>A variants were investigated by luciferase reporter assays.

Results: No associations of *ATG5* and *ATG7* polymorphisms with asthma susceptibility and severity were found. *ATG5* -769T>C and -335G>A were in complete linkage disequilibrium. In the asthma group, GA/AA genotypes at *ATG5* -335G>A were associated with higher neutrophil counts in sputum ($p < 0.05$); CC/TT genotype at *ATG5* 8830C>T associated with lower FEV1% predicted value ($p < 0.05$). DNA fragments containing *ATG5* -769T and -335G alleles had higher promoter activities compared to those with -769C and -335A in both human airway epithelial cells (A549, $p < 0.01$) and human mast cell (HMC-1, $p < 0.001$). GG and CC genotype at *ATG7* -100A>G and 25108G>C were significantly associated with high serum levels of IL-8 ($p < 0.05$ for both variants).

Conclusions: Genetic polymorphisms of *ATG5* and *ATG7* could contribute to neutrophilic airway inflammation in the pathogenesis of adult asthma.

Keywords: Asthma; Autophagy; Polymorphism, genetic; Neutrophils

INTRODUCTION

Asthma is a chronic airway inflammatory disease characterized by airflow obstruction and bronchial hyperresponsiveness [1,2]. Asthmatic airway inflammation has heterogeneous clinical phenotypes which can depend on

different cellular inflammatory pathogenic mechanisms [3]. Eosinophil is considered as a key effector cell in asthma [4,5]; however, abundant published data reported strong associations of airway neutrophilic inflammation with severe asthma, corticosteroid treatment responses, asthma exacerbations and asthma phenotypes/

endotypes [6]. The roles of neutrophil in asthma pathogenesis are not completely understood and the genetic predictive factor for neutrophilic inflammatory phenotypes in asthma has not been identified.

Macroautophagy (referred as autophagy in this report) is a cellular process important to cell survival, by which damaged organelles and proteins are sequestered to autophagosomes and delivered to lysosomes for degradation [7]. Among the proteins that take part in autophagy pathway, autophagy related (ATG) 5 protein in the conjugation with ATG12, and ATG7 acting as E1-like enzyme play important roles in the maturation of autophagosomes by facilitating the elongation of isolation membranes (phagophore). Recently, autophagy has been implicated in inflammation and immune responses of airway diseases such as asthma and chronic obstructive pulmonary disease [8]. Autophagy plays crucial roles in innate as well as adaptive immune responses by supporting the proliferation, activation and function of various immune cells such as macrophages, neutrophils, mast cells, T and B cells [9-11]. Regarding neutrophils, autophagy was found to regulate cell differentiation, formation of neutrophil extracellular trap (NET) and cell death [12], which could contribute to the pathogenesis of immune diseases such as asthma. In asthma patients autophagosomes were detected more frequently in epithelial cells compared to those in healthy subjects [13]. In addition, single nucleotide polymorphisms (SNPs) of ATG5 such as rs510432 and rs12212740 were found to associate with asthma susceptibility and lung function [13,14]. These findings suggest an association of autophagy and ATG genetic polymorphisms in asthma pathogenesis.

Based on the findings, we investigated the association of ATG5 and ATG7 genetic polymorphisms with asthma susceptibility, severity and clinical features with a focus on neutrophilic inflammation in the present study.

METHODS

Study subjects recruitment

We enrolled 408 asthmatic patients and 201 healthy normal controls (NCs) from Ajou University Hospital (Suwon, Korea) to a case-controlled study. Asthma was diagnosed at the first evaluation based on a history of respiratory symptoms as well as the results of airway

reversibility and hyperresponsiveness to methacholine following the Global Initiative for Asthma guideline. Severe asthma was defined as asthma that requires treatment with high dose inhaled corticosteroids plus a second controller and/or systemic corticosteroids to control it or that remains “uncontrolled” despite this therapy, follows American Thoracic Society guidelines [15]. All asthma patients were recruited when the disease was stable on their regular medications without any viral or bacterial airway infection. NCs were healthy individuals who had no history of asthma symptoms. All the subjects were provided written informed consents prior to participating in this study.

Atopy was defined as one or more positive reactions on a skin prick test with 55 common inhalant allergens (Bencard Co., Brentford, UK) with histamine and saline controls. Methacholine bronchial challenge tests were performed as previously described using doubled doses of methacholine (0.075 to 25 mg/mL) [16]. The methacholine PC₂₀ value (the concentration of methacholine needed to produce a 20% decrease in forced expiratory volume in 1 second [FEV₁]) was determined by interpolation from a dose-response curve.

Sputum induction and blood collection

Sputum induction was performed as previously described [17]. Briefly, asthma subjects were pretreated with 200 µg salbutamol through a metered dose inhaler. The subjects then were inhaled nebulized sterile 3% saline solution for 20 minutes through an ultrasonic nebulizer (Omron Co., Kyoto, Japan). Expecterated sputum was collected into a petri dish after excluding the saliva. Simultaneously with sputum collection, venous blood was collected into acid citric dextrose containing tubes (BD Falcon, Franklin Lakes, NJ, USA) for genomic DNA preparation and Vacuette tubes (Greiner Bio-One, Monroe, NC, USA) for serum collection.

Assessment of sputum neutrophil count

Each sputum sample in petri dish was weighted and transferred into a 50 mL polystyrene tube. Four times volume (v/w) of freshly prepared dithiothreitol 0.1% (DTT, Sigma, St. Louis, MO, USA) diluted in distilled water was added to each sputum tube followed by incubation at 37°C for 20 minutes with occasionally gentle vortex to dissociate the disulfide bonds of the mucus.

Table 1. The single nucleotide polymorphisms of ATG₅ and ATG₇ in this study

Gene	SNP	Chromosome	Location	MAF	HWE (p value)	
ATG ₅	-769 T>C	rs506027	6	Promoter	0.419	0.961
	-335 G>A	rs510432	6	Promoter	0.419	0.961
	8830 C>T	rs573775	6	Intron 1	0.331	0.530
ATG ₇	-100 A>G	rs2594971	3	Promoter	0.32	0.530
	25108 G>C	rs1375206	3	Intron 1	0.342	0.012

p values were obtained by chi-square tests in normal control group.

SNP, single nucleotide polymorphism; MAF, minor allele frequency; HWE, Hardy-Weinberg equilibrium.

Table 2. The amplifying and extension primers and probes for ATG₅ and ATG₇ polymorphisms genotyping

Gene	Locus (SNP ID)	Primer
ATG ₅	-769T>C (rs506027)	Forward: AGTCCAACCTCCAAGAAGATTT Reverse: GTCCCAGCTACTGGAGAGT Genotyping: TTATTAGTAAGTTTACCAATTATTA
	-335G>A (rs510432)	TaqMan assay, predesigned
	8830C>T (rs573775)	TaqMan assay, predesigned
ATG ₇	-100A>G (rs2594971)	TaqMan assay, predesigned
	25108G>C (rs1375206)	TaqMan assay, predesigned

SNP, single nucleotide polymorphism.

The reaction was stopped by added phosphate buffer saline (PBS) in a volume equal to the sputum plus DTT solution. The tubes were centrifuged at 1,500 rpm for 5 minutes. The cell pellet was resuspended in 50 mL PBS and filtered through a 40 µm nylon filter (Millipore, Bedford, MS, USA). Total cell count and cell viability were determined by staining with Trypan Blue (Sigma) and observing under an inverted microscope. To assess differential cell count, a cytospin slide of each sample was prepared, fixed with methanol and performed hematoxylin and eosin stain. All slides were overread (total 500 cells) and the samples with more than 80% squamous cells were excluded from analysis. Sputum neutrophil count was expressed as percentage of neutrophils among non-squamous cells in each sample.

SNP selection and genotyping

Genomic DNA was prepared from peripheral blood samples using Puregene Blood Core Kit (Qiagen, Germantown, MD, USA) following the manufacturer's protocol. We screened for SNPs located on promoter and 5'-untranslated region (UTR) of ATG₅ and ATG₇ from 40

healthy Korean volunteers by direct sequencing. Three SNPs of ATG₅ including -769T>C, -335G>A in promoter region and 8830C>T in 5'UTR, were selected. Two SNPs of ATG₇, including -100 A>G in promoter and 25108G>C in 5'UTR, were selected. ATG₅ and ATG₇ are located on chromosome 6 and 3, respectively. The information of ATG₅ and ATG₇ polymorphisms are shown in Table 1. Minor allele frequency of all selected SNPs was over than 5%. Target SNPs were genotypes from all genomic DNA samples of study subjects using a primer extension method and the SNaPshot ddNTP primer extension kit (Applied Biosystems, Foster City, CA, USA) with primers shown in Table 2.

Preparation of ATG₅ constructs for promoter assays

A 3201 base-pair DNA fragment that contains ATG₅ -769T>C and -335G>A was amplified using the following primers: forward primer 5'-CTCTATCGA-TAGGTACCAAGTGATGGTTAGGGTTCATGG-3' and reverse primer 5'-GATCGCAGATCTCGAGGCCCTCCGTGTTCTGCCTAA-3'. The 3' end of each primer (underlined) contains homologous sequences with ATG₅

and the 5' ends contain 15-16 bases homologous to each end of the linear pGL3-basic vector (Promega, Madison, WI, USA). Linear vector was prepared using Kpn1 and Xho1 restriction enzymes (Takara, Shuzo, Japan). The PCR amplified fragments were gel-purified using Expin Gel kit (GeneAll, Seoul, Korea) and then cloned into the linear pGL3-basic vectors using In-fusion HD cloning kit (Clontech Laboratories Inc., Mountain View, CA, USA) following the manufacturer's manual.

Cell culture

Human airway epithelial cell (A549) and human mast cell (HMC-1) were purchased from American Type Culture Collection (Manassas, VA, USA). A549 cells were cultured in RPMI-1640 and HMC-1 cells were cultured in Iscove's modified Dulbecco's medium (Gibco, Grand Island, NY, USA) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 100 U/mL penicillin G sodium and 100 µg/mL of streptomycin sulfate (Gibco). The cells were maintained at 37°C in 5% CO₂ humidified atmosphere.

Endogenous mRNA expressions of ATG5 and ATG7

Total mRNA was extracted from various cell lines using Easy-Blue reagent (Intron Biotechnology, Seongnam, Korea) according to the manufacturer's protocol. cDNA was synthesized by reverse transcription (RT)-PCR from 3 µg of total mRNA using MMLV-RTase (Promega) under optimized reaction condition (reaction buffer: 50 mM Tris-HCl pH 8.3, 17 mM KCl, 3 mM MgCl₂, 10 mM DTT, 2 mM dNTP, 0.5 U/mL RNase inhibitors [Promega] at 42°C for 1 hour 10 minutes and extended at 70°C for 10 minutes). The mRNA expression levels of ATG5 and ATG7 was evaluated by PCR using the specific primers (ATG5: forward 5'-TCTCTTGACTGCAGTACCTTTTC-3' and reverse 5'-CGTGTCTGATACTGTAAGCCTTT-3'; ATG7: forward 5'-CGACGCTCAGTAGCCTGTAG-3' and reverse 5'-TTTCTATCCCCTCCCCTGG-3'), which was normalized by beta-actin mRNA expression level using specific primers (forward 5'-TCCTTCTGCATCCTGTC-GGC-3' and reverse 5'-CAAGAGATGGCCACGGCTGC-3').

Transient transfection and dual luciferase assay

A549 cells were transfected with constructed plasmid DNA using lipofectamine (Invitrogen, Carlsbad, CA, USA) following the manufacturer's protocol. Briefly, 1 ×

10⁵ cells/well were seeded onto 12-well plate to obtain 70% to 80% confluence. Cells were transfected with 1 µg of constructed plasmid DNA, 5 ng of *Renilla* plasmid DNA, 5 µL lipofectamine and grown for 48 hours. HMC-1 cells were transfected with plasmid constructs using a microporator (MP-100, Digital Bio Technology, Seoul, Korea) and Neon transfection system (Invitrogen). Each well of 12-well plate was filled with 1 mL of appropriate medium supplemented with 10% FBS. For each well, 5 × 10⁵ cells mixed with constructed plasmid DNA (1 µg) and *Renilla* plasmid DNA (5 ng) were suspended in 10 µL of solution R. The mixture then was shocked at 1,650 V/20 mA/plus width 1 in the microporator pipette station containing 3 mL of solution E. Cells were then transferred into 12-well plate and incubated for 24 hours at 37°C in 5% CO₂ humidified atmosphere. Transfected cells were lysed with passive lysis buffer (Promega). 10 µL of each lysate was applied to a dual-luciferase reporter assay (Promega) using a Clarity Luminescence microplate reader (BioTek, Winooski, VT, USA) following the manufacturer's protocol. The pGL3-control (Promega) and promoter-less pGL3-basic vectors were used as positive and negative control, respectively. Transfection and luciferase assays were replicated in three independent experiments.

Measurement of serum interleukin 8 by ELISA

Levels of serum interleukin 8 (IL-8) in were measured by using an enzyme-linked immunosorbent assay (ELISA) kit (Endogen, Woburn, MA, USA) according to the manufacturer's protocol.

Statistical analysis

The chi-square test with one degree of freedom was used to test the frequency of each SNP for significant departure from the Hardy-Weinberg equilibrium in NC group; and *p* values of less than 0.01 were considered statistically significant [18]. The haplotype block pattern was constructed using Haploview version 4.2 (<http://haploview.software.informer.com/4.2/>). Haplotypes that had frequency of more than 5% were selected for statistical analysis.

Clinical characteristics were compared between the two study groups using Student *t* test and chi-square test. Differences in genotype and haplotype frequencies between the study groups were examined by a lo-

Table 3. Clinical characteristics of study subjects

Characteristic	Asthma (n = 408)	NC (n = 201)	p value
Age, yr	42.4 ± 13.3	28.6 ± 12.3	< 0.001 ^a
Male sex	164/399 (41.1)	91/180 (50.6)	0.018 ^b
Smoking	60/136 (44.1)	6/48 (12.5)	0.003 ^b
Atopy	215/353 (60.9)	24/104 (23.1)	< 0.001 ^b
Severe asthma	64/369 (17.3)	NA	NA
FEV ₁ % predicted	84.2 ± 21.9	NA	NA
Metacholine PC ₂₀ , mg/mL	8.1 ± 16.3	NA	NA
Sputum neutrophil count, %	53.6 ± 34.8	NA	NA
Total IgE, IU/L	402.8 ± 710.9	NA	NA

Values are presented as mean ± SD or number (%).

NC, normal control; FEV₁, forced expiratory volume in 1 second; NA, not available.

^ap values were obtained using one way analysis of variance (ANOVA).

^bp values were obtained using chi-square test.

gistic regression analysis with codominant, dominant and recessive models after accounting for age and sex as covariates. Generalized linear models were applied to compare sputum neutrophil count and serum IL-8 level according to genotypes and haplotypes distribution in the asthma group. The statistical analyses used IBM SPSS version 20 (IBM Co., Armonk, NY, USA) and p values of less than 0.05 were considered statistically significant.

RESULTS

Clinical characteristics of study subjects

Table 3 demonstrates the clinical characteristics of study subjects. Asthma patients were older than NCs (42.4 ± 13.3 vs. 28.6 ± 12.3, $p < 0.001$). Smoking and atopy rate were significantly higher in the asthma group compared to those of NC group ($p = 0.003$ and $p < 0.001$, respectively). Among asthma patients, 17.3% were classified as severe asthma. Mean sputum neutrophil count of the asthma group was 53.6% ± 34.8%.

Allele, genotype and haplotype frequency of ATG5 and ATG7 polymorphisms

The distributions of the three SNPs of ATG5 and two SNPs of ATG7 did not depart significantly from the Hardy-Weinberg equilibrium in NC group ($p > 0.01$ for all SNPs) (Table 1). All the SNPs of each ATG5 and ATG7




Figure 1. Linkage disequilibrium plot of ATG5 (A) and ATG7 (B) genetic polymorphisms using Haploview with confidence bounds color scheme. All single nucleotide polymorphisms were in strong linkage disequilibrium.

were in strong linkage disequilibrium (Fig. 1). Interestingly, the two promoter SNPs of ATG5 gene (−769T>C and −335G>A) were in a complete linkage disequilibrium in our study population; consequently, the statistically analyzed data of −335G>A were shown as the representatives for the two ATG5 promoter SNPs. Three common haplotypes (HT1 [AC], HT2 [GT], HT3 [GC]) for ATG5 −335G>A and 8830C>T; and two common haplotypes (HT1 [AG], HT2 [GC]) for the two SNPs of ATG7 were constructed using Haploview software. The haplotypes that have a frequency more than 5% were selected for analysis. The distribution of genotypes and haplotypes of ATG5 and ATG7 polymorphisms in the two study groups was not significantly different (Tables 4 and 5); and no significant differences in those distributions were

Table 4. Distribution of genotypes and haplotypes of ATG5 genetic polymorphisms into the two study groups

SNP/HT	Genotype	Asthma (n = 408)	Normal control (n = 201)	p value
-335 G>A	GG	133/404 (32.9)	66/200 (33.0)	0.238
	GA	175/404 (43.3)	100/100 (50.0)	0.275
	AA	96/404 (23.8)	34/200 (17.0)	0.386
8830 C>T	CC	177/406 (43.6)	93/201 (46.3)	0.442
	CT	182/406 (44.8)	84/201 (41.8)	0.097
	TT	47/406 (11.6)	24/201 (11.9)	0.998
HT1 [AC]	HT1/HT1	96/396 (24.2)	34/198 (17.2)	0.197
	HT1/-	172/396 (43.4)	98/198 (49.5)	0.339
	-/-	128/396 (32.3)	66/198 (33.3)	0.237
HT2 [GT]	HT2/HT2	46/396 (11.6)	24/198 (12.1)	0.337
	HT2/-	175/396 (44.2)	83/198 (41.9)	0.776
	-/-	175/396 (44.2)	91/198 (46.0)	0.108
HT3 [GC]	HT3/HT3	15/396 (3.8)	16/198 (8.1)	0.631
	HT3/-	131/396 (33.1)	67/198 (33.8)	0.918
	-/-	250/396 (63.1)	115/198 (58.1)	0.314

Values are presented as number (%). Each p value was calculated using the codominant, dominant and recessive models. Logistic regression analysis was applied to control age and sex as covariates. SNP, single nucleotide polymorphism; HT, haplotype.

Table 5. Distribution of genotypes and haplotypes of ATG7 genetic polymorphisms into the two study groups

SNP/HT	Genotype	Asthma (n = 408)	Normal control (n = 201)	p value
-100 A>G	AA	185/402 (46.0)	86/201 (42.8)	0.427
	AG	189/402 (47.0)	100/201 (49.8)	0.999
	GG	28/402 (7.0)	15/201 (7.5)	0.313
25108 G>C	GG	168/403 (41.7)	78/200 (39.0)	0.467
	GC	202/403 (50.1)	107/200 (53.5)	0.785
	CC	33/403 (8.2)	15/200 (7.5)	0.274
HT1 [AG]	HT1/HT1	166/399 (41.6)	78/200 (39.0)	0.503
	HT1/-	200/399 (50.1)	107/200 (53.5)	0.747
	-/-	33/399 (8.3)	15/200 (7.5)	0.294
HT2 [GC]	HT2/HT2	28/399 (7.0)	14/200 (7.0)	0.490
	HT2/-	188/399 (47.1)	100/200 (50.0)	0.327
	-/-	183/399 (45.9)	86/200 (43.0)	0.845

Values are presented as number (%). Each p value was calculated using the codominant, dominant and recessive models. Logistic regression analysis was applied to control age and sex as covariates. SNP, single nucleotide polymorphism; HT, haplotype.

noted according to severity of asthma (Tables 6 and 7).

Clinical characteristics according to ATG5 and ATG7 polymorphisms

Fig. 2A shows the comparison of clinical characteristics

according to the distributions of ATG5 genotypes in the asthma group. Asthmatics that carried GA/AA genotype at ATG5 -335G>A had significantly higher sputum neutrophil counts compared to those carrying GG genotypes (59.5% ± 33.5% vs. 39.8% ± 34.9%, p = 0.025). Asth-

Table 6. Distribution of genotypes and haplotypes of ATG5 genetic polymorphisms in asthma group according to asthma severity

SNP/HT	Genotype	Severe asthma (n = 64)	Non-severe asthma (n = 305)	p value
-335 G>A	GG	21/64 (32.8)	99/301 (32.9)	0.773
	GA	26/64 (40.6)	130/301 (43.2)	0.609
	AA	17/64 (26.6)	72/301 (23.9)	0.994
8830 C>T	CC	29/64 (45.3)	133/303 (43.9)	0.659
	CT	24/64 (37.5)	138/303 (45.5)	0.178
	TT	11/64 (17.2)	32/303 (10.6)	0.767
HT1 [AC]	HT1/ HT1	17/62 (27.4)	72/295 (24.4)	0.622
	HT1/-	26/62 (41.9)	127/295 (43.1)	0.800
	-/-	19/62 (30.6)	96/295 (32.5)	0.560
HT2 [GT]	HT2/ HT2	11/62 (17.7)	31/295 (10.5)	0.687
	HT2/-	22/62 (35.5)	133/295 (45.1)	0.656
	-/-	29/62 (46.8)	131/295 (44.4)	0.136
HT3 [GC]	HT3/ HT3	2/62 (3.2)	10/295 (3.4)	0.241
	HT3/-	16/62 (25.8)	104/295 (35.3)	0.163
	-/-	44/62 (71.0)	181/295 (61.4)	0.867

Values are presented as number (%). Each p value was calculated using the codominant, dominant and recessive models. Logistic regression analysis was applied to control age and sex as covariates.

SNP, single nucleotide polymorphism; HT, haplotype.

Table 7. Distribution of genotypes and haplotypes of ATG7 genetic polymorphisms in asthma group according to asthma severity

SNP/HT	Genotype	Severe asthma (n = 64)	Non-severe asthma (n = 305)	p value
-100 A>G	AA	27/64 (42.2)	138/299 (46.2)	0.397
	AG	30/64 (46.9)	141/299 (47.2)	0.175
	GG	7/64 (10.9)	20/299 (6.7)	0.740
25108 G>C	GG	23/64 (35.9)	126/300 (42.0)	0.151
	GC	32/64 (50.0)	151/300 (50.3)	0.069
	CC	9/64 (14.1)	23/300 (7.7)	0.439
HT1 [AG]	HT1/HT1	23/64 (35.9)	124/296 (41.9)	0.155
	HT1/-	32/64 (50.0)	149/296 (50.3)	0.075
	-/-	9/64 (14.1)	23/296 (7.8)	0.441
HT2 [GC]	HT2/HT2	7/64 (10.9)	20/296 (6.8)	0.418
	HT2/-	30/64 (46.9)	140/296 (47.3)	0.769
	-/-	27/64 (42.2)	136/296 (45.9)	0.182

Values are presented as number (%). Each p value was calculated using the codominant, dominant and recessive models. Logistic regression analysis was applied to control age and sex as covariates.

SNP, single nucleotide polymorphism; HT, haplotype.

atics carrying CC/CT genotype at ATG5 8830C>T had significantly lower FEV1% predicted values compared to the carriers of TT genotype (83.3% ± 22.1% vs. 92.3% ± 19.2%, p = 0.021). Additionally, the prevalence of asth-

atics with FEV1% predicted value < 80% tended to be higher in those carrying ATG5 8830 CC/CT genotype than in those carrying TT genotype (41.0% vs. 22.6%, p = 0.053). Fig. 2B shows that asthmatics carrying ATG5 HT1




Figure 2. Comparison of clinical parameters according to the genotypes of *ATG5* and *ATG7* polymorphisms in asthmatic patients. (A) Clinical parameters according to the genotypes of *ATG5* -335 G>A and 8830C>T. (B) Clinical parameters according to the HT1 [AC] and HT3 [GC] of *ATG5* polymorphisms. (C) Serum levels of interleukin (IL) 8 according to *ATG7* -100A>G and 25108G>C. A *p* value was obtained by generalized linear models adjusted for age and sex as covariates.

[AC] haplotype had a significantly higher sputum neutrophil count compared to non-carriers ($58.9\% \pm 33.6\%$ vs. $39.6\% \pm 35.5\%$, $p = 0.037$); while asthmatic carriers of HT3 [GC] haplotype had a slightly but significantly lower sputum neutrophil count compared to non-carriers ($43.5\% \pm 35.1\%$ vs. $60.1\% \pm 33.6\%$, $p = 0.049$). No associations of HT2 [GT] with clinical parameters were noted.

Regarding *ATG7* polymorphisms (Fig. 2C), serum IL-8 levels were significantly higher in asthmatic carriers of GG genotype at -100A>G as well as those carrying CC




Figure 3. Functional promoter activities of *ATG5* polymorphisms. DNA fragments containing Ht1 [CA] or Ht2 [TG] of *ATG5* -769T>C and -335G>A were cloned into pGL3-basic vectors. Plasmid constructs were transfected into A549 cells (A) and HMC-1 cells (B). Promoter activities of the DNA fragments were evaluated by dual-luciferase report assays. Transfections and luciferase assays were performed in three independent experiments (total $n = 9$). Values are presented as mean \pm SD; A *p* value was obtained by one way ANOVA.

genotype at 25108G>C compared to the carriers of AA/AG and GG/GC genotypes (137 ± 445 pg/mL vs. 24.2 ± 55.5 pg/mL, $p = 0.001$; 123 ± 418 pg/mL vs. 24.5 ± 55.9 pg/mL, $p = 0.003$; respectively).

Transcriptional activity of *ATG5* promoter polymorphisms

In order to select the target cell lines to investigate the functional promoter activity of *ATG5* promoter polymorphisms, we evaluated the endogenous expression of *ATG5* in various human cell lines including lung epithelial cell (A549), bronchial epithelial cell (Beas-2B), promyelocytic leukemia cell (HL-60), leukemic monocyte lymphoma cell (U937) and mast cell (HMC-1). A549 and HMC-1 cell line, known to play role in asthma pathogenesis, were selected based on their high expressions of *ATG5* (data not shown).

We constructed two DNA fragments that contained Ht1 [CA] with two minor alleles, and Ht2 [TG] with two major alleles of *ATG5* promoter SNPs as described in the Method section for dual luciferase assays. Plasmids with DNA fragments containing major haplotype exhibited significantly higher luciferase activity compared to those containing minor haplotype in both A549 ($p < 0.01$) (Fig. 3A) and HMC-1 cells ($p < 0.001$) (Fig. 3B).

DISCUSSION

Previous reports demonstrated the associations of *ATG5*,

but not *ATG7*, polymorphisms with asthma susceptibility and lung function [13,14]. In the present study with Korean adult asthmatics, we identified two promoter SNPs of *ATG5*, -769T>C and -335G>A, associated with sputum neutrophil count; and two SNPs of *ATG7* gene, -100A>G and 25108G>A, associated with serum level of IL-8.

In a previous genome wide association study with White/Caucasian children asthmatics, allele G of *ATG5* -335G>A (rs510432) was found to be associated with increased asthma risk. In the present study, the association of *ATG5* -335G>A with asthma susceptibility was not noted; however, GA/AA genotypes at *ATG5* -335G>A was associated with a higher sputum neutrophil count in adult asthmatic patients. Intriguingly, the two *ATG5* promoter SNPs were in a complete linkage equilibrium in our study population, which explains the same finding observed in regards to the sputum neutrophil count in association with *ATG5* -769T>C. These findings suggested a possibly functional effect of the two *ATG5* promoter SNPs on transcriptional activity.

Martin et al. [14] demonstrated that *ATG5* -335G allele increased a higher promoter activity compared to A allele; and two asthma-associated transcription factors, STAT1 and C-Fos, were assumed to bind to *ATG5* -335G>A. Consistently, we found that Ht1 [TG] of *ATG5* -769T>C and -335G>A enhanced significantly higher promoter activity compared to Ht2 [CA]. We do not have a direct evidence to explain for that the major haplotype of *ATG5* promoter polymorphisms was associated with lower sputum neutrophil counts but enhanced the higher promoter activity. However, Yousefi et al. [19] identified a calpain-dependently cleaved fragment of mouse *ATG5* protein related to spontaneous neutrophil apoptosis, which was independent to autophagy process. In addition, *ATG5* could contribute to cell death by interaction with Fas-associated protein with death domain under the stimulation of interferon gamma (INF- γ) [20]. Because the formation of autophagosome in mouse only needs a small amount of *ATG5* [21], the overexpression of *ATG5* in human, which is affected by *ATG5* promoter polymorphisms, probably increases intracellular autophagy levels and/or elicits neutrophil apoptosis mediated by the *ATG5* cleaved fragment.

Asthmatics carrying C allele of *ATG5* 8830C>T had a lower FEV1% predicted value than carriers of the TT

genotype. However, we did not observe a difference in the transcriptional activity of the two alleles of this SNP (data not shown). In the present study population, we observed that all subjects carrying *ATG5* -335A allele concomitantly carried *ATG5* 8830C allele; however the reverse linkage was not found. This linkage explained for that HT1 [AC] and HT3 [GC] of *ATG5* -335G>A and 8830C>T were associated only with sputum neutrophil count, but not FEV1% predicted value. This finding suggests that the two SNPs of *ATG5* could play role in asthma pathogenesis by different mechanisms, which needs to be revealed by further studies.

GG genotype at *ATG7* -100A>G and CC genotype at *ATG7* 25108G>C was also found to be associated with high serum levels of IL-8, a major chemoattractant and stimulator of neutrophils. However, these two variants did not exhibit different transcriptional activities in our transfection and luciferase assay system (data not shown).

The fact that the two promoter SNPs of *ATG5* and the two SNPs of *ATG7* were all associated with neutrophilic inflammatory markers suggested an interaction between the two genes. However, we did not find the associations of *ATG5* and *ATG7* polymorphisms in gene-gene interactions with asthma susceptibility as well as asthma clinical parameters in the present study (data not shown). This finding suggests independent roles of the two genes in neutrophilic inflammation of asthma.

Autophagy machinery was identified in murine as well as human neutrophils [22,23]. Reactive oxygen species (ROS) possibly trigger and regulate autophagy in neutrophils and other cell types [23,24]. Inflammatory cells including neutrophils that are recruited into the asthmatic airways produced greater ROS levels compared to those in normal subjects [25]. Concomitantly, greater number of double membrane autophagosomes were depicted in airway epithelial cells and fibroblast from asthmatics compared to those from healthy subjects [13]. These findings suggested a high level of autophagy occurring in cells that are involved in asthma pathogenesis, including neutrophils.

It is not fully understood if autophagy is a harmful or protective mechanism in asthma. However, autophagy has been known to protect the cells from damage or even cell death, as well as be essential for the survival, differentiation and function of various kinds of cell in-

involved in inflammatory and immune responses [10,26-28]. Consequently, autophagy is able to augment the effects of immune cells to prolong airway inflammation that increase asthma severity. In addition, respiratory viral infection, which could lead to asthma exacerbation, also could activate ATG genes and autophagy [29]. Concurrently, Remijnsen et al. [12] demonstrated that autophagy is crucial for the formation of NET which was identified in airways of allergic asthma patients and could be triggered by IL-8 [30,31]. NET is a structure composed by DNA net with histones and coated by several proteins such as neutrophil elastase and myeloperoxidase released from activated neutrophils. The role of NET in asthma pathogenesis is under investigations; however, neutrophil enzymes that coat the NET may damage the airway epithelia [32]. IL-8 levels, sputum neutrophil counts and NETs are all increased in neutrophilic asthmatics, and likely contribute to asthma severity [32-35]. Additionally, both neutrophils and IL-8 are features of difficult-to-treat asthma [36]. These findings suggest a potent role of autophagy in neutrophil activation which may enhance the severity of asthma.

We had a limitation in investigating the gene-environment interaction in the present study, which could be strictly evaluated by a further prospective study. In addition, larger cohorts and further functional studies are needed to reveal the role of autophagy and ATG genes in the pathogenesis of asthma.

We provide evidence showing the association of ATG5 and ATG7 gene polymorphisms with several markers of neutrophilic asthma phenotype, and the different functional effects of ATG5 promoter SNPs on transcriptional activities. The findings suggest a role of ATG genetic polymorphisms and autophagy machinery in neutrophil survival and function, which may contribute to the neutrophilic airway inflammation in asthma.

KEY MESSAGE

1. ATG5 and ATG7 genetic polymorphisms were associated with several makers for neutrophilic asthma phenotype such as sputum neutrophil counts and serum interleukin 8 levels.
2. ATG gene polymorphisms and autophagy machinery could contribute to neutrophilic airway inflammation in asthma pathogenesis.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

This study was supported by a grant of the Korean Health 21 R&D Project, Ministry of Health and Welfare, Republic of Korea (HI14C2628).

REFERENCES

1. Lotvall J, Akdis CA, Bacharier LB, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011;127:355-360.
2. Holgate ST. Mechanisms of asthma and implications for its prevention and treatment: a personal journey. *Allergy Asthma Immunol Res* 2013;5:343-347.
3. Fahy JV. Eosinophilic and neutrophilic inflammation in asthma: insights from clinical studies. *Proc Am Thorac Soc* 2009;6:256-259.
4. Jo EJ, Kim MY, Lee SE, et al. Eosinophilic airway inflammation and airway hyperresponsiveness according to aeroallergen sensitization pattern in patients with lower airway symptoms. *Allergy Asthma Immunol Res* 2014;6:39-46.
5. Schleich FN, Manise M, Sele J, Henket M, Seidel L, Louis R. Distribution of sputum cellular phenotype in a large asthma cohort: predicting factors for eosinophilic vs neutrophilic inflammation. *BMC Pulm Med* 2013;13:11.
6. Douwes J, Gibson P, Pekkanen J, Pearce N. Non-eosinophilic asthma: importance and possible mechanisms. *Thorax* 2002;57:643-648.
7. Deretic V, Saitoh T, Akira S. Autophagy in infection, inflammation and immunity. *Nat Rev Immunol* 2013;13:722-737.
8. Ryter SW, Choi AM. Autophagy in the lung. *Proc Am Thorac Soc* 2010;7:13-21.
9. Bhattacharya A, Eissa NT. Autophagy and autoimmunity crosstalks. *Front Immunol* 2013;4:88.
10. Mihalache CC, Simon HU. Autophagy regulation in macrophages and neutrophils. *Exp Cell Res* 2012;318:1187-1192.
11. Ushio H, Ueno T, Kojima Y, et al. Crucial role for autophagy in degranulation of mast cells. *J Allergy Clin Immunol*

- nol 2011;127:1267-1276.e6.
12. Remijsen Q, Vanden Berghe T, Wirawan E, et al. Neutrophil extracellular trap cell death requires both autophagy and superoxide generation. *Cell Res* 2011;21:290-304.
 13. Poon AH, Chouiali F, Tse SM, et al. Genetic and histologic evidence for autophagy in asthma pathogenesis. *J Allergy Clin Immunol* 2012;129:569-571.
 14. Martin LJ, Gupta J, Jyothula SS, et al. Functional variant in the autophagy-related 5 gene promotor is associated with childhood asthma. *PLoS One* 2012;7:e33454.
 15. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014;43:343-373.
 16. Park HS, Kim HY, Nahm DH, Son JW, Kim YY. Specific IgG, but not specific IgE, antibodies to toluene diisocyanate-human serum albumin conjugate are associated with toluene diisocyanate bronchoprovocation test results. *J Allergy Clin Immunol* 1999;104(4 Pt 1):847-851.
 17. Nahm DH, Park HS. Analysis of induced sputum for studying allergen-specific IgE antibodies in airway secretion from asthmatic patients. *Clin Exp Allergy* 1998;28:686-693.
 18. Lunetta KL. Genetic association studies. *Circulation* 2008;118:96-101.
 19. Yousefi S, Perozzo R, Schmid I, et al. Calpain-mediated cleavage of Atg5 switches autophagy to apoptosis. *Nat Cell Biol* 2006;8:1124-1132.
 20. Pyo JO, Jang MH, Kwon YK, et al. Essential roles of Atg5 and FADD in autophagic cell death: dissection of autophagic cell death into vacuole formation and cell death. *J Biol Chem* 2005;280:20722-20729.
 21. Hosokawa N, Hara Y, Mizushima N. Generation of cell lines with tetracycline-regulated autophagy and a role for autophagy in controlling cell size. *FEBS Lett* 2006;580:2623-2629.
 22. Huang J, Canadien V, Lam GY, et al. Activation of antibacterial autophagy by NADPH oxidases. *Proc Natl Acad Sci U S A* 2009;106:6226-6231.
 23. Mitroulis I, Kourtzelis I, Kambas K, et al. Regulation of the autophagic machinery in human neutrophils. *Eur J Immunol* 2010;40:1461-1472.
 24. Scherz-Shouval R, Elazar Z. Regulation of autophagy by ROS: physiology and pathology. *Trends Biochem Sci* 2011;36:30-38.
 25. Dworski R. Oxidant stress in asthma. *Thorax* 2000;55 Suppl 2:S51-S53.
 26. He MX, McLeod IX, Jia W, He YW. Macroautophagy in T lymphocyte development and function. *Front Immunol* 2012;3:22.
 27. Pua HH, Dzhagalov I, Chuck M, Mizushima N, He YW. A critical role for the autophagy gene Atg5 in T cell survival and proliferation. *J Exp Med* 2007;204:25-31.
 28. Conway KL, Kuballa P, Khor B, et al. ATG5 regulates plasma cell differentiation. *Autophagy* 2013;9:528-537.
 29. Jyothula SS, Eissa NT. Autophagy and role in asthma. *Curr Opin Pulm Med* 2013;19:30-35.
 30. Dworski R, Simon HU, Hoskins A, Yousefi S. Eosinophil and neutrophil extracellular DNA traps in human allergic asthmatic airways. *J Allergy Clin Immunol* 2011;127:1260-1266.
 31. Brinkmann V, Reichard U, Goosmann C, et al. Neutrophil extracellular traps kill bacteria. *Science* 2004;303:1532-1535.
 32. Cheng OZ, Palaniyar N. NET balancing: a problem in inflammatory lung diseases. *Front Immunol* 2013;4:1.
 33. Jatakanon A, Uasuf C, Maziak W, Lim S, Chung KF, Barnes PJ. Neutrophilic inflammation in severe persistent asthma. *Am J Respir Crit Care Med* 1999;160(5 Pt 1):1532-1539.
 34. Kikuchi S, Nagata M, Kikuchi I, Hagiwara K, Kanazawa M. Association between neutrophilic and eosinophilic inflammation in patients with severe persistent asthma. *Int Arch Allergy Immunol* 2005;137 Suppl 1:7-11.
 35. Silvestri M, Bontempelli M, Giacomelli M, et al. High serum levels of tumour necrosis factor-alpha and interleukin-8 in severe asthma: markers of systemic inflammation? *Clin Exp Allergy* 2006;36:1373-1381.
 36. Chun YH, Park JY, Lee H, et al. Rhinovirus-infected epithelial cells produce more IL-8 and RANTES compared with other respiratory viruses. *Allergy Asthma Immunol Res* 2013;5:216-223.