

- logica. 2004;89:348-56.
2. De Franceschi L, Cappellini MD, Olivieri O. Thrombosis and sickle cell disease. *Semin Thromb Hemost.* 2011; 37:226-36.
  3. Uwaezuoke SN, Ayuk AC, Ndu IK, et al. Vaso-occlusive crisis in sickle cell disease: Current paradigm on pain management. *J Pain Res.* 2018;11:3141-50.
  4. Rees DC, Olujohungbe AD, Parker NE, et al. Guidelines for the Management of the Acute Painful Crisis in Sickle Cell Disease. *Br J Haematol.* 2003;120:744-52.
  5. Okpala I, Tawil A. Management of pain in sickle-cell disease. *J R Soc Med.* 2002;95:456-58.
  6. Stinson J, Naser B. Pain management in children with sickle cell disease. *Pediatr Drugs.* 2003;5:229-41.
  7. National Institute of Health and Care Excellence (NICE). Guidance – sickle cell acute painful episode. Accessed February 28, 2020. Available from: <http://www.nice.org.uk/>
  8. Gopinath R. Suggesting the value of diluted Diclofenac infused over a longer period of time as a method for reducing venous sequelae of IV diclofenac. *Br J Anaesth.* 1991;67:803.
  9. Strumer T, Erb A, Keller F, Gunther KP, Brenner H. Determinants of impaired renal functions with non-steroidal anti-inflammatory drugs. *Am J Med.* 2001;111:521-7.
  10. Shastri N. Intravenous acetaminophen use in pediatrics. *Pediatr Emerg Care.* 2015;31:444-50.
  11. Dampier CD, Setty BN, Logan J, Ioli JG, Dean R. Intravenous morphine pharmacokinetics in pediatric patients with sickle cell disease. *J Pediatr.* 1995;126:461-7.
  12. Jahr JS, Lee VK. IV Acetaminophen. *Anaesthesiol Clin.* 2010; 28:619-45.
  13. Jacob E, Miaskowski C, Savedra M, et al. Management of vaso-occlusive pain in children with sickle cell disease. *J Pediatr Hematol Oncol.* 2003;25:307-11.
  14. Baichoo P, Asuncion A, El-Chaar G. Intravenous acetaminophen for the management of pain during vaso-occlusive crises in pediatric patients. *PT.* 2019;44:5-8.
  15. Ceelie I, de Wildt SN, van Dijk M, et al. Effect of intravenous paracetamol on postoperative morphine requirements in neonates and infants undergoing major noncardiac surgery: A randomized controlled trial. *JAMA.* 2013;309:149-54.
  16. Lee CKK, Engorn B, Drug dosages. *In: Engorn B, Flerlage J, editor. The Harriet Lane Hand Book: A manual for pediatric house officers, 20<sup>th</sup>ed. Elsevier; 2015.p.667-8.*
  17. Standing JF, Savage I, Pritchard D, Waddington M. Diclofenac for Acute pain in children. *Cochrane Database Syst Rev.* 2009;4:CD005538.
  18. Merkel SI, Voepel-Lewis T, Shayevitz JR, Malviya S. The FLACC: A behavioral scale for scoring postoperative pain in young children. *Pediatr Nurs.* 1997;23:293-7.
  19. Wong DL, Baker CM. Pain in children: comparison of assessment scales. *Pediatr Nurs.* 1988;14:9-17.
  20. Shields BJ, Cohen DM, Harbeck-Weber C, et al. Pediatric pain measurement using a visual analogue scale: A comparison of two teaching methods. *Clin Pediatr (Phila).* 2003;42:227-34.
  21. Cooper TE, Hambleton IR, Ballas SK, Wiffen PJ. Pharmacological interventions for painful sickle cell vaso-occlusive crises in adults. *Cochrane Database. Syst Rev.* 2016, CD012187.
  22. Varrassi G, Marinangeli F, Agro FA, et al. A double-blinded evaluation of propacetamol versus ketorolac in combination with patient-controlled analgesia morphine: Analgesic efficacy and tolerability after gynecologic surgery. *Anesth Analg.* 1999;88:611-16.
  23. Paul D, Mathews J, Kulkarni SN. Comparative evaluation of efficacy of intravenous paracetamol and intravenous diclofenac as post-operative analgesia in laparoscopic cholecystectomy. *International J Biomedical Res.* 2015;6:482-87.
  24. Shaffer EE, Pham A, Woldman RL, et al. Estimating the effect of intravenous acetaminophen for postoperative pain management on length of stay and inpatient hospital costs. *Adv Ther.* 2016;33:2211-28.
  25. Sinatra RS, Jahr JS, Reynolds LW, et al. Efficacy and safety of single and repeated administration of 1 gram intravenous acetaminophen injection (paracetamol) for pain management after major orthopedic surgery. *Anesthesiology.* 2005; 102:822-31.
  26. Tariq Haider S, Anwar A, Sultan T. Effectiveness and safety of Intravenous water based diclofenac sodium for post-operative pain management. *Anaesthesia, Pain and Intensive Care.* 2000;4:27-30.

### ERRATUM

Please note the following correction in the recommendation titled “Association of Child Neurology (AOCN) – Indian Epilepsy Society (IES) consensus guidelines for the diagnosis and management of West syndrome” published in *Indian Pediatr.* 2020;58:54-66.

The first sentence of last paragraph, column 1, page 61 should read: “In the 2004 United Kingdom Infantile Spasms study, spasm freedom was achieved in 70% of children taking high dose oral prednisolone (40-60 mg/day) and 76% of children taking ACTH (40 IU/alternate day) [21]” instead of “In the 2004 United Kingdom Infantile Spasms study, spasm freedom was achieved in 70% of children taking high dose oral prednisolone (40-60 mg/kg/day) and 76% of children taking ACTH (40 IU/alternate day) [21].”

Appropriate corrections have already been done in the web version at <https://www.indianpediatrics.net/jan2021/jan-54-66.htm> on January 9, 2021.