



Association of Exposure to Ambient Air Pollution With Thyroid Function During Pregnancy

Akhgar Ghassabian, MD; Livia Pierotti, MSc; Mikel Basterrechea, PhD; Leda Chatzi, MD; Marisa Estarlich, PhD; Ana Fernández-Somoano, PhD; Abby F. Fleisch, MD; Diane R. Gold, MD; Jordi Julvez, PhD; Polyxeni Karakosta, MD; Aitana Lertxundi, PhD; Maria-Jose Lopez-Espinosa, PhD; Tessa A. Mulder, BSc(Med); Tim I. M. Korevaar, MD; Emily Oken, MD; Robin P. Peeters, MD; Sheryl Rifas-Shiman, MPH; Euripides Stephanou, PhD; Adonina Tardón, MD; Henning Tiemeier, MD; Martine Vrijheid, PhD; Tanja G. M. Vrijkotte, PhD; Jordi Sunyer, MD; Mònica Guxens, MD

Abstract

IMPORTANCE Air pollutants interact with estrogen nuclear receptors, but their effect on thyroid signaling is less clear. Thyroid function is of particular importance for pregnant women because of the thyroid's role in fetal brain development.

OBJECTIVE To determine the short-term association of exposure to air pollution in the first trimester with thyroid function throughout pregnancy.

DESIGN, SETTING, AND PARTICIPANTS In this cohort study, 9931 pregnant women from 4 European cohorts (the Amsterdam Born Children and Their Development Study, the Generation R Study, Infancia y Medio Ambiente, and Rhea) and 1 US cohort (Project Viva) with data on air pollution exposure and thyroid function during pregnancy were included. The recruitment period for the Amsterdam Born Children and Their Development Study was January 2003 to March 2004; for Generation R, April 2002 to January 2006; for Infancia y Medio Ambiente, November 2003 to January 2008; for Rhea, February 2007 to February 2008; and for Project Viva, April 1999 to November 2002. Statistical analyses were conducted from January 2018 to April 2019.

MAIN OUTCOMES AND MEASURES Residential air pollution concentrations (ie, nitrogen oxide and particulate matter [PM]) during the first trimester of pregnancy were estimated using land-use regression and satellite-derived aerosol optical depth models. Free thyroxine, thyrotropin, and thyroid peroxidase antibody levels were measured across gestation. Hypothyroxinemia was defined as free thyroxine below the fifth percentile of the cohort distribution with normal thyrotropin levels, following the American Thyroid Association guidelines.

RESULTS Among 9931 participants, the mean (SD) age was 31.2 (4.8) years, 4853 (48.9%) had more than secondary educational levels, 5616 (56.6%) were nulliparous, 404 (4.2%) had hypothyroxinemia, and 506 (6.7%) tested positive for thyroid peroxidase antibodies. Concentrations of nitrogen dioxide and PM with an aerodynamic diameter of 2.5 μm or less ($\text{PM}_{2.5}$) were lower and had less variation in women in the US cohort than those in European cohorts. No associations of nitrogen dioxide with thyroid function were found. Higher exposures to $\text{PM}_{2.5}$ were associated with higher odds of hypothyroxinemia in pregnant women (odds ratio per 5- $\mu\text{g}/\text{m}^3$ change, 1.21; 95% CI, 1.00-1.47). Although exposure to PM with an aerodynamic diameter of 10 μm or less was not significantly associated with hypothyroxinemia, the coefficient was similar to that for the association of $\text{PM}_{2.5}$ with hypothyroxinemia (odds ratio per 10- $\mu\text{g}/\text{m}^3$ change, 1.18; 95% CI, 0.93-1.48). Absorbances of $\text{PM}_{2.5}$ and PM with aerodynamic diameter from 2.5 to 10 μm and were not associated with hypothyroxinemia. There was substantial heterogeneity among cohorts with respect to thyroid

(continued)

Key Points

Question Is exposure to ambient air pollution in the first trimester associated with thyroid function throughout pregnancy?

Findings Among 9931 pregnant women in 4 European cohorts and 1 US cohort, an increase of 5 $\mu\text{g}/\text{m}^3$ in exposure to particulate matter with an aerodynamic diameter of 2.5 μm or less was associated with 20% higher odds of hypothyroxinemia.

Meaning The findings of this study raise the possibility that exposure to particulate matter might disrupt thyroid function in pregnant women.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Abstract (continued)

peroxidase antibodies (*P* for heterogeneity, $<.001$), showing associations of nitrogen oxide and PM with thyroid autoimmunity only in the women in the Generation R Study.

CONCLUSIONS AND RELEVANCE The findings of this study suggest that first-trimester exposures to $PM_{2.5}$ were associated with mild thyroid dysfunction throughout pregnancy. The association of $PM_{2.5}$ exposure with thyroid function during pregnancy is of global health importance because air pollution exposure is widespread and hypothyroxinemia may adversely influence the brain development of offspring.

JAMA Network Open. 2019;2(10):e1912902. doi:10.1001/jamanetworkopen.2019.12902

Introduction

Exposure to ambient air pollution is a leading contributor to the burden of disease globally. Exposure to air pollutants, such as nitrogen dioxide (NO_2), nitrogen oxides (NO_x), particulate matter (PM), and polycyclic aromatic hydrocarbons (PAHs), during pregnancy is associated with brain structural alterations, impaired executive function, learning disabilities, and behavioral problems in offspring.¹⁻⁴ Oxidative stress, neuroinflammation, and the disruption of the hypothalamus-pituitary-adrenal axis are some underlying factors.^{5,6} Endocrine disruption might be among other mechanisms; for example, PAHs induce activation of the estrogen receptor gene, and PAHs and PM can interfere with nuclear receptors, such as estrogen receptor signaling.^{7,8} Experimental and epidemiological studies have also shown associations between PAH exposure and thyroid function.^{9,10} Nonetheless, evidence on whether exposure to air pollution might also disrupt thyroid signaling and thyroid function is limited.

Thyroid function is of particular importance for pregnant women because of the critical role of thyroid hormones in fetal brain development. Because the fetal thyroid gland achieves its full function only from midgestation on, undetected or inadequately treated thyroid insufficiency in pregnant women adversely influences growth and development of offspring, even in the absence of neonatal hypothyroidism.¹¹ Recent evidence suggests that mild thyroid insufficiency, ie, hypothyroxinemia, defined as low free thyroxine (T_4) levels with normal thyrotropin (TSH) concentrations, during pregnancy may also contribute to impaired cognition and neurodevelopmental disorders in offspring.¹²⁻¹⁴ While inadequate iodine intake is a common cause of thyroid insufficiency worldwide, exposure to environmental contaminants is increasingly considered important.¹⁵ Some studies have examined the association of exposure to ambient and traffic-related air pollution with thyroid function in pregnant women. A 2017 study¹⁶ of 499 mother-child pairs in Belgium examined the association of third-trimester exposure to PM with an aerodynamic diameter of $2.5\ \mu m$ or less ($PM_{2.5}$) with maternal and fetal thyroid function. This study found inverse associations of $PM_{2.5}$ exposure with levels of free T_4 , free triiodothyronine, and TSH in fetal cord blood and maternal free T_4 levels.¹⁶ In 8077 pregnant women in Shanghai,¹⁷ exposure to higher concentrations of $PM_{2.5}$ but not NO_2 in the first and second trimesters was associated with hypothyroxinemia in midgestation. A study in California with 2050 mother-child pairs¹⁸ showed that newborns who were prenatally exposed to $PM_{2.5}$ and PM with an aerodynamic diameter of $10\ \mu m$ or less (PM_{10}) had higher concentrations of total T_4 levels in heel-stick blood spots.

A potential association of air pollution exposure with thyroid function during gestation might further clarify underlying mechanisms of hypothyroxinemia in pregnant women. Accordingly, using 5 birth cohorts (4 in Europe and 1 in the United States), we examined whether measures of air pollutants, ie, NO_2 , NO_x , and PM, averaged in the first trimester were associated with thyroid function and thyroid autoimmunity throughout the pregnancy. We primarily investigated the short-term associations of air pollutants with mild thyroid hormone insufficiency, ie, hypothyroxinemia and high TSH, because of emerging evidence suggesting their implications for fetal development.^{12-14,19}

We also examined thyroid peroxidase antibody (TPOAb) positivity on the basis of evidence suggesting the association of air pollution exposure with inflammation²⁰ and our earlier findings on the association of TPOAb positivity in pregnancy with maternal and offspring health outcomes.²¹⁻²³ We specifically focused on air pollution exposure during the first trimester to ensure that air pollution exposure preceded thyroid function measurement.

Methods

Participants

This analysis used data from 5 birth cohorts with prenatal recruitment, including the Amsterdam Born Children and Their Development Study (ABCD; Amsterdam, the Netherlands; n = 3867),²⁴ the Generation R Study (Rotterdam, the Netherlands; n = 2605),²⁵ Infancia y Medio Ambiente (INMA; including the regions of Sabadell, Gipuzkoa, Valencia, and Asturias, Spain; n = 2239),²⁶ Rhea (island of Crete, Greece; n = 483),²⁷ and Project Viva (eastern Massachusetts, United States; n = 737),²⁸ yielding a total sample of 9931. Recruitment periods for the cohorts were as follows: ABCD, January 2003 to March 2004; Generation R, April 2002 to January 2006; INMA, November 2003 to January 2008; Rhea, February 2007 to February 2008; and Project Viva, April 1999 to November 2002.

Iodine status of participants varied among cohorts. Median urinary iodine concentrations of pregnant women in Generation R and Rhea were optimal according to World Health Organization references.^{29,30} In contrast, mild to moderate iodine insufficiency was observed in INMA (except for women in the Gipuzkoa region).³¹ Urinary iodine concentrations were not measured in ABCD or Project Viva. However, reports from pregnant women in Massachusetts showed optimal iodine status,³² and iodine status of participants in Amsterdam and Rotterdam are expected to be comparable. There was no pattern suggesting associations of the iodine status of cohorts with air pollution exposure.

In each cohort, we included pregnant women who had data on first-trimester air pollution exposure and gestational thyroid function. For each woman, thyroid assessments were performed once throughout pregnancy, with most measures collected in the first half of pregnancy (ie, gestational age, <17 weeks). We excluded women with twin pregnancies, women with a history of thyroid disease, and women who reported taking medication affecting thyroid function. There were no data available on medication use in INMA. We did not exclude women who tested positive for TPOAb because conditioning on autoimmune processes, an intermediate factor, might introduce bias in the analysis. This study was granted an exception from ethics review and informed consent because we used deidentified data in each cohort. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

Measurements

In ABCD, Generation R, INMA, and Rhea, 3 two-week air pollution monitoring campaigns were performed within 1 year (January 2009 to April 2011). Air pollution concentrations (ie, NO₂; NO_x; PM_{2.5} absorbance, determined as the reflectance of PM_{2.5} filters; PM_{2.5}; PM₁₀; and PM with aerodynamic diameters between 2.5 and 10 μm [PM_{2.5-10}]) at the participants' home addresses were estimated on a daily basis for the whole period of pregnancy using land-use regression following a standardized procedure.^{33,34} Among the INMA regions, data on PM were only available in Sabadell. In Rhea, only PM₁₀, PM_{2.5}, and PM_{2.5-10} concentrations were estimated. Land-use regression models were developed for each pollutant metric using all measurement sites. We used a back-extrapolation procedure to estimate exposure concentrations for each woman, averaged across the first trimester at the participant's home address using daily concentrations from routine background monitoring network sites. Land-use regression models explained a large fraction of the spatial variance in measured annual average air pollutant concentrations.^{33,34} In Project Viva, validated prediction models were used to obtain spatially and temporally resolved estimates of daily PM_{2.5} exposure at each participant's residential address, following a method described elsewhere.³⁵ Briefly,

this method combined the satellite aerosol optical depth data at a 10 km × 10 km spatial grid with the spatiotemporal land-use regression models based on monitored ground PM_{2.5} measurements. Satellite remote sensing provides an important tool for monitoring aerosols when surface monitors are not available. For measurement of NO₂ in Project Viva, we calculated hourly ambient concentrations of NO₂ by averaging data from the Massachusetts Department of Environmental Protection's greater Boston monitoring sites,^{36,37} then calculated daily and first-trimester NO₂ exposure. In ABCD, Generation R, INMA, and Project Viva, if more than 1 address was collected during the first trimester, we calculated the weighted average concentrations of all addresses according to the time spent at each address.

Serum concentrations of free T₄ and TSH were measured at the median gestational age of 13 weeks in ABCD (range, 5-37 weeks), Generation R (range, 6-18 weeks), INMA (range, 7-33 weeks) and Rhea (range, 6-27 weeks). In Project Viva, TSH, total T₄, and triiodothyronine resin uptake were measured at the median gestational age of 10 weeks (range, 6-21 weeks), allowing for the calculation of the free T₄ index, an estimate of circulating free T₄ levels from total T₄ × triiodothyronine uptake. Cohorts used different assays for measurements of thyroid hormones and TPOAb (eTable 1 in the Supplement).

To define *hypothyroxinemia*, we followed the recommendations of the American Thyroid Association guideline, which describes hypothyroxinemia as free T₄ concentrations in the lower 2.5th to 5th percentile of the population, despite a normal TSH level.³⁸ Following this guideline, we calculated population-specific cutoffs in individuals without a history of thyroid disease or thyroid medication use and those who tested negative for TPOAb. In INMA, TPOAb was not measured, and thus, no exclusions were made based on TPOAb results. We defined *high TSH* as concentrations higher than the 95th percentile of the cohort. Cutoffs were population-specific, and corresponding cutoffs in each cohort appear in eTable 2 in the Supplement. We also tested alternative cutoffs for free T₄ as well as the cutoff of 0.03 to 2.5 mIU/L for a normal TSH level to examine whether any observed associations were independent of cutoff choice. We used the laboratory recommended cutoffs for TPOAb positivity (eTable 1 in the Supplement). Details on measurements of covariates appear in the eMethods in the Supplement.

Statistical Analysis

We used Spearman correlations to examine the correlations between concentrations of air pollutants. We performed logistic regression models to assess the short-term association of air pollutants (averaged in the first trimester) with hypothyroxinemia, high TSH, and TPOAb positivity in pregnant women throughout pregnancy. First, we explored any indication of nonlinearity in the associations of air pollutants with thyroid parameters using generalized additive models. Results confirmed the linearity of associations and indicated no threshold effect. Next, associations of air pollutants with hypothyroxinemia, high TSH, and TPOAb positivity were examined in each cohort. Cohort-specific effect estimates from regression models were then combined using a random-effects meta-analysis after exploring the heterogeneity in the estimates among cohorts. We assessed heterogeneity in the estimates using the *Q* test and the *I*² statistic. Similar to previous studies, we reported the odds ratios (ORs) of hypothyroxinemia, high TSH, and TPOAb positivity per 10-μg/m³ change in NO₂ and PM₁₀ levels, 20-μg/m³ change for NO_x levels, 5-μg/m³ change for PM_{2.5} and PM_{2.5-10} levels, and 10⁻⁵ × m⁻¹ change for PM_{2.5} absorbance based on the distribution of pollutants.¹⁷

Selection of confounders was a priori and based on the direct acyclic graph of the study question and factors associated with air pollution exposure and thyroid function.³⁹⁻⁴¹ We did not adjust for season because, to our knowledge, no evidence exists on the association of season of pregnancy with free T₄ or TSH levels (despite associations with air pollution).^{42,43} Models included information on maternal age at enrollment; education; country of birth; smoking and alcohol intake during pregnancy; parity; prepregnancy body mass index, calculated as weight in kilograms divided by height in meters squared; gestational age at thyroid measurement; socioeconomic status, defined

using information on household income or occupation obtained from self-administered questionnaires; and marital status. We adjusted the analyses in INMA for 4 regions.

Pregnant women who were included in the analysis were different from those excluded because of missing data on exposure and outcome (eTable 3 in the [Supplement](#)). To address the selective nonresponse arising from these differences, we used inverse probability weighting. Briefly, we used the available information for eligible women (eTable 4 in the [Supplement](#)) to estimate the probability of participation in the study and used the inverse of those probabilities as weights in the analyses so that the results would be representative of the full cohort.

Among participants with exposure and outcome data, information on covariates was missing in less than 10% of participants, except for smoking during pregnancy (266 of 2605 [10.2%] in Generation R), any alcohol intake during pregnancy (76 of 483 [15.7%] in Rhea), socioeconomic status (626 of 3867 [16.2%] in ABCD, 651 of 2605 [25.0%] in Generation R, and 146 of 483 [30.2%] in Rhea), and prepregnancy body mass index (458 of 2605 [17.6%] in Generation R). We addressed missing data in covariates by imputing data using the Stata `ice` command for chained equations imputation. We created 25 data sets with complete observations, in which analyses were performed using standard combination rules for multiple imputations (eTable 5 in the [Supplement](#)).

In a sensitivity analysis, we ran the meta-analysis for hypothyroxinemia excluding Project Viva because a measure of free T_4 index instead of free T_4 level was included in that cohort. Additionally, in Project Viva, NO_2 was measured using central monitors; therefore, we also reran the NO_2 meta-analysis excluding Project Viva.

All analyses were performed in Stata statistical software version 14.0 (StataCorp) between January 2018 and April 2019. All statistical tests were 2-sided with a significance threshold of $P < .05$.

Results

Table 1 summarizes participant characteristics. Of 9931 women in the analysis, the mean (SD) age of participants at enrollment was 31.2 (4.8) years, 4853 (48.9%) had more than a secondary education, 5616 (56.6%) were nulliparous, and 584 (5.8%) were single. Overall, 7568 (76.2%) did not smoke during pregnancy, and 2911 (29.3%) reported alcohol intake in pregnancy. A smaller proportion of participants in Project Viva had a low socioeconomic status compared with women in other cohorts (10 [1.4%] in Project Viva, 464 [12.0%] in ABCD, 102 [3.9%] in Generation R, 1189 [53.1%] in INMA, and 40 [8.3%] in Rhea). Women in INMA and Rhea had lower educational levels compared with women in other cohorts (no more than elementary education: 544 [24.4%] in INMA, 97 [20.1%] in Rhea, 683 [17.8%] in ABCD, 428 [17.2%] in Generation R, and 9 [1.2%] in Project Viva). In Project Viva, 103 women (13.9%) tested positive for TPOAb, which was modestly higher than in ABCD (213 [5.5%]), Generation R (145 [6.0%]), and Rhea (45 [9.3%]). Median (interquartile range) concentrations of $PM_{2.5}$ were between 11.5 (10.7-12.3) $\mu g/m^3$ in Project Viva and 20.6 (19.7-24.2) $\mu g/m^3$ in ABCD (**Figure 1**; eTable 6, eFigure 1, and eFigure 2 in the [Supplement](#)). The median (interquartile range) concentration of NO_2 was 21.6 (20.7-23.3) $\mu g/m^3$ in Project Viva and 41.6 (34.7-49.4) $\mu g/m^3$ in ABCD. Concentrations of NO_2 and $PM_{2.5}$ were lower and had less variation in Project Viva compared with the 4 European cohorts. The correlations between air pollutants in each cohort reflected high correlations between NO_2 and NO_x (correlation coefficients varied between 0.88 and 0.92) and among PM concentrations (correlation coefficients varied between 0.59 and 0.95) (eTable 7 in the [Supplement](#)). Overall, 404 women (4.2%) had hypothyroxinemia, and 506 (6.7%) tested positive for TPOAb.

The short-term associations of air pollutants (concentrations averaged in the first trimester) with hypothyroxinemia and high TSH throughout pregnancy are presented in **Table 2** and **Figure 2**. Unadjusted analyses appear in eTable 8 in the [Supplement](#), and complete-case analysis without imputation of covariates appears in eTable 9 in the [Supplement](#). We found no associations of NO_2 and NO_x concentrations with hypothyroxinemia during pregnancy in the meta-analysis of estimates in cohorts with available data. When we ran the meta-analysis excluding Project Viva, results

remained unchanged (data not shown). Exposures to NO₂ and NO_x were not associated with high TSH during pregnancy.

Women with higher exposures to PM_{2.5} in the first trimester had higher odds of hypothyroxinemia during pregnancy (OR, 1.21; 95% CI, 1.00-1.47) (Table 2 and **Figure 3**; eTable 8 and eTable 9 in the **Supplement**). Exposure to PM₁₀ was not associated with higher odds of hypothyroxinemia (OR, 1.18; 95% CI, 0.93-1.48). When we used the 2.5th percentile as the cutoff for free T₄ level to define hypothyroxinemia, the results did not change (data not shown). The associations of PM exposures and high TSH (concentrations >95th percentile) during pregnancy were not significant (Table 2), but when we used the clinical cutoff of 0.03 to 2.5 mIU/l for a normal TSH level, we found significant associations of exposures to PM_{2.5} and PM₁₀ with high TSH (OR per 5-μg/m³ change of PM_{2.5} level, 1.23; 95% CI, 1.09-1.39; OR per 10-μg/m³ change of PM₁₀ level, 1.24; 95% CI, 1.02-1.51). Levels of PM_{2.5-10} and PM_{2.5} absorbance were not associated with hypothyroxinemia (PM_{2.5-10} level: OR, 1.05; 95% CI, 0.76-1.45; PM_{2.5} absorbance level: OR, 1.05; 95% CI, 0.88-1.26) or high TSH (PM_{2.5-10} level: OR, 1.18; 95% CI, 0.88-1.57; PM_{2.5} absorbance level: OR, 1.10; 95% CI, 0.95-1.26), with effect estimates close to null (Table 2). Examination of the association of air pollution exposure with TPOAb positivity showed large heterogeneity among cohorts (*P* values for heterogeneity for NO₂, PM₁₀, PM_{2.5}, and PM_{2.5-10}, <.001; for NO_x, *P* = .01). Therefore, we only performed cohort-specific analysis for TPOAb. In Generation R, higher concentrations of air pollutants were associated with TPOAb positivity (OR per 10-μg/m³ change of NO₂ level, 1.22; 95% CI, 1.11-1.34; OR per 5-μg/m³ change of PM_{2.5} level, 1.76; 95% CI, 1.51-2.04; OR per 10-μg/m³ change of PM₁₀ level, 1.96; 95% CI, 1.64-2.35). There was no association of air pollution exposure with TPOAb positivity in the other cohorts with available data.

Table 1. Participant Characteristics

Characteristic	No. (%)					
	ABCD, the Netherlands (n = 3867)	Generation R, the Netherlands (n = 2605)	INMA, Spain (n = 2239)	Rhea, Greece (n = 483)	Project Viva, United States (n = 737)	Total (N = 9931)
Age at enrollment, mean (SD), y	33.6 (3.9)	30.8 (4.7)	31.4 (4.2)	29.3 (4.9)	32.5 (4.7)	31.2 (4.8)
Educational levels						
Elementary	683 (17.8)	428 (17.2)	544 (24.4)	97 (20.1)	9 (1.2)	1761 (18.0)
Secondary	1059 (27.6)	756 (30.4)	921 (41.2)	256 (53.0)	179 (24.3)	3171 (32.4)
Higher	2098 (54.6)	1307 (52.4)	769 (34.4)	130 (26.9)	549 (74.5)	4853 (49.6)
Nulliparous	2278 (57.4)	1574 (60.7)	1266 (56.6)	192 (40.4)	366 (49.7)	5616 (56.7)
Foreign country of birth	1210 (31.3)	1115 (42.8)	188 (8.4)	43 (8.9)	119 (16.2)	2675 (27.0)
Single	173 (4.5)	271 (10.9)	38 (1.7)	62 (12.8)	40 (5.4)	584 (6.0)
Never smoked during pregnancy	3503 (90.7)	1767 (75.5)	1485 (68.2)	302 (66.7)	511 (69.5)	7568 (79.1)
Drank alcohol during pregnancy	932 (24.1)	1130 (46.7)	206 (9.6)	102 (25.1)	541 (73.6)	2911 (30.4)
Low socioeconomic status	464 (12.0)	102 (3.9)	1189 (53.1)	40 (8.3)	10 (1.4)	1805 (20.0)
Prepregnancy BMI, mean (SD)	21.9 (3.5)	22.6 (4.4)	22.5 (4.4)	23.4 (5.3)	23.5 (5.3)	23.4 (4.2)
Thyroid function in pregnancy						
Free T ₄ , median (IQR), ng/dL	0.7 (0.7-0.8)	1.2 (1.0-1.3)	0.8 (0.7-0.8)	1.2 (1.1-1.3)	2.1 (1.9-2.3) ^a	0.8 (0.7-1.1) ^b
TSH, median (IQR), mIU/L	1.2 (0.8-1.7)	1.7 (1.0-2.5)	1.3 (0.8-1.8)	1.1 (0.7-1.6)	1.2 (0.7-1.9)	1.3 (0.8-1.9)
Hypothyroxinemia ^c	158 (4.1)	112 (4.4)	88 (4.2)	23 (4.8)	23 (3.2)	404 (4.2)
TPOAb positive	213 (5.5)	145 (6.0)	NA	45 (9.3)	103 (13.9)	506 (6.7)
Gestational age at thyroid measurement, median (IQR), wk	13.0 (11.9-14.0)	13.1 (12.1-16.8)	13.0 (12.3-14.0)	13.0 (12.0-15.0)	9.6 (8.7-10.7)	12.9 (11.9-12.3)

Abbreviations: ABCD, Amsterdam Born Children and Their Development; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); INMA, Infancia y Medio Ambiente; IQR, interquartile range; NA, not available; TPOAb, thyroid peroxidase antibodies; TSH, thyrotropin; T₄, thyroxine.

SI conversion factor: To convert free T₄ to pmol/L, multiply by 12.871.

^a Calculated from total T₄ × triiodothyronine resin uptake.


^b Excluding Project Viva.

^c Defined as free T₄ below the fifth percentile of cohort distribution despite normal TSH level.

Discussion

In a large sample from 5 cohorts in Europe and the United States, we found that first-trimester exposures to PM_{2.5} were associated with mild thyroid dysfunction throughout pregnancy. Exposures

Figure 1. Distribution of Air Pollutants Averaged Across the First Trimester of Pregnancy



Data on particulate matter (PM) were only available in the Sabadell subcohort of Infancia y Medio Ambiente (INMA). Center line indicates 50th percentile; upper and lower boundaries of boxes, 75th and 25th percentile, respectively; upper whisker, 75th percentile plus 1.5 times the interquartile range; lower whisker, 25th percentile minus 1.5 times the interquartile range; and circles, any value above or below the whiskers. ABCD indicates the Amsterdam Born Children and Their Development Study; Gen R, the Generation R Study; NO₂, nitrogen dioxide; NO_x, nitrogen oxides; PM_{2.5}, particulate matter less than 2.5 µm; PM₁₀, particulate matter less than 10 µm; and Viva, Project Viva.

Table 2. Associations of Exposure to Ambient Air Pollutants in the First Trimester With Thyroid Function During Pregnancy

Exposure	Hypothyroxinemia				High TSH			
	Cohorts Included, No. ^a	OR (95% CI) ^b	P _h	I ² , %	Cohorts Included, No. ^a	OR (95% CI) ^b	P _h	I ² , %
NO ₂ , per 10-µg/m ³ change	4	0.96 (0.82-1.12) ^c	.16	41.25	4	1.02 (0.94-1.12)	.83	0
NO _x , per 20-µg/m ³ change	3	0.95 (0.87-1.03)	.60	0	3	0.99 (0.93-1.06)	.80	0
PM _{2.5} , per 5-µg/m ³ change	5	1.21 (1.00-1.47) ^d	.37	6.65	5	1.14 (0.88-1.48)	.12	45.58
PM ₁₀ , per 10-µg/m ³ change	4	1.18 (0.93-1.48)	.33	13.02	4	1.17 (0.87-1.58)	.09	53.72
PM _{2.5-10} , per 5-µg/m ³ change	4	1.05 (0.76-1.45)	.17	40.63	4	1.18 (0.88-1.57)	.12	47.79
PM _{2.5} absorbance, per 10 ⁻⁵ × m ⁻¹ change	4	1.05 (0.88-1.26)	.40	0	4	1.10 (0.95-1.26)	.84	0

Abbreviations: NO₂, nitrogen dioxide; NO_x, nitrogen oxides; OR, odds ratio; P_h, P value of heterogeneity; PM_{2.5-10}, particulate matter between 2.5 and 10 µm; PM₁₀, particulate matter less than 10 µm; PM_{2.5}, particulate matter less than 2.5 µm; TSH, thyrotropin.

^a Data on PM was only available in Sabadell region of Infancia y Medio Ambiente.

^b Estimated using random-effects meta-analysis by cohort (Amsterdam Born Children and Their Development, Generation R, Infancia y Medio Ambiente, Rhea, and Project Viva). The median gestational age at thyroid measurement was at 13 weeks in all cohorts, except for Project Viva, in which measurement was at median gestational age of 10 weeks. Models were adjusted for maternal age at enrollment, educational level,

country of birth, gestational age at thyroid measurement, smoking and alcohol intake during pregnancy, socioeconomic status, marital status, parity, and prepregnancy body mass index (calculated as weight in kilograms divided by height in meters squared). Analysis in Infancia y Medio Ambiente was adjusted for region (ie, Sabadell, Gipuzkoa, Valencia, or Asturias).


^c Excluding participants of Project Viva: OR per 10-µg/m³ change of NO₂ level, 0.97; 95% CI, 0.87-1.08.

^d Excluding participants of Project Viva: OR per 5-µg/m³ change of PM_{2.5} level, 1.23; 95% CI, 1.04-1.47.

to NO_x and NO₂ were not associated with hypothyroxinemia or high TSH during pregnancy. In the Generation R cohort, we observed that pregnant women with higher exposures to NO and PM were more likely to be TPOAb positive.

Studies have found associations of PAH exposure with thyroid dysfunction in nonpregnant populations⁴⁴ and of cigarette smoking with thyroid dysfunction in pregnant women.⁴¹ Three observational studies¹⁶⁻¹⁸ have specifically examined the association of air pollutants with the thyroid function of pregnant women and their neonates. Howe et al¹⁸ showed that prenatal exposures to PM_{2.5} and PM₁₀ but not NO₂ and ozone were associated with higher neonatal total T₄. Zhao et al¹⁷ reported positive associations between residential PM_{2.5} concentrations and maternal hypothyroxinemia during midgestation. Janssen et al¹⁶ found that third trimester PM_{2.5} exposure was negatively associated with free T₄ in maternal serum. Our results extend these observations and showed that the association of PM_{2.5} concentrations with hypothyroxinemia were present in the first trimester, the period when the fetus is most sensitive to maternal thyroid dysfunction. In addition, we found no associations of NO₂ and NO_x exposures with thyroid function in pregnancy, similar to a

Figure 2. Association of Exposure to Nitrogen Dioxide (NO₂) and Nitrogen Oxides (NO_x) in the First Trimester With Thyroid Function During Pregnancy




Odds ratios (ORs) were estimated using random-effects meta-analysis by cohort (the Amsterdam Born Children and Their Development Study [ABCD], Generation R, Infancia y Medio Ambiente [INMA], Rhea, and Project Viva [Viva]). Models were adjusted for pregnant maternal age at enrollment, educational level, country of birth, gestational age at thyroid measurement, smoking and alcohol intake during pregnancy, socioeconomic status, marital status, parity, and prepregnancy body mass index (calculated as weight in kilograms divided by height in meters squared). In addition, analysis in INMA was adjusted for region (Sabadell, Gipuzkoa, Valencia, and Asturias). Data are presented as available in each cohort. Hypothyroxinemia was defined as free thyroxine below the fifth percentile of cohort distribution despite normal thyrotropin (TSH) levels. High TSH concentration was defined as levels higher than the 95th percentile. Size of box indicates weight.

previous study in China.¹⁷ Consistent findings on null associations of NO₂ with thyroid function and our findings on null associations of NO_x and thyroid function, combined with observed associations of PM with thyroid function, suggest that the association of air pollutants with thyroid function may be mostly associated with PM. Importantly, our findings also confirm that the associations of air pollution exposure with thyroid dysfunction in pregnant women are present with concentrations of pollutants at levels much lower than the study in China,¹⁷ as shown in Belgium and California.^{16,18}

While our findings indicate a short-term association of PM exposure with thyroid function, the mechanisms of this association are not fully understood and need further investigation. Although speculative, direct interference in the intracellular action of deiodinase enzymes and the induction of oxidative stress and inflammation are likely among the short-term mechanisms.^{5,45} Autoimmune

Figure 3. Association of Exposure to Particulate Matter With an Aerodynamic Diameter of 2.5 μm or Less (PM_{2.5}) and Particulate Matter With an Aerodynamic Diameter of 10 μm or Less (PM₁₀) in the First Trimester With Thyroid Function During Pregnancy



Odds ratios (ORs) were estimated using random-effects meta-analysis by cohort (the Amsterdam Born Children and Their Development Study [ABCD], Generation R, Infancia y Medio Ambiente [INMA], Rhea, and Project Viva [Viva]). Models were adjusted for pregnant maternal age at enrollment, educational level, country of birth, gestational age at thyroid measurement, smoking and alcohol intake during pregnancy, socioeconomic status, marital status, parity, and prepregnancy body mass index (calculated as weight in kilograms divided by height in meters squared). Data on particulate matter (PM) was only available in the Sabadell region of INMA. Data are presented as available in each cohort. Hypothyroxinemia was defined as free thyroxine below the fifth percentile of cohort distribution despite normal thyrotropin (TSH) levels. High TSH concentration was defined as levels higher than the 95th percentile. Size of box indicates weight.

processes might act on thyroid function in a longer period. In women of reproductive age, autoimmunity is a common cause of thyroid dysfunction in iodine-sufficient areas. Earlier studies among pregnant women have shown that prolonged lead exposure is associated with TPOAb positivity and subsequently low thyroid function.⁴⁶ We found that exposure to PM during early pregnancy was associated with higher odds of thyroid autoimmunity in Generation R, an iodine-sufficient cohort in the Netherlands. The observed associations between PM exposures and low thyroid function and the null association with NO₂, NO_x, and PM_{2.5} absorbance, a measure of black carbon, suggest that the composition of PM rather than the general markers of traffic-related pollution may be responsible for thyroid disruption. One hypothesis—supported by in vitro studies⁴⁷—is that PAHs are associated with the thyroid toxic effects of PM exposure, but we cannot rule out the role of other components, such as trace elements.

The cohorts in our study varied with respect to the concentrations of air pollution exposure. For example, ABCD and Generation R had higher concentrations of PM_{2.5} exposure compared with other cohorts. Exposure to concentrations of NO₂ and PM measures in the European cohorts were positive and moderate to strong, but there was a negative and small correlation between NO₂ and PM_{2.5} in the US cohort Project Viva. These differences can be explained by varying sources for exposure to pollutants in different regions as well as different exposure assessment methods in European cohorts and in Project Viva. Nonetheless, examination of the heterogeneity in estimates for analysis of the association of NO₂ and PM_{2.5} with thyroid function confirmed that estimates among cohorts could be combined in the meta-analysis. There was an inverse but imprecise association of PM_{2.5} with hypothyroxinemia in Project Viva, the smallest cohort with the lowest concentrations of PM_{2.5}, suggesting that the association might be present at a threshold of exposure. Nonetheless, results of analyses using generalized additive models to create the smoothing curve spline confirmed no threshold effect of pollutants. With regard to TPOAb positivity, appropriate testing showed large heterogeneity among cohorts. In particular, a larger number of women were positive for TPOAb in Project Viva compared with other cohorts, potentially explained by natural variation, use of different assays, slightly higher mean age in Project Viva at the time of assessment, and earlier measurement during pregnancy. Also, the association of air pollution exposure with TPOAb positivity was only present in Generation R. Further investigation is needed to determine whether the iodine status of the cohort population or other characteristics might explain the differences in the association of air pollution exposure with thyroid autoimmunity.

Strengths and Limitations

This study has several strengths, including a large number of participants from regions with different iodine status and diverse sociodemographic characteristics in Europe and the United States. Nonetheless, this study had important limitations. We had only measures of TSH, free T₄, and TPOAb during pregnancy (mostly in the first half of pregnancy), and measurements were performed using different assays among cohorts. While the absolute concentrations of TSH and free T₄ can vary between assays, we defined the outcomes by population- and assay-specific cutoffs to overcome any issues related to the interchangeability of absolute-concentration assay results. We adjusted the models for history of smoking in pregnancy, but we did not have data on secondhand tobacco smoke exposure in all cohorts. We relied on residential addresses to estimate air pollution exposure, without consideration for within-individual spatial variation in exposure. Furthermore, estimation of air pollution exposure in European cohorts did not account for temporal variation during the first trimester, and estimation of NO₂ exposure in Project Viva did not account for spatial variation between individuals. Another limitation concerning the European cohorts is that air pollution exposure models were developed based on monitoring campaigns performed between 2009 and 2011 and used to estimate exposures of pregnant women in preceding years. We used routine monitoring data to back-extrapolate the concentrations to the exact first trimester of pregnancy. Therefore, we assumed that the spatial distribution of the sources and predictors of air pollution levels remained stable over time, as previous research has shown.³⁶ We did not have history of

addresses in pregnant women and thus could not examine the association of air pollution exposure prior to pregnancy with thyroid function. Subsequently, no conclusion on the critical window of exposure or accumulation of risk can be drawn from this analysis.

Conclusions

The findings of this study suggest that first-trimester exposures to PM_{2.5} were associated with mild thyroid dysfunction throughout pregnancy. The association of PM_{2.5} exposure with thyroid function in pregnant women is of global health importance because air pollution exposure is widespread and hypothyroxinemia may adversely influence offspring brain development.

ARTICLE INFORMATION

Accepted for Publication: August 15, 2019.

Published: October 16, 2019. doi:10.1001/jamanetworkopen.2019.12902

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2019 Ghassabian A et al. *JAMA Network Open*.

Corresponding Author: Mònica Guxens, MD, ISGlobal, Barcelona Institute for Global Health, Campus MAR, Barcelona Biomedical Research Park (PRBB) (Office 190), Doctor Aiguader, 88, 08003 Barcelona, Spain (monica.guxens@isglobal.org).

Author Affiliations: Departments of Pediatrics, Environmental Medicine, and Population Health, School of Medicine, New York University, New York (Ghassabian); ISGlobal, Barcelona, Spain (Pierotti, Julvez, Vrijheid, Sunyer, Guxens); Spanish Consortium for Research on Epidemiology and Public Health, Madrid, Spain (Pierotti, Basterrechea, Estarlich, Fernández-Somoano, Julvez, Lertxundi, Lopez-Espinosa, Tardón, Vrijheid, Sunyer, Guxens); Universitat Pompeu Fabra, Barcelona, Spain (Pierotti, Julvez, Vrijheid, Sunyer, Guxens); Biodonostia Health Research Institute, San Sebastian, Spain (Basterrechea, Lertxundi); Public Health Division of Gipuzkoa, Basque Government, San Sebastian, Spain (Basterrechea); Department of Social Medicine, University of Crete, Heraklion, Greece (Chatzi, Karakosta); Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles (Chatzi); Epidemiology and Environmental Health Joint Research Unit, The Foundation for the Promotion of Health and Biomedical Research of Valencia Region, Universitat Jaume I-Universitat de València, Valencia, Spain (Estarlich, Lopez-Espinosa); Instituto Universitari de Oncología del Principado de Asturias, Department of Medicine, University of Oviedo, Oviedo, Spain (Fernández-Somoano, Tardón); Department of Pediatric Endocrinology and Diabetes, Maine Medical Center, Portland (Fleisch); Center for Outcomes Research and Evaluation, Maine Medical Center Research Institute, Portland (Fleisch); Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts (Gold); Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Gold); Hospital del Mar Research Institute, Barcelona, Spain (Julvez, Sunyer); Clinical Microbiology Laboratory, Attikon University Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece (Karakosta); Department of Public Health and Preventive Medicine, University of Basque Country, Bilbao, Spain (Lertxundi); Academic Center for Thyroid Diseases, Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands (Mulder, Korevaar, Peeters); Department of Child and Adolescent Psychiatry, Erasmus Medical Center–Sophia Children's Hospital, Rotterdam, the Netherlands (Mulder, Tiemeier, Guxens); Division of Chronic Disease Research Across the Lifecourse, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institution, Boston, Massachusetts (Oken, Rifas-Shiman); Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Oken); Department of Chemistry, University of Crete, Heraklion, Greece (Stephanou); Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Tiemeier); Department of Public Health, Amsterdam Public Health Research Institute, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, the Netherlands (Vrijkotte).

Author Contributions: Dr Guxens had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Ghassabian, Lertxundi, Lopez-Espinosa, Peeters, Guxens.

Acquisition, analysis, or interpretation of data: Ghassabian, Pierotti, Basterrechea, Chatzi, Estarlich, Fernández-Somoano, Fleisch, Gold, Julvez, Karakosta, Lopez-Espinosa, Mulder, Korevaar, Oken, Peeters, Rifas-Shiman, Stephanou, Tardón, Tiemeier, Vrijheid, Vrijkotte, Sunyer, Guxens.

Drafting of the manuscript: Ghassabian, Pierotti, Gold, Lopez-Espinosa, Stephanou, Guxens.

Critical revision of the manuscript for important intellectual content: Basterrechea, Chatzi, Estarlich, Fernández-Somoano, Fleisch, Julvez, Karakosta, Lertxundi, Lopez-Espinosa, Mulder, Korevaar, Oken, Peeters, Rifas-Shiman, Stephanou, Tardón, Tiemeier, Vrijheid, Vrijkotte, Sunyer, Guxens.

Statistical analysis: Ghassabian, Pierotti, Estarlich, Korevaar, Rifas-Shiman, Guxens.

Obtained funding: Chatzi, Gold, Lopez-Espinosa, Oken, Tardón, Vrijheid, Vrijkotte, Sunyer, Guxens.

Administrative, technical, or material support: Fernández-Somoano, Karakosta, Lopez-Espinosa, Korevaar, Oken, Peeters, Tardón, Vrijkotte.

Supervision: Basterrechea, Chatzi, Estarlich, Lertxundi, Peeters, Stephanou, Tiemeier, Vrijheid, Vrijkotte, Guxens.

Conflict of Interest Disclosures: Dr Basterrechea reported receiving grants from the Instituto de Salud Carlos III and the Department of Health of the Basque Government during the conduct of the study. Dr Fleisch reported receiving grants from the National Institutes of Health during the conduct of the study. Dr Gold reported receiving grants from the National Institutes of Health during the conduct of the study. Dr Lopez-Espinosa reported receiving grants and personal fees from the Instituto de Salud Carlos III and grants from the Alicia Koplowitz Foundation 2017 during the conduct of the study. Dr Oken reported receiving grants from the National Institutes of Health during the conduct of the study. Dr Vrijkotte reported receiving grants from the Netherlands Institute for Research and Development during the conduct of the study and outside the submitted work. Dr Sunyer reported receiving grants from the Instituto de Salud Carlos III, the EU Commission, the Centro de Investigación Biomedica en Red Epidemiología y Salud Pública, and the Generalitat de Catalunya during the conduct of the study. Dr Guxens reported receiving grants from the Instituto de Salud Carlos III, the EU Commission, and the Centro de Investigación Biomedica en Red Epidemiología y Salud Pública during the conduct of the study. No other disclosures were reported.

Funding/Support: The Amsterdam Born Children and Their Development Study was supported by the Netherlands Organization for Health Research and Development, Public Health Service Amsterdam, and the Amsterdam UMC. The Generation R Study was supported by the Erasmus Medical Center, Rotterdam; Erasmus University, Rotterdam; the Netherlands Organization for Health Research and Development; the Netherlands Organization for Scientific Research; and the Ministry of Health, Welfare and Sport. Nederlandse Organisatie voor Toegepast Natuurwetenschappelijk Onderzoek received funding from the Netherlands Ministry of Infrastructure and the Environment to support exposure assessment. Drs Mulder, Korevaar, Peeters, and Tiemeier were supported by grant S17-19 from the Sophia Children's Hospital Foundation. Dr Tiemeier was supported by grant 016.VICI.170.200 from the Netherlands Organization for Health Research and Development. The Generation R Study was conducted by the Erasmus Medical Center in close collaboration with the Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, Rotterdam, and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond, Rotterdam. Infancia y Medio Ambiente was funded by grants Red INMA G03/176 and CBO6/02/0041 FIS-FEDER P103/1615, P104/1509, P104/1112, P104/1931, P1042018, P105/1079, P105/1052, P106/0867, P106/1213, P107/0314, P109/02311, P109/02647, P111/02591, P111/02038, P113/1944, P113/2032, P113/02429, P114/00891, P114/01687, P116/1288, and P117/00663, and Miguel Servet-FEDER CP11/00178, CP15/00025, MS116/00051, MS13/00054, and CP118/00018 from the Instituto de Salud Carlos III; grant CIRIT 1999SGR 00241 from the Generalitat de Catalunya; grants FISABIO UGP 15-230, UGP-15-244, and UGP-15-249 from the Generalitat Valenciana; grant 2005111093 from the Department of Health of the Basque Government; grant DFG06/002 from the Provincial Government of Gipuzkoa, annual agreements with the municipalities in the study area (Zumarraga, Urretxu, Legazpi, Azkoitia, Azpeitia, and Beasain); grants FP7/2007-2011-GA#211250, FP7-ENV-2011 cod 282957, HEALTH.2010.2.4.5-1 from the EU Commission; the Obra Social Cajastur and Fundación Liberbank; the Universidad de Oviedo; the Centro de Investigación Biomedica en Red Epidemiología y Salud Pública. ISGlobal is a member of the CERCA Programme, Generalitat de Catalunya. The Rhea project was supported by European projects (EU FP6-2003-Food-3-A NewGeneris, EU FP6. STREP Hiwate, EU FP7 ENV.2007.1.2.2.2. Project No 211250 Escape, EU FP7-2008-ENV-1.2.1.4 Envirogenomarkers, EU FP7-HEALTH-2009-single-stage CHICOS, EU FP7 ENV.2008.1.2.1.6. Proposal No 226285 ENRIECO) and the Greek Ministry of Health (Program of Prevention of Obesity and Neurodevelopmental Disorders in Preschool Children, in Heraklion district, Crete, Greece, 2011-2014). Project Viva was funded by grants R01 HD 034568, UH3 ODO23286, K23 ESO24803 from the National Institutes of Health.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We gratefully acknowledge the contribution of children and parents, general practitioners, hospitals, midwives, and pharmacies in Rotterdam.

REFERENCES

1. Guxens M, Garcia-Esteban R, Giorgis-Allemand L, et al. Air pollution during pregnancy and childhood cognitive and psychomotor development: six European birth cohorts. *Epidemiology*. 2014;25(5):636-647. doi:10.1097/EDE.000000000000133
2. Suades-González E, Gascon M, Guxens M, Sunyer J. Air pollution and neuropsychological development: a review of the latest evidence. *Endocrinology*. 2015;156(10):3473-3482. doi:10.1210/en.2015-1403
3. Sunyer J, Esnaola M, Alvarez-Pedrerol M, et al. Association between traffic-related air pollution in schools and cognitive development in primary school children: a prospective cohort study. *PLoS Med*. 2015;12(3):e1001792. doi:10.1371/journal.pmed.1001792
4. Guxens M, Lubczyńska MJ, Muetzel RL, et al. Air pollution exposure during fetal life, brain morphology, and cognitive function in school-age children. *Biol Psychiatry*. 2018;84(4):295-303. doi:10.1016/j.biopsych.2018.01.016
5. Block ML, Calderón-Garcidueñas L. Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci*. 2009;32(9):506-516. doi:10.1016/j.tins.2009.05.009
6. Thomson EM. Neurobehavioral and metabolic impacts of inhaled pollutants: a role for the hypothalamic-pituitary-adrenal axis? *Endocr Disruptors (Austin)*. 2013;1(1):e27489. doi:10.4161/endo.27489
7. Klein GP, Hodge EM, Diamond ML, et al. Gas-phase ambient air contaminants exhibit significant dioxin-like and estrogen-like activity in vitro. *Environ Health Perspect*. 2006;114(5):697-703. doi:10.1289/ehp.8496
8. Clemons JH, Allan LM, Marvin CH, et al. Evidence of estrogen- and TCDD-like activities in crude and fractionated extracts of PM10 air particulate material using in vitro gene expression assays. *Environ Sci Technol*. 1998;32(12):1853-1860. doi:10.1021/es971124n
9. Kelishadi R, Sobhani P, Poursafa P, et al. Is there any association between urinary metabolites of polycyclic aromatic hydrocarbons and thyroid hormone levels in children and adolescents? *Environ Sci Pollut Res Int*. 2018;25(2):1962-1968. doi:10.1007/s11356-017-0577-y
10. Movahedinia A, Salamat N, Kheradmand P. Effects of the environmental endocrine disrupting compound benzo[a]pyrene on thyroidal status of abu mullet (*Liza abu*) during short-term exposure. *Toxicol Rep*. 2018;5:377-382. doi:10.1016/j.toxrep.2018.02.018
11. Haddow JE, Palomaki GE, Allan WC, et al. Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child. *N Engl J Med*. 1999;341(8):549-555. doi:10.1056/NEJM199908193410801
12. Levie D, Korevaar TIM, Bath SC, et al. Thyroid function in early pregnancy, child IQ, and autistic traits: a meta-analysis of individual participant data. *J Clin Endocrinol Metab*. 2018;103(8):2967-2979. doi:10.1210/jc.2018-00224
13. Ghassabian A, El Marroun H, Peeters RP, et al. Downstream effects of maternal hypothyroxinemia in early pregnancy: nonverbal IQ and brain morphology in school-age children. *J Clin Endocrinol Metab*. 2014;99(7):2383-2390. doi:10.1210/jc.2013-4281
14. Modesto T, Tiemeier H, Peeters RP, et al. Maternal mild thyroid hormone insufficiency in early pregnancy and attention-deficit/hyperactivity disorder symptoms in children. *JAMA Pediatr*. 2015;169(9):838-845. doi:10.1001/jamapediatrics.2015.0498
15. Zoeller RT, Brown TR, Doan LL, et al. Endocrine-disrupting chemicals and public health protection: a statement of principles from The Endocrine Society. *Endocrinology*. 2012;153(9):4097-4110. doi:10.1210/en.2012-1422
16. Janssen BG, Saenen ND, Roels HA, et al. Fetal thyroid function, birth weight, and in utero exposure to fine particle air pollution: a birth cohort study. *Environ Health Perspect*. 2017;125(4):699-705. doi:10.1289/EHP508
17. Zhao Y, Cao Z, Li H, et al. Air pollution exposure in association with maternal thyroid function during early pregnancy. *J Hazard Mater*. 2019;367:188-193. doi:10.1016/j.jhazmat.2018.12.078
18. Howe CG, Eckel SP, Habre R, et al. Association of prenatal exposure to ambient and traffic-related air pollution with newborn thyroid function: findings from the Children's Health Study. *JAMA Netw Open*. 2018;1(5):e182172. doi:10.1001/jamanetworkopen.2018.2172
19. Ghassabian A, Bongers-Schokking JJ, Henrichs J, et al. Maternal thyroid function during pregnancy and behavioral problems in the offspring: the Generation R Study. *Pediatr Res*. 2011;69(5, pt 1):454-459. doi:10.1203/PDR.Ob013e3182125b0c
20. Gawda A, Majka G, Nowak B, Marcinkiewicz J. Air pollution, oxidative stress, and exacerbation of autoimmune diseases. *Cent Eur J Immunol*. 2017;42(3):305-312. doi:10.5114/ceji.2017.70975
21. Derakhshan A, Korevaar TIM, Taylor PN, et al. The association of maternal thyroid autoimmunity during pregnancy with child IQ. *J Clin Endocrinol Metab*. 2018;103(10):3729-3736. doi:10.1210/jc.2018-00743

22. Ghassabian A, Bongers-Schokking JJ, de Rijke YB, et al. Maternal thyroid autoimmunity during pregnancy and the risk of attention deficit/hyperactivity problems in children: the Generation R Study. *Thyroid*. 2012;22(2):178-186.
23. Korevaar TI, Schalekamp-Timmermans S, de Rijke YB, et al. Hypothyroxinemia and TPO-antibody positivity are risk factors for premature delivery: the Generation R Study. *J Clin Endocrinol Metab*. 2013;98(11):4382-4390. doi:10.1210/jc.2013-2855
24. van Eijsden M, Vrijkotte TG, Gemke RJ, van der Wal MF. Cohort profile: the Amsterdam Born Children and Their Development (ABCD) Study. *Int J Epidemiol*. 2011;40(5):1176-1186. doi:10.1093/ije/dyq128
25. Kooijman MN, Kruijthof CJ, van Duijn CM, et al. The Generation R Study: design and cohort update, 2017. *Eur J Epidemiol*. 2016;31(12):1243-1264. doi:10.1007/s10654-016-0224-9
26. Guxens M, Ballester F, Espada M, et al; INMA Project. Cohort profile: the INMA—Infancia y Medio Ambiente—(Environment and Childhood) Project. *Int J Epidemiol*. 2012;41(4):930-940. doi:10.1093/ije/dyr054
27. Chatzi L, Leventakou V, Vafeiadi M, et al. Cohort profile: the Mother-Child Cohort in Crete, Greece (Rhea Study). *Int J Epidemiol*. 2017;46(5):1392-1393k. doi:10.1093/ije/dyx084
28. Oken E, Baccarelli AA, Gold DR, et al. Cohort profile: Project Viva. *Int J Epidemiol*. 2015;44(1):37-48. doi:10.1093/ije/dyu008
29. Kippler M, Bottai M, Georgiou V, et al. Impact of prenatal exposure to cadmium on cognitive development at preschool age and the importance of selenium and iodine. *Eur J Epidemiol*. 2016;31(11):1123-1134. doi:10.1007/s10654-016-0151-9
30. Ghassabian A, Steenweg-de Graaff J, Peeters RP, et al. Maternal urinary iodine concentration in pregnancy and children's cognition: results from a population-based birth cohort in an iodine-sufficient area. *BMJ Open*. 2014;4(6):e005520. doi:10.1136/bmjopen-2014-005520
31. Murcia M, Rebagliato M, Espada M, et al; INMA Study Group. Iodine intake in a population of pregnant women: INMA mother and child cohort study, Spain. *J Epidemiol Community Health*. 2010;64(12):1094-1099. doi:10.1136/jech.2009.092593
32. Pearce EN, Bazrafshan HR, He X, Pino S, Braverman LE. Dietary iodine in pregnant women from the Boston, Massachusetts area. *Thyroid*. 2004;14(4):327-328. doi:10.1089/105072504323031013
33. Eeftens M, Beelen R, de Hoogh K, et al. Development of land use regression models for PM(2.5), PM(2.5) absorbance, PM(10) and PM(coarse) in 20 European study areas: results of the ESCAPE project. *Environ Sci Technol*. 2012;46(20):11195-11205. doi:10.1021/es301948k
34. Beelen R, Hoek G, Vienneau D, et al. Development of NO2 and NOx land use regression models for estimating air pollution exposure in 36 study areas in Europe: the ESCAPE project. *Atmos Environ*. 2013;72:10-23. doi:10.1016/j.atmosenv.2013.02.037
35. Kloog I, Coull BA, Zanobetti A, Koutrakis P, Schwartz JD. Acute and chronic effects of particles on hospital admissions in New-England. *PLoS One*. 2012;7(4):e34664. doi:10.1371/journal.pone.0034664
36. Eeftens M, Beelen R, Fischer P, Brunekreef B, Meliefste K, Hoek G. Stability of measured and modelled spatial contrasts in NO(2) over time. *Occup Environ Med*. 2011;68(10):765-770. doi:10.1136/oem.2010.061135
37. Department of Environmental Protection. MassAir Online. <http://eeaonline.eea.state.ma.us/dep/massair/web/#/pollution/map/max>. Accessed August 27, 2019.
38. Alexander EK, Pearce EN, Brent GA, et al. 2017 guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and the postpartum. *Thyroid*. 2017;27(3):315-389. doi:10.1089/thy.2016.0457
39. Chi GC, Hajat A, Bird CE, et al. Individual and neighborhood socioeconomic status and the association between air pollution and cardiovascular disease. *Environ Health Perspect*. 2016;124(12):1840-1847. doi:10.1289/EHP199
40. Alderete TL, Chen Z, Toledo-Corral CM, et al. Ambient and traffic-related air pollution exposures as novel risk factors for metabolic dysfunction and type 2 diabetes. *Curr Epidemiol Rep*. 2018;5(2):79-91. doi:10.1007/s40471-018-0140-5
41. Männistö T, Hartikainen A-L, Vääräsmäki M, et al. Smoking and early pregnancy thyroid hormone and anti-thyroid antibody levels in euthyroid mothers of the Northern Finland Birth Cohort 1986. *Thyroid*. 2012;22(9):944-950. doi:10.1089/thy.2011.0377
42. Mahwi TO, Abdulateef DS. Relation of different components of climate with human pituitary-thyroid axis and FT3/FT4 ratio: a study on euthyroid and SCH subjects in two different seasons. *Int J Endocrinol*. 2019;2019:2762978. doi:10.1155/2019/2762978

43. Ren C, Tong S. Temperature modifies the health effects of particulate matter in Brisbane, Australia. *Int J Biometeorol*. 2006;51(2):87-96. doi:10.1007/s00484-006-0054-7
44. Jain RB. Association between polycyclic aromatic hydrocarbons and thyroid function among males and females: data from NHANES 2007-2008. *Int J Environ Health Res*. 2016;26(4):405-419. doi:10.1080/09603123.2015.1135311
45. Mancini A, Di Segni C, Raimondo S, et al. Thyroid hormones, oxidative stress, and inflammation. *Mediators Inflamm*. 2016;2016:6757154. doi:10.1155/2016/6757154
46. Kahn LG, Liu X, Rajovic B, et al. Blood lead concentration and thyroid function during pregnancy: results from the Yugoslavia Prospective Study of Environmental Lead Exposure. *Environ Health Perspect*. 2014;122(10):1134-1140. doi:10.1289/ehp.1307669
47. Sun H, Shen OX, Xu XL, Song L, Wang XR. Carbaryl, 1-naphthol and 2-naphthol inhibit the beta-1 thyroid hormone receptor-mediated transcription in vitro. *Toxicology*. 2008;249(2-3):238-242. doi:10.1016/j.tox.2008.05.008

SUPPLEMENT.

eMethods. Measurements of Covariates

eTable 1. Immunoassays Used for Measurement of Thyroid Parameters in 5 Participating Cohorts

eTable 2. Cohort-Specific Cutoffs for Free Thyroxine (T₄) and Thyrotropin (TSH) in 5 Cohorts

eTable 3. Participants' Characteristics in Pregnant Women Included in the Analysis and Those Women Excluded Because of Missing Data in Exposure and/or Outcome

eTable 4. Variables Used in Forward Selection Logistic Regression Model to Calculate Inverse Probability of Attrition Weights

eTable 5. Imputation Modelling

eTable 6. Concentrations of Air Pollutants Averaged Over the First Trimester of Pregnancy Based on Residential Addresses in 5 Cohorts

eTable 7. Spearman Correlations Between Concentrations of Traffic-Related Air Pollutants in 5 Cohorts

eTable 8. Associations Between Exposure to Ambient Air Pollutants in the First Trimester and Thyroid Function During Pregnancy: Unadjusted Analysis

eTable 9. Associations Between Exposure to Ambient Air Pollutants in the First Trimester and Thyroid Function During Pregnancy: Complete Case Analysis

eFigure 1. Distribution of PM_{2.5-10} and PM_{2.5} Absorbance Averaged Across the First Trimester of Pregnancy in 5 Birth Cohorts

eFigure 2. Exposure to PM_{2.5-10} and PM_{2.5} Absorbance in the First Trimester and Thyroid Function During Pregnancy