POSTER PRESENTATION





Association of HLA DRB variants to genetic predisposition to rheumatoid arthritis

Bushra S. Jarullah

From 2nd International Genomic Medical Conference (IGMC 2013) Jeddah, Kingdom of Saudi Arabia. 24-27 November 2013

Background

Genetic predisposition to rheumatoid arthritis related to the presence of specific polymorphic HLA sequences has been frequently reported [1,2]. HLA class II is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Association studies have provided evidence that a sequence of amino acids, termed the shared epitope, are involved in the disease process [3].

Materials and methods

In order to understand the influence of variation in HLA DRB genes on rheumatoid arthritis sequence comparison of 21 allelic variants of HLA DRB gene was carried out.

Results

Although a lot of variation was observed in the 5' ends, almost all sequences showed alignment scores of above 200 for 30-40% of the sequence towards the 3' end. High levels of variations were however observed in two alleles; HLA-DRB1- 1318 and HLA-B*5904. Both these alleles have been recently reported as novel variants of HLA [4,5].

Conclusions

Our results strongly support the shared epitope hypothesis presenting an explanation for the susceptibility to rheumatoid arthritis. Yet considering the length of similar sequence at the 3' end and the hyper variable region at the 5' end, we conclude the hypothesis may not completely explain the course of disease severity and emphasize the need to develop superior biomarkers to predict the course of the disease.

Department of Biotechnology, Kadi Sarva Vishwavidyalaya, Gandhinagar, Gujarat, India

Published: 2 April 2014

References

- Deighton CM, Wentzel J, Cavanagh G, Roberts DF, Walker DJ: Contribution of inherited factors to rheumatoid arthritis. Ann Rheum Dis 1992, 51:182-5.
- Weyand CM, Hicok KC, Conn DL, Goronzy JJ: The influence of HLADRB1 genes on disease severity in rheumatoid arthritis. Ann Intern Med 1992, 117:801-806.
- 3. Gregersen PK, Silver J, Winchester RJ: The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 1987, **30**:1205-1213.
- Zetterquist H, Olerup O: A novel DRB1 allele (HLA-DRB1*1318) featuring a DR8-associated sequence motif on a DR52 haplotype. *Tissue Antigens* 1995, 46:337-339.
- Ueta M, Matsushita M, Sotozono C, Kinoshita S, Tokunaga K: Identification of a novel HLA-B allele, HLA-B*5904. *Tissue Antigens* 2009, 73:612-614.

doi:10.1186/1471-2164-15-S2-P32

Cite this article as: Jarullah: Association of HLA DRB variants to genetic predisposition to rheumatoid arthritis. *BMC Genomics* 2014 15(Suppl 2): P32.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

) BioMed Central

Submit your manuscript at www.biomedcentral.com/submit



© 2014 Jarullah; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correspondence: bjarullah@yahoo.com