

SCIENTIFIC REPORTS



OPEN

Association of Prehospital Epinephrine Administration With Survival Among Patients With Traumatic Cardiac Arrest Caused By Traffic Collisions

Makoto Aoki¹ , Toshikazu Abe^{2,3} & Kiyohiro Oshima¹

For traumatic cardiac arrest (TCA), the effect of prehospital epinephrine administration was unclear. The aim of this study was to evaluate the relationship between prehospital epinephrine administration and survival in patients with TCA caused by traffic collisions. We conducted a nationwide, prospective, population-based observational study involving patients who experienced out-of-hospital cardiac arrest (OHCA) by using the All-Japan Utstein Registry. Blunt trauma patients with TCA who received prehospital epinephrine were compared with those who did not receive prehospital epinephrine. The primary outcome was 1-month survival of patients. The secondary outcome was prehospital return of spontaneous circulation (ROSC). A total of 5,204 patients with TCA were analyzed. Of those, 758 patients (14.6%) received prehospital epinephrine (Epinephrine group), whereas the remaining 4,446 patients (85.4%) did not receive prehospital epinephrine (No epinephrine group). Eleven (1.5%) and 41 (0.9%) patients in the Epinephrine and No epinephrine groups, respectively, survived for 1 month. In addition, 74 (9.8%) and 40 (0.9%) patients achieved prehospital ROSC in the Epinephrine and No epinephrine groups, respectively. In multivariable logistic regression models, prehospital epinephrine administration was not associated with 1-month survival (odds ratio [OR] 1.495, 95% confidence interval [CI] 0.758 to 2.946) and was associated with prehospital ROSC (OR 3.784, 95% CI 2.102 to 6.812). A propensity score-matched analysis showed similar results for 1-month survival (OR 2.363, 95% CI 0.606 to 9,223) and prehospital ROSC (OR 6.870, 95% CI 3.326 to 14.192). Prehospital epinephrine administration in patients with TCA was not associated with 1-month survival, but was beneficial in regard to prehospital ROSC.

Various treatments have been developed to decrease the mortality caused by trauma¹. However, the mortality rate related to traumatic cardiac arrest (TCA) remains high and the latest systematic review reported the mortality was 97.6%²⁻⁹. Prehospital advanced life support (ALS) is currently widely used. An observational study reported that prehospital ALS was associated with survival of patients with TCA¹⁰. However, the component of prehospital ALS associated with survival is unknown¹¹.

Recently, numerous observational studies and the latest randomized control trial of endogenous cardiac arrest reported that prehospital epinephrine administration was associated with patient survival¹²⁻¹⁴. However, few TCA patients were included in this randomized control trial.

Resuscitation algorithm for TCA was naturally introduced from the immediate treatment of reversible causes such as resuscitative thoracotomy and did not provide obvious opinion of epinephrine administration in patients with TCA¹⁵⁻¹⁷. There is limited evidence regarding the effectiveness of prehospital epinephrine administration in patients with TCA, except one small observational study¹¹. The study revealed that prehospital administration of epinephrine in patients with TCA was associated with increased survival-to-discharge. However, half of the

¹Department of Emergency Medicine, Gunma University Graduate School of Medicine, Gunma, Japan. ²Department of General Medicine, Juntendo University, Tokyo, Japan. ³Department of Health Services Research, Faculty of Medicine, University of Tsukuba, Ibaraki, Japan. Correspondence and requests for materials should be addressed to M.A. (email: aokimakoto@gunma-u.ac.jp)




Figure 1. Flow chart of patients included in this study. ALS, advanced life support; CPR, cardiopulmonary resuscitation; ROSC, return of spontaneous circulation.

patients who survived in this study were penetrating trauma patients. Thus, the efficacy of prehospital epinephrine administration in blunt trauma patients with TCA was unknown.

The objective of the present study was to investigate the efficacy of prehospital epinephrine administration in blunt trauma patients with TCA.

Methods

Study design and setting. This study was a post-hoc analysis of data from the All-Japan Utstein Registry – a prospective, nationwide, population-based out-of-hospital cardiac arrest (OHCA) registry system based on the Utstein style – established by the Fire and Disaster Management Agency^{18,19}. This OHCA registry includes all patients with TCA caused by traffic collisions, who were treated by the emergency medical service (EMS) from January 1, 2012 to December 31, 2015. The analysis protocol of this study was approved by the Ethics Committee of the Gunma University Hospital.

Cardiac arrest was defined as the cessation of cardiac mechanical activity, confirmed by the absence of signs of circulation^{18,19}. The cause of arrest was presumed to be of traumatic origin based on circumstantial evidence. Confirmation of the traumatic origin of cardiac arrest was clinically determined by the treating physicians who collaborated with the EMS personnel.

EMS system in Japan and data collection. The free emergency telephone number 1-1-9 is used to call an ambulance from anywhere in Japan. Emergency services are provided 24 hours per day. Following a call, an ambulance is dispatched from the nearest fire station. In general, an ambulance includes a crew of three EMS personnel. Most ambulances include at least one emergency life-saving technician (ELST), certified to insert an intravenous catheter and an adjunct airway and use semiautomated external defibrillators in patients with OHCA. In addition, since July 2004 and April 2006, specially trained ELST are certified to insert an endotracheal tube and administer intravenous epinephrine. In 2014, almost all (97.4%) ambulances included at least one ELST and the vast majority of those (82.9%) were specially trained ELSTs²⁰. However, these ALS procedures cannot be performed without the instruction of a medical director in each municipality. The treatment of cardiac arrest is based on the Japanese cardiopulmonary resuscitation (CPR) guidelines, derived from those established by the American Heart Association, the European Resuscitation Council, and the International Liaison Committee on Resuscitation guidelines. EMS providers are not permitted to terminate resuscitation in the field²¹. Therefore, most patients with OHCA are treated by EMS personnel, transported to a hospital, and registered in the All-Japan Utstein Registry.

Patient selection. The patient flow chart is shown in Fig. 1. Exclusion criteria were: no resuscitation, ALS performed by a physician (i.e., bias from another ALS procedure such as blood transfusion, insertion of chest tube, thoracotomy, resuscitative endovascular occlusion of the aorta, etc.), unknown adrenaline use, CPR duration time < 1 min, CPR duration time > 30 min (i.e., unsurvivable long time), and prehospital return of spontaneous circulation (ROSC) within 10 min from CPR (i.e., bias caused by inclusion of patients successfully resuscitated prior to epinephrine administration).

Data collection. Data regarding patient age, gender, type of bystander witness status, first recorded cardiac rhythm, life support by EMS personnel (i.e., use of ALS devices, insertion of an intravenous line), time course of resuscitation and epinephrine administration, prehospital ROSC, and 1-month survival were obtained. Data regarding EMS times of call receipt, arrival of the ambulance at the scene of the accident, contact with patients, initiation of CPR, defibrillation performed by EMS personnel, and arrival at the hospital were recorded using the clock of each EMS system. In cases of shock delivery by bystanders using a public-access automated external defibrillator (AED), the patient's first recorded rhythm was regarded as ventricular fibrillation or pulseless

ventricular tachycardia. The estimated times of collapse, initiation of public-access AED shocks, and initiation of bystander CPR were obtained through an interview with the bystander performed by the EMS personnel. The type of bystander CPR was obtained through observation and interview with the bystander performed by the EMS personnel before leaving the scene of the accident. A set of specific questions on the presence or absence of chest compressions and rescue breathing was used for this purpose. The time interval from collapse to shocks using public-access AEDs was replaced with time to bystander CPR, unless the time of shocks delivered using public access AEDs was available. Patients who survived OHCA were followed up for a maximum of 1 month after the event by the EMS personnel in charge. The data forms were completed by the EMS personnel in cooperation with the treating physicians. The data were integrated into the All-Japan Utstein Registry database server and logically checked by the computer system. In cases of incomplete data forms, the Fire and Disaster Management Agency requested the provision of missing data from the respective fire station.

Outcomes. The primary outcome was 1-month survival. The secondary outcome was prehospital return of spontaneous circulation (ROSC).

Statistical analysis. Continuous variables were expressed as medians (interquartile range). The patients were divided into two groups, namely Epinephrine and No epinephrine. Comparisons of continuous variables between the Epinephrine and No epinephrine groups were performed using the Mann–Whitney U test. Categorical variables were expressed as counts and percentages, and comparisons of each categorical variable between groups were performed using the chi-squared test. Outcomes were evaluated using univariate and multiple logistic regression analyses to assess the independent effect of epinephrine administration. Confounders for multiple logistic regression models were selected on the basis of the assumption that these were directly affected by epinephrine administration (according to previous reports) and their clinical importance¹¹. Covariates of the primary outcome included age, gender, first rhythm and time from the call to arrival at the scene of the accident. The presence of a witness was not included in the covariates because this was evaluated in the subgroup analysis. We set two multiple logistic regression models of the secondary outcome. In the first model we selected the covariates of the primary analysis, while in the second model we included age, gender, first rhythm, use of ALS devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident to adjust the differences between two groups more precisely. In addition, multicollinearity was assessed using a variance inflation factor and the tolerance value was set at <2 . Moreover, the goodness-of-fit was assessed using the Hosmer–Lemeshow test. Statistical significance was defined as a two-sided $P < 0.05$ or assessed using a 95% confidence interval (CI) in all statistical analyses. All statistical analyses, excluding the propensity score (PS) matching were performed using the IBM SPSS Statistics version 23.0 (SPSS Inc, Chicago, Illinois, USA).

Propensity score. We chose the PS-matching analysis because prehospital epinephrine administration was not randomly assigned. A logistic regression analysis was performed to estimate a PS for the prediction of prehospital epinephrine administration from the available predictors. Confounders for the PS were selected according to a previous report¹¹. Clinically important confounders were included in the calculation of the PS. These variables were age, gender, presence of a witness, bystander CPR, first rhythm, use of ALS devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident. PS matching extracted 1:1 matched pairs of patients using a caliper with 0.0005 with administration/or not of epinephrine based on the averaged PS. The absolute standardized difference of variables for the estimation of PS was used to assess the match balance, whereby an absolute standardized difference >0.1 represented a meaningful imbalance. In the PS-matched cohort, a univariate logistic regression analysis was performed to evaluate the association between epinephrine administration and outcomes. PS matching was performed using the R software for Windows (version 3.4.0; R Foundation for Statistical Computing, Vienna, Austria).

Subgroup analysis. Subgroup analysis was performed to identify the potential benefits and demerits of prehospital epinephrine administration. Patients with TCA and presence of a witness were selected. In the subgroup analysis, we performed PS matching. The variables used to calculate the PS were age, gender, bystander CPR, first rhythm, use of ALS devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident. In the PS-matched cohort, a univariate logistic regression analysis was performed.

Sensitivity analysis. We performed a sensitivity analysis involving patients with TCA who achieved prehospital ROSC within 10 min from CPR to verify that the results were not altered by the selection of patients. In the sensitivity analysis, we performed PS matching. The variables used to calculate the PS were age, gender, bystander CPR, first rhythm, use of ALS devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident. In the PS-matched cohort, a univariate logistic regression analysis was performed.

Ethics approval and consent to participate. The analysis protocol of this study was approved by the Ethics Committee of the Gunma University Hospital. Because of the anonymous and retrospective nature of the study, the need for informed consent was waived.

Results

A total of 465,932 patients with OHCA were documented from January 1, 2012 to December 31, 2015. Among those, cardiac arrest was presumed to be of traumatic origin caused by traffic collision in 8,111 patients. We identified a total of 5,204 patients with TCA eligible for analysis (Fig. 1). Of those, 758 patients received prehospital epinephrine (Epinephrine group), whereas the remaining 4,446 patients did not receive prehospital epinephrine (No epinephrine group). Patient characteristics are shown in Table 1. The median age of patients was 61 years (40–75 years). Age was significantly higher in the Epinephrine group versus the No epinephrine group (64 [44–77]

Characteristics	Epinephrine (n = 758)	No epinephrine (n = 4,446)	P value
Age, median (IQR), y	64 (44–77)	61 (39–75)	0.004
≤17	19 (2.5%)	211 (4.7%)	0.006
18–64	370 (49%)	2258 (51%)	
≥65	369 (49%)	1977 (45%)	
Gender			0.277
Male	537 (71%)	3062 (69%)	
Female	221 (29%)	1384 (31%)	
Witness			0.001
Unwitnessed	191 (25%)	1393 (31%)	
Witnessed	567 (75%)	3053 (69%)	
Bystander CPR			<0.001
Any CPR	195 (26%)	796 (18%)	
No CPR	563 (74%)	3650 (82%)	
First rhythm			<0.001
Ventricular fibrillation	21 (2.8%)	82 (1.8%)	
Pulseless ventricular tachycardia	2 (0.3%)	9 (0.2%)	
Pulseless electrical activity	341 (45%)	1482 (33%)	
Asystole	354 (47%)	2618 (59%)	
Other	40 (5.3%)	255 (5.7%)	
Life support by emergency medical personnel			
Use of advanced life support devices	469 (62%)	1049 (24%)	<0.001
Insertion of intravenous line	739 (98%)	480 (11%)	<0.001
Time from call to arrival at scene	8 (6–11)	8 (6–10)	<0.001
Time from call to arrival at hospital	38 (30–48)	32 (25–43)	<0.001

Table 1. Baseline characteristics of the patients with traumatic cardiac arrest according to epinephrine administration (n = 5,204). IQR; interquartile range, CPR; cardiopulmonary resuscitation. Missing; Time from call to arrival at scene = 13 and Time from call to arrival at hospital = 13.

years vs. 61 [39–75] years, respectively; $P = 0.004$). The majority of patients were male (69%; 3,599/5,204 patients). The percentages of witness presence and bystander CPR were significantly higher in the Epinephrine group versus the No epinephrine group (witness: 567/758 patients [75%] vs. 3,053/4,446 patients [69%], respectively; $P = 0.001$ and bystander CPR: 195/758 patients (26%) vs. 796/4,446 patients (18%), respectively; $P < 0.001$). Approximately 92% of the patients exhibited a non-shockable first rhythm. There were significantly more non-shockable first rhythms in Epinephrine group than No epinephrine group. Regarding life support provided by EMS personnel, use of ALS devices and insertion of an intravenous line were more frequent in the Epinephrine group (use of ALS devices: 469/758 patients (62%) vs. 1,049/4,446 patients (24%), respectively; $P < 0.001$ and insertion of an intravenous line: 739/758 patients (98%) vs. 480/4,446 patients (11%), respectively; $P < 0.001$). Both times from the call to arrival at the scene of the accident and at the hospital were significantly longer in the Epinephrine group (call to arrival at the scene of the accident: 8 (6–11) min vs. 8 (6–10) min, respectively; $P < 0.001$ and call to arrival at the hospital: 38 (30–48) min vs. 32 (25–43) min, respectively; $P < 0.001$).

Figures 2 and 3 show the comparison of outcomes between the Epinephrine and the No epinephrine groups. There was no significant difference in 1-month survival between the groups (1.5% [11/758 patients] vs. 0.9% [41/4,446 patients], respectively; $P = 0.176$). However, the rate of prehospital ROSC was higher in the Epinephrine group (9.8% [74/758 patients] vs. 0.9% [40/4,446 patients], respectively; $P < 0.001$). The multiple logistic regression analysis showed that prehospital epinephrine administration was not associated with 1-month survival (odds ratio [OR] 1.495, 95% CI 0.758 to 2.946); however, it was associated with prehospital ROSC (OR 3.784, 95% CI 2.102 to 6.812) (Figs 2 and 3).

Table 2 shows the baseline characteristics of PS-matched patients. The characteristics of PS-matched patients were balanced in terms of absolute standardized mean difference (< 0.1 between the groups). In PS-matched patients, there was no significant difference in 1-month survival between the Epinephrine and No epinephrine groups (2.2% [7/317 patients] vs. 0.9% [3/317 patients], respectively; $P = 0.340$). However, the rate of prehospital ROSC was higher in the Epinephrine group (16.7% [53/317 patients] vs. 2.8% [9/317 patients], respectively; $P < 0.001$). Prehospital epinephrine administration was not associated with 1-month survival (OR 2.363, 95% CI 0.606 to 9.223); however, it was associated with prehospital ROSC (OR 6.870, 95% CI 3.326 to 14.192) (Figs 2 and 3).

The subgroup and sensitivity analyses corroborated the results of the main analyses (Supplemental Tables 1 and 2, Figs 2 and 3).

Discussion

Brief summary. This study demonstrated that prehospital epinephrine administration in patients with TCA caused by traffic collisions was not associated with 1-month survival, whereas it was associated with prehospital ROSC.




Figure 2. Effect of prehospital epinephrine administration on 1-month survival. (a) The adjusted confounders included age, gender, first rhythm, and time from the call to arrival at the scene of the accident. (b) The covariates used to estimate the PS were age, gender, witness, bystander CPR, first rhythm, advanced life support devices, insertion of an intravenous line, and time from call to arrival at the scene of the accident. CI, confidence interval; CPR, cardiopulmonary resuscitation; PS, propensity score.




Figure 3. Effect of prehospital epinephrine administration on return of spontaneous circulation. (a) The adjusted confounders included age, gender, first rhythm, and time from the call to arrival at the scene of the accident. (b) The adjusted confounders included age, gender, first rhythm, advanced life support devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident. (c) The covariates used to estimate the PS were age, gender, witness, bystander CPR, first rhythm, advanced life support devices, insertion of an intravenous line, and time from the call to arrival at the scene of the accident. CI, confidence interval; ROSC, return of spontaneous circulation; CPR, cardiopulmonary resuscitation; PS, propensity score.

Comparison with previous studies. The results of the present study are consistent with those of a previous study, demonstrating that prehospital epinephrine administration in patients with TCA is associated with temporary ROSC¹¹. Adrenergic system in the regulation of the cardiovascular system was well studied²²⁻²⁴. In patients with endogenous cardiac arrest, epinephrine administration has been reported to potentially increase the chance of ROSC through the arterioles mediated by α -adrenergic receptors²⁵. This effect augmented temporal coronary blood flow and increased prehospital ROSC in patients with TCA^{14,26}. However, the main cause of TCA was hemorrhagic shock²⁷. A previous observational study reported that use of a vasopressor was associated with mortality in patients with traumatic hemorrhagic shock²⁸. This study showed that the effect of epinephrine was limited and did not reach 1-month survival without definitive hemostatic treatment and/or blood transfusion management for hemorrhagic shock.

Previous article was that half of the patients who survived were penetrating trauma patients and this constituted a major bias¹¹. The present study focused on blunt trauma patients with TCA caused by traffic collisions. In addition, our study analyzed data from a nationwide registry. Therefore, the sample size and generalizability of the findings are major strengths of this study. Besides, there were no biases related to hemostatic treatment because Japanese EMS systems could not do definitive hemostatic treatment.

In this study we used PS matching to control the variables including the time from the call to arrival at the scene of the accident and the time from the call to arrival at the hospital. Regarding clinical study in resuscitation area, the time dependent propensity score is thought to be more robust analysis^{29,30} and we included the time from the call to arrival at scene as one covariate for investigating the true effect of prehospital epinephrine administration for TCA.

Possible explanations and implications. This study assessed the circumscription of prehospital management for patients with TCA in Japan. The mortality of patients with TCA varied depending on the previous reports²⁻⁹. The 1-month survival reported in the present study cohort was approximately 1%. This rate was lower compared with that reported in a recent systematic review³. Treatment options for EMS personnel in other countries extend to insertion of a chest tube, insertion of an intravenous line, and use of ALS devices³¹. Regarding fluid management, the usefulness of prehospital administration of red blood cells and blood plasma has been reported. Moreover, aggressive and varying prehospital management has been related to survival³²⁻³⁵. A previous report demonstrated that ALS performed by a treating physician was associated with survival among patients with TCA¹⁰. Therefore, it is recommended to use aggressive therapeutic options at the prehospital setting to achieve more favorable outcomes in patients with TCA.

Characteristics	Epinephrine (n = 317)	No epinephrine (n = 317)	Standardized difference
Age, median (IQR), y	67 (47–77)	65 (43–77)	
≤17	3 (0.9%)	5 (1.6%)	−0.07
18–64	147 (46%)	152 (48%)	−0.03
≥65	167 (53%)	160 (51%)	0.04
Gender			0.01
Male	230 (73%)	232 (73%)	
Female	87 (27%)	85 (27%)	
Witness			−0.01
Unwitnessed	87 (27%)	89 (28%)	
Witnessed	230 (73%)	228 (72%)	
Bystander CPR			0.02
Any CPR	61 (19%)	64 (20%)	
No CPR	256 (81%)	253 (80%)	
First rhythm			
Ventricular fibrillation	2 (0.6%)	4 (1.3%)	−0.08
Pulseless ventricular tachycardia	1 (0.3%)	2 (0.6%)	−0.06
Pulseless electrical activity	127 (40%)	123 (39%)	0.03
Asystole	177 (56%)	180 (57%)	−0.02
Other	10 (3.2%)	8 (2.5%)	0.04
Life support emergency medical personnel			
Use of advanced life support devices	185 (58%)	188 (59%)	−0.02
Insertion of intravenous line	302 (95%)	302 (95%)	0
Time from call to arrival at scene	8 (6–10)	8 (6–11)	−0.08
Time from call to arrival at hospital	35 (28–44)	34 (27–44)	

Table 2. Baseline characteristics of the propensity score-matched patients with traumatic cardiac arrest according to epinephrine administration (n = 634). IQR; interquartile range, CPR; cardiopulmonary resuscitation.

The authors acknowledge the following limitations of this study. Firstly, as expected, survival from TCA was rare. First, the cause of mortality was not registered in the All-Japan Utstein Registry, therefore, we could not discuss what is the most severe injured organ and what in-hospital treatment was performed. In addition, there may be some medical caused cardiac arrest patients. These were strong bias of this study. Second, the rate of 1-month survival was approximately 1.0%. Considering that the goal of CPR is a favorable neurological outcome, we were unable to set the neurological outcome as an outcome of this study.

Conclusion

Prehospital epinephrine administration in patients with TCA caused by traffic collisions was not associated with 1-month survival, but was beneficial in regards to prehospital ROSC. Currently, options for the prehospital treatment of trauma are limited.

References

- Davidson, G. H. *et al.* Long-term survival of adult trauma patients. *JAMA*. **305**, 1001–1017 (2011).
- Shimazu, S. & Shatney, C. H. Outcomes of trauma patients with no vital signs on hospital admission. *J Trauma*. **23**, 213–216 (1983).
- Zwingmann, J. *et al.* Survival and neurologic outcome after traumatic out-of-hospital cardiopulmonary arrest in a pediatric and adult population: a systematic review. *Crit Care*. **16**, R117 (2012).
- Hopson, L. R. *et al.* Guidelines for withholding or termination of resuscitation in prehospital traumatic cardiopulmonary arrest: joint position statement of the National Association of EMS Physicians and the American College of Surgeons Committee on Trauma. *J Am Coll Surg*. **196**, 106–112 (2003).
- Beck, B. *et al.* Trends in traumatic out-of-hospital cardiac arrest in Perth, Western Australia from 1997 to 2014. *Resuscitation*. **98**, 79–84 (2016).
- Deasy, C. *et al.* Traumatic out-of-hospital cardiac arrests in Melbourne, Australia. *Resuscitation*. **83**, 465–470 (2012).
- Lockey, D., Crewdson, K. & Davies, G. Traumatic cardiac arrest: who are the survivors? *Ann Emerg Med*. **48**, 240–244 (2006).
- Huber-Wagner, S. *et al.* Outcome in 757 severely injured patients with traumatic cardiorespiratory arrest. *Resuscitation*. **75**, 276–285 (2007).
- Leis, C. C. *et al.* Traumatic cardiac arrest: should advanced life support be initiated? *J Trauma Acute Care Surg*. **74**, 634–638 (2013).
- Fukuda, T., Ohashi-Fukuda, N., Kondo, Y., Hayashida, K. & Kukita, I. Association of Prehospital Advanced Life Support by Physician With Survival After Out-of-Hospital Cardiac Arrest With Blunt Trauma Following Traffic Collisions: Japanese Registry-Based Study. *JAMA Surg*. **153**, e180674 (2018).
- Chiang, W. C. *et al.* Prehospital intravenous epinephrine may boost survival of patients with traumatic cardiac arrest: a retrospective cohort study. *Scand J Trauma Resusc Emerg Med*. **23**, 102 (2015).
- Hagihara, A. *et al.* Prehospital epinephrine use and survival among patients with out-of-hospital cardiac arrest. *JAMA*. **307**, 1161–1168 (2012).
- Hansen, M. *et al.* Time to epinephrine administration and survival from nonshockable out-of-hospital cardiac arrest among children and adults. *Circulation*. **137**, 2032–2040 (2018).
- Perkins, G. D. *et al.* A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Eng J Med*. **379**, 711–721 (2018).

15. Monsieurs, K. G. ERC Guidelines 2015 Writing Group, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 1. *Executive summary. Resuscitation* **95**, 1–80 (2015).
16. Locke, D. J., Lyon, R. M. & Davies, G. E. Development of a simple algorithm to guide the effective management of traumatic cardiac arrest. *Resuscitation*. **84**(6), 738–42 (2013).
17. Sherrin, P. B., Reid, C., Habig, K. & Burns, B. J. Algorithm for the resuscitation of traumatic cardiac arrest patients in a physician-staffed helicopter emergency medical service. *Crit Care*. **17**(2), 308 (2013).
18. Cummins, R. O. et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation*. **84**, 960–975 (1991).
19. Jacobs, I. et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Councils of Southern Africa). *Circulation*. **110**, 3385–3397 (2004).
20. Fire and Disaster Management Agency. *Kyukyukyuujo-no genkyo 2014* [Current Situations of Emergency Medical Services and Rescue Services 2014]. Tokyo, Japan: Fire and Disaster Management Agency; 2015.
21. Kitamura, T. et al. Nationwide public access defibrillation in Japan. *N Engl J Med*. **362**, 994–1004 (2010).
22. Santulli, G. et al. G protein-coupled receptor kinase 2 in patients with acute myocardial infarction. *Am J Cardiol*. **15**, 107 (2011).
23. Santulli, G. & Iaccarino, G. Adrenergic signaling in heart failure and cardiovascular aging. *Maturitas* **93**, 65–72 (2016).
24. Ciccarelli, M., Santulli, G., Pascale, V., Trimarco, B. & Iaccarino, G. Adrenergic receptors and metabolism: role in development of cardiovascular disease. *Front Physiol* **4**, 265 (2013).
25. Sorriento, D. et al. Endothelial cells are able to synthesize and release catecholamines both *in vitro* and *in vivo*. *Hypertension*. **60**, 129–136 (2012).
26. Perkins, G. D., Cottrell, P. & Gates, S. Is adrenaline safe and effective as a treatment for out of hospital cardiac arrest? *BMJ*. **348**, g2435 (2014).
27. Kauvar, D. S., Lefering, R. & Wade, C. E. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma*. **60**, S3–11 (2006).
28. Aoki, M., Abe, T., Saitoh, D., Hagiwara, S. & Oshima, K. Use of vasopressor increases the risk of mortality in traumatic hemorrhagic shock: A nationwide cohort study in Japan. *Crit Care Med*. **46**, e1145–e1151 (2018).
29. Nakahara, S. et al. Evaluation of pre-hospital administration of adrenaline (epinephrine) by emergency medical services for patients with out of hospital cardiac arrest in Japan: controlled propensity matched retrospective cohort study. *BMJ*. **347**, f6829 (2013).
30. Izawa, J. et al. Pre-hospital advanced airway management for adults with out-of-hospital cardiac arrest: nationwide cohort study. *BMJ*. **364**, l430 (2019).
31. Dickson, R. L. et al. Emergency medical services simple thoracostomy for traumatic cardiac arrest: Postimplementation experience in a ground-based suburban/rural emergency medical services agency. *J Emerg Med*. **55**, 366–371 (2018).
32. O'Reilly, D. J. et al. Prehospital blood transfusion in the en route management of severe combat trauma: a matched cohort study. *J Trauma Acute Care Surg*. **77**, S114–120 (2014).
33. Shackelford, S. A. et al. Association of prehospital blood product transfusion during medical evacuation of combat casualties in Afghanistan with acute and 30-day survival. *JAMA*. **318**, 1581–1591 (2017).
34. Sperry, J. L. et al. Prehospital plasma during air medical transport in trauma patients at risk for hemorrhagic shock. *N Engl J Med*. **379**, 315–326 (2018).
35. Van Oostendorp, S. E., Tan, E. C. & Geeraedts, L. M. Jr. Prehospital control of life-threatening truncal and junctional haemorrhage is the ultimate challenge in optimizing trauma care; a review of treatment options and their applicability in the civilian trauma setting. *Scand J Trauma Resusc Emerg Med*. **24**, 110 (2016).

Acknowledgements

We thank all of the EMS personnel and physicians in Japan and the Fire and Disaster Management Agency and Institute for Fire Safety and Disaster Preparedness of Japan for their generous cooperation in establishing and maintaining the Utstein database.

Author Contributions

M.A. and T.A. conceived of and designed this study, interpreted the data, drafted the manuscript, and revised the manuscript for important intellectual content. K.O. interpreted the data and revised the manuscript for important intellectual content.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-019-46460-w>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019