

SFRP1 shapes astrocyte to microglia cross-talk in acute and chronic neuroinflammation

Javier Rueda-Carrasco^{1,2}, María Inés Mateo^{1,2}, Aldo Borroto¹, María Jesús Martín-Bermejo^{1,2}, Markus P. Kummer^{3,4}, Stephanie Schwartz^{3,4}; José P. López-Atalaya⁵; Balbino Alarcon¹, Michael T. Heneka^{3,4}, Pilar Esteve^{1,2} and Paola Bovolenta^{1,2*}

¹Centro de Biología Molecular Severo Ochoa, CSIC-UAM, c/ Nicolás Cabrera, 1, Campus de la Universidad Autónoma de Madrid, Madrid 28049, Spain; ²CIBER de Enfermedades Raras (CIBERER) Madrid 28049, Spain; ³Universitätsklinikum Bonn, Neurology; ⁴German Center for Neurodegenerative Diseases (DZNE) 53127 Bonn, Germany. ⁵Instituto de Neurociencias, CSIC-UMH, Sant Joan d'Alacant, Spain

Running title: SFRP1 enhances microglial activation

Key words: Neuroinflammation; microglia, HIF pathway; LPS; EAE; reactive astrocytes; Multiple Sclerosis; Alzheimer's disease

* corresponding author at pbovolenta@cbm.csic.es

Abstract

Neuroinflammation is a common feature of many neurodegenerative diseases, which fosters a dysfunctional neuron-microglia-astrocyte crosstalk that, in turn, maintains microglial cells into a perniciously reactive state that often enhance neuronal damage. The molecular components that mediate this critical communication are however not fully explored. Here, we have asked whether Secreted-Frizzled-Related-Protein-1 (SFRP1), a multifunctional regulator of cell-to-cell communication, is part of the cellular crosstalk underlying neuroinflammation. We show that in mouse models of acute and chronic neuroinflammation, astrocyte-derived SFRP1 is sufficient to promote and sustain microglial activation, and thus a chronic inflammatory state. SFRP1 allows the upregulation of components of Hypoxia Induced Factors-dependent inflammatory pathway and, to a lower extent, of those downstream of the Nuclear Factor-kappaB. We thus propose that SFRP1 acts as a critical astrocyte to microglia amplifier of neuroinflammation, representing a potential valuable therapeutic target for counteracting the harmful effect of chronic inflammation present in several neurodegenerative diseases.

Introduction

Degeneration of neurons occurs in a variety of rare and common pathological conditions of the CNS including, for example, Alzheimer's disease (AD) or Multiple Sclerosis (MS). CNS function is supported by a robust neurons-glia crosstalk [1,2], so that neuronal damage is almost invariably associated with the activation of two types of glial cells: microglia and astrocytes. The prompt glial response to insults brings about an inflammatory reaction that favors tissue healing and helps restoring CNS homeostasis [3,4]. However, excessive glial activation, as often occurs in neurodegeneration, is itself a cause of neuronal loss, which, in turn, establishes a state of pernicious chronic neuroinflammation [4–7]. Consistent with an important cellular crosstalk, glial cell dysfunction, either as a consequence of hyper- or hypo-functionality, can also be the cause of neuronal damage, rather than its consequence, strongly contributing to the progression of neurodegenerative diseases [3,4,8,9]. Although this is nowadays a widely accepted idea, there is still only partial information on the molecular components that sustain a dys- or hyper-functional state of glial cells and an abnormal glia/neuron crosstalk. Here we have investigated whether SFRP1 may represent one of such components.

SFRP1 is a small, secreted and dispersible protein with a dual function: modulation of Wnt signaling [10] and inhibition of ADAM10 activity [11]. The latter function is particularly relevant in the context of neurodegeneration because ADAM10, a member of the A Disintegrin and Metalloprotease family of plasma-membrane proteins [12], acts as an α -secretase (or sheddase; [13] for many substrates expressed in both neurons and glial cells. In microglial cells, ADAM10-mediated shedding of TREM2 (Triggering Receptor Expressed on Myeloid) is a relevant mechanism for controlling their activation [14]. In neurons, ADAM10 activity releases the ectodomain of CD200 and CX₃CL1 [15,16], two components of the mechanism by which neurons maintain microglial cells in a surveying state [2]. Consistent with these roles and the additional regulation of proteins involved in synapse formation [12,17], genetic inactivation of *Adam10* in mice causes neuroinflammation and loss of synaptic plasticity [18]. Interest in ADAM10 is further increased by genetic studies supporting a link between impaired ADAM10 activity and AD [19,20]. Thus, and based on its non-amyloidogenic processing of the Amyloid Precursor Protein (APP) [21], ADAM10 may have a protective role against AD [22]. Consistent with this possibility, we have recently shown that elevated levels of SFRP1, its endogenous negative modulator, contribute to AD pathogenesis [23]. SFRP1 upregulation correlates with poor ADAM10-mediated processing

of APP and of the synaptic protein N-Cadherin, whereas neutralization of its activity prevents the appearance of AD pathological traits, including glial cell activation [23].

Sfrp1 is abundantly expressed in mammalian radial glial progenitors of the developing CNS [11,24,25] but is largely down-regulated in the adult brain with the exception of restricted neurogenic areas [25–27]. Besides in human neurodegenerative diseases [23,28–30], in which the mRNA localizes to glial cells [23,27], SFRP1 upregulated expression has been reported in several other inflammatory conditions such as periodontitis, rheumatoid arthritis, uropathies or pulmonary emphysema [10,31]. An upregulation has been observed also in the aged brain [29], in which a low-grade chronic inflammation is present [32]. Notwithstanding, whether SFRP1 is directly involved in the modulation of neuroinflammation remains unexplored.

By addressing this issue, here we show that *Sfrp1* is a novel mediator of the astrocyte to microglia crosstalk that underlies mammalian CNS inflammation. In mice, astrocyte-derived SFRP1 is sufficient to activate microglial cells and to amplify their response to distinct acute and chronic neuroinflammatory challenges, sustaining their chronic activation. From a molecular point of view, SFRP1 allows for the full expression of down-stream targets of the transcription factors hypoxia induced factors (HIF) and, to a lesser extent, nuclear factor-kappa B (NF- κ B), which are mediators of neuroinflammatory responses [33,34]. Thus, neutralizing SFRP1 function may represent a strategy to counteract pernicious chronic neuroinflammation that contributes to many human neurodegenerative conditions.

Results

Acute brain neuroinflammation elevates astrocyte-specific levels of SFRP1 expression

To evaluate a possible link between SFRP1 and neuroinflammation, we induced acute brain inflammation by injecting bacterial lipopolysaccharides (LPS; [35] or control saline into the somatosensory cortex of 3 months-old *Sfrp1*^{+/ β gal}; *CX₃CR1*^{+/*GFP*} mice (n=4), which allow the simultaneous identification of microglial (GFP; [36] and *Sfrp1* producing cells (nuclear β gal-positive; [37]. Immunofluorescence of the brains three days after injection, when inflammation is at its peak [38], revealed a broader β gal immunoreactivity (reporter of *Sfrp1* expression) in LPS vs saline treated animals (Fig. 1A), largely localized in GFAP⁺ astrocytes but not in Iba1⁺ microglial cells (Fig. 1A; S1A). Immunodetection of SFRP1 with specific antibodies [23] confirmed an increased SFRP1 production after LPS compared to saline injections (Fig. 1B). Notably, SFRP1 protein distribution in saline injected animals was comparable to that of non-injected mouse brains [23]. Consistent with the secreted and dispersible nature of SFRP1 [39,40], immuno-signal was widely distributed in the brain parenchyma (arrows, Fig. 1B). The use of a highly specific ELISA [23] further confirmed a significant increase of SFRP1 levels in extracts of small cortical tissue samples from LPS treated animals (Fig. 1C) as compared to that present in equivalent brain regions from non-injected or saline injected animals (Fig 1C).

Together these data show that astrocytes produce and secrete increased level of SFRP1 in response to a bacterial lipopolysaccharide.

In vivo Sfrp1 gene addition is sufficient to trigger and sustain glial cell activation

We next reasoned that if SFRP1 is indeed associated with neuroinflammation, its forced expression should be sufficient to activate glial cells. To test this possibility, we infused lentiviral particles (LV) containing *Sfrp1*-IRES-*Gfp* or control IRES-*Gfp* into the lateral ventricle of 10 weeks-old wt mice (n=8; Fig. 2). As expected by the injection site, GFP⁺ (used to determine infection efficiency) LV-transduced cells were largely found lining along the wall of the lateral ventricle, in the choroid plexus and, to a less extent, along the rostral migratory stream (Fig. S1B). Immunohistochemistry of cortical sections 1 month after injection showed a significantly higher presence of SFRP1 protein at the infected site (Fig 2A) associated with GFAP⁺ reactive astrocytes and Iba1⁺ microglial cells as compared to LV-IRES-*Gfp* control animals (Fig. 2A,B; quantifications performed in 3 of the injected animals). Iba1 immunoreactivity was accumulated around the injection site (Fig. 2A,B) and many cells

had a round amoeboid morphology and a significantly higher CD45 expression (Fig. 2B; S1C), two characteristics of reactive microglia [41]. CD45^{hi} round cells could also represent blood-borne infiltrating macrophages (Fig. S1C), although the molecular and functional difference between the two cell types is currently a matter of debate [42]. A wider distribution of hyper-phosphorylated Tau, which often appears as a brain response to inflammation and degeneration [43], was also detected in LV-*Sfrp1*-IRES-*Gfp* transduced vs control brains with a significantly wider distribution (Fig. 2A,B). Analysis of a different set of LV-transduced animals (n=4) 5 months after LV delivery showed a persistent microglial activation in SFRP1- vs GFP-treated animals, with more CD45⁺ cells than those observed at 1-month post-injection (Fig. S1C). Furthermore, several CD45^{hi}-round-shaped microglia/macrophages were detected in the parenchyma, especially in the proximity of transduced cells (white arrowheads in Fig. S1C).

Together these results indicate that SFRP1 is sufficient to trigger an inflammatory response of glial cells, which persists for prolonged periods of time.

Sfrp1 is required for amplifying CNS inflammatory response

Given that SFRP1 was sufficient to induce glial cell activation, we next asked whether it was also a necessary component of the CNS inflammatory response. To this end, we took advantage of *Sfrp1*^{-/-} mice, in which SFRP1 protein is completely undetectable [23] (Fig 1C). These mice have a slightly shorter and thicker cortex that however does not affect their life span, reproduction rate or cognitive and motor behavior and present no evident neuronal defects [23,25]. Furthermore, their content of astrocytes and microglial cells was undistinguishable from that of wt mice (Fig. 2C-F; Fig S1D). We thus compared the effect of intra-cortical LPS infusion into the brains of 3-months-old wt and *Sfrp1*^{-/-} mice (Fig. 2C,D). In the somatosensory cortex of wt brains (Fig. S1D), LPS but not saline treatment caused the appearance of GFAP⁺ reactive astrocytes (Fig. 2C) and CD45⁺ reactive microglia (Fig. 2D). In contrast, *Sfrp1*^{-/-} littermates presented fewer and less immuno-positive astrocytes and a significant reduction of CD45⁺ reactive microglia/macrophages (Fig. 2C,D), suggesting that SFRP1 is relevant for the activation of both astrocytes and microglial cells. Quantitation of immunoreactivity among different genotypes and treatments confirmed no significant differences between the two saline-treated genotypes but showed a significantly lower response of *Sfrp1*^{-/-} mice to LPS as compared to wt (Fig. 2E,F). Notably however, *Sfrp1*^{-/-} mice do not completely lose their respond to LPS, given that this is significantly different

from that observed in saline injected mice (Fig. 2E,F). This suggests that SFRP1 is involved in boosting inflammation.

To evaluate a possible specificity of this response, we induced Experimental Autoimmune Encephalomyelitis (EAE) in wt and *Sfrp1*^{-/-} mice. This is a widely used experimental model for MS, a human inflammatory demyelinating disease. In EAE, CNS inflammation and gliosis occur as a consequence of a strong autoimmune response against the peripheral exposure to myelin components [44], thus representing a neuroinflammatory paradigm quite different from the direct LPS intra-cerebral infusion. Female littermates of the two genotypes were immunized following a standard protocol and animals were scored for the development/remission of their clinical symptoms over the course of a month [45]. Animals were classified with a standard 0 to 5 rank based on their paralysis degree, with 0 corresponding to absence of symptoms and 5 to a moribund condition [45]. In wt mice (n=19), tail limping -the first symptom of the disease- became apparent around 8 days after immunization with a subsequent rapid progression, so that, by 16 days, most of the wt animals presented hind limb paralysis followed by a slow recovery (Fig. 3A). Notably, 47% of the immunized wt mice developed extreme and protracted symptoms (Fig 3A,B). The response of *Sfrp1*^{-/-} mice (n=19) to immunization was instead slower and milder: only 16% of them developed extreme symptoms and their recovery was significantly faster (Fig. 3A,B). Immunostaining of spinal cord sections before immunization showed no differences between wt and *Sfrp1*^{-/-} in astrocytes, microglia or myelin distribution (not shown). In contrast, in animals (n=4) sacrificed 16 days after immunization, there was a significant reduction of pathological signs in *Sfrp1*^{-/-} vs wt mice, including infiltration of CD4⁺ lymphocytes, presence of Iba1⁺ macrophages/activated microglial cells and GFAP⁺ reactive astrocytes, pial surface disruption and loss of MBP⁺ myelin (Fig. 3C,D).

All in all, these data indicate that SFRP1 is commonly required for a robust neuroinflammatory response, likely mediating astrocyte to microglial crosstalk.

Astrocyte-derived Sfrp1 is required for a robust response of microglial cells to damage

The latter possibility found support in SFRP1 localization around microglial cells (Fig. 1B) and in the remarkably poor microglial activation (Fig. 2; Fig. 3) and myeloid cell recruitment (Fig. 3C) observed in the absence of *Sfrp1*. To further explore this possibility we used flow cytometry analysis to determine the proportion of the different CD11b⁺ myeloid populations [46] in the cortex of *CX3CRI*^{+/GFP} and *CX3CRI*^{+/GFP};*Sfrp1*^{-/-} mice 3 days after intra-cerebroventricular LPS or saline administration (Fig. 4A; Fig S2A). In the absence of *Sfrp1*, there

was a significant decrease of LPS-induced infiltration of CD11b⁺/CD45⁺/GFP⁻ monocytes and of the proportion of microglial cells that passed from a CD11b⁺/CD45^{lo}/GFP⁺ surveying to a CD11b⁺/CD45⁺/GFP⁺ activated state (Fig. 4A,B). Furthermore, FACS analysis of isolated GFP-positive microglial cells (Fig. S3A) revealed that, upon infusion of pHrodo *E.coli* bioparticles [47] into the third ventricle, microglial cells derived from *CX3CRI^{+/GFP};Sfrp1^{-/-}* mouse brains showed less cumulative phagocytized fluorescent signal than those isolated from *CX3CRI^{+/GFP}* brains at both 40 and 65 hr post infusion (Fig. 4C; Fig. S3). Notably, at 40 hr the percentage of phagocytizing microglial cells in *CX3CRI^{+/GFP};Sfrp1^{-/-}* mice was significantly higher than in controls (Fig. 4C) but this proportion underwent a drastic reduction at 65 hr (Fig. 4C). These data suggest that in SFRP1 absence microglial cells present a less efficient phagocytizing activity that, at earlier time points, might be compensated by increasing cell recruitment. A differential rate of phagocytosis vs degradation or the acidity of the lysosomal compartment [48] may explain the time course of phagocytosis observed in mutant microglia.

Collectively, these data support the contention that SFRP1, mainly derived from astrocytes, modulates microglial activation in response to an inflammatory challenge and favors their efficient function. To assess that indeed there is an astrocyte-to-microglial flux of information, we next established microglial or astrocyte/microglia mixed cultures (1:1 ratio) from wt and *Sfrp1^{-/-}* neonatal mice and used a multiplex ELISA to determine their cytokines' release (IFN γ , TNF α , IL1- β , IL4, IL6 and IL10) in the culture medium, as a measure of their LPS-induced activation (Heneka et al., 2014). At 24 hr, LPS but not saline treatment enhanced the accumulation of all tested cytokines in the culture media of purified microglial cultures with no significant differences between genotypes (Fig. 4D). In astrocytes/microglia cultures derived from *Sfrp1^{-/-}* LPS-induced cytokines' release was instead significantly reduced, as compared to that detected in similar wt-derived cultures (Fig. 4D). ELISA determination confirmed the presence of SFRP1 only in culture media of astrocytes/microglia mixed cultures derived from wt pups (Fig. S2B), strongly supporting that astrocyte-derived SFRP1 enhances microglial activation.

Sfrp1 allows for the full expression of down-stream targets of HIF transcription factors

We next reasoned that if SFRP1 enhances microglial activation, its absence should modify their transcriptomic profiling towards a less activated or surveying state. To test this possibility, we used fluorescence-associated cell sorting to purify GFP⁺ microglial cells from the brain of *CX3CRI^{+/GFP}* and *CX3CRI^{+/GFP};Sfrp1^{-/-}* mice (n=4). Cells were purified three

days after the injection of LPS or saline and used to obtain the corresponding transcriptomic profiles. We initially compared the gene-expression signature of *CX3CR1*^{+GFP} and *CX3CR1*^{+GFP};*Sfrp1*^{-/-}-derived microglial cells in response to LPS (Fig. 5). Principal-component analysis demonstrated LPS as the main source of variation (72% variance), whereas only 6% of the total variations could be attributed to the genotype after secondary-component analysis (Fig. 5A, Table S1A-D), well in line with the observation that SFRP1 is poorly expressed in the adult brain under homeostatic conditions (Fig. 1; Esteve et al., 2019b). The relatively central position of LPS-treated *Sfrp1*^{-/-} microglia supported their milder inflammatory response (Fig. 2, 3), which was further confirmed when overall gene expression variations of LPS-treated control and *Sfrp1*^{-/-} microglia were plotted and compared (Fig. 5B,C). Linear regression analysis of this comparison demonstrated a 30% reduction of the global transcriptomic response induced by LPS in the absence of *Sfrp1* (slope of red dotted line, Fig. 5D). LPS treatment in control microglia induced the upregulation of 1128 genes, 487 of which were shared with *Sfrp1*^{-/-} microglia (Table S1A-B). LPS treatment also induced the upregulation of 121 and the down-regulation of 167 microglial genes in the absence of SFRP1 (Fig. 5E; Table S1A-D). Genes associated with metabolic pathways (i.e. *AldoA*, *Ldha* or *Pygl*), the cell cycle (i.e. *Gas6*) or immune regulators (i.e. *Mif*, *Mefv*) were highly upregulated in controls but not in *Sfrp1*^{-/-} microglia (Table S1A-D), whereas genes downstream of the Toll-like 4 receptor (TLR4), such as *Tlr2*, were upregulated at similar levels in both genotypes (Table S1A-D). TLR4 is fundamental for LPS recognition and the activation of the immediate inflammatory response (Ransohoff and Perry, 2009), strongly suggesting that, in the absence of *Sfrp1*, microglial cells retain their prompt response to damage, as also shown by their cytokines' release upon LPS treatment (Fig. 4D).

Consistent with this observation, hierarchical clustering of the samples demonstrated treatment-dependent similarities, but with *Sfrp1*^{-/-} microglia showing an attenuated response to LPS treatment as compared to wt (Fig. 5F). Hierarchical clustering of the differentially expressed genes presented a pattern of covariance that was further analysed by Z-score covariance unsupervised clustering (Fig. 5F). This analysis generated 4 different clusters of upregulated genes, which were analysed for enrichment of regulatory elements and gene ontology annotations (Fig. S4). Genes regulated by E2F transcription factors and implicated in cell cycle regulation composed the less abundant 3 and 4 clusters. Genes belonging to cluster 3, involved in the regulation of chromosomal segregation and spindle organization, were slightly more upregulated in control microglia. On the contrary, genes belonging to

cluster 4 and involved in chromatin assembly and DNA packing were upregulated in *Sfrp1*^{-/-} microglia, suggesting possible differences in the length of the cell cycle (Fig. S4).

Cluster 1 and 2 were the largest clusters and included genes that mediate the inflammatory response and regulators of the defense response (Fig. S4A,B). Both clusters were strongly upregulated in control microglia in response to LPS (Fig. S4A): genes in cluster 1 showed significant enrichment in NF-κB TF binding sites at the promoter regions, whereas those putatively controlled by Hif transcription factors were enriched in cluster 2 (Fig. S4C; Table S1E,F). In microglia derived from *Sfrp1*^{-/-} mice, the expression of a few genes belonging to cluster 1 was decreased (Fig. S5A; Table S6), whereas that of genes belonging to cluster 2 was basically abrogated (Fig. S4A; Table S1F). Ingenuity Pathway Analysis network representation of the NF-κB and HIF downstream targets showed that the absence of SFRP1 has a greater effect on downstream targets of HIF (Fig. 6A). Genes downstream NF-κB, such as *Tlr2* and *ApoE* showed LPS-dependent up-regulation with levels almost similar in the two genotypes, whereas the expression of the homeostatic *Cx3cr1* and *Trem2* genes was unchanged (Fig. 6B; Table S1E). Of note, a number of genes shared between the NF-κB and HIF pathways, such as *Cxcl2*, *Cxcl3* and *Ptgs2*, were strongly up-regulated in LPS-treated control microglia but not in those derived from *Sfrp1*^{-/-} brains (Fig. 6A). Similarly, other LPS-induced HIF targets failed to be up-regulated in the absence of *Sfrp1* and maintained levels of expression comparable to those of saline-treated control microglia (Fig. 6A,B; Table S1F). These included *Vegfa*, *Sod2*, *Mif*, *Aldoa* and *Ldha*. Notably, *Hif1α* expression was similar to that observed in LPS treated controls (Table S1F), possibly in line with the notion that Hif1 levels and activity are largely modulated by post-translational modifications and proteolysis [49].

All in all, these results support the idea that in the absence of *Sfrp1* microglial cells can sense and respond to an inflammatory insult undertaking an initial response. SFRP1 is however required to enhance this response allowing for the full activation of microglial-mediated inflammation in the brain.

Discussion

Microglial cells transit from a surveying to an activated state in response to adverse signals derived from the surrounding environment [50]. However, both insufficient or prolonged microglia activation is harmful to the brain so that an elaborated neuron-microglia-astrocyte cross talk is in charge of shaping the brain immune response to damage [51]. The molecular components that mediate the flux of information from neurons to microglia (i.e. the CD200/CD200R signaling system) [50] or from microglia to astrocytes (i.e. cytokines and NO) [1] have been in part identified. In contrast, there is perhaps less information on how astrocytes communicate with microglial cells [1]. Our study unveils that SFRP1 is part of the molecular signals that astrocytes provide to microglial cells to enhance their inflammatory response. In response to damage, reactive astrocytes produce and secrete SFRP1, which, in turn, increases the number of activated microglial cells and fosters the expression of HIF and, to a much lower extent, NF- κ B down-stream targets in microglial cells. SFRP1 persistency is however pernicious as it sustains a chronic inflammatory state. Indeed, its inactivation significantly reduces the prolonged neuroinflammation associated with EAE. These observations indicate that SFRP1 is a potential valuable target to counteract the harmful effect of prolonged inflammatory conditions, such as those present in MS or AD.

SFRP1 function has been studied in many developmental contexts largely linked to the control of cell specification, proliferation and differentiation [52] but its homeostatic role in adult tissues is less clear. Its poor or absent expression (or that of other SFRP family members) due to promoter hypermethylation is considered a sign of bad prognosis in different types of tumor [10,53,54], although *Sfrp1*^{-/-} mice are viable, fertile and do not form tumor spontaneously [55], indicating that SFRP1 absence per se is not tumorigenic. On the other hand, our finding unveils that elevation of SFRP1 levels has also pathological consequences but related to neuroinflammation (this study) and neurodegeneration [23]. As an advantage, SFRP1 neutralization in AD-like mice counteracts disease pathology without any apparent side effect [23]. In the specific case of neuroinflammation, neutralization of SFRP1 function should have less effect on the initial acute inflammatory phase, which is a necessary step towards pathogen elimination, tissue repair and homeostasis restoration [51]. Abrogation of SFRP1 function does not prevent this acute response given that, in its absence, both LPS-infusion and MOG-immunization induce an initial inflammatory reaction, albeit at somewhat lower levels. Furthermore, activated microglial cells and infiltrated monocytes can still be isolated from the brain of *Sfrp1*^{-/-} mice, although at significantly decreased numbers. These

microglial cells retain phagocytic competence, although with a significant less efficiency and no significant difference was observed in the LPS-induced cytokines release from purified microglial cells obtained from *Sfrp1*^{-/-} and wt brains. This agrees with the observation that in homeostatic conditions SFRP1 is very poorly expressed in the adult mammalian brain [23] and this study). Our transcriptomic analysis shows that *Sfrp1*^{-/-} microglia cells retain their capacity of sensing LPS, providing strong support to the implication of SFRP1 in the amplification of the inflammatory response. Indeed, *Tlr2* expression, a read-out of LPS-mediated TLR4 activation [38], is increased (and to a comparable extent) in microglial cells from both genotypes. A similar consideration applies to a large fraction of the genes related to the NFκB-dependent pathway, which is largely related to the acute inflammatory response [34].

Brain response to LPS, but also to other harmful signals, depends on the fast reaction of functional microglial cells [56], whereas astrocytes and neurons do not initially and directly participate in this response despite their reported expression of the TLR4 receptor [57,58]. Reactive microglial cells thereafter induce astrocytes activation that, in turn, feeds back on microglial function [1]. Fitting our data into this loop, we propose that up-regulation and release of SFRP1 from astrocytes occurs as part of their microglia-mediated early activation. Secreted SFRP1 then acts on microglial cells amplifying a HIF-dependent inflammatory response and thus a significant increase in cytokine secretion, further impinging upon inflammation. Supporting SFRP1 implication in this astrocyte-microglia loop, astrocytes *Sfrp1*^{-/-} mice show decreased GFAP expression and thus are also less reactive, likely producing lower levels of other microglia-activating molecules. This implies that, as long as SFRP1 up-regulation persists, neuroinflammation persists, contributing to its chronicization. This idea is well in agreement with our observation that genetic inactivation of *Sfrp1* significantly limits the severity and progression of EAE, a condition that, as MS, is characterized by persistent microglial activation. Both EAE and MS are driven by the infiltration of peripheral macrophages and lymphocytes [59]. We cannot exclude that the lack of peripherally expressed *Sfrp1* may limit this infiltration and thus the disease, especially because SFRP1 has been shown to influence lymphocytes' differentiation [60]. Notwithstanding there is strong evidence that endogenous microglial activation is critical for EAE onset and maintenance [61,62] and the *SFRP1* gene is hypomethylated in brain samples from MS patients [63], likely promoting an abnormal protein increase. Thus, reduced microglial activation is a plausible cause of the milder EAE symptoms observed in *Sfrp1*^{-/-} mice.

SFRP1 up-regulation in other neurodegenerative diseases characterized by the presence of chronic inflammation further supports a role of SFRP1 microglial activation. An example is Glaucoma [64] or AD [23,30], in which antibody-mediated neutralization of SFRP1 strongly decreases different AD pathological features, including neuroinflammation [23]. Notably, SFRP1-mediated inhibition of ADAM10, a metalloprotease responsible for non-amyloidogenic processing of APP, contributes to the generation of toxic amyloid peptides in AD patients, whereas SFRP1 deficiency decreases amyloid burden [23]. Low neuroinflammation upon SFRP1 neutralization could therefore be secondary to this reduction [23]. The present study shows that this is not necessarily the case as SFRP1 acts directly on microglial cells, suggesting that SFRP1 simultaneously impinges upon multiple pathological events in AD. A possible SFRP1-mediated activation of microglial cells in AD finds further support in our transcriptomic analysis that links SFRP1 with HIF signaling. Indeed, many LPS-induced genes regulated or co-regulated by Hif were not activated in the absence of *Sfrp1*, although *Hif1 α* expression was not modified, possibly because its levels are largely post-translationally regulated [49]. Consistent with our observations, a number of recent studies have shown that microglial cells isolated from AD-like mouse models undergo important metabolic changes with the activation of the HIF pathway [65–67]. Furthermore, *Hif1 α* is involved in astrocytes proinflammatory activity [68] and its activity seems to be dysregulated in association with other genes genetically linked to AD risk, suggesting that HIF-1 α may be detrimental in AD pathology [66].

The precise molecular interactions underlying SFRP1-mediated HIF pathway activation are at the moment an open question. Microglial cells express members of the ADAM family of metalloproteases [14], including ADAM10 which has been shown to be upregulated in both neurons and microglial cells during microglial mediated synaptic pruning [69]. Furthermore, SFRP1 effectively inhibits ADAM enzymatic activity in cultured microglial cells (our own observations). Given that TREM2 is a proven substrate of ADAM10/17 [14], it seems plausible to postulate that SFRP1 may modulate the shedding of TREM2 on the surface of microglial cells. This shedding occurs, for example, in response to LPS induced activation of TLR4 [70,71], thereby attenuating microglial activation [70]. Notably, microglial cells deficient in TREM2 undergo only a partial and abortive activation and remain locked in an almost homeostatic state [72,73]. It is thus tempting to speculate that in SFRP1 absence enhanced shedding of TREM2 may reduce microglial activation as we have observed in *Sfrp1*^{-/-} mice, whereas high SFRP1 levels, by interfering with ADAM function, may enhance TREM2 signaling. In addition (or alternatively), SFRP1 may interfere with ADAM10-

mediated shedding of the neuronal ligands CX3CL1 and CD200 preventing the generation of their soluble forms [15,16] and thus the activation of their respective microglial receptors, CX3CR1 and CD200R [36,74]. Unfortunately, lack of appropriate biochemical tools has prevented us from verifying these possibilities in our mouse models, leaving this question open for future studies.

An alternative and non-mutually exclusive possibility involves SFRP1 function as a Wnt signaling modulator. The role of this pathway in microglial activation is however somewhat controversial, perhaps in line with the notion that Wnt pathway activity is highly context dependent. For example, studies *in vitro* have shown that Wnt3a and Wnt5a can counteract LPS-induced microglia activation [75]. In contrast, in preterm-born infants and in different postnatal animal models of injured-induced neuroinflammation, downregulation of the expression of Wnt signaling components seems a prerequisite for pro-inflammatory activation of microglia [76]. SFRP1 upregulation could lead to the same net effect, inducing at least the down-regulation of the expression of *Axin2* or *Lef1*, genes that targets and read-out of Wnt/ β catenin pathway activation. Notably however, we found no evidence of changes in the expression of these genes in our RNaseq analysis, making us favor the possibility that SFRP1 acts through ADAM10.

In conclusion, we have shown that astrocyte-derived SFRP1 plays an important role in shaping microglia response to CNS damage, sustaining chronic neuroinflammation. This effect might not be limited to the brain, as SFRP1 upregulation has been reported in different pathological conditions associated with inflammation or fibrosis such as periodontitis, rheumatoid arthritis, uropathies or pulmonary emphysema [10,31]. Therefore, once analyzed the possible side effect, SFRP1 neutralization [23] may represent a promising therapeutic avenue to treat a wide variety of chronic pathological conditions.

Material and Methods

Animals. Newborn and adult mice of both sexes were used, unless otherwise indicated. Mice were maintained in pathogen-free conditions at the CBMSO animal facilities, following current national and European guidelines (Directive 2010/63/EU). Experimental procedures were approved by the CBMSO and Comunidad Autónoma de Madrid ethical committees. *Sfrp1*^{-/-} mice were generated by inter-cross of the *Sfrp1*^{-/-};*Sfrp2*^{+/-} mice in a mixed 129/C57BL/6 background as described [11] and then back-crossed at least 4X with C57BL/6J to unify the background. Wild type (wt) animals were littermates selected from heterozygous crosses. Breeding pairs of *CX₃CR1::GFP* mice [36] were kindly provided by Prof. J Avila, CBMSO. Mice were further crossed with the *Sfrp1*^{-/-} to obtain *CX₃CR1::GFP*;*Sfrp1*^{-/-}.

Stereotactic LPS infusion. LPS or saline were infused into the brain parenchyma of 10-12 weeks-old littermates from wt and *Sfrp1*^{-/-} or *CX₃CR1*^{+/*GFP*} and *CX₃CR1*^{+/*GFP*};*Sfrp1*^{-/-} mice. Animals were anaesthetized with 4% Isoflurane (Forane, AbbVie Farmacéutica) vaporized into a sealed anesthetic induction chamber (SurgiVet, Smiths Medical) and placed in a stereotaxic apparatus (Stoelting). Anesthesia was maintained at 2.5% in 250ml/min oxygen flow. Saline (2.5µl) or LPS (5µg; Escherichia coli 0111:B4; Sigma Aldrich) were delivered through a small skull window using a Quintessential Stereotaxic Injector (Stoelting) coupled to a 10µl syringe with a 34G needle (Hamilton) at the rate of 0.5µl/min using the following bregma coordinates: 0.0mm A-P; -1.0 mm lateral, and -1.5mm D/V. Mice were let survive for three days and then sacrificed and processed for further analysis. Delivery of pHrodo Red *E.coli* BioParticles Conjugate (Molecular Probes) was performed with a similar procedure at -2.5 mm A-P; 0.0 mm lateral, and -2.3 mm D-V from bregma. Lentiviral vectors (2.5µl), generated as described below, were delivered 0.5 mm A-P; 1.0 mm lateral and -2.3 mm D-V from bregma. Mice were let survive 1-5 months and then analyzed.

Lentiviral (LV) particle generation. LV particles carrying IRES-*Gfp* or *Sfrp1*-IRES-*Gfp* were obtained by transient transfection of mycoplasma-free HEK-293T cells [77,78]. Cells were transfected employing a three-plasmids HIV-derived and VSV pseudotyped LV system kindly provided by M.K. Collins, University College London, UK; A. Thrasher, Institute of Child Health, UK; and D. Trono, Ecole Polytechnique Fédérale de Lausanne, Switzerland. Culture supernatants were collected two and three days after transfection and ultra-centrifuged. Pellets containing the concentrated particles were resuspended in PBS. Functional titer of the viral

preparations was determined by transfection of HEK-293T cells (1×10^8 TU/ml, in both cases).

Experimental Autoimmune Encephalomyelitis (EAE). Chronic EAE was induced as described [45]. Briefly, 8-10 weeks-old female C57BL/6J and *Sfrp1*^{-/-} littermates were injected bilaterally in subcutaneous femoral region with 150 mg of MOG35–55 (Espikem) emulsified with Freund's complete adjuvant (Sigma Aldrich), supplemented with Mycobacterium tuberculosis (1mg/ml; H37Ra strain from Difco), followed by two intraperitoneal injections of pertussis toxin (200 ng; Sigma Aldrich) separated by 48h. An observer blind to the animals' genotype weighed and inspected the animals daily to detect the appearance of clinical signs according to the following classification: 0) no overt signs of disease; 1) weakness at the tail distal portion; 1.5) complete tail flaccidity; 2) moderate hind limb weakness; 2.5) severe hind limb weakness; 3) ataxia; 3.5) partial hind limb paralysis; 4) complete hind limb paralysis; 4.5) complete hind limb paralysis with muscle stiffness; 5) moribund state and hence sacrificed according to ethical procedures. A representative pool of mice was anesthetized and perfused intracardially sixteen days after immunization, when symptoms picked, for histological analysis. Other animals were let recover. Spinal cords were dissected and processed.

Primary cultures. Glial primary cultures were established from cerebral cortices of C57BL/6J or *Sfrp1*^{-/-} 1-3 days-old pups dissected in Ca²⁺/Mg²⁺-free Hank's Balanced Salt Solution (HBSS, Invitrogen) and processed following standard procedures [79]. Cells were plated in Dulbecco's modified Eagle medium and F-12 nutrient mixture (DMEM/F12, Invitrogen) containing 10% FCS (Invitrogen), and gentamycin (Sigma Aldrich). Cortices from two pups were plated in a 75cm² flask pre-treated with Poly-D-Lys (P7280, Sigma Aldrich) and cultured at 37°C in a humidified 5% CO₂ incubator. After 24h, the medium was refreshed and supplemented with m-CSF1 to improve microglial survival. Cultures were let reach confluency (about 2 weeks) without further changes. For mixed culture analysis, cells were detached by washing with warm PBS and then incubated with 0.25% trypsin (Invitrogen), 1mM EDTA in PBS at 37°C for 15min. After adding DMEM/F12 with 10% FCS, cells were centrifuged for 5 min at 1000 rpm, and re-suspended in the same medium. Microglial cells were purified by mechanical detachment of mixed glial cultures, recovered by medium centrifugation (5 min; 1000 rpm) [79] and plated replacing only 50% of the medium to promote microglial proliferation. Cells were seeded on multi-well culture plates (Falcon) at a density of 10⁵ cells/cm². In mixed cultures, the astrocyte/microglia ratio was 1:1. Cells were

let settle for 48h and treated with either LPS (1µg/ml) or saline. Cell media were then collected and processed.

Immunohistochemistry. Newborn and adult mice were perfused transcardially with 4% PFA in PB 0.1M (wt/vol). Brains were removed, post-fixed by immersion overnight and then washed for 24h in PBS, on a rocking platform at 4°C. Spinal cords were extracted after body post-fixation and washing. Tissues were incubated in a 30% sucrose-PB solution (wt/vol) for 24h, embedded in a 7.5% gelatin; 15% sucrose solution (wt/vol), frozen on dry ice and, if necessary, stored at -80°C until serially sectioned coronally at 15µm of thickness using a cryostat (Leica). Histological and cytological immunostaining was performed following standard protocols, after antigen retrieval (10mM citrate buffer, pH6, for 5min at 110°C in a boiling chamber, Biocaremedical). Primary antibodies (Table S2) were incubated ON at RT. The following secondary antibodies were incubated for 1h at RT: Alexa488 or Alexa594-conjugated donkey anti-rabbit or anti-mouse (1:1000); goat anti-rat (1:3000; Molecular Probes, Invitrogen) or anti-chicken, (1:2000, AbCAM); biotin conjugated goat anti-mouse or anti-rabbit (1:500, Jackson Lab). Alexa488, Alexa594 (1:500; Molecular Probes, Invitrogen) or POD conjugated (Jackson Lab) streptavidin followed by reaction with 3,3-diaminobenzidine (0.05%; Sigma) and 0.03% H₂O₂. For immunofluorescence, sections were counterstained with Hoechst (Sigma Aldrich). Tissue was analysed with a DMCTR5000 microscope equipped with a DFC350Fx monochrome camera or a DFC500 color camera (Leica Microsystems) or with a LSM710 confocal imaging system coupled to an AxioImager.M2 vertical microscope (Zeiss) or a LSM800 coupled to an Axio Observer inverted microscope (Zeiss). Fluorescence was quantified with ImageJ software (National Institute of Health) using 12-14 sections per analyzed brain.

ELISA. The amount of cytokines' present in the glial culture media was determined with electro-chemo-luminescence in a MSD MULTI-SPOT Assay System using 96 well V-PLEX plates for pro-inflammatory mouse panel 1 or custom mouse cytokine V-PLEX plates for IFN γ , IL1 β , IL4, IL6, IL10 and TNF α (Meso Scale Discovery), following the manufacturer's indications and using a SECTOR Imager 2400 reader (Meso Scale Discovery). SFRP1 presents in glial culture medium or in the RIPA fraction of brain lysates was determined with a capture ELISA [23]. Culture media were diluted five folds, and brain lysates used at 0.1 µg/µl protein concentration determined with BCA protein assay (Thermo Scientific). Values were determined at 450 nm wavelength using a FLUOstar OPTIMA micro-titter plate reader (BMG LABTECH).

Fluorescence Activated Cell Sorting (FACS) and Flow Cytometry Analysis. *CX₃CR1^{+GFP}* and *CX₃CR1^{+GFP};Sfrp1^{-/-}* 10-12 weeks-old male littermates were treated with LPS or saline as described. After 3 days, animals were perfused with ice-cold saline and brains collected on ice-cold, HBSS Ca²⁺, Mg²⁺-free (Invitrogen). After meninges' removal, cortices were isolated, finely chopped and digested for 20 min in DMEM with GlutMAX (Invitrogen) containing papain 20U/ml (Worthington Biochemical Corporation), DNase (50U/ml, Sigma Aldrich) and L-Cysteine (1mM, MERCK). After addition of 20% FCS (Invitrogen), the tissue was mechanically dissociated and filtered through a 35µm nylon strainer (Falcon). Microglial cells were separated from myelin/debris by Isotonic Percoll gradient centrifugation (35% Fisher Scientific) at 2000 rpm for 45min at 4°C. The pellet was recovered and sequentially incubated with anti-mouse CD16/CD32 (1:250, BD Pharmingen; for 15min at 4°C) followed by rat anti-CD11b and PerCP-Cy5.5 rat anti-mouse CD45 (1:200, BD Pharmingen; for 30min at 4°C), all in 2% BSA, 5mM EDTA in PBS. After washing, cells were sorted using a BD FACS Aria Fusion Flow Cytometer and their signal, size and complexity acquired with DiVA8 Software (BD Pharmingen). Analysis was performed using FlowJo v10.0.7 Software (BD Pharmingen).

Genomic Data Processing and Access. RNA from sorted microglia was extracted with the TRI reagent (MERCK) according to the manufacturer's instructions, purified with RNeasy Lipid Mini Kit (Qiagen) following the manufacturer's protocol and the resulting total RNA was treated with RNase-Free DNase Set (Qiagen). RNA quality was assessed with a Bioanalyzer 2100 system (Aligent), obtaining RIN (RNA integrity number) values between 8.8 and 10. Each sample (microglia sorted from a single mouse brain) was processed to obtain a RiboZero Stranded Gold Library (Illumina) and sequenced in HiSeq 4000 sequencer in paired-end configuration with 150bp sequence reads (Illumina). Sequence quality was determined with FastQC (Babraham Bioinformatics) revealing more than 32 million mean cluster reads of over 38 quality score and 93% mean Q30. Reads were aligned with HISAT2 v2.1.0 [80] to Ensemble *Mus musculus* (GRCm38.94). Aligned reads were further processed using Samtools v1.9 [81] and quantified to gene level using HTseq v0.11.2 [82]. Whole genome alignments were visualized with Integrative Genomics Viewer (IGV v2.5) [83]. Differential expression analysis (DGE) was performed using the Bioconductor package DESeq2 v1.10.0 [84]. DGE data were processed with custom R scripts (R version 3.5.1, 2018) considering genes with adjusted p-value < 0.05 and log₂ Fold Change > +/- 0.8 as significantly up- or down-regulated. Analysis of GO terms was performed with the

Bioconductor package clusterProfiler [85] and that of promoter-based motif enrichment with HOMER [86]. RNAseq data sets can be accessed at the European Nucleotide Archive public repository under the following accession numbers (ERP119668 / PRJEB3647).

Statistical analysis was performed using Prism v7 software (GraphPad). Different statistical tests were used as indicated in each figure footnote and represented by *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Acknowledgements We wish to thank J Avila, CBM, for providing a breeding pair of the *CX₃CRI::GFP* mice and the CBM image analysis, genomic and cytometry facilities for their support and advices during the course of this study. This work was supported by grants from the Spanish AEI (BFU2013-43213-P; BFU2016-75412-R with FEDER support), Fundación Tatiana Perez de Guzman el Bueno and CIBERER to PB and PE. JRC (BES-2011-047189) and MIM (BES-2014-068797) were supported by FPI fellowships from the MINECO. We also acknowledge a CBM Institutional grant from the Fundación Ramon Areces.

Author contributions JRC, PE and PB conceptualized and designed the research study. JRC, MIM and MJMB conducted the experiments and acquired the data. AB, JRC and BA designed, the EAE study and JRC and AB conducted and analyzed it. JRC, MIM, MTH, PE and PB analysed the data. MPK and SS provided support with cytokine analysis. JRC and PB wrote the paper. JPLA supervised RNAseq analysis. All authors read and approved the manuscript.

Conflict of interest: The authors declare no conflict of interest

Data Availability Section: Materials will be available upon request

RNAseq data can be accessed at the European Nucleotide Archive public repository: ERP119668 / PRJEB3647.

References

1. Jha MK, Jo M, Kim JH, Suk K (2019) Microglia-Astrocyte Crosstalk: An Intimate Molecular Conversation. *Neuroscientist* 25: 227–240.
2. Marinelli S, Basilico B, Marrone MC, Ragozzino D (2019) Microglia-neuron crosstalk: Signaling mechanism and control of synaptic transmission. *Semin Cell Dev Biol*.
3. Ransohoff RM (2016) How neuroinflammation contributes to neurodegeneration. *Science* 353: 777–783.
4. Escartin C, Guillemaud O, Carrillo-de Sauvage MA (2019) Questions and (some) answers on reactive astrocytes. *Glia*.
5. Frank-Cannon TC, Alto LT, McAlpine FE, Tansey MG (2009) Does neuroinflammation fan the flame in neurodegenerative diseases? *Mol Neurodegener* 4: 47.
6. Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH (2010) Mechanisms underlying inflammation in neurodegeneration. *Cell* 140: 918–934.
7. Perry VH, Holmes C (2014) Microglial priming in neurodegenerative disease. *Nat Rev Neurol* 10: 217–224.
8. Sarlus H, Heneka MT (2017) Microglia in Alzheimer’s disease. *J Clin Invest* 127: 3240–3249.
9. Hickman S, Izzy S, Sen P, Morsett L, El Khoury J (2018) Microglia in neurodegeneration. *Nat Neurosci* 21: 1359–1369.
10. Esteve P, Bovolenta P (2010) The advantages and disadvantages of sfrp1 and sfrp2 expression in pathological events. *Tohoku J Exp Med* 221: 11–17.
11. Esteve P, Sandonis A, Cardozo M, Malapeira J, Ibanez C, Crespo I, Marcos S, Gonzalez-Garcia S, Toribio ML, Arribas J, et al. (2011) SFRPs act as negative modulators of ADAM10 to regulate retinal neurogenesis. *Nat Neurosci* 14: 562–569.
12. Saftig P, Lichtenthaler SF (2015) The alpha secretase ADAM10: A metalloprotease with multiple functions in the brain. *Prog Neurobiol* 135: 1–20.
13. Lichtenthaler SF, Lemberg MK, Fluhrer R (2018) Proteolytic ectodomain shedding of membrane proteins in mammals—hardware, concepts, and recent developments. *EMBO J* 37:.
14. Kleinberger G, Yamanishi Y, Suarez-Calvet M, Czirr E, Lohmann E, Cuyvers E, Struyfs H, Pettkus N, Wenninger-Weinzierl A, Mazaheri F, et al. (2014) TREM2 mutations implicated in neurodegeneration impair cell surface transport and phagocytosis. *Sci Transl Med* 6: 243ra86.
15. Hundhausen C, Schulte A, Schulz B, Andrzejewski MG, Schwarz N, von Hundelshausen P, Winter U, Paliga K, Reiss K, Saftig P, et al. (2007) Regulated Shedding of Transmembrane Chemokines by the Disintegrin and Metalloproteinase 10 Facilitates Detachment of Adherent Leukocytes. *The Journal of Immunology* 178: 8064–8072.
16. Wong KK, Zhu F, Khatri I, Huo Q, Spaner DE, Gorczynski RM (2016) Characterization of CD200 Ectodomain Shedding. *PLoS One* 11: e0152073.
17. Musardo S, Marcello E, Gardoni F, Di Luca M (2014) ADAM10 in synaptic physiology and pathology. *Neurodegener Dis* 13: 72–74.
18. Prox J, Bernreuther C, Altmeyen H, Grendel J, Glatzel M, D’Hooge R, Stroobants S, Ahmed T, Balschun D, Willem M, et al. (2013) Postnatal disruption of the disintegrin/metalloproteinase ADAM10 in brain causes epileptic seizures, learning deficits, altered spine morphology, and defective synaptic functions. *J Neurosci* 33: 12915–12928, 12928a.
19. Suh J, Choi SH, Romano DM, Gannon MA, Lesinski AN, Kim DY, Tanzi RE (2013) ADAM10 missense mutations potentiate beta-amyloid accumulation by impairing prodomain chaperone function. *Neuron* 80: 385–401.
20. Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC, Boland A, Vronskaya M, van der Lee SJ, Amlie-Wolf A, et al. (2019) Genetic meta-analysis of diagnosed Alzheimer’s disease identifies new risk loci and implicates Abeta, tau, immunity and lipid processing. *Nat Genet* 51: 414–430.
21. Kuhn PH, Wang H, Dislich B, Colombo A, Zeitschel U, Ellwart JW, Kremmer E, Rossner S, Lichtenthaler SF (2010) ADAM10 is the physiologically relevant, constitutive alpha-secretase of the amyloid precursor protein in primary neurons. *EMBO J* 29: 3020–3032.

22. Marcello E, Borroni B, Pelucchi S, Gardoni F, Di Luca M (2017) ADAM10 as a therapeutic target for brain diseases: from developmental disorders to Alzheimer's disease. *Expert Opin Ther Targets* 21: 1017–1026.
23. Esteve P, Rueda-Carrasco J, Ines Mateo M, Martin-Bermejo MJ, Draffin J, Pereyra G, Sandonis A, Crespo I, Moreno I, Aso E, et al. (2019) Elevated levels of Secreted-Frizzled-Related-Protein 1 contribute to Alzheimer's disease pathogenesis. *Nat Neurosci* 22: 1258–1268.
24. Campanelli JT, Sandrock RW, Wheatley W, Xue H, Zheng J, Liang F, Chesnut JD, Zhan M, Rao MS, Liu Y (2008) Expression profiling of human glial precursors. *BMC Dev Biol* 8: 102.
25. Esteve P, Crespo I, Kaimakis P, Sandonis A, Bovolenta P (2019) Sfrp1 Modulates Cell-signaling Events Underlying Telencephalic Patterning, Growth and Differentiation. *Cereb Cortex* 29: 1059–1074.
26. Augustine C, Gunnarsen J, Spirkoska V, Tan SS (2001) Place- and time-dependent expression of mouse sFRP-1 during development of the cerebral neocortex. *Mech Dev* 109: 395–397.
27. Zhang Y, Sloan SA, Clarke LE, Caneda C, Plaza CA, Blumenthal PD, Vogel H, Steinberg GK, Edwards MS, Li G, et al. (2016) Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* 89: 37–53.
28. Blalock EM, Geddes JW, Chen KC, Porter NM, Markesbery WR, Landfield PW (2004) Incipient Alzheimer's disease: microarray correlation analyses reveal major transcriptional and tumor suppressor responses. *Proc Natl Acad Sci U S A* 101: 2173–2178.
29. Folke J, Pakkenberg B, Brudek T (2019) Impaired Wnt Signaling in the Prefrontal Cortex of Alzheimer's Disease. *Mol Neurobiol* 56: 873–891.
30. Bai B, Wang X, Li Y, Chen PC, Yu K, Dey KK, Yarbrow JM, Han X, Lutz BM, Rao S, et al. (2020) Deep Multilayer Brain Proteomics Identifies Molecular Networks in Alzheimer's Disease Progression. *Neuron* 105: 975-991 e7.
31. Claudel M, Jouzeau JY, Cailotto F (2019) Secreted Frizzled-related proteins (sFRPs) in osteo-articular diseases: much more than simple antagonists of Wnt signaling? *FEBS J*.
32. Youm YH, Grant RW, McCabe LR, Albarado DC, Nguyen KY, Ravussin A, Pistell P, Newman S, Carter R, Laque A, et al. (2013) Canonical Nlrp3 inflammasome links systemic low-grade inflammation to functional decline in aging. *Cell Metab* 18: 519–532.
33. Helton R, Cui J, Scheel JR, Ellison JA, Ames C, Gibson C, Blouw B, Ouyang L, Dragatsis I, Zeitlin S, et al. (2005) Brain-specific knock-out of hypoxia-inducible factor-1alpha reduces rather than increases hypoxic-ischemic damage. *J Neurosci* 25: 4099–4107.
34. Kaltschmidt B, Kaltschmidt C (2009) NF-kappaB in the nervous system. *Cold Spring Harb Perspect Biol* 1: a001271.
35. Wright SD (1999) Toll, a new piece in the puzzle of innate immunity. *J Exp Med* 189: 605–609.
36. Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, Littman DR (2000) Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol Cell Biol* 20: 4106–4114.
37. Satoh W, Gotoh T, Tsunematsu Y, Aizawa S, Shimono A (2006) Sfrp1 and Sfrp2 regulate anteroposterior axis elongation and somite segmentation during mouse embryogenesis. *Development* 133: 989–999.
38. Rivest S (2009) Regulation of innate immune responses in the brain. *Nat Rev Immunol* 9: 429–439.
39. Mii Y, Taira M (2009) Secreted Frizzled-related proteins enhance the diffusion of Wnt ligands and expand their signalling range. *Development* 136: 4083–4088.
40. Esteve P, Sandonis A, Ibanez C, Shimono A, Guerrero I, Bovolenta P (2011) Secreted frizzled-related proteins are required for Wnt/beta-catenin signalling activation in the vertebrate optic cup. *Development* 138: 4179–4184.
41. Heneka MT, Kummer MP, Latz E (2014) Innate immune activation in neurodegenerative disease. *Nat Rev Immunol* 14: 463–477.
42. Grassivaro F, Menon R, Acquaviva M, Ottoboni L, Ruffini F, Bergamaschi A, Muzio L, Farina C, Martino G (2020) Convergence between Microglia and Peripheral Macrophages Phenotype during Development and Neuroinflammation. *J Neurosci* 40: 784–795.
43. Ising C, Venegas C, Zhang S, Scheiblich H, Schmidt SV, Vieira-Saecker A, Schwartz S, Albasset S, McManus RM, Tejera D, et al. (2019) NLRP3 inflammasome activation drives tau pathology. *Nature* 575: 669–673.

44. Constantinescu CS, Farooqi N, O'Brien K, Gran B (2011) Experimental autoimmune encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *Br J Pharmacol* 164: 1079–1106.
45. Borroto A, Reyes-Garau D, Jimenez MA, Carrasco E, Moreno B, Martinez-Pasamar S, Cortes JR, Perona A, Abia D, Blanco S, et al. (2016) First-in-class inhibitor of the T cell receptor for the treatment of autoimmune diseases. *Sci Transl Med* 8: 370ra184.
46. Greter M, Lelios I, Croxford AL (2015) Microglia Versus Myeloid Cell Nomenclature during Brain Inflammation. *Front Immunol* 6: 249.
47. Kapellos TS, Taylor L, Lee H, Cowley SA, James WS, Iqbal AJ, Greaves DR (2016) A novel real time imaging platform to quantify macrophage phagocytosis. *Biochem Pharmacol* 116: 107–119.
48. Sole-Domenech S, Cruz DL, Capetillo-Zarate E, Maxfield FR (2016) The endocytic pathway in microglia during health, aging and Alzheimer's disease. *Ageing Res Rev* 32: 89–103.
49. Choudhry H, Harris AL (2018) Advances in Hypoxia-Inducible Factor Biology. *Cell Metab* 27: 281–298.
50. Kierdorf K, Prinz M (2013) Factors regulating microglia activation. *Front Cell Neurosci* 7: 44.
51. Colonna M, Butovsky O (2017) Microglia Function in the Central Nervous System During Health and Neurodegeneration. *Annu Rev Immunol* 35: 441–468.
52. Bovolenta P, Esteve P, Ruiz JM, Cisneros E, Lopez-Rios J (2008) Beyond Wnt inhibition: new functions of secreted Frizzled-related proteins in development and disease. *Journal of cell science* 121: 737–746.
53. Dahl E, Wiesmann F, Woenckhaus M, Stoehr R, Wild PJ, Veeck J, Knuchel R, Klopocki E, Sauter G, Simon R, et al. (2007) Frequent loss of SFRP1 expression in multiple human solid tumours: association with aberrant promoter methylation in renal cell carcinoma. *Oncogene* 26: 5680–5691.
54. Baharudin R, Tieng FYF, Lee LH, Ab Mutalib NS (2020) Epigenetics of SFRP1: The Dual Roles in Human Cancers. *Cancers (Basel)* 12:.
55. Trevant B, Gaur T, Hussain S, Symons J, Komm BS, Bodine PV, Stein GS, Lian JB (2008) Expression of secreted frizzled related protein 1, a Wnt antagonist, in brain, kidney, and skeleton is dispensable for normal embryonic development. *J Cell Physiol* 217: 113–126.
56. Holm TH, Draeby D, Owens T (2012) Microglia are required for astroglial Toll-like receptor 4 response and for optimal TLR2 and TLR3 response. *Glia* 60: 630–638.
57. Bowman CC, Rasley A, Tranguch SL, Marriott I (2003) Cultured astrocytes express toll-like receptors for bacterial products. *Glia* 43: 281–291.
58. Rolls A, Shechter R, London A, Ziv Y, Ronen A, Levy R, Schwartz M (2007) Toll-like receptors modulate adult hippocampal neurogenesis. *Nat Cell Biol* 9: 1081–1088.
59. Ajami B, Bennett JL, Krieger C, McNagny KM, Rossi FM (2011) Infiltrating monocytes trigger EAE progression, but do not contribute to the resident microglia pool. *Nat Neurosci* 14: 1142–1149.
60. Lee YS, Lee KA, Yoon HB, Yoo SA, Park YW, Chung Y, Kim WU, Kang CY (2012) The Wnt inhibitor secreted Frizzled-Related Protein 1 (sFRP1) promotes human Th17 differentiation. *Eur J Immunol* 42: 2564–2573.
61. Becher B, Durell BG, Miga AV, Hickey WF, Noelle RJ (2001) The clinical course of experimental autoimmune encephalomyelitis and inflammation is controlled by the expression of CD40 within the central nervous system. *J Exp Med* 193: 967–974.
62. Heppner FL, Greter M, Marino D, Falsig J, Raivich G, Hovelmeyer N, Waisman A, Rulicke T, Prinz M, Priller J, et al. (2005) Experimental autoimmune encephalomyelitis repressed by microglial paralysis. *Nat Med* 11: 146–152.
63. Chomyk AM, Volsko C, Tripathi A, Deckard SA, Trapp BD, Fox RJ, Dutta R (2017) DNA methylation in demyelinated multiple sclerosis hippocampus. *Sci Rep* 7: 8696.
64. Wang WH, McNatt LG, Pang IH, Millar JC, Hellberg PE, Hellberg MH, Steely HT, Rubin JS, Fingert JH, Sheffield VC, et al. (2008) Increased expression of the WNT antagonist sFRP-1 in glaucoma elevates intraocular pressure. *J Clin Invest* 118: 1056–1064.
65. Ulland TK, Song WM, Huang SC, Ulrich JD, Sergushichev A, Beatty WL, Loboda AA, Zhou Y, Cairns NJ, Kambal A, et al. (2017) TREM2 Maintains Microglial Metabolic Fitness in Alzheimer's Disease. *Cell* 170: 649–663 e13.
66. Wendeln AC, Degenhardt K, Kaurani L, Gertig M, Ulas T, Jain G, Wagner J, Hasler LM, Wild K, Skodras A, et al. (2018) Innate immune memory in the brain shapes neurological disease hallmarks. *Nature* 556: 332–338.

67. Baik SH, Kang S, Lee W, Choi H, Chung S, Kim JI, Mook-Jung I (2019) A Breakdown in Metabolic Reprogramming Causes Microglia Dysfunction in Alzheimer's Disease. *Cell Metab* 30: 493-507 e6.
68. Wheeler MA, Jaronen M, Covacu R, Zandee SEJ, Scalisi G, Rothhammer V, Tjon EC, Chao C-C, Kenison JE, Blain M, et al. (2019) Environmental Control of Astrocyte Pathogenic Activities in CNS Inflammation. *Cell* 176: 581-596.e18.
69. Gunner G, Cheadle L, Johnson KM, Ayata P, Badimon A, Mondo E, Nagy MA, Liu L, Bemiller SM, Kim KW, et al. (2019) Sensory lesioning induces microglial synapse elimination via ADAM10 and fractalkine signaling. *Nat Neurosci* 22: 1075–1088.
70. Turnbull IR, Gilfillan S, Cella M, Aoshi T, Miller M, Piccio L, Hernandez M, Colonna M (2006) Cutting Edge: TREM-2 Attenuates Macrophage Activation. *The Journal of Immunology* 177: 3520–3524.
71. Piccio L, Buonsanti C, Mariani M, Cella M, Gilfillan S, Cross AH, Colonna M, Panina-Bordignon P (2007) Blockade of TREM-2 exacerbates experimental autoimmune encephalomyelitis. *Eur J Immunol* 37: 1290–1301.
72. Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, David E, Baruch K, Lara-Astaiso D, Toth B, et al. (2017) A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell* 169: 1276-1290 e17.
73. Mazaheri F, Snaidero N, Kleinberger G, Madore C, Daria A, Werner G, Krasemann S, Capell A, Trumbach D, Wurst W, et al. (2017) TREM2 deficiency impairs chemotaxis and microglial responses to neuronal injury. *EMBO Rep* 18: 1186–1198.
74. Wright GJ, Puklavec MJ, Willis AC, Hoek RM, Sedgwick JD, Brown MH, Barclay AN (2000) Lymphoid/neuronal cell surface OX2 glycoprotein recognizes a novel receptor on macrophages implicated in the control of their function. *Immunity* 13: 233–242.
75. Halleskog C, Schulte G (2013) WNT-3A and WNT-5A counteract lipopolysaccharide-induced pro-inflammatory changes in mouse primary microglia. *J Neurochem* 125: 803–808.
76. Van Steenwinckel J, Schang AL, Krishnan ML, Degos V, Delahaye-Duriez A, Bokobza C, Csaba Z, Verdonk F, Montane A, Sigaut S, et al. (2019) Decreased microglial Wnt/beta-catenin signalling drives microglial pro-inflammatory activation in the developing brain. *Brain* 142: 3806–3833.
77. Marin V, Stornaiuolo A, Piovan C, Corna S, Bossi S, Pema M, Giuliani E, Scavullo C, Zucchelli E, Bordignon C, et al. (2016) RD-MolPack technology for the constitutive production of self-inactivating lentiviral vectors pseudotyped with the nontoxic RD114-TR envelope. *Mol Ther Methods Clin Dev* 3: 16033.
78. Merten OW, Hebben M, Bovolenta C (2016) Production of lentiviral vectors. *Mol Ther Methods Clin Dev* 3: 16017.
79. Bovolenta P, Wandosell F, Nieto-Sampedro M (1993) Neurite outgrowth inhibitors associated with glial cells and glial cell lines. *Neuroreport* 5: 345–348.
80. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907–915.
81. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–2079.
82. Anders S, Pyl PT, Huber W (2015) HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31: 166–169.
83. Thorvaldsdottir H, Robinson JT, Mesirov JP (2013) Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 14: 178–192.
84. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550.
85. Yu G, Wang LG, Han Y, He QY (2012) clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* 16: 284–287.
86. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* 38: 576–589.

Figure legends

Figure 1. Astrocytes upregulate SFRP1 expression upon LPS stimulation. A, B) Confocal image analysis of cryostat sections from adult $CX_3CRI^{+/GFP};Sfrp1^{+/βgal}$ mouse brains three days after intra-cortical infusion of saline or LPS. Sections were immunostained for βgal (magenta) and Iba1 (green, A) or Sfrp1 (magenta) and GFP (green, B), and GFAP (cyan). Arrowheads indicate βgal/GFAP (A) and Sfrp1/GFP co-localization (B). Scale bar 25μm. C) ELISA determination of Sfrp1 levels in brain extracts from 3 months-old wt and $Sfrp1^{-/-}$ mice. WT mice were injected either with saline or LPS. Three days after injection the region around the injected side (10 mm³ cortical cube) was isolated and SFRP1 content compared with that present in similar region of non-injected or $Sfrp1$ null mice used as negative control (n=5 mice for each group). Error bars represent Standard Error. Statistical significance: ****P<0.0001; One-way ANOVA followed by Bonferroni test.

Figure 2. SFRP1 is sufficient and required to enhance glial cell activation upon LPS treatment. A) Coronal cryostat sections of LV-IRES-*Gfp* or LV-*Sfrp1*-IRES-*Gfp* infected brains 1-month post-infusion, immunostained for SFRP1 (magenta) Iba1 (green) GFAP (cyan), or TauP (red). Scale bar 100μm. B) The graph shows the level of GFAP, Iba1 and CD45 immunoreactivity (IR) and the area occupied by TauP signal (n=24 acquisitions, white dots; N=3 mice, black dots, for each group), normalized to LV-IRES-*Gfp* infected brains. Error bars represent Standard Error. Statistical significance: ****P<0.0001 by two-sided Student's t-test. C, D) Coronal sections from wt and $Sfrp1^{-/-}$ mouse brains three days after infusion of saline or LPS, immunostained for GFAP (cyan, A) or CD45 (magenta, B). The images are high power views of the somatosensory cortex (lower power view in Fig S2). Scale bar 60μm. E, F) The graphs show the levels of immunoreactivity (IR) for GFAP (C) and CD45 (D, P=0,006) present in cortical sections (n=24 acquisitions white dots; from N=3 animals, black dots, per group) from wt and $Sfrp1^{-/-}$ animals infused with saline or LPS. Bars represent Standard Error. Statistical significance: ** or ### P<0.01, **** or ##### P<0.0001 by two-way ANOVA followed by Bonferroni's multiple comparisons test. * and # indicate significance between genotypes and treatments, respectively.

Figure 3. *Sfrp1*^{-/-} mice develop a milder form of EAE. A, B) Time course analysis and severity of the symptoms in wt and $Sfrp1^{-/-}$ mice after EAE induction. In A means are depicted with black (wt) and cyan ($Sfrp1^{-/-}$) lines. In B data are expressed as % of the total number of analyzed animals (n=19 per genotype). B) Wt and $Sfrp1^{-/-}$ mice immunized with

MOG and sacrificed 16 days post immunization. Cryostat sections of the thoracic spinal cords were stained with antibodies against CD4, Iba1, GFAP and MBP. Images show the region dorsal fasciculus. Scale bar 200 μ m. C) Quantification of CD4⁺ infiltrated lymphocytes, Iba1 immunoreactivity and MBP immunoreactive area in the dorsal fasciculus of spinal cord sections from wt and *Sfrp1*^{-/-} mice 16 days after immunization (n=16 acquisitions, white dots; from N=4 animals, black dots, per genotype). Statistical significance: *P<0.05, **P<0.01, ***P<0.001 by Kolmogorov-Smirnov test followed by Mann–Whitney U nonparametric test comparing mice of same day after immunization (a) or two-sided Student's t-test (c).

Figure 4. SFRP1 amplifies LPS-induced microglial activation and enhances their cytokine secretion.

A) Representative cytometry plots of CD11b⁺ populations present in the cortex of *CX₃CR1*^{GFP/+} or *CX₃CR1*^{GFP/+};*Sfrp1*^{-/-} mice 3 days after saline or LPS intraventricular infusion. Gating was set for isolating the following myeloid cell populations: CD11b⁺;CD45^{lo};GFP⁺ surveying microglia, CD11b⁺;CD45⁺;GFP⁺ activated microglia and CD11b⁺;CD45⁺;GFP⁻ infiltrated monocytes. B) Quantification of the percentage of activated microglia and infiltrated monocyte present in *Sfrp1*^{-/-} and controls brains 3 days after infusion of saline or LPS. C) Quantification of the total fluorescence from phagocytized pHrodo-labelled *E. coli* bioparticles and percentage of phagocytizing cells in the CD11b⁺;GFP⁺ microglial population. Error bars represent Standard Error. Statistical significance: **P<0.01, ***P<0.001 by two-way ANOVA followed by Bonferroni's multiple comparisons test. D) Secretory profile of cytokines present in the medium of primary microglia or microglia and astrocytes cultures from wt and *Sfrp1*^{-/-} pups exposed for 24h to saline or LPS (1 μ g/ml). The content of IFN γ , TNF α , IL1 β , IL4, IL6 and IL10 present in the culture media was determined by ELISA. Values, normalized to those of untreated wt cultures, are represented in Log scale. Error bars represent Standard Error. Statistical significance: ns P>0.05; *P<0.05; ***P<0.001; ****P<0.0001 by two-way ANOVA followed by Bonferroni's multiple comparisons test.

Figure 5. SFRP1 enhances the transcriptional response of microglial cells to LPS treatment.

A) Principal component analysis with the 1000 most variable genes from microglial cells isolated from *CX₃CR1*^{GFP/+} and *CX₃CR1*^{GFP/+};*Sfrp1*^{-/-} mouse brains three days after saline or LPS intra-cerebro-ventricular infusion. The analysis depicts sample clusterization by genotype and treatment. B, C) Volcano plots of differential gene expression from *CX₃CR1*^{GFP/+} (B) or *CX₃CR1*^{GFP/+};*Sfrp1*^{-/-} (C) microglial cell in response to LPS. Data are represented as Log₂ Fold Change vs -Log₁₀ adjusted p-value. Blue vertical lines represent

an increase of 75 and 400% in the expression levels respectively. Green horizontal lines represent a 0.05 or 0.01 adjusted p-value of statistical significance respectively. D) Scatterplot of differential expressed genes in $CX_3CRI^{GFP/+}$ and $CX_3CRI^{GFP/+};Sfrp1^{-/-}$ microglia; the red line represents linear regression (slope 0.704014 p-value: $< 2.2e-16$) of the differential expression, indicating response attenuation. Colored dots represent response which are exclusive for each genotype as represented in E. E) Venn diagram showing the extent of differential gene expression between $CX_3CRI^{GFP/+}$ and $CX_3CRI^{GFP/+};Sfrp1^{-/-}$ microglial cells. Fold change (75%) and adjusted p-value < 0.05 cut-off. F) Heatmap showing fold changes of regularized log transformed gene-level RNA-seq counts with hierarchical clustering of samples and differentially expressed genes

Figure 6. SFRP1 induces the up-regulation of HIF target genes during neuroinflammation. A) Ingenuity Pathway Analysis network representation of NF- κ B and HIF downstream targets for $CX_3CRI^{GFP/+}$ and $CX_3CRI^{GFP/+};Sfrp1^{-/-}$ microglial cells. Colored by Z-score alteration of their expression after LPS stimulus. B) Integrative Genomics Viewer transcription profile of represented genes for saline and LPS treatments of $CX_3CRI^{GFP/+}$ and $CX_3CRI^{GFP/+};Sfrp1^{-/-}$ microglial cells. Scale bar: 2 Kbp.

Supplementary Figure Legends

Supplementary Figure 1. SFRP1 enhances glial cell activation. A) Confocal image analysis of cryostat sections from adult *Sfrp1*^{+/ β gal} mouse brains three days after intra-cortical infusion of LPS. Sections were immunostained for β gal (green) and GFP (red) or Iba1 (red). Note that astrocytes but not microglial cells express the β gal/*Sfrp1* reporter. B) On the left, diagram of the observed GFP distribution (green dots) after lentiviral (LV) particles' delivery into the lateral ventricle (*Lv*). Images show examples of immunostaining against GFP protein (GFP signal was no longer visible, unless detected with immunohistochemistry) in the *Lv* of brains transduced with LV-*IRE5-Gfp* or LV-*Sfrp1-IRE5-Gfp* 5 months (m) post-infusion. C) Coronal sections from LV-*IRE5-Gfp* or LV-*Sfrp1-IRE5-Gfp* infected brains 1 or 5 months (m) post-infusion (PI). Sections were immunostained for CD45. Arrowheads indicate CD45^{hi}-positive macrophages or lymphocytes infiltrated in the parenchyma after prolonged LV-*Sfrp1-IRE5-Gfp* infection. High power images 1 and 2 were taken from the regions indicated with grey dotted lines in A. D) Coronal sections from wt and *Sfrp1*^{-/-} mouse brains three days after infusion of saline or LPS, immunostained for GFAP. The white arrows indicate the injection site; the area boxed with white lines is represented at high power in Fig 3.

Supplementary Figure 2. Astrocytes-derived SFRP1 enhances glial cell activation upon LPS treatment. A) The graph shows the % CD11b⁺ populations present in the cortex of *CX₃CR1*^{GFP/+} or *CX₃CR1*^{GFP/+};*Sfrp1*^{-/-} mice 3 days after saline or LPS intraventricular infusion. Gating was set for isolating the following myeloid cell populations: CD11b⁺;CD45^{lo};GFP⁺ surveying microglia, CD11b⁺;CD45⁺;GFP⁺ activated microglia and CD11b⁺;CD45⁺;GFP⁻ infiltrated monocytes. B) ELISA determination of SFRP1 levels present in the media of microglia or microglial and astrocytes culture derived from wt or *Sfrp1*^{-/-} pups. Note that SFRP1 is detected only in cultures in which astrocytes are present. Note also that LPS induces a significant increase of secreted SFRP1 protein. Error bars represent Standard Error. Statistical significance: ****P<0.0001; One way ANOVA followed by Bonferroni Test

Supplementary Figure 3. Gating strategy to assess phagocytosis. A) Representative cytometry plots of brain cells suspension showing gating of individualized live cells CD11b⁺;CX3CR1⁺ recognized as microglial cells for pHrodo labeled *E. coli* bio-particles quantitation. Comparison of microglial cells exposed (*E.coli* Ctrl+) or not exposed (*E.coli* Ctrl-) to pHrodo labeled *E.coli* shows specific pHrodo recognition. B) Representative

confocal image of FACS sorted GFP⁺ microglia. Note specific pHodo⁺ lysosomal inclusion of *E.coli* particles in phagocytosing vs non-phagocytosing microglia.

Supplementary Figure 4. SFRP1 microglial inflammatory response depends on HIF factors. A) Representation of the four up-regulated k-means unsupervised clusters of up-regulated genes in response to LPS in *CX₃CR1^{GFP/+}* or *CX₃CR1^{GFP/+};Sfrp1^{-/-}* microglial cells. Eight different clusters were generated by Z-scores co-variation. The clusters are indicated in the heatmap of Fig. 5F. B) Gene Ontology Enrichment Analysis of different clusters, GO biological process annotated are represented by fold enrichment and color-coded by adjusted p-value of enrichment and number of genes of each cluster in that term. C) Regulatory Elements Analysis of different clusters. Specifically enriched transcription factors are represented by fold enrichment relative to the overall transcripts present in the microglia transcriptome. Colored by Log odds detection threshold and the percentage of total targets for each transcription factor present in the cluster.











