

 Open access • Journal Article • DOI:10.1111/J.1466-8238.2010.00594.X

Asymmetric boundary shifts of tropical montane Lepidoptera over four decades of climate warming — [Source link](#)

I-Ching Chen, Jane K. Hill, Hau-Jie Shiu, Jeremy D. Holloway ...+4 more authors

Institutions: University of York, National University of Tainan, Natural History Museum, Universiti Malaysia Sabah

Published on: 01 Jan 2011 - Global Ecology and Biogeography (Blackwell Publishing Ltd)

Topics: Range (biology) and Extinction

Related papers:

- [Rapid Range Shifts of Species Associated with High Levels of Climate Warming](#)
- [A globally coherent fingerprint of climate change impacts across natural systems](#)
- [Ecological and Evolutionary Responses to Recent Climate Change](#)
- [Elevation increases in moth assemblages over 42 years on a tropical mountain](#)
- [Global Warming, Elevational Range Shifts, and Lowland Biotic Attrition in the Wet Tropics](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/asymmetric-boundary-shifts-of-tropical-montane-lepidoptera-2139k47z76>

Asymmetric boundary shifts of tropical montane Lepidoptera over four decades of climate warming

Abstract

Aim To estimate whether species have shifted at equal rates at their leading edges (cool boundaries) and trailing edges (warm boundaries) in response to climate change. We provide the first such evidence for tropical insects, here examining elevation shifts for the upper and lower boundaries shifts of montane moths. Threats to species on tropical mountains are considered. **Location** Mount Kinabalu, Sabah, Malaysia. **Methods** We surveyed Lepidoptera (Geometridae) on Mount Kinabalu in 2007, 42 years after the previous surveys in 1965. Changes in species upper and lower boundaries, elevational extents and range areas were assessed. We randomly subsampled the data to ensure comparable datasets between years. Estimated shifts were compared for endemic versus more widespread species, and for species that reached their range limits at different elevations. **Results** Species that reached their upper limits at 2500-2700 m (n= 28 species, 20% of those considered) retreated at both their lower and upper boundaries, and hence showed substantial average range contractions (-300 m in elevational extent and -45 km² in estimated range area). These declines may be associated with changes in cloud cover and the presence of ecological barriers (geological and vegetation transitions) which impede uphill movement. Other than this group, most species (n= 109, 80% of the species considered) expanded their upper boundaries upwards (by an average of 152 m) more than they retreated at their lower boundaries (77 m). **Main conclusions** Without constraints, leading margins shifted uphill faster than trailing margins retreated, such that many species increased their elevational extents. However, this did not result in increases in range area because the area of land available declines with increasing elevation. Species close to a major ecological/geological transition zone on the mountain flank declined in their range

areas. Extinction risk may increase long before species reach the summit, even when undisturbed habitats are available.