



Published in final edited form as:

J Pers Disord. 2018 January ; 32(Suppl): 93–114. doi:10.1521/pedi.2018.32.supp.93.

Attachment-related regulatory processes moderate the impact of adverse childhood experiences on stress reaction in Borderline Personality Disorder

Johannes C. Ehrenthal^{1,2}, Kenneth N. Levy¹, Lori N. Scott³, and Douglas A. Granger^{4,5,6}

¹Department of Psychology, Pennsylvania State University

²Department of Psychology, Klagenfurt University, Austria

³Department of Psychiatry, University of Pittsburgh School of Medicine

⁴Institute for Interdisciplinary Salivary Bioscience Research, University of California at Irvine

⁵Johns Hopkins University School of Medicine; Johns Hopkins University Bloomberg School of Public Health; Johns Hopkins University School of Nursing

⁶Department of Psychology and Salivary Bioscience Laboratory, University of Nebraska

Abstract

In this study, we explored whether attachment insecurity moderates the effects of adverse childhood experiences on stress-reactivity in the context of Borderline Personality Disorder (BPD). Participants were 113 women (39 with BPD, 15 with some BPD criteria present, 59 without any BPD symptoms) who participated in the Trier Social Stress Test. Saliva samples were collected before and after the stressor, and assayed for alpha-amylase and cortisol. Adverse childhood experiences were measured using the Childhood Trauma Questionnaire, and Attachment by the Experiences in Close Relationships-Revised questionnaire. Results revealed that attachment avoidance, and a combination of more adverse childhood experiences and attachment insecurity (anxiety and avoidance) resulted in higher salivary alpha-amylase (sAA) levels as well as differences in reactivity to the stressor. Interactions between attachment anxiety and adverse childhood experiences were related to blunted cortisol reactivity. The results suggest that the influence of adverse childhood experiences on stress-regulation in BPD may be moderated by attachment-related regulatory processes.

Keywords

Borderline Personality Disorder; attachment; childhood trauma; cortisol; TSST; alpha-amylase

Correspondence concerning this article should be addressed to Johannes C. Ehrenthal or Kenneth N. Levy, Pennsylvania State University, Department of Psychology, 362 Bruce V. Moore Building, University Park, PA 16802. johannes.ehrenthal@aau.at or klevy@psu.edu.

Disclosure Statement

In the interest of full disclosure, DAG is the founder of Salimetrics LLC and Salivabio LLC (Carlsbad, CA), serves as the companies' chief scientific and strategy advisor, and these relationships are managed by the policies of the committees on conflict of interest at the Johns Hopkins University School of Medicine and University of California at Irvine. Drs. Johannes C. Ehrenthal, Kenneth N. Levy, and Lori N. Scott and report no competing interests

Stress physiology offers a unique approach to the study of psychopathology, as it has the potential to provide information over and above self-report measures of distress and self-regulation (Campbell & Ehlert, 2012). Adding biomarkers to psychological measures may improve diagnostic accuracy, or increase predictive validity (Halford, Anderzén, & Arnetz, 2003). At the same time, individuals react very differently to stress on the level of biomarkers, and these interindividual differences can be related a variety of developmental and psychological factors (Granger, Kivlighan, El-Sheikh, Gordis, & Stroud, 2007). Integrating models from psychology and stress physiology has a twofold potential: On the one hand, it increases our general understanding of biobehavioral response to stress; on the other, it helps to understand disorder-specific and transdiagnostic regulatory processes. This may be especially relevant for mental health conditions such as Borderline Personality Disorder (BPD), where diagnostic heterogeneity is a common phenomenon due to 126 different ways to fulfill the minimal requirement for a valid diagnosis (American Psychiatric Association, 2013).

At the same time, BPD has been conceptually linked to stress. Several contemporary models of BPD see heightened reactivity to challenges, high levels of arousal and negative affect, and difficulties in emotional self-regulation as key features of the disorder (Linehan, 1993; Tragesser, Solhan, Schwartz-Mette, & Trull, 2007; Zanarini & Frankenburg, 2007). However, current research indicates that these factors may rather be transdiagnostic variables that are important for individuals with BPD, but do not explain the disorder sufficiently. For example, ecological momentary assessment studies reveal that neither affective instability nor switching of emotional states seem specific for BPD (Houben et al., 2016; Köhling et al., 2015; Santangelo et al., 2016). The same is true when looking at direct measures of psychophysiological arousal. Findings on reactions of the hypothalamic–pituitary–adrenal (HPA) axis and the autonomic nervous system (ANS) in individuals with BPD are mixed (Scott, Levy, & Granger, 2013; Wingenfeld, Spitzer, Rullkotter, & Lowe, 2010; Wingenfeld et al., 2010; Zimmerman & Choi-Kain, 2009). This indicates a need for studies that help explain these diverse findings on stress physiology in BPD. Some researchers specifically recommend investigating moderating constructs from developmental psychopathology (Wingenfeld et al., 2010; Zimmerman & Choi-Kain, 2009).

Two prominent developmental variables that have received considerable attention in BPD are early life stress and attachment. As we will explain in later sections of the manuscript, an advantage of these variables for the study of biomarkers is that both have been linked to stress physiology. Similar to the data on psychological reactivity mentioned above, current research suggests that both childhood trauma and insecure attachment are important for understanding individuals with borderline conditions. Early research suggested a strong relationship between BPD and developmental trauma (e.g., Herman, Perry, & van der Kolk, 1989). However, subsequent studies were not able to support this finding as particularly disorder-specific (for a discussion, see Ball & Links, 2009; Ford & Courtois, 2014; Fossati, Madeddu, & Maffei, 1999). Another research tradition conceptualized BPD as an attachment disorder (Fonagy, Gergely, Jurist, & Target, 2002; Liotti, Pasquini, & Cirrincione, 2000). Again, while attachment insecurity seems to be an important risk factor for psychopathology in general, including BPD, it is probably more a transdiagnostic than a BPD-specific construct (Levy, 2005; Levy, Johnson, Clouthier, Scala, & Temes, 2015). In other words,

developmental variables such as early traumatic experiences and insecure attachment patterns are relevant for BPD, but may rather be helpful for understanding interindividual differences within the diagnosis than distinguishing it from other psychiatric disease conditions. When considering developmental variables to understand stress regulation in BPD, it is important to adopt an approach that tries to take into account their dynamic interplay. Most of the current studies utilize either main-effect models or simple diathesis-stress models on the influence of a developmental variable on human stress-physiology. This suggests a linear influence of early priming on adult stress-systems (Carrion & Kletter, 2012; Radley et al., 2011), or a biological model of state-related influence by a current mental disorder as biomarker or endophenotype (Goldstein & Klein, 2014; Lenzenweger, 2013). These approaches have the potential of underestimating the complexity of developmental trajectories (Haltigan, Roisman, & Fraley, 2013). For example, as specified below, different attachment styles are associated with specific self-regulatory strategies, while childhood trauma seems to be a general risk factor that influences early priming of the biological stress systems.

The aim of this study is to understand stress regulation in individuals with varying levels of BPD in the context of the impact of childhood trauma and attachment styles. Reaching beyond main effect models, we focus on the dynamic interplay between the two variables in predicting levels of psychobiological arousal, reactivity, and recovery to a social-evaluative stressor. In the following sections, we will first describe contemporary concepts of stress regulation, then models and findings of the impact of adverse childhood experiences and attachment on stress regulation, before describing the study protocol.

Interindividual differences – A stress model approach

A fundamental component of stress research refers to the adaptiveness of regulatory biological systems. Expanding on ideas from classical concepts of homeostasis regulation, one of the most influential contribution to the understanding of stress and disease stems from the model of allostasis and allostatic load (Karatsoreos & McEwen, 2011; McEwen & Wingfield, 2010). Allostasis refers to adaptive processes of body and brain to changing internal or external demands by providing optimal levels of anticipation and energy mobilization. In addition, the effective down-regulation of arousal systems in the absence of demands is considered central for biological functioning. Chronic or repeated stress results in high, sometimes deleterious demand on the organism to adapt, called ‘allostatic load’ (Koolhaas et al., 2011). While general mechanisms from the allostatic load model have stimulated a large body of research, empirical results regarding psychophysiology depend on a variety of known and yet to explore predictor variables (Kudielka, Hellhammer, & Wust, 2009). In addition, while the model focuses on mechanisms of adaptation and disease development, it underemphasizes possible developmental trajectories of stress response (Juster et al., 2011). Some of these issues are addressed by current approaches, such as the adaptive calibration model (Del Giudice, Ellis, & Shirtcliff, 2011; Del Giudice, Hinnant, Ellis, & El-Sheikh, 2012; Essex et al., 2011). Key features of the adaptive calibration model are the focus on the integration of allostasis, life history, and the functioning of long-term, regulatory feedback loops. Furthermore, it describes different physiological ‘phenotype’ patterns in the context of their developmental history and biological responses. The model

seems especially relevant for the study of developmental variables in BPD, as it adds to previous ideas on stress-regulation the perspective of interindividual differences related to adverse childhood experiences in a testable form. We will summarize specific findings regarding the impact of childhood trauma on psychobiological arousal in the following section.

The impact of adverse childhood experiences on biobehavioral regulatory function

The detrimental impact of adverse childhood experiences and trauma on adult health places a major burden on individuals and society (Edwards, Holden, Felitti, & Anda, 2003; Felitti et al., 1998). A significant part of this strain is attributed to altered stress sensitivity by early life stress, influencing susceptibility to later life stressors (Harkness, Bruce, & Lumley, 2006), as well as to alterations of psychophysiological pathways (Ehlert, 2013). The general hypothesis of alterations of psychophysiological pathways by a negative impact of early life stress as well as adult trauma has been tested in a growing number of studies over the last 20 years (Heim & Nemeroff, 2001; Ehlert, 2013). Results on the impact of adult traumatic experiences on HPA-axis function are diverse. Although differences in HPA functioning have been described in studies using healthy control-groups (Zoladz & Diamond, 2013), evidence suggests that it is not just state PTSD symptoms that drive changes in HPA reaction. For example, a meta-analysis by Klaassens and colleagues (2012) found no difference in overall cortisol levels between individuals exposed to a traumatic event vs. those without. The results of another meta-analysis, incorporating studies with childhood trauma in addition to adult exposure to adverse experiences, show higher cortisol output in close temporal proximity to the adverse situation, but lower levels of cortisol secretion when more time elapsed since the traumatic event (Morris, Compas, & Garber, 2012). The influence of adverse childhood experiences, but not current PTSD diagnosis, on cortisol levels is also supported by a study by van Voorhees and colleagues (2014). Similarly, recent traumatic events were related to hypercortisolism, but more time elapsed since the event was associated with relatively lower levels of cortisol in children (Carrion & Kletter, 2012; Bellis & Zisk, 2014). In a mixed sample, Carvalho Fernando et al. (2012) found specific adverse childhood experiences to be more predictive of salivary cortisol levels and reaction after than other variables, including depression or BPD symptoms, indicating a transdiagnostic psychopathological process. However, a large study by Holleman and colleagues (2012) saw no clear main effect of childhood trauma on cortisol level or reaction after adjusting for covariates. Taken together, the majority of the available studies suggest that adverse childhood experiences may impact adult HPA stress-reactions in at-risk samples, and that developmental timing, time elapsed since trauma, and measurement issues are important variables to consider when trying to generalize the results. Nonetheless, contradictory findings indicate that the exact influence of childhood trauma on stress sensitivity in adulthood remains insufficiently understood.

Despite its solid empirical base, research studies on associations between childhood trauma and adult stress reaction place less emphasis on how social-cognitive mechanisms influence psychophysiological reactions. For example, stress-responses are associated with

psychopathological states (Ehrental, Herrmann-Lingen, Fey, & Schauenburg, 2010; Llera & Newman, 2014), which in turn are prone to be influenced by mechanisms of worry and rumination (Ottaviani, Shapiro, & Couyoumdjian, 2013). As a substantial proportion of social-cognitive mechanisms influence stress reaction outside of conscious processes (Brosschot, 2010), or through challenges to the self-concept (Papousek, Paechter, & Lackner, 2011), there is a need for models that helps to predict which psychological processes individuals use for self-regulation. A cogent developmental model of interindividual differences in psychological regulatory capacity is described by attachment theory.

Attachment and psychological regulatory function

The attachment motivational system emerges in early childhood, depending on contingency and co-regulatory quality of interactions with the primary caregivers (Bretherton & Munholland, 2008). Main goals of attachment system activation in childhood are the (re-) establishment of proximity with an attachment figure, contact maintenance, and a state of felt security. So-called ‘internal working models’ of attachment are social-cognitive representational structures which serve to integrate and interpret relational information, and regulate attachment-oriented motivation, cognitions, emotions, and behavior (Dykas & Cassidy, 2011). In children and adults, there are three organized attachment styles: secure, anxious, and avoidant. Prototypically secure individuals enjoy close relationships, and are able to adequately regulate proximity to significant others. Prototypically avoidant persons are usually uncomfortable relying on others in times of distress, deny interpersonal needs, and deactivate attachment-related cognitions. Anxious individuals see themselves as insufficient with respect to self-regulatory competence, and have a tendency for a hyperactivation of the attachment system. Their attempts to secure the attention and proximity of others are often accompanied by emotional hyperarousal, clinging or controlling behavior. In addition to the three organized styles, a fourth category is related to unresolved attachment trauma and loss (e.g., Bernier & Meins, 2008; Madigan et al., 2006). Individuals with this kind of attachment classification experience seemingly contradictory relationship needs, wishes, and behaviors, resulting from the impact of incoherent, fragmented representational structures of the self and others in, parallel to relationship patterns observed in patients with BPD (Levy, 2005). A central theoretical aspect of internal working models is the conceptualization of organized insecure attachment strategies as defensive regulatory processes, which are “mental mechanisms aimed at adaptation and self-regulation” (Mikulincer, Shaver, Cassidy, & Berant, 2009, p. 294). Attachment-related regulatory strategies are usually activated to regulate distress resulting from rejection, loneliness, fear, and further consequences of unaccomplished relational needs. Based on these strategies, contemporary models of attachment (Mikulincer, Shaver, and Pereg, 2003), allow to predict dynamic aspects of attachment security and insecurity (e.g., Coan, 2010; Ein-Dor, Mikulincer, & Shaver, 2011; Rusk & Rothbaum, 2010; Sadikaj, Moskowitz, & Zuroff, 2011). In addition, attachment-related excitatory and inhibitory feedback-loops can be used to understand how co- and self-regulatory processes lead to stability or change of attachment representations (Diamond, 2001; Sbarra & Hazan, 2008). Attachment-related regulatory function depends on an individual’s predominant attachment style. Anxious

attachment is associated with the abovementioned hyperactivation and hypersensitivity to attachment related cues, and a negative view of the self. Attachment avoidance and its related downregulation of relational cognitions and emotions, on the other hand, seems to be associated with a more positive view of the self (e.g., Ehrental, Dinger, Lamla, Funken, & Schauenburg, 2009). However, avoidant regulatory functioning is more fragile and prone to failure. Initially adaptive avoidant strategies may break down under conditions of emotional or cognitive load (Gillath, Giesbrecht, & Shaver, 2009). Also, while individuals with attachment anxiety are challenged by relational threats such as separation, avoidant individuals react more strongly to evaluative, self-challenging situations, such as real or imagined failure in an achievement task (Mikulincer, 1998). It remains an open question if the model by Mikulincer et al. (2003) may be limited for research in personality disorders, as it was tested mostly in relatively healthy student samples, and does not incorporate assumptions on the level of integration and coherence of the attachment system. This could be relevant for the study of BPD, as individuals with a very low level of integration and coherence of internal working models, related to high levels of unresolved relational childhood trauma, may react differently to stressors than individuals with higher levels of integration of the attachment system.

The impact of attachment strategies on biobehavioral regulatory function

While attachment insecurity is a normal facet of development, theoretical as well as empirical evidence suggests that insecure attachment may narrow regulatory strategies and therefore serve as a risk factor for health-related outcome. Known pathways include impaired biological stress-physiology as well as reduced abilities for an adequate social modulation of stress (Maunder and Hunter, 2001, 2008). Although stress regulation is a central function of attachment theory (Gunnar, Brodersen, Nachmias, Buss, & Rigatuso, 1996), systematic psychophysiological in this area started to unfold just over the last two decades (Diamond, 2001). Central to all cited approaches is the emphasis on attachment-related interindividual differences, importance of interpersonal co-regulation, and the intertwinement of psychological factors and psychophysiology, such as reaction of the HPA-axis (Dykas & Cassidy, 2011; Sbarra & Hazan, 2008).

Research on HPA-functioning and adult attachment differs widely with regard to methods used and results (Diamond & Fagundes, 2010). Existing studies can be grouped according to their way of assessing attachment (e.g., by questionnaire or interview). A number of articles on attachment and HPA-function are published that make use of self-report measures of attachment styles. A large study by Kidd, Hamer, and Steptoe (2011) found a blunted cortisol response to two non-social, behavioural tasks for individuals classified as “fearful” (individuals reporting high levels of both attachment/separation anxiety as well as attachment avoidance). In a reanalysis of cortisol patterns from the Whitehall II cohort, preoccupied attachment was related to higher levels of cortisol as well as less decrease of cortisol output over the day (Kidd, Hamer, & Steptoe, 2013). Quirin, Pruessner, and Kuhl (2008) reported a higher cortisol response to an acute stressor, but lower cortisol awakening response to be associated with higher attachment anxiety in healthy women. Gordon et al. (2008) found attachment anxiety to be related to cortisol in a sample of young adults. Cortisol reaction during a group-related stress paradigm was greater in women with higher

attachment anxiety scores (Smyth et al., 2015). In a small study with healthy women by Tops, van Peer, Korf, Wijers, and Tucker (2007), the attachment subscale of the Temperament and Character Inventory, which may serve as a proxy for tendencies to communicate and relate to friends with regard to own emotional states and needs, was negatively related to higher plasma cortisol levels. Smeets (2010), however, found no associations between attachment and cortisol reaction in a sample of 68 healthy participants. Fewer studies related interview-based attachment classifications to cortisol reaction. Scheidt et al. (2000) found an association between a more pronounced cortisol response during the administration of an Adult Attachment Interview (AAI) and higher levels of dismissing attachment in patients with idiopathic spasmodic torticollis, but not for healthy control participants. Rifkin-Graboi (2008) reported no evidence for a correlation between attachment insecurity and ambulatory cortisol responses, but there was an association between the AAI subscale “dismissing idealization” and higher cortisol levels after an attachment-activating memory task. And finally, Pierrehumbert et al. (2009) compared women with and without sexual abuse in childhood or adolescence, and simultaneously applied the AAI to measure the individuals’ attachment states of mind. They found the most suppressed cortisol reaction to a standardized laboratory stress induction task in a subgroup of women with a combination of abuse experience as well as a classification as unresolved with regard to attachment trauma in the AAI.

To sum up, research using attachment questionnaires found some evidence for an impact of attachment insecurity on cortisol reaction. However, although more studies report a specific impact of attachment anxiety, there are data favouring attachment avoidance as well. In addition, there is evidence that with a lower level of general integration of internal working models of attachment, often seen in individuals with repeated traumatic childhood experiences, cortisol reaction may be suppressed.

The current study

Although research on the impact of adverse childhood experiences and trauma on BPD and adult stress physiology has advanced over the last decades, conflicting findings call for further research on individual differences in biobehavioral regulatory function. Recent models of stress-regulation propose a non-linear, moderated impact of childhood trauma on the HPA axis. At the same time, developmental models of social cognition, such as attachment theory, provide a framework to understand the moderating impact of hyperactivation and downregulation in the face of threat. Therefore, our study aims at testing the moderating influence of adult attachment strategies on the impact of adverse childhood experiences on psychophysiological stress-regulation. In other words, we hypothesized that interactions between childhood trauma and attachment insecurity would significantly predict psychophysiological stress reaction in individuals with varying levels of BPD symptoms.

Methods

Participants

Participants were recruited in the greater State College area, either from an outpatient community mental health clinic, or from a pool of university students and community

residents following screening procedures described by Scott et al. (2013). Individuals diagnosed with psychotic disorders, Bipolar I disorder, delirium, dementia, mental retardation, heart disease, endocrinological diseases (with the exception of diabetes and thyroid disorders; Jones et al., 2004), pregnancy during the preceding six months, or current lactation were excluded.

Participants' characteristics are displayed in Table 1. The sample consisted of 118 women who were on average in their mid-twenties, ranging from 18 to 48 years, and predominantly white/Caucasian, with about ten percent identifying as African American. A little over 40 % had achieved a high-school degree; the majority of the other participants (41.6 % of the total sample) had finished partial 4-year or standard college. Eight participants had completed a master's level, and 2 a doctoral level degree. Only 17.7 % worked full-time. Thirty-four participants had a diagnosis of BPD, and five a sub-threshold BPD, but exhibited prominent symptoms that would justify a BPD diagnosis. The number of additional, non-BPD DSM-IV Axis-II diagnosis ranged from zero to two, the number of current Axis-I diagnoses from zero to six. On average, participants took about one medication at the time of the study, ranging from zero to ten (more data on medication is available on request).

Psychosocial Stressor

Stress induction was performed with a social-evaluative stress protocol, the Trier Social Stress test (TSST; Kirschbaum, Pirke, & Hellhammer, 2004). It comprises two stressful tasks (public speaking, mental arithmetic) in front of a stern and evaluative jury of confederate "judges". The tasks are preceded by a standardized baseline resting phase, and followed by a recovery phase. The TSST is known to reliably elicit HPA reaction (Foley & Kirschbaum, 2010).

Measures

Adverse childhood experiences—To assess adverse childhood experiences, we administered the Childhood Trauma Questionnaire (CTQ). The CTQ measures self-reported physical, sexual, and emotional abuse, as well as physical and emotional neglect on 28 items (Bernstein & Fink, 1998; Scher, Stein, Asmundson, McCreary, & Forde, 2001). We used the total sum score of the abuse and neglect scales as a measure of general adverse childhood experiences.

Attachment—Attachment styles were measured with the Experiences in Close Relationships – Revised questionnaire (ECR-R). It assesses attachment related anxiety and avoidance on 36 items addressing romantic relationship expectations and experiences (Fraley, Waller, & Brennan, 2000), and has shown good psychometric properties (Fairchild, 2006; Sibley, Fischer, & Liu, 2005).

Trait negative affect—To control for possible impacts of trait negative affect and impulsivity, the facet scales Angry Hostility, Impulsivity, Depression, and Anxiety from the Revised NEO Personality Inventory (Costa & McCrae, 1992) were administered to all participants.

Diagnostic status—Psychological diagnoses were assessed using the Structural Clinical Interview for DSM-IV (SCID-I-CV; (First, Spitzer, Gibbon, & Williams, 1997), and the International Personality Disorder Examination (IPDE; Loranger, 1999), to generate reliable diagnosis of DSM-IV personality disorders. Both interviews were administered by experienced, trained raters at graduate student level who were blind to the initial recruitment background.

Other covariates—Other questions aimed to assess additional possible covariates, such as food and beverage intake, socio-demographical data, body mass index (BMI), dental hygiene, sleep patterns, physical activity, and menstrual cycle, but also overall substance use or recent life stress.

Procedure

At the first appointment, all participants provided written informed consent, thereafter completing the diagnostic interviews and questionnaires. On a second appointment, participants returned for the TSST. They were instructed to neither drink alcohol 24 hours before the TSST nor use any non-prescribed medication or drugs six hours before examination. In addition, they were asked to restrain from caffeine or tobacco consumption as well as dental work for four hours prior to the test. A standard regime regarding consumption of foods and other beverages, dairy, citrus fruits, and sleep was administered. All participants were scheduled during the follicular phase of their menstrual cycle. The measurements took place during the afternoon for each individual participant in order to control for daily fluctuation of cortisol output. Before each testing session, these prerequisites were monitored; if participants had not been able to comply, their session was rescheduled.

After arriving at the lab, participants rinsed their mouth with water, completed some basic questionnaires, provided the first saliva sample, then sat quietly for the baseline period. Eight saliva samples per person were collected during the assessment by the passive drool method: -40 minutes, -30 minutes, directly before the TSST, directly after the TSST, + 10 minutes, + 20 minutes, + 30 minutes, + 40 minutes. From the samples, cortisol as a measure of HPA reaction, and alpha-amylase as a marker of ANS function were derived (Granger, Hibel, Fortunato, & Kapelewski, 2009; Kudielka, Gierens, Hellhammer, Wust, & Schlotz, 2012; Nater & Rohleder, 2009).

Saliva Collection, Cortisol, and salivary Alpha-Amylase Assay

Saliva samples were assayed for cortisol in duplicate using a commercially available enzyme immunoassay (Salimetrics, LLC). The assay uses 25 microliters of saliva per determination, has a lower limit of detection of 0.003 µg/dl, standard curve range from 0.012 to 3.0 µg/dl, and average intra- and inter-assay coefficients of variation less than 5 % and 10 %, respectively. Samples were assayed for alpha-amylase using a commercially available kinetic reaction assay (Salimetrics, LLC), which employs a chromagenic substrate, 2-chloro-p-nitrophenol, linked to maltotriose. The enzymatic action of alpha-amylase on this substrate yields 2-chloro-p-nitrophenol, which can be measured spectrophotometrically at 405 nm using a standard laboratory plate reader. The amount of alpha-amylase activity present in the

sample is directly proportional to the increase (over a 2-minute period) in absorbance at 405 nm. The intra-assay variation (CV) based on 30 replicate tests was less than 7.5%. The inter-assay variation based on 16 separate runs was less than 6%.

Data Preparation

Cortisol and salivary alpha-amylase (sAA) data were inspected for distribution and outliers. Outliers, defined as ± 3 *SD* away from the group mean, were winzorized (Tukey, 1977), relating to 5.7% of the cortisol as well as the sAA data. Prior to all analyses due to their skewness, sAA values were square-root-, and cortisol values log-transformed.

Statistical analyses

To assess cortisol and sAA during resting, reactivity, and recovery, we implemented a coding procedure described by Llabre and colleagues (2001). Eight units of time were constructed by dividing the minutes passed since first measurement by 10. Slower bio-reactivity of cortisol compared to sAA was accounted for by implementing different coding schemes as described in more detail elsewhere (Scott et al., 2013).

For each biomarker, we applied three mixed models analyses to separately assess effects of psychological predictors for resting phase, reactivity, or recovery, while simultaneously controlling for the impact of the other two phases. Time (resting phase, reactivity, recovery) served as a level-1 (i.e., within-person) predictor, and CTQ and ECR-R as level-2 (i.e., between-person) predictors (Garson, 2013). Intercept and slopes for each time-phase were treated as random effects with an unstructured covariance matrix. We examined both main and interaction effects of the CTQ and ECR-R anxiety and avoidance subscales for predicting intercepts and slopes (i.e., change) during each phase. Covariates such as age, medication, day in menstrual cycle, time of day, or NEO-PI-R subscales were included if they were at least marginally significant ($p < .10$) and contributed to better fit indices of the particular model. We also checked whether the expected associations between childhood trauma, attachment, and biomarkers were not just artefacts of BPD diagnosis by adding the number of BPD symptoms, and their interaction with time, to the analyses. All analyses were conducted with IBM SPSS 22 (IBM Corp., Amonk, NY)

Results

All baseline-models indicated a significant amount of variance to be explained at the participants' level (Hox, 2002). For cortisol, time was a significant level-1 predictor, indicating a relative change in cortisol from the beginning of each phase to the end of each phase. Specifically, there was a decline in cortisol levels during rest, followed by an increase in cortisol during reactivity to the TSST, and a decrease during recovery after the TSST (see Table 2). Time was a significant level-1 predictor for sAA levels during reactivity to, and a significant negative predictor for recovery from the TSST, with the expected pattern of an increase during, and a decrease after the stressor (see Table 3).

Childhood Trauma

There were no significant associations between levels of childhood trauma and cortisol or sAA during resting state, reactivity, or recovery.

Attachment

There was no main effect of attachment anxiety on cortisol or sAA during resting state, reactivity, or recovery. ECR-R attachment avoidance was significantly associated with higher levels of sAA in all three phases.

Interaction Childhood Trauma by Attachment

Cortisol reaction to the TSST interacted significantly with CTQ and ECR-R anxiety: For individuals with higher levels of adverse childhood experiences, higher attachment anxiety was associated with a less steep increase in cortisol during the reactivity phase.

Regarding sAA, there was a significant positive interaction effect between CTQ and attachment avoidance during all three phases, and a significant positive interaction effect between CTQ and ECR-R anxiety on sAA during reactivity and recovery. We also found a three-way interaction between time, CTQ, and attachment anxiety, in that for individuals with higher CTQ values, more attachment anxiety was associated with less reactivity, and more recovery of sAA (see Table 3).

Covariates and impact of BPD symptoms

Regarding cortisol, for all three phases the time the participant had been awake that day was the only significant covariate. Running the models with a full set of all possible covariates did not change the results. The number of IPDE BPD criteria did not show any significant associations with cortisol levels or time-related change in cortisol output. Furthermore, it did not change any of the other results. For sAA, significant covariates were the NEO anxiety and depression subscale, and the number of days in menstrual cycle. Running the models with a full set of all possible covariates did not alter the results. Entering number of IPDE BPD criteria, or BPD-criteria by time interactions into the model, did neither change the results, nor display any significant associations with the biomarkers.

Discussion

In a study on 113 women with varying levels BPD pathology, there was an impact of psychosocial variables on HPA as well as SNS reaction to a social-evaluative stressor. Especially attachment avoidance was related to generally elevated levels of salivary alpha-amylase. Interactions between higher levels of childhood trauma and attachment insecurity (avoidance, and, to a lesser degree anxiety) were associated with higher levels of SNS activation as well. Under conditions of higher levels of childhood trauma and adversities, attachment anxiety was related to dampened HPA reactivity, reduced SNS reactivity, and more SNS recovery.

An important finding of our study is that variables from the tradition of developmental psychopathology were more impactful for predicting HPA and SNS activity in a sample of

women with varying levels of personality functioning than BPD symptoms. In other words, adding the number of BPD symptoms or their interaction with time to the statistical model did neither add to, nor change the pattern of results. Instead, attachment avoidance, but also the interaction between adverse childhood experiences and attachment insecurity were related to higher levels of ANS functioning, especially regarding the SNS.

From a perspective of attachment theory, this can be understood as a breakdown of attachment regulatory functioning, either under conditions of the social-evaluative stressor, or the moderating influence of an internal risk-factor, childhood trauma. Although especially attachment avoidance may be helpful in the down-regulation of stressful memory content (Dykas & Cassidy, 2011), this effect disappears under conditions of stress and psychological load (Gillath et al., 2009). Similarly, the evaluative aspect of the TSST could be especially stressful for individuals with high levels of avoidance (Mikulincer, 1998). This could also explain the larger influence on sAA, which is more prone to react rapidly to, and be prolonged through psychological processes such as rumination and perseverative cognitions (e.g., Ottaviani et al., 2013). More overall activation of the SNS could lead to a psychobiological restriction of reactivity of the ANS, but also to a general blunting of the stress-response due to prolonged allostatic load (Juster, McEwen, & Lupien, 2010). Especially the developmental combination of attachment hyperactivation and risk factor childhood trauma would lead to a constant hyperactivation of the stress systems over the life span. The lower levels of HPA reactivity could then be a consequence of a counter-regulatory mechanisms, a preventive dampening of stress response, also described by Scott et al. (2013) for individuals with BPD, or Kidd et al. (2013) for preoccupied attachment. Interestingly, this would fit to the ‘vigilant pattern’ of the adaptive calibration model described by Del Giudice and colleagues (2011), that is assumed to develop in unpredictable or dangerous childhood conditions. At the same time, heightened levels of cortisol or sAA may in turn negatively influence encoding and processing of emotional and trauma-related stimuli, resulting in a deleterious regulatory feedback loop (Holz, Lass-Hennemann, Streb, Pfaltz, & Michael, 2014; Nicholson, Bryant, & Felmingham, 2014).

Our results do not support previous findings on a possible main effect of attachment anxiety on cortisol reaction. On the one hand, this could be due to sample effects. In contrast to other studies, we included women with low levels of personality dysfunction as well as women fulfilling full criteria for DSM-IV BPD diagnoses. By broadening the range of psychopathology, we also intended to broaden the variability of attachment and CTQ ratings. Studies with exclusively non-clinical or clinical samples are at a risk of over- or underestimating true effects of attachment and childhood trauma on stress reaction, especially when taking into account that individuals can have different ranges of psychobiological responsiveness (Del Giudice et al., 2011). For example, according to a perspective of allostatic load, for individuals in the upper range of attachment security, attachment anxiety could stimulate a larger cortisol reaction, as the organism can rely on adaptive downregulation on a psychological as well as biological level.

Strengths of the study are a reasonably large, well diagnosed sample of women with varying degrees of BPD symptoms, attachment insecurity, and childhood trauma. This is important not only for ensuring enough variance to be explained in the models, but also for addressing

these questions from an perspective that explicitly tries to capture dimensions of not only psychopathology, but also normal behavior and regulatory processes (Cuthbert & Kozak, 2013). We also made an effort to control for covariates that possibly influence cortisol or sAA measurements. And lastly, by using a mixed model approach, we were able to test the influence of our psychological predictors while simultaneously controlling for the influence of possibly competing variables. This is relevant, as early life stress and attachment insecurity share some conceptual overlap, namely the influence of actual experiences in an interpersonal context. At the same time, attachment formation needs the activation of the attachment system, and focuses on repeated micro-interactions (Beebe et al., 2010), while for the occurrence of adverse childhood experiences is not bound to attachment-related situations, and single events such as accidents are possible. In that regard, studying the impact of both variables simultaneously could help to disentangle their specific influences, especially when using a cross-sectional design. In our study, the risk factor childhood trauma was less impactful than the psychological variable attachment for HPA and ANS reaction.

Possibly relevant limitations of the study are the exclusive focus on women, and the use of questionnaire data for assessing attachment. While excluding men may help to reduce gender-related variance in the results (Bangasser & Valentino, 2012), it affects the generalizability of our results. Applying questionnaire assessment of attachment instead of interview-based data could also limit the understanding of the data by not being able to assess different levels of personality functioning of attachment directly (Blatt & Levy, 2003; Roisman et al., 2007). In other words, individuals with disorganized internal working models of attachment may not be able to rely on stable strategies to regulate themselves in comparison to individuals with organized attachment styles. Future studies, especially in the context of BPD, should apply both self-report and interview measures of attachment. Furthermore, while our predictor variables come from a tradition of developmental psychopathology, they measure current state of mind with regard to attachment, and current representations of childhood trauma. It would be interesting to address these questions in a sample from a longitudinal study with actual data from early childhood and adolescence. And last, but not least, did we simultaneously test the impact of our predictors on psychobiological functioning, controlling for effects of all the other variables, which may reduce the variance explained by each predictor if entered separately.

Our results allow for some tentative comments on possible implications for the study of BPD and its treatments. Firstly, as mentioned above, it may be helpful to complement classical BPD diagnosis according to DSM by other approaches, at least when looking at biological markers. The NIMH Research Domain Criteria (RDoC; Insel et al., 2010) initiative of studying proposed subsystems of psychobiological behavior is obviously one way of organizing research in the hope of reducing or explaining diagnostic heterogeneity. RDoC proposes a set of fundamental, transdiagnostic constructs hypothesized to be at the basis of human behavior and psychopathology, namely negative and positive valence systems, systems related to cognitive and social processes, and arousal and regulatory systems. While the inconsistencies regarding BPD described in the introduction can be seen as a prime example for the necessity of the RDoC approach, it seems important to exercise caution and not draw premature conclusion (Lilienfeld, 2014). Despite the attempt to extract empirically driven dimensions of personality and psychopathology, its current empirical

basis is heavily influenced by major research traditions from biological psychiatry, but also other areas of psychopathology. In fact, our results indicate that at this moment, a viable approach to making sense of RDoC for psychopathology and psychotherapy is to combine it with strong theoretical models from psychology. In the RDoC terminology, we used the ‘attachment’ construct from the domain ‘social processes’ together with the developmental risk factor adverse childhood experiences to understand interindividual psychophysiological differences in the construct ‘arousal’ of the domain ‘arousal and regulatory systems’. This theory driven approach is of special relevance for psychosocial treatments of BPD, as in the RDoC approach there is a tendency to favor possible biological substrates (Cuthbert & Kozak, 2013) over well-known psychological theories, but also over therapy-oriented diagnostic models (e.g., Oldham, 2015; Skodol et al., 2011). For future studies it might be helpful to integrate these and other approaches (Zimmermann et al., 2012) wherever possible, rather than to repeat old research programs under a new nomenclature.

At the same time, adding RDoC-oriented measures to existing paradigms from psychopathology can help to clarify disease mechanisms and mechanisms of change as targets for psychotherapy interventions. For example, Transference Focused Psychotherapy (TFP) has been shown to increase attachment security and reflective functioning (Fischer-Kern et al., 2015; Levy et al., 2006). Although other approaches to the treatment of BPD, for example Dialectical Behavior Therapy (DBT), are generally open to understanding interpersonal patterns and processes, for example regarding changes in patient introject (Bedics, Atkins, Comtois, & Linehan, 2012), they yet failed to show changes in attachment representations. The increase in coherence of the narratives on early attachment experiences in TFP may be directly related to more integrated representations of the self and others, which is central to the treatment model. A next step could be to delineate to what extent the increase in attachment security is related to short-or long-term increase in psychosocial functioning. Integrating this idea into the context of our study, it would be possible to test if an increase in attachment security is related to better psychophysiological recovery from a stressor after experimental activation of the representation of an attachment figure (Bryant & Chan, 2015). Similarly, multiple measurement points of attachment representations, but also attachment related stress reaction, over the course of interventions are needed to delineate if increase in attachment security precedes symptom change in TFP or other treatments, or vice versa (e.g., Daniel, Poulsen, & Lunn, 2016). While both TFP and attachment theory share the view that the current representation of traumatic interactional experiences is central for the understanding of regulatory psychodynamics, and results from our study indicate the importance of childhood trauma for stress-regulation, it would be interesting to see if and how TFP changes trauma-related symptoms. Here DBT has already started to combine its core program with features of other trauma-related interventions (Bohus et al., 2013; Harned, 2014). Especially in the light of promising results of non-exposure based treatments for trauma-related disorders (e.g., Markowitz et al., 2015; Steinert et al., 2016), further research is needed to show if TFP may be a viable option for the treatment of comorbid complex PTSD in BPD as well.

Acknowledgments

This research was supported by grants to Kenneth N. Levy from the Pennsylvania State University Social Science Research Institute, International Psychoanalytic Association, and American Psychoanalytic Association, and by grants to Lori N. Scott from the National Institute of Mental Health (F31MH081395, PI: Lori N. Scott), Pennsylvania State University College of Liberal Arts, and American Psychological Association. Parts of the sample were used in the doctoral dissertation of the Lori N. Scott. We would like to thank Joseph E. Beeney, Ph.D., William D. Ellison, Ph.D., Ann B. Stonebraker, M.S., Christina M. Temes, M.S., Rachel H. Wasserman, Ph.D., and Samantha Bernecker, M.S., for their assistance in conducting interview assessments. We would also like to thank Amber L. Walser, Psy.D., Rachel L. Tomko, Stevie N. Grasseti, M.A., Samantha Bernecker, and Laura Moser for their assistance in recruiting participants and technical assistance, and the members of the lab who contributed to the study by serving as judges in the TSST and coding data.

Reference List

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th. Washington, DC: American Psychiatric Association; 2013.
- Ball JS, Links PS. Borderline personality disorder and childhood trauma: Evidence for a causal relationship. *Current Psychiatry Reports*. 2009; 11(1):63–68. [PubMed: 19187711]
- Bangasser DA, Valentino RJ. Sex differences in molecular and cellular substrates of stress. *Cellular and Molecular Neurobiology*. 2012; 32(5):709–723. [PubMed: 22488525]
- Bedics JD, Atkins DC, Comtois KA, Linehan MM. Weekly therapist ratings of the therapeutic relationship and patient introject during the course of dialectical behavioral therapy for the treatment of borderline personality disorder. *Psychotherapy*. 2012; 49(2):231–240. [PubMed: 22642526]
- Beebe B, Jaffe J, Markese S, Buck K, Chen H, Cohen P, et al. The origins of 12-month attachment: a microanalysis of 4-month mother-infant interaction. *Attachment & Human Development*. 2010; 12(1–2):3–141. [PubMed: 20390524]
- de Bellis MD, Zisk A. The biological effects of childhood trauma. *Child and Adolescent Psychiatric Clinics of North America*. 2014; 23(2):185–222. vii. [PubMed: 24656576]
- Bernier A, Meins E. A threshold approach to understanding the origins of attachment disorganization. *Developmental Psychology*. 2008; 44(4):969–982. [PubMed: 18605828]
- Bernstein, DP., Fink, LA. CTQ: Childhood Trauma Questionnaire: a retrospective self-report. San Antonio, TX: Psychological Corp; 1998.
- Blatt SJ, Levy KN. Attachment theory, psychoanalysis, personality development, and psychopathology. *Psychoanalytic Inquiry*. 2003; 23(1):102–150.
- Bohus M, Dyer AS, Priebe K, Krüger A, Kleindienst N, Schmahl C, Niedtfeld I, Steil R. Dialectical behaviour therapy for post-traumatic stress disorder after childhood sexual abuse in patients with and without borderline personality disorder: a randomised controlled trial. *Psychotherapy and Psychosomatics*. 2013; 82(4):221–233. [PubMed: 23712109]
- Bretherton, I., Munholland, KA. Internal working models in attachment relationships: Elaborating a central construct in attachment theory. In: Cassidy, Shaver, editor. *Handbook of Attachment: Theory, research, and Clinical Applications*. 2nd. 2008. p. 102-130.
- Brosschot JF. Markers of chronic stress: prolonged physiological activation and (un)conscious perseverative cognition. *Neuroscience and Biobehavioral Reviews*. 2010; 35(1):46–50. [PubMed: 20096302]
- Bryant RA, Chan L. Thinking of attachments reduces noradrenergic stress response. *Psychoneuroendocrinology*. 2015; 60:39–45. [PubMed: 26115145]
- Campbell J, Ehlert U. Acute psychosocial stress: does the emotional stress response correspond with physiological responses? *Psychoneuroendocrinology*. 2012; 37(8):1111–1134. [PubMed: 22260938]
- Carrion VG, Kletter H. Posttraumatic stress disorder: shifting toward a developmental framework. *Child and Adolescent Psychiatric Clinics of North America*. 2012; 21(3):573–591. [PubMed: 22800995]

- Carvalho Fernando S, Beblo T, Schlosser N, Terfehr K, Otte C, Löwe B, et al. Associations of childhood trauma with hypothalamic-pituitary-adrenal function in borderline personality disorder and major depression. *Psychoneuroendocrinology*. 2012; 37(10):1659–1668. [PubMed: 22444624]
- Coan JA. Adult attachment and the brain. *Journal of Social and Personal Relationships*. 2010; 27(2): 210–217.
- Costa PT, McCrae RR. Normal personality assessment in clinical practice: The NEO Personality Inventory. *Psychological Assessment*. 1992; 4(1):5–13.
- Cuthbert BN, Kozak MJ. Constructing constructs for psychopathology: the NIMH research domain criteria. *Journal of Abnormal Psychology*. 2013; 122(3):928–937. [PubMed: 24016027]
- Daniel SIF, Poulsen S, Lunn S. Client attachment in a randomized clinical trial of psychoanalytic and cognitive-behavioral psychotherapy for bulimia nervosa: Outcome moderation and change. *Psychotherapy*. 2016; 53(2):174–184. [PubMed: 26950692]
- Del Giudice M, Ellis BJ, Shirtcliff EA. The Adaptive Calibration Model of stress responsivity. *Neuroscience and Biobehavioral Reviews*. 2011; 35(7):1562–1592. [PubMed: 21145350]
- Del Giudice M, Hinnant JB, Ellis BJ, El-Sheikh M. Adaptive patterns of stress responsivity: a preliminary investigation. *Developmental Psychology*. 2012; 48(3):775–790. [PubMed: 22148947]
- Diamond LM, Fagundes CP. Psychobiological research on attachment. *Journal of Social and Personal Relationships*. 2010; 27(2):218–225.
- Diamond LM. Contributions of Psychophysiology to Research on Adult Attachment: Review and Recommendations. *Personality and Social Psychology Review*. 2001; 5(4):276–295.
- Dykas MJ, Cassidy J. Attachment and the processing of social information across the life span: theory and evidence. *Psychological Bulletin*. 2011; 137(1):19–46. [PubMed: 21219056]
- Edwards VJ, Holden GW, Felitti VJ, Anda RF. Relationship between multiple forms of childhood maltreatment and adult mental health in community respondents: results from the adverse childhood experiences study. *The American Journal of Psychiatry*. 2003; 160(8):1453–1460. [PubMed: 12900308]
- Ehler U. Enduring psychobiological effects of childhood adversity. *Psychoneuroendocrinology*. 2013; 38(9):1850–1857. [PubMed: 23850228]
- Ehrenthal JC, Dinger U, Lamla A, Funken B, Schauenburg H. Evaluation of the German version of the attachment questionnaire “Experiences in Close Relationships– Revised” (ECR-RD). *Psychotherapie, Psychosomatik, medizinische Psychologie*. 2009; 59(6):215–223.
- Ehrenthal JC, Herrmann-Lingen C, Fey M, Schauenburg H. Altered cardiovascular adaptability in depressed patients without heart disease. *The World Journal of Biological Psychiatry*. 2010; 11(3): 586–593. [PubMed: 20218928]
- Ein-Dor T, Mikulincer M, Shaver PR. Attachment insecurities and the processing of threat-related information: Studying the schemas involved in insecure people’s coping strategies. *Journal of Personality and Social Psychology*. 2011; 101(1):78–93. [PubMed: 21355660]
- Essex MJ, Shirtcliff EA, Burk LR, Ruttle PL, Klein MH, Slattery MJ, et al. Influence of early life stress on later hypothalamic-pituitary-adrenal axis functioning and its covariation with mental health symptoms: a study of the allostatic process from childhood into adolescence. *Development and Psychopathology*. 2011; 23(4):1039–1058. [PubMed: 22018080]
- Fairchild AJ. Investigating Validity Evidence for the Experiences in Close Relationships-Revised Questionnaire. *Educational and Psychological Measurement*. 2006; 66(1):116–135.
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, et al. Relationship of Childhood Abuse and Household Dysfunction to Many of the Leading Causes of Death in Adults. *American Journal of Preventive Medicine*. 1998; 14(4):245–258. [PubMed: 9635069]
- First, MB., Spitzer, RL., Gibbon, M., Williams, JB. Structured clinical interview for DSM-IV clinical version (SCID-I/CV). New York, NY: Biometrics Research Department, New York State Psychiatric; 1997.
- Fischer-Kern M, Doering S, Taubner S, Horz S, Zimmermann J, Rentrop M, et al. Transference-focused psychotherapy for borderline personality disorder: Change in reflective function. *The British Journal of Psychiatry*. 2015; 207(2):173–174. Retrieved April 12, 2016. [PubMed: 25999334]

- Foley P, Kirschbaum C. Human hypothalamus-pituitary-adrenal axis responses to acute psychosocial stress in laboratory settings. *Neuroscience and Biobehavioral Reviews*. 2010; 35(1):91–96. [PubMed: 20109491]
- Fonagy, Peter, Gergely, György, Jurist Elliot, J., Target, Mary. *Affect Regulation, Mentalization and the Development of the Self*. New York, NY: Karnac Books; 2002.
- Ford JD, Courtois CA. Complex PTSD, affect dysregulation, and borderline personality disorder. *Borderline Personality Disorder and Emotion Dysregulation*. 2014; 1:9. [PubMed: 26401293]
- Fossati A, Madeddu F, Maffei C. Borderline Personality Disorder and Childhood Sexual Abuse: A Meta-Analytic Study. *Journal of Personality Disorders*. 1999; 13(3):268–280. [PubMed: 10498039]
- Fraley RC, Waller NG, Brennan KA. An item response theory analysis of self-report measures of adult attachment. *Journal of Personality and Social Psychology*. 2000; 78(2):350–365. [PubMed: 10707340]
- Garson, GD. *Hierarchical Linear Modeling: Guide and Applications*. Thousand Oaks, CA: Sage Publications; 2013.
- Gillath O, Giesbrecht B, Shaver PR. Attachment, attention, and cognitive control: Attachment style and performance on general attention tasks. *Journal of Experimental Social Psychology*. 2009; 45(4):647–654.
- Goldstein BL, Klein DN. A review of selected candidate endophenotypes for depression. *Clinical Psychology Review*. 2014; 34(5):417–427. [PubMed: 25006008]
- Gordon I, Zagoory-Sharon O, Schneiderman I, Leckman JF, Weller A, Feldman R. Oxytocin and cortisol in romantically unattached young adults: associations with bonding and psychological distress. *Psychophysiology*. 2008; 45(3):349–352. [PubMed: 18266803]
- Granger DA, Hibel LC, Fortunato CK, Kapelewski CH. Medication effects on salivary cortisol: tactics and strategy to minimize impact in behavioral and developmental science. *Psychoneuroendocrinology*. 2009; 34(10):1437–1448. [PubMed: 19632788]
- Granger DA, Kivlighan KT, El-Sheikh M, Gordis EB, Stroud LR. Salivary alpha-amylase in biobehavioral research: recent developments and applications. *Annals of the New York Academy of Sciences*. 2007; 1098:122–144. [PubMed: 17332070]
- Gunnar MR, Brodersen L, Nachmias M, Buss K, Rigatuso J. Stress reactivity and attachment security. *Developmental Psychobiology*. 1996; 29(3):191–204. [PubMed: 8666128]
- Halford C, Anderzén I, Arnetz B. Endocrine measures of stress and self-rated health: A longitudinal study. *Journal of Psychosomatic Research*. 2003; 55(4):317–320. [PubMed: 14507542]
- Haltigan JD, Roisman GI, Fraley RC. The predictive significance of early caregiving experiences for symptoms of psychopathology through midadolescence: enduring or transient effects? *Development and Psychopathology*. 2013; 25(1):209–221. [PubMed: 22800694]
- Harkness KL, Bruce AE, Lumley MN. The role of childhood abuse and neglect in the sensitization to stressful life events in adolescent depression. *Journal of Abnormal Psychology*. 2006; 115(4):730–741. [PubMed: 17100530]
- Harned MS. The Combined Treatment of PTSD with Borderline Personality Disorder. *Current Treatment Options in Psychiatry*. 2014; 1(4):335–344.
- Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*. 2001; 49(12):1023–1039. [PubMed: 11430844]
- Herman JL, Perry JC, van der Kolk BA. Childhood trauma in borderline personality disorder. *The American Journal of Psychiatry*. 1989; 146(4):490–495. [PubMed: 2929750]
- Holleman M, Vreeburg SA, Dekker JJM, Penninx BWJH. The relationships of working conditions, recent stressors and childhood trauma with salivary cortisol levels. *Psychoneuroendocrinology*. 2012; 37(6):801–809. [PubMed: 22000684]
- Holz E, Lass-Hennemann J, Streb M, Pfaltz M, Michael T. Effects of acute cortisol administration on perceptual priming of trauma-related material. *PloS one*. 2014; 9(9):e104864. [PubMed: 25192334]

- Houben M, Bohus M, Santangelo PS, Ebner-Priemer U, Trull TJ, Kuppens P. The specificity of emotional switching in borderline personality disorder in comparison to other clinical groups. *Personality Disorders*. 2016; 7(2):198–204. [PubMed: 26882282]
- Hox, J. *Multilevel analysis*. Mahwah, NJ: Lawrence Erlbaum; 2002.
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *The American Journal of Psychiatry*. 2010; 167(7):748–751. [PubMed: 20595427]
- Jones DR, Macias C, Barreira PJ, Fisher WH, Hargreaves WA, Harding CM. Prevalence, severity, and co-occurrence of chronic physical health problems of persons with serious mental illness. *Psychiatric Services*. 2004; 55(11):1250–1257. [PubMed: 15534013]
- Juster RP, Bizik G, Picard M, Arseneault-Lapierre G, Sindi S, Trepanier L, et al. A transdisciplinary perspective of chronic stress in relation to psychopathology throughout life span development. *Development and Psychopathology*. 2011; 23(3):725–776. [PubMed: 21756430]
- Juster RP, McEwen BS, Lupien SJ. Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience and biobehavioral reviews*. 2010; 35(1):2–16. [PubMed: 19822172]
- Karatsoreos IN, McEwen BS. Psychobiological allostasis: resistance, resilience and vulnerability. *Trends in Cognitive Sciences*. 2011; 15(12):576–584. [PubMed: 22078931]
- Kidd T, Hamer M, Steptoe A. Examining the association between adult attachment style and cortisol responses to acute stress. *Psychoneuroendocrinology*. 2011; 36(6):771–779. [PubMed: 21106296]
- Kidd T, Hamer M, Steptoe A. Adult attachment style and cortisol responses across the day in older adults. *Psychophysiology*. 2013; 50(9):841–847. [PubMed: 23808770]
- Kirschbaum C, Pirke KM, Hellhammer DH. The ‘Trier Social Stress Test’ – A Tool for Investigating Psychobiological Stress Responses in a Laboratory Setting. *Neuropsychobiology*. 2004; 28(1–2): 76–81.
- Klaassens ER, Giltay EJ, Cuijpers P, van Veen T, Zitman FG. Adulthood trauma and HPA-axis functioning in healthy subjects and PTSD patients: a meta-analysis. *Psychoneuroendocrinology*. 2012; 37(3):317–331. [PubMed: 21802212]
- Köhling J, Moessner M, Ehrenthal JC, Bauer S, Cierpka M, Kammerer A, et al. Affective Instability and Reactivity in Depressed Patients With and Without Borderline Pathology. *Journal of Personality Disorders*. 2015:1–20.
- Koolhaas JM, Bartolomucci A, Buwalda B, de Boer SF, Flugge G, Korte SM, et al. Stress revisited: a critical evaluation of the stress concept. *Neuroscience and Biobehavioral Reviews*. 2011; 35(5): 1291–1301. [PubMed: 21316391]
- Kudielka BM, Gierens A, Hellhammer DH, Wust S, Schlotz W. Salivary cortisol in ambulatory assessment—some dos, some don'ts, and some open questions. *Psychosomatic Medicine*. 2012; 74(4):418–431. [PubMed: 22582339]
- Kudielka BM, Hellhammer DH, Wust S. Why do we respond so differently? Reviewing determinants of human salivary cortisol responses to challenge. *Psychoneuroendocrinology*. 2009; 34(1):2–18. [PubMed: 19041187]
- Lenzenweger MF. Thinking clearly about the endophenotype-intermediate phenotype-biomarker distinctions in developmental psychopathology research. *Development and Psychopathology*. 2013; 25(4 Pt 2):1347–1357. [PubMed: 24342844]
- Levy KN. The implications of attachment theory and research for understanding borderline personality disorder. *Development and Psychopathology*. 2005; 17(04)
- Levy KN, Johnson BN, Clouthier TL, Scala JW, Temes CM. An attachment theoretical framework for personality disorders. *Canadian Psychology/Psychologie Canadienne*. 2015; 56(2):197–207.
- Levy KN, Meehan KB, Kelly KM, Reynoso JS, Weber M, Clarkin JF, Kernberg OF. Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. *Journal of Consulting and Clinical Psychology*. 2006; 74(6):1027–1040. [PubMed: 17154733]
- Lilienfeld SO. The Research Domain Criteria (RDoC): an analysis of methodological and conceptual challenges. *Behaviour Research and Therapy*. 2014; 62:129–139. [PubMed: 25156396]
- Linehan, MM. *Cognitive-behavioral treatment of borderline personality disorder*. New York, NY: Guilford Press; 1993.

- Liotti G, Pasquini P, Cirrincione R. Predictive factors for borderline personality disorder: Patients' early traumatic experiences and losses suffered by the attachment figure. *Acta Psychiatrica Scandinavica*. 2000; 102(4):282–289. [PubMed: 11089728]
- Llabre MM, Spitzer SB, Saab PG, Schneiderman N. Piecewise latent growth curve modeling of systolic blood pressure reactivity and recovery from the cold pressor test. *Psychophysiology*. 2001; 38(6):951–960. [PubMed: 12240671]
- Llera SJ, Newman MG. Rethinking the role of worry in generalized anxiety disorder: evidence supporting a model of emotional contrast avoidance. *Behavior Therapy*. 2014; 45(3):283–299. [PubMed: 24680226]
- Loranger, AW. *International Personality Disorder Examination (IPDE) manual*. Odessa, FL: Psychological Assessment Resources, Inc; 1999.
- Markowitz JC, Petkova E, Neria Y, van Meter PE, Zhao Y, Hembree E, et al. Is Exposure Necessary? A Randomized Clinical Trial of Interpersonal Psychotherapy for PTSD. *The American Journal of Psychiatry*. 2015; 172(5):430–440. [PubMed: 25677355]
- McEwen BS, Wingfield JC. What is in a name? Integrating homeostasis, allostasis and stress. *Hormones and Behavior*. 2010; 57(2):105–111. [PubMed: 19786032]
- Mikulincer M. Adult attachment style and affect regulation: Strategic variations in self-appraisals. *Journal of Personality and Social Psychology*. 1998; 75(2):420–435. [PubMed: 9731317]
- Mikulincer, M., Shaver, PR., Cassidy, J., Berant, E. Attachment-related defensive processes. In: Obegi, JH., Berant, E., editors. *Attachment theory and research in clinical work with adults*. New York: Guilford Press; 2009. p. 293-327.
- Mikulincer M, Shaver PR, Pereg D. Attachment theory and affect regulation: The dynamics, development, and cognitive consequences of attachment-related strategies. *Motivation and Emotion*. 2003; 27(2):77–102.
- Morris MC, Compas BE, Garber J. Relations among posttraumatic stress disorder, comorbid major depression, and HPA function: a systematic review and meta-analysis. *Clinical Psychology Review*. 2012; 32(4):301–315. [PubMed: 22459791]
- Nater UM, Rohleder N. Salivary alpha-amylase as a non-invasive biomarker for the sympathetic nervous system: current state of research. *Psychoneuroendocrinology*. 2009; 34(4):486–496. [PubMed: 19249160]
- Nicholson EL, Bryant RA, Felmingham KL. Interaction of noradrenaline and cortisol predicts negative intrusive memories in posttraumatic stress disorder. *Neurobiology of Learning and Memory*. 2014; 112:204–211. [PubMed: 24296460]
- Ottaviani C, Shapiro D, Couyoumdjian A. Flexibility as the key for somatic health: From mind wandering to perseverative cognition. *Biological Psychology*. 2013; 94(1):38–43. [PubMed: 23680439]
- Papousek I, Paechter M, Lackner HK. Delayed psychophysiological recovery after self-concept-inconsistent negative performance feedback. *International Journal of Psychophysiology*. 2011; 82(3):275–282. [PubMed: 21963528]
- Pierrehumbert B, Torrisi R, Glatz N, Dimitrova N, Heinrichs M, Halfon O. The influence of attachment on perceived stress and cortisol response to acute stress in women sexually abused in childhood or adolescence. *Psychoneuroendocrinology*. 2009; 34(6):924–938. [PubMed: 19217212]
- Quirin M, Pruessner JC, Kuhl J. HPA system regulation and adult attachment anxiety: Individual differences in reactive and awakening cortisol. *Psychoneuroendocrinology*. 2008; 33(5):581–590. [PubMed: 18329180]
- Radley JJ, Kabbaj M, Jacobson L, Heydendael W, Yehuda R, Herman JP. Stress risk factors and stress-related pathology: neuroplasticity, epigenetics and endophenotypes. *Stress*. 2011; 14(5):481–497. [PubMed: 21848436]
- Rifkin-Graboi A. Attachment status and salivary cortisol in a normal day and during simulated interpersonal stress in young men. *Stress*. 2008; 11(3):210–224. [PubMed: 18465468]
- Roisman GI, Holland A, Fortuna K, Fraley RC, Clausell E, Clarke A. The Adult Attachment Interview and self-reports of attachment style: an empirical rapprochement. *Journal of Personality and Social Psychology*. 2007; 92(4):678–697. [PubMed: 17469952]

- Rusk N, Rothbaum F. From stress to learning: Attachment theory meets goal orientation theory. *Review of General Psychology*. 2010; 14(1):31–43.
- Sadikaj G, Moskowitz DS, Zuroff DC. Attachment-related affective dynamics: differential reactivity to others' interpersonal behavior. *Journal of Personality and Social Psychology*. 2011; 100(5):905–917. [PubMed: 21443369]
- Santangelo PS, Limberger MF, Stiglmayr C, Houben M, Coosemans J, Verleysen G, et al. Analyzing subcomponents of affective dysregulation in borderline personality disorder in comparison to other clinical groups using multiple e-diary datasets. *Borderline Personality Disorder and Emotion Dysregulation*. 2016; 3:5. [PubMed: 27386138]
- Sbarra DA, Hazan C. Coregulation, dysregulation, self-regulation: an integrative analysis and empirical agenda for understanding adult attachment, separation, loss, and recovery. *Personality and Social Psychology Review*. 2008; 12(2):141–167. [PubMed: 18453476]
- Scheidt CE, Waller E, Malchow H, Ehler U, Becker-Stoll F, Schulte-Mönting J, Lücking CH. Attachment Representation and Cortisol Response to the Adult Attachment Interview in Idiopathic Spasmodic Torticollis. *Psychotherapy and Psychosomatics*. 2000; 69(3):155–162. [PubMed: 10773780]
- Scher CD, Stein MB, Asmundson GJ, McCreary DR, Forde DR. The childhood trauma questionnaire in a community sample: psychometric properties and normative data. *Journal of Traumatic Stress*. 2001; 14(4):843–857. [PubMed: 11776429]
- Scott LN, Levy KN, Granger DA. Biobehavioral reactivity to social evaluative stress in women with borderline personality disorder. *Personality Disorders*. 2013; 4(2):91–100. [PubMed: 23244772]
- Sibley CG, Fischer R, Liu JH. Reliability and validity of the revised experiences in close relationships (ECR-R) self-report measure of adult romantic attachment. *Personality & Social Psychology Bulletin*. 2005; 31(11):1524–1536. [PubMed: 16207771]
- Skodol AE, Clark LA, Bender DS, Krueger RF, Morey LC, Verheul R, et al. Proposed changes in personality and personality disorder assessment and diagnosis for DSM-5 Part I: Description and rationale. *Personality Disorders*. 2011; 2(1):4–22. [PubMed: 22448687]
- Smeets T. Autonomic and hypothalamic-pituitary-adrenal stress resilience: Impact of cardiac vagal tone. *Biological Psychology*. 2010; 84(2):290–295. [PubMed: 20206227]
- Smyth N, Thorn L, Oskis A, Hucklebridge F, Evans P, Clow A. Anxious attachment style predicts an enhanced cortisol response to group psychosocial stress. *Stress*. 2015; 18(2):143–148. [PubMed: 25758939]
- Steinert C, Bumke PJ, Hollekamp RL, Larisch A, Leichsenring F, Matthes H, et al. Resource activation for treating post-traumatic stress disorder, co-morbid symptoms and impaired functioning: a randomized controlled trial in Cambodia. *Psychological Medicine*. 2016:1–12.
- Tops M, van Peer JM, Korf J, Wijers AA, Tucker DM. Anxiety, cortisol, and attachment predict plasma oxytocin. *Psychophysiology*. 2007; 44(3):444–449. [PubMed: 17371496]
- Tragesser SL, Solhan M, Schwartz-Mette R, Trull TJ. The role of affective instability and impulsivity in predicting future BPD features. *Journal of Personality Disorders*. 2007; 21(6):603–614. [PubMed: 18072862]
- Tukey, JW. *Exploratory data analysis*. Don Mills, Ontario: Addison-Wesley; 1977.
- van Voorhees EE, Dennis MF, Calhoun PS, Beckham JC. Association of DHEA, DHEAS, and cortisol with childhood trauma exposure and post-traumatic stress disorder. *International Clinical Psychopharmacology*. 2014; 29(1):56–62. [PubMed: 23907073]
- Wingenfeld K, Spitzer C, Rullkötter N, Lowe B. Borderline personality disorder: hypothalamus pituitary adrenal axis and findings from neuroimaging studies. *Psychoneuroendocrinology*. 2010; 35(1):154–170. [PubMed: 19837517]
- Zanarini MC, Frankenburg FR. The essential nature of borderline psychopathology. *Journal of Personality Disorders*. 2007; 21(5):518–535. [PubMed: 17953504]
- Zimmerman DJ, Choi-Kain LW. The hypothalamic-pituitary-adrenal axis in borderline personality disorder: a review. *Harvard Review of Psychiatry*. 2009; 17(3):167–183. [PubMed: 19499417]
- Zoladz PR, Diamond DM. Current status on behavioral and biological markers of PTSD: a search for clarity in a conflicting literature. *Neuroscience and Biobehavioral Reviews*. 2013; 37(5):860–895. [PubMed: 23567521]

Table 1

Participants Characteristics

Variables	M (sd) or %
Age	25.86 (8.33)
Race white/caucasian	77.9 %
Highest level of education	
High School Graduate or GED	42.5%
> High School Graduate	57.5 %
Currently employed (full or part-time)	40.7 %
Relationship status single	75.2 %
Number of medication	1.21 (1.96)
Number of BPD criteria	2.10 (2.74)
Number of non-BPD Axis-II diagnoses	.15 (.36)
Number of Axis-I diagnoses	.46 (.87)
NEO Angry hostility	56.97 (13.42)
NEO Anxiety	55.01 (12.13)
NEO Depression	55.29 (15.18)
NEO Impulsiveness	51.96 (12.66)
CTQ Total score	42.42 (19.58)
ECR-R Attachment anxiety	3.18 (1.37)
ECR-R Attachment avoidance	3.06 (1.22)

Note. GED = General Educational Development Test; BPD = Borderline Personality Disorder; NEO = Revised NEO Personality Inventory; CTQ = Childhood Trauma Questionnaire; ECR-R = Experiences in Close Relationships - Revised Questionnaire.

Table 2

Cortisol Reaction to the TSST

	Resting		Reactivity		Recovery	
	Coeff.	t	Coeff.	T	Coeff.	t
Level 1						
Time	-.032	-3.351**	.210	7.422***	-.075	-5.672***
Level 2						
CTQ	.004	1.213	.004	1.318	.004	1.243
ECR-R Anxiety	.023	.468	.041	.876	.034	.734
ECR-R Avoidance	.006	.126	.003	.055	.001	.025
CTQ*ECR-R Anxiety	-.0002	-.097	-.001	-.380	-.002	-.556
CTQ*ECR-R Avoidance	-.001	-.470	-.0001	-.042	<.0001	-.005
Time*CTQ	-.0002	-.397	-.001	-1.322	-.001	-1.328
Time*ECR-R Anxiety	.004	.499	-.030	-1.565	-.008	-.911
Time*ECR-R Avoidance	-.003	-.347	-.006	-.301	-.001	-.139
Time*CTQ*ECR-R Anxiety	-.001	-1.625	-.002	-2.031*	-.001	-1.444
Time*CTQ*ECR-R Avoidance	.001	1.484	.0005	.450	.0001	.365

Note: , p < .10;

* p < .05;

** p < .01;

*** p < .001.

CTQ = Childhood Trauma Questionnaire; ECR-R = Experiences in Close Relationships - Revised Questionnaire. All models controlled for the respective other two time-variables (i.e., resting, reactivity, recovery), and specific covariates that contributed significantly (p < .10) to the model as specified in the methods and results section.

Table 3

Alpha-Amylase Reaction to the TSSST

	Resting		Reactivity		Recovery	
	Coeff.	T	Coeff.	t	Coeff.	T
Level 1						
Time	-.115	-1.328	.963	7.893***	-.788	-10.134***
Level 2						
CTQ	.015	.708	.0001	.006	-.0001	-.008
ECR-R Anxiety	.016	.527	.243	.859	.164	.550
ECR-R Avoidance	.679	2.339*	.692	2.622*	.676	2.174*
CTQ*ECR-R Anxiety	.021	1.615	.025	2.081*	.028	2.168*
CTQ*ECR-R Avoidance	.041	2.686**	.044	3.149**	.043	2.825**
Time*CTQ	-.007	-1.548	-.002	-.405	-.001	-.260
Time*ECR-R Anxiety	-.001	-.019	-.050	-.817	-.002	-.047
Time*ECR-R Avoidance	-.002	-.029	-.0011	-.174	.016	.369
Time*CTQ*ECR-R Anxiety	-.001	-1.548	-.008	-2.795**	-.005	-2.526*
Time*CTQ*ECR-R Avoidance	-.004	-1.503	.004	1.148	.002	1.055

Note: , p < .10;

* p < .05;

** p < .01;

*** p < .001.

CTQ = Childhood Trauma Questionnaire; ECR-R = Experiences in Close Relationships - Revised Questionnaire. All models controlled for the respective other two time-variables (i.e., resting, reactivity, recovery), and specific covariates that contributed significantly (p < .10) to the model as specified in the methods and results section.