

Attachment tests of *Pasteuria penetrans* to the cuticle of plant and animal parasitic nematodes, free living nematodes and *srf* mutants of *Caenorhabditis elegans*

P. Mendoza de Gives^{1,2,3*}, K.G. Davies¹, M. Morgan²
and J.M. Behnke²

¹Entomology and Nematology Department, IACR-Rothamsted, Harpenden, Herts, AL5 2JQ, UK; ²Department of Life Science, University of Nottingham, University Park, Nottingham, NG2 2RD, UK; ³Centro Nacional de Investigaciones Disciplinarias en Parasitología Veterinaria, CENID-PAVET, Km 11.5 Carr. Fed. Cuernavaca-Cuautla, Col. Progreso, Municipio de Jiutepec, Estado de Morelos, Ap. Postal 206, CIVAC, Morelos, México, 62500

Abstract

Populations of *Pasteuria penetrans* isolated from root-knot nematodes (*Meloidogyne* spp.) and cyst nematodes (*Heterodera* spp.) were tested for their ability to adhere to a limited selection of sheathed and exsheathed animal parasitic nematodes, free living nematodes, including *Caenorhabditis elegans* wild type and several *srf* mutants, and plant parasitic nematodes. The attachment of spores of *Pasteuria* was restricted and no spores were observed adhering to any of the animal parasitic nematodes either with or without their sheath or to any of the free living nematodes including *C. elegans* and the *srf* mutants. All spore attachment was restricted to plant parasitic nematodes; however, spores isolated from cyst nematodes showed the ability to adhere to other genera of plant parasitic nematodes which was not the case with spores isolated from root-knot nematodes. The results are discussed in relationship to cuticular heterogeneity.

Introduction

The control of human and animal nematode infections is largely based on the administration of anthelmintic drugs. However, in response to the intensive use of anthelmintics, resistance has been reported (Waller, 1990; Jackson, 1993; De Clercq *et al.*, 1997; Reynoldson *et al.*, 1998) and this has led to a search for alternative strategies.

Nematophagous fungi are currently being evaluated for their potential to control plant parasitic (Kerry, 1993; Kerry & Bourne, 1996) and animal parasitic nematodes (Waller, 1993; Mendoza de Gives *et al.*, 1994; Wolstrup *et al.*, 1996; Morgan *et al.*, 1997; Llerandi-Juarez & Mendoza de Gives, 1998). However, to date, there are very few reports evaluating the use of bacteria to control these animal parasites. The *Pasteuria* group of Gram-positive endospore-forming bacteria are parasites of nematodes and water fleas (*Daphnia* spp.). Certain basic morphological types of spore, each with a variety of sub-types, have been identified and are found almost exclusively in Tylenchida and Dorylaimida; however, others have recently been found associated with Araeolaimida, Chromadorida, and Enoplida

*Address for correspondence: Centro Nacional de Investigaciones Disciplinarias en Parasitología Veterinaria, CENID-PAVET, Km 11.5 Carr. Fed. Cuernavaca-Cuautla, Col. Progreso, Municipio de Jiutepec, Estado de Morelos, Ap. Postal 206, CIVAC, Morelos, México, 62500.
Fax: 73 20 55 44

(Sturhan, 1996). Four species of *Pasteuria* have been described so far: (i) *P. penetrans* parasitic on *Meloidogyne incognita* and probably other root-knot nematodes (Sayre & Starr, 1985); (ii) *P. thornei* on *Pratylenchus brachyurus* (Sayre & Starr, 1988); (iii) *P. nishizawae* which parasitizes cyst nematodes (Sayre *et al.*, 1991); and (iv) *P. ramosa*, a parasite of water fleas *Daphnia* spp. (Sayre *et al.*, 1983; Ebert *et al.*, 1996).

Most research on *P. penetrans* has concentrated on its interaction with root-knot nematodes (*Meloidogyne* spp.) and it has been shown to have potential to control root-knot nematodes (Stirling, 1991). The life cycle of *P. penetrans* commences when endospores attach to the cuticle of motile larvae as they migrate through the soil. Spores either germinate and penetrate the second-stage juvenile before the nematode has infected a plant root, as in the case of spores adhering to the cuticle of *Heterodera avenae* (Davies *et al.*, 1990), or after the nematode has infected a plant root and started feeding (Sayre & Starr, 1985, 1988). In both cases, the spores each produce a germ tube which penetrates the nematode cuticle and produces a dichotomously branched microcolony. These microcolonies subsequently divide and proliferate throughout the pseudocoelom eventually killing the nematode and producing a cadaver filled with spores (Sayre & Starr, 1988). *Pasteuria* spores have been shown to differ in their ability to adhere to the cuticle of plant parasitic nematodes (Davies *et al.*, 1988; Stirling, 1991) and the interaction between the nematode and *Pasteuria* is thought to involve a protein/carbohydrate like mechanism between the spore and the nematode cuticle (Davies *et al.*, 1994). There are no reports of isolates of the bacteria being tested against either animal parasitic or free living nematodes, and *Caenorhabditis elegans* represents the latter category where mutants are available which differ solely in the surface characteristics of their cuticle.

This paper reports the results of tests to study the attachment of spores, of selected *Pasteuria* isolates, to different species of animal parasitic (with and without sheath) and free living nematodes, to plant parasitic nematodes and surface (*srf*) mutants of *Caenorhabditis elegans*, the cuticles of which react differently to either antibodies and/or lectins.

Material and methods

Nematodes

The plant-parasitic nematodes *Meloidogyne* spp. and *Rotylenchulus reniformis* were obtained from plant cultures maintained in the glasshouse at 25°C on tomato plants, cv. Pixie, grown in a peat/sand (1:1, v/v) compost. *Pratylenchus* spp. and *Radopholus similis* were obtained from axenic maize root cultures (Hooper, 1986b). Nematodes were hatched from infected root material by placing small samples of infected root material in tap water on a small sieve in a tray of water at room temperature (Hooper, 1986a). The juveniles of cyst forming nematodes were obtained by incubating cysts at optimum temperatures in tap water and, in the case of the two species of potato cyst nematode, *Globodera pallida* and *G. rostochiensis*, in the presence of potato root diffusate. *Aphelenchoides*

spp. and *Ditylenchus* spp. were obtained from axenic Petri dish cultures of *Botrytis cinerea* maintained at room temperature in the laboratory by washing the surface of the agar with water (Hooper, 1986b). Samples of *Anguina tritici* were obtained by breaking open infected grains of wheat in a small drop of tap water to release the nematodes (Hooper, 1986a).

Cultures of the animal parasitic nematodes *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta* and *Trichostrongylus axei* were provided by Drs E. Munn, Babraham Institute, Cambridge, and R. Coop, The Moredun Research Institute, Edinburgh. *Ancylostoma ceylanicum* and *Heligmosomoides polygyrus* were obtained from the faecal material of infected hamsters and mice respectively (Garside & Behnke, 1989). *Steinernema* and *Heterorhabditis* were cultured in *Galleria* larvae. The free-living nematodes *Panagrellus redivivus*, *Pelodera strongyloides*, *Diplogaster* sp., *Mesodiplogaster* sp., *Panagrolaimus* sp., and *Rhabditis* sp. were obtained from axenic Petri dish cultures maintained at room temperature in the laboratory. The wild type culture of *Caenorhabditis elegans* (N2) was provided by Dr Julie Arhinger, Medical Research Council, Cambridge and the surface mutants AT6, AT10 and CL261 (table 2) were obtained from Dr Theresa Stiernagle, *Caenorhabditis* Genetics Center, University of Minnesota and were maintained in Petri dishes seeded with *E. coli* strain OP50 following the method of Wood (1988).

Bacterial cultures

Populations of *Pasteuria* were obtained from the species of nematode from which they were originally isolated growing on a suitable host plant. Either, the infected roots were dried and the powder produced following the method of Stirling & Wachtel (1980), or *Pasteuria* infected nematode cadavers were collected from field soils. The latter were recognized using a dissecting microscope and identifying females present on or in roots but not producing egg masses (Sharma & Davies, 1996). Suspensions of spores were prepared by grinding either *Pasteuria* infested root powder, or infected cadavers, in tap water with a pestle and mortar. The spores were filtered with a 10 µm sieve, counted using a haemocytometer slide and the concentrations of suspensions were adjusted to 10⁶ spores/ml. Stock suspensions were stored frozen at -20°C.

Attachment tests

Samples (250 µl) of spore suspensions of each of the stock *Pasteuria* populations were placed in separate siliconized Eppendorf tubes together with a 250 µl of a suspension of the test nematode population containing approximately 500 individuals. The nematodes and spores were thoroughly mixed and an attachment test performed by centrifugation (10 000 g for 5 min) following the method of Hewlett & Dickson (1993). A semi-quantitative score (0, no spores per nematode; +, 1-10 spores per nematode; ++, 11-40 spores per nematode) was given for each population of nematode tested, assessing a minimum of 25 nematodes for each nematode population, using a light microscope (×400).

Table 1. Animal parasitic nematodes, third stage larvae with and without sheath, to which no spores of the bacterial hyperparasite *Pasteuria penetrans* adhered.

Genus/species	Origin ¹	<i>Pasteuria</i> populations tested
<i>Haemonchus contortus</i>	BI	PP1 ² , PP3O ³
<i>Heligmosomoides polygyrus</i>	UN	PP1, PP3A ⁴ , PP3O, B7 ² , PA ⁴
<i>Ostertagia circumcincta</i>	MRI	PP1, PP3O
<i>Trichostrongylus axei</i>	MRI	PP1, PP3O
<i>Ancylostoma ceylanicum</i>	UN	PP1, PP3A, PP3O, B7, PA
<i>Steinernema feltiae</i>	IACR	PP1, PPJ ³ , PPC ⁵
<i>Heterorhabditis megidis</i>	IACR	PP1, PPJ, PPC

¹BI, Babraham Institute, Cambridge; UN, University of Nottingham; MRI, Moredun Research Institute, Edinburgh; IACR, Institute of Arable Crops Research.
²Spores originating from *Meloidogyne incognita*; ³spores originating from *M. javanica*; ⁴spores originating from *M. arenaria*; ⁵spores originating from *Heterodera cajani*.

Results and discussion

No *Pasteuria* spores were observed adhering to any of the 3rd stage infective larvae of the animal parasitic nematodes either with or without their sheath (table 1) or to any of the free living nematodes including *C. elegans* and three *srf* mutants (table 2). All populations of *Pasteuria* used in these experiments had been isolated from plant parasitic nematodes and their attachment was restricted to plant parasitic nematodes (table 3). Attachment of those populations of spores isolated from root-knot nematodes (*Meloidogyne* spp.) was similarly restricted to root-knot nematodes, however, those isolated from the genus *Heterodera* appeared to have a broader range of hosts and all three *Pasteuria* populations, PPC, PPN and PPW also attached to *Globodera*. One population of spores, PPC, was also observed attaching to *Pratylenchus*, *Radopholus*, *Rotylenchulus* and *Aphelenchoides*; the attachment of these spores also exhibited interspecific variation between species within genera (table 3). It is interesting to note that the populations of *Pasteuria* from the apomictic populations of nematodes, i.e. the root-knot populations, appear to have a more restricted host range

than those isolated from the cyst nematode populations which are amphimictic.

There are two fundamental problems in the deployment of *Pasteuria* as a biological nematicide, firstly, the inability to culture large populations of spores (Williams *et al.*, 1989; Bishop and Ellar, 1991) and secondly, its host specificity (Stirling, 1985; Channer & Gowen, 1992; Davies *et al.*, 1988). Populations of *Pasteuria* are found which parasitize all the major genera of plant parasitic nematodes (Sayre and Starr, 1988) and there have recently been reports of other populations which parasitize nematodes in other families and even orders (Sturhan, 1996). Monoclonal antibodies have shown that the surface of the spores of a *Pasteuria* isolate originating from *M. incognita* race 2 was highly heterogeneous, and baiting experiments showed that different sub-populations of spores adhere to different species and races of nematode (Davies *et al.*, 1994). These and subsequent studies (Davies & Redden, 1997) have suggested that the surface properties of the spore are responsible for the virulence of the bacterium and suggest that similar heterogeneity will also be present in the nematode cuticle. As the bacterium infects other invertebrates such as the cladoceran *Moina*

Table 2. Free living nematodes, mixed stages, to which no spores of the bacterial hyperparasite *Pasteuria penetrans* adhered.

Genus/species	Origin ¹	<i>Pasteuria</i> populations tested
<i>Caenorhabditis elegans</i> N2	MRC	PP1 ² , PP3O ³ , B7 ² , PNG ³ , PPN ⁵
Surface mutants	CGC	PP1, PP3O
AT6	CGC	PP1, PP3O
AT10	CGC	PP1, PP3O
CL261	CGC	PP1, PP3O
<i>Panagrellus redivivus</i>	IACR	PP1, PP3O, B7, PNG, PPN
<i>Pelodera strongyloides</i>	IACR	PP1, PP3O, B7, PNG, PPN
<i>Diplogaster</i> sp.	IACR	PP1, PP3O, B7, PNG, PPN
<i>Mesodiplogaster</i> sp.	IACR	PP1, PP3O, B7, PNG, PPN
<i>Panagrolaimus</i> sp.	IACR	PP1, PPA ⁴ , B7, PNG, PPN
<i>Rhabditis</i> sp.	IACR	PP1, PPA, B7, PNG, PPN

¹MRC, Medical Research Council, Cambridge; CGC, *Caenorhabditis* Genetics Center, Minnesota.

²Spores originating from *Meloidogyne incognita*; ³spores originating from *M. javanica*; ⁴spores originating from *M. arenaria*; ⁵spores originating from *Heterodera glycines*.

Table 3. Attachment of spores of six populations of *Pasteuria* (PP1, PPA, PPJ, PPC, PPN, PPW) to the cuticle of second-stage juveniles of plant parasitic nematodes (0, no attachment; +, 1–10 spores; ++, >10 spores; –, not available; based on a mean of 25 individual nematodes).

Genus/species	Origin ¹	PP1 ²	PPA ²	PPJ ³	PPC ⁴	PPN ⁵	PPW ⁶
<i>Meloidogyne incognita</i>	IACR	++	+	++	0	0	0
<i>M. javanica</i>	IACR	++	++	++	0	+	+
<i>M. arenaria</i>	IACR	+	++	+	0	0	0
<i>M. hapla</i>	IACR	+	+	0	0	0	0
<i>Heterodera avenae</i>	IACR	0	0	0	–	–	++
<i>H. schachtii</i>	IACR	0	0	0	++	+	++
<i>H. glycines</i>	IACR	0	0	–	+	+	+
<i>H. cajani</i>	ICRISAT	0	0	0	++	+	–
<i>Globodera rostochiensis</i>	IACR	0	0	0	++	+	+
<i>G. pallida</i>	IACR	0	0	0	++	–	+
<i>Pratylenchus crenatus</i>	IACR	0	0	0	0	0	0
<i>P. neglectus</i>	IACR	0	0	0	0	0	0
<i>P. coffeae</i>	IIP	0	0	0	+	0	–
<i>Radopholus similis</i>	IIP	0	0	0	+	–	–
<i>Rotylenchulus reinformis</i>	ICRISAT	0	0	0	+	–	–
<i>Anguina tritici</i>	IACR	0	0	0	0	–	–
<i>Aphelenchoides</i> sp.	IACR	0	0	0	+	–	–
<i>Ditylenchus</i> sp.	IACR	0	0	0	0	–	–

¹IACR, Institute of Arable Crops Research; ICRISAT, International Crop Research Institute for the Semi-Arid Tropics; IIP, International Institute of Parasitology.

²Spores originating from *Meloidogyne incognita*; ³spores originating from *M. javanica*; ⁴spores originating from *H. cajani*; ⁵spores, *P. nishizawae*, originating from *Heterodera glycines*; ⁶spores originating from *H. avenae*

(Sayre *et al.*, 1977; Ebert *et al.*, 1996) it would seem likely that similar bacteria will be found infecting animal parasitic nematodes, especially those which have to spend prolonged periods of their life cycle in soil before infecting their respective animal hosts, and these infective stages will also exhibit a high level of cuticular heterogeneity. The challenge for the future will therefore be to isolate such bacteria targeting animal and human parasitic nematodes, and to evaluate their potential as tools for the biological control of important human and livestock diseases.

Acknowledgements

P. Mendoza de Gives and M. Morgan were financially supported by Consejo Nacional de Ciencia y Tecnología, México and Lawes Agricultural Trust respectively. IACR-Rothamsted receives grant-aided support from the Biotechnological and Biological Sciences Research Council of the United Kingdom.

References

- Bishop, A.H. & Ellar, D.J. (1991) Attempts to culture *Pasteuria penetrans* *in vitro*. *Biocontrol Science and Technology* **1**, 101–114.
- Channer, A.G. De R. & Gowen, S.R. (1992) Selection for increased resistance and increased pathogen specificity in the *Meloidogyne*–*Pasteuria penetrans* interaction. *Fundamental and Applied Nematology* **15**, 331–339.
- Davies, K.G. & Redden, M. (1997) Diversity and partial characterisation of putative virulence determinants in *Pasteuria penetrans*, the hyperparasitic bacterium of root-knot nematodes (*Meloidogyne* spp.). *Journal of Applied Microbiology* **83**, 227–235.
- Davies, K.G., Kerry, B.R. & Flynn, C.A. (1988) Observations on the pathogenicity of *Pasteuria penetrans*, a parasite of root-knot nematodes. *Annals of Applied Biology* **112**, 491–501.
- Davies, K.G., Flynn, C.A., Laird, V. & Kerry, B.R. (1990) The life-cycle, population dynamics and host specificity of a parasite of *Heterodera avenae*, similar to *Pasteuria penetrans*. *Revue de Nematologie* **13**, 303–309.
- Davies, K.G., Pearson, T. & Redden, M. (1994) Endospore heterogeneity in *Pasteuria penetrans* related to adhesion to plant parasitic nematodes. *Letters in Applied Microbiology* **19**, 370–373.
- De Clercq, D., Sacko, M., Behnke, J., Gilbert, F., Dorny, P. & Vercruyse, J. (1997) Failure of mebendazole in treatment of human hookworm infections in the southern region of Mali. *American Journal of Tropical Medicine and Hygiene* **57**, 25–50.
- Ebert, D., Rainey, P., Embley, M. & Scholz, D. (1996) Development, life cycle, ultrastructure and phylogenetic position of *Pasteuria ramosa* Metchnikoff 1888: rediscovery of an obligate endoparasite of *Daphnia magna* Straus. *Philosophical Transactions of the Royal Society London B* **351**, 1689–1701.
- Garside, P. & Behnke, J.M. (1989) *Ancylostoma ceylanicum*: observations on host–parasite relationship during primary infection. *Parasitology* **98**, 283–289.
- Hewlett, T.E. & Dickson, D.W. (1993) A centrifugation method for attaching endospores of *Pasteuria* spp. to nematodes. *Journal of Nematology* **25** (Supplement), 785–788.
- Hooper, D.J. (1986a) Extraction of nematodes from plant

- material. pp. 51–58 in Southey, J.F. (Ed.) *Laboratory methods for work with plant and soil nematodes*. Ministry of Agriculture, Fisheries and Food, London, HMSO.
- Hooper, D.J.** (1986b) Culturing nematodes and related experimental techniques. pp. 133–137 in Southey, J.F. (Ed.) *Laboratory methods for work with plant and soil nematodes*. Ministry of Agriculture Fisheries and Food, London, HMSO.
- Jackson, F.** (1993) Anthelmintic resistance – the state of play. *British Veterinary Journal* **149**, 123–138.
- Kerry, B.R.** (1993) The use of microbial agents for the biological control of plant parasitic nematodes. pp. 81–104 in Jones, D.G. (Ed.) *Exploitation of microorganisms*. London, Chapman & Hall.
- Kerry, B.R. & Bourne, J.M.** (1996) The importance of rhizosphere interactions in the biological control of plant parasitic nematodes a case study using *Verticillium chlamyosporium*. *Pesticide Science* **47**, 69–75.
- Llerandi-Juarez, R.D. & Mendoza-de Gives, P.** (1998) Resistance of chlamydospores of nematophagous fungi to the digestive processes of sheep in Mexico. *Journal of Helminthology* **72**, 155–158.
- Mendoza de Gives, P., Zavaleta-Mejia, E., Herrera-Rodriguez, D. & Quiroz-Romero, H.** (1994) *In vitro* trapping capability of *Arthrobotrys* spp. on infective larvae of *Haemonchus contortus* and *Nacobus aberrans*. *Journal of Helminthology* **68**, 223–229.
- Morgan, M., Behnke, J.M., Lucas, J.A. & Peberdy, J.F.** (1997) *In vitro* assessment of the influence of nutrition, temperature and larval density on trapping of the infective larvae of *Heligmosomoides polygyrus* by *Arthrobotrys oligospora*, *Duddingtonia flagrans* and *Monacrosporium megalosporum*. *Parasitology* **115**, 303–310.
- Reynoldson, J.A., Behnke, J.M., Pallant, L.J., MacNish, M.G., Gilbert, F. & Thompson, R.C.A.** (1998) Failure of pyrantel to control hookworms in Australia. *Acta Tropica* (in press).
- Sayre, R.M. & Starr, M.P.** (1985) *Pasteuria penetrans* (ex Thorne, 1940) nom. rev., comb. n., sp. n., a mycelial endospore-forming bacterium parasitic in plant-parasitic nematodes. *Proceedings of the Helminthological Society of Washington* **52**, 149–165.
- Sayre, R.M. & Starr, M.P.** (1988) Bacterial diseases and antagonisms of nematodes. pp. 69–101 in Poinar, G.O., Jr. & Jansson, H.B. (Eds) *Diseases of nematodes*, Vol. 1. Boca Raton, USA, CRC Press.
- Sayre, R.M., Wergin, W.P. & Davis, R.E.** (1977) Occurrence in *Moina rectirostis* (Cladocera: Daphnidae) of a parasite morphologically similar to *Pasteuria ramosa* (Metchnikoff, 1988). *Canadian Journal of Microbiology* **23**, 1573–1579.
- Sayre, R.M., Gherna, R.L. & Wergin, W.P.** (1983) Morphological and taxonomic reevaluation of *Pasteuria ramosa* Metchnikoff 1888 and *Bacillus penetrans* Mankau 1975. *International Journal of Systematic Bacteriology* **33**, 636–649.
- Sayre, R.M., Wergin, W.P., Schmidt, J.M. & Starr, M.P.** (1991) *Pasteuria nishizawae* sp. nov., a mycelial and endospore-forming bacterium parasitic on cyst nematodes of genera *Heterodera* and *Globodera*. *Research in Microbiology* **142**, 551–564.
- Sharma, R. & Davies, K.G.** (1996) A comparison of two sympatryx species of *Pasteuria* isolated from a tropical vertisol soil. *World Journal of Microbiology and Biotechnology* **12**, 361–366.
- Stirling, G.R.** (1985) Host specificity of *Pasteuria penetrans* within the genus of *Meloidogyne*. *Nematologica* **31**, 203–209.
- Stirling, G.R.** (1991) *Biological control of plant-parasitic nematodes: progress, problems and prospects*. 282 pp. Wallingford, CAB International.
- Stirling, G.R. & Wachtel, M.F.** (1980) Mass production of *Bacillus penetrans* for the biological control of root-knot nematodes. *Nematologica* **26**, 308–312.
- Sturhan, D.** (1996) Diversity and host specificity in *Pasteuria* bacteria. *Nematropica* **26**, 226, 1996.
- Waller, P.J.** (1990) Resistance in nematode parasites of livestock to the benzimidazole anthelmintics. *Parasitology Today* **6**, 127–129.
- Waller, P.J.** (1993) Nematophagous fungi: prospective biological control agents of animal parasitic nematodes? *Parasitology Today* **9**, 429–431.
- Williams, A.B., Stirling, G.R., Hayward, A.C. & Perry, J.** (1989) Properties and attempted culture of *Pasteuria penetrans*, a bacterial parasite of root-knot nematodes (*Meloidogyne javanica*). *Journal of Applied Bacteriology* **67**, 145–156.
- Wolstrup, P., Nansen, P., Grønvold, J., Henriksen, S.A. & Larsen, M.** (1996) Toward practical biological control of parasitic nematodes in domestic animals. *Journal of Nematology* **28**, 129–132.
- Wood, W.B.** (1988) *The nematode Caenorhabditis elegans*. Cold Spring Harbor, New York.

(Accepted 15 July 1998)

© CAB INTERNATIONAL, 1999