

## Augmented Zika and Dengue Neutralizing Antibodies Are Associated With Guillain-Barré Syndrome

Rebecca M. Lynch,<sup>1</sup> Grace Mantus,<sup>1</sup> Liliana Encinales,<sup>2</sup> Nelly Pacheco,<sup>2</sup> Guangzhao Li,<sup>1</sup> Alexandra Porras,<sup>3</sup> Alejandro R. Mendoza,<sup>3</sup> Jin Peng,<sup>1</sup> Monica Rengifo-Pardo,<sup>4</sup> Magelda Montoya Cruz,<sup>5</sup> Eva Harris,<sup>5</sup> Jeff M. Bethony,<sup>1</sup> Gary L. Simon,<sup>6</sup> and Aileen Y. Chang<sup>6</sup>

<sup>1</sup>Department of Microbiology, Immunology, and Tropical Medicine, George Washington University, Washington, District of Columbia; <sup>2</sup>Department of Medicine, Allied Research Society LLC, Baranquilla, and <sup>3</sup>Grupo de Medicina Comunitaria, Universidad El Bosque, Bogotá, Colombia; <sup>4</sup>Department of Dermatology, George Washington University, Washington, District of Columbia; <sup>5</sup>Division of Infectious Diseases and Vaccinology, School of Public Health, University of California, Berkeley; and <sup>6</sup>Department of Medicine, George Washington University, Washington, District of Columbia

The role of neutralizing antibodies in Zika-induced Guillain-Barré syndrome (GBS) has not yet been investigated. We conducted a case-control study using sera from the 2016 Zika epidemic in Colombia to determine the neutralizing antibody activity against Zika virus (ZIKV) and dengue virus serotype 2 (DENV2). We observed increased neutralizing antibody titers against DENV2 in ZIKV-infected individuals compared with uninfected controls and higher titers to both ZIKV and DENV2 in ZIKV-infected patients diagnosed with GBS compared with non-GBS ZIKV-infected controls. These data suggest that high neutralizing antibody titers to DENV and to ZIKV are associated with GBS during ZIKV infection.

**Keywords.** Guillain-Barré syndrome; neutralizing antibody; flavivirus; Zika; dengue.

Zika virus (ZIKV) is a flavivirus spread mainly by the *Aedes aegypti* mosquito. In 2015–2016, an epidemic of Zika in the Americas was accompanied by severe neurologic complications including microcephaly in babies born to mothers infected with ZIKV during pregnancy and Guillain-Barré syndrome (GBS) in adults [1]. GBS is a disorder of the peripheral nervous system often triggered by a preceding viral or bacterial infection or vaccination [2]. Although the exact cause of most GBS cases remains unknown, several studies have demonstrated that for some pathogens, such as *Campylobacter jejuni*,

an infection-induced antibody cross-reacts with the ganglioside surface components of peripheral nerves [2]. Although the mechanism whereby Zika is associated with GBS has not been clearly elucidated, it is likely that there is a similar pathogenesis. As many vaccines currently under development for Zika are designed to elicit protective titers of neutralizing antibodies, it is critical to define the role of ZIKV antibodies in the development of GBS.

During the 2015–2016 Zika epidemic in Colombia, there was a simultaneous increase in the number of neuroinflammatory disorders reported [3]. Specifically, there was an increase in GBS cases in individuals found to be ZIKV positive (ZIKV<sup>+</sup>) by reverse-transcription polymerase chain reaction, lending support to the role of ZIKV infection in GBS pathogenesis [3]. To investigate the relationship between ZIKV infection and GBS, anti-ZIKV neutralizing antibodies were assayed in plasma samples obtained from ZIKV-infected patients and controls collected during the 2016 outbreak in Barranquilla, Colombia.

### METHODS

#### Ethics Statement

This study was approved by the ethics committee of the Universidad El Bosque, and a nonhuman subjects determination was made by the George Washington University Institutional Review Board for analysis of de-identified data. All participants received written informed consent.

#### Participants and Setting

Adult patients with a clinical diagnosis of Zika and Zika-related GBS were referred to this study from the Atlántico Department and Bolívar Department, Colombia, while asymptomatic participants from Bogotá, Cundinamarca Department, a mountainous region without endemic ZIKV transmission, were enrolled as Zika-negative controls.

#### Case and Control Definitions

##### Zika-Related Guillain-Barré Syndrome Case

A Zika-related GBS (ZGBS) case was defined as a participant with clinically diagnosed ZIKV infection, confirmed by serologic analysis as described below, and GBS, as diagnosed and reported by a local neurologist. The Brighton criteria for the level of GBS diagnosis certainty was determined if documentation was available [4].

##### Zika-Positive Control

Participants with clinical symptoms of ZIKV infection and serological ZIKV confirmation (ZIKV<sup>+</sup> control) were matched by age and sex using simple stratified random sampling from patients from the Atlántico and Bolívar departments with clinical ZIKV infection.

Received 6 April 2018; editorial decision 20 July 2018; accepted 2 August 2018; published online August 3, 2018.

Correspondence: R. M. Lynch, PhD, George Washington University, 2300 Eye St NW, Washington, DC 20037 (rmlynch@email.gwu.edu).

The Journal of Infectious Diseases® 2019;219:26–30

© The Author(s) 2018. Published by Oxford University Press for the Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com DOI: 10.1093/infdis/jiy466

### Zika-Negative Control

A Zika-negative control (ZIKV<sup>-</sup>) was defined as an asymptomatic participant confirmed to be negative for ZIKV infection by serology.

### Design

Each ZGBS case was age- and sex- matched to 2 ZIKV<sup>+</sup> controls and 1 ZIKV<sup>-</sup> control by simple random sampling within age and sex strata. All patients with a clinical diagnosis of ZIKV or ZGBS completed a brief symptom questionnaire prior to blood sample collection. A retrospective chart review was performed for cases of ZGBS cases where the medical records were available.

### Serologic ZIKV Infection Determination

Participants were considered to be positive for a ZIKV infection if they fulfilled the following diagnostic criteria: ZIKV nonstructural protein 1 (NS1) antibody positive by the previously described Zika NS1 blockade-of-binding assay [5] or a reciprocal 50% neutralizing titer (NT<sub>50</sub>) against ZIKV strain H/PF/2013 that was at least 2-fold greater than the NT<sub>50</sub> titer against dengue virus serotype 2 (DENV2) 16681.

### Reporter Virus Particle Neutralization Assays

Neutralization of ZIKV H/PF/2013 and DENV2 16681 by plasma samples was measured using a reporter virus particle assay as described previously [6]. In brief, heat-inactivated plasma was serially diluted 5-fold from 1:50 and incubated with 100 µL of virus for 1 hour at 37°C, after which 50 µL of target Vero cells (400 000 cells/mL) was added. Input virus dilution was calculated from titration experiments to ensure sufficient luciferase output within the linear portion of the titration curve. Cell-only and virus-only controls were included on each plate, and all serum samples (and virus only) were run in triplicate. After a 48-hour incubation, luciferase activity was measured, and neutralization curves were calculated by averaging luciferase units from triplicates, subtracting cell-only control background and calculating the percentage difference in serum samples to virus-only controls. Data was fit by nonlinear regression using the asymmetric 5-parameter logistic function in GraphPad Prism. The 50%, 80%, and 90% neutralizing titers (NT<sub>50</sub>, NT<sub>80</sub>, and NT<sub>90</sub>, respectively) were defined as the reciprocal serum dilution resulting in a 50%, 80%, or 90% reduction in infectivity.

### Statistical Analysis

Nonparametric Mann–Whitney *U* tests were performed to determine if there were differences in reciprocal dilutions between ZGBS and ZIKV<sup>+</sup> groups for neutralization of DENV and ZIKV, respectively. Samples with no neutralization at a dilution of 1:50 were assigned a titer of 49 for statistical analysis. The differences between the mean reciprocal dilution vectors for neutralization of DENV2 and ZIKV in these groups (ZGBS and ZIKV<sup>+</sup>) were further assessed with Hotelling T<sup>2</sup> test and

graphically with 95% probability confidence ellipses. Spearman rank correlation was used to determine the association between neutralization of DENV and ZIKV for each group. Statistical analyses were conducted using SAS software, version 9.4 (SAS Institute, Cary, North Carolina), and tests were considered statistically significant with a *P* value ≤.05.

## RESULTS


The clinical and serological factors associated with ZGBS were studied in 23 patients with a clinical diagnosis of Zika and GBS in Barranquilla, Colombia, from December 2015 through May 2016. Six participants were excluded from further analysis because their clinical Zika diagnosis was not serologically confirmed. Seventeen ZGBS cases, 34 age- and sex-matched ZIKV<sup>+</sup> controls, and 17 age- and sex-matched ZIKV<sup>-</sup> controls were included in the current analysis. The ZGBS cases were adults with median age of 49 years, and 47% were male (Supplementary Table 1). Two patients reported a history of a previous suspected DENV infection, and 2 patients of a suspected prior chikungunya virus infection. All patients reported viral symptoms during ZIKV infection including arthralgias (94%), fever (88%), and myalgias (88%). The median time from onset of ZIKV symptoms to neurologic symptoms was 10 days (interquartile range [IQR], 7–19; Supplementary Table 1). Access to medical records allowed Brighton criteria GBS classification in 8 of the 17 patients, demonstrating certainty of diagnosis level 1 (based on both nerve conduction studies and cerebrospinal fluid [CSF] analysis) in 18% of cases, level 2 in 18% of cases based on either nerve conduction studies or CSF analysis, and level 3 (based on clinical features) in 12% of cases [4]. One patient was diagnosed with Miller–Fisher syndrome (Supplementary Table 1). Two patients demonstrated demyelination and axonal involvement based on nerve conduction studies.

The most common neurologic symptoms were lower extremity weakness (100%), inability to walk (88%), and paresthesias (100%). The great majority of patients were cared for in the intensive care unit (88%). Half the patients had difficulty breathing, and 38% had respiratory failure requiring intubation. Most patients were treated with intravenous immunoglobulin (63%) or plasmapheresis (25%), and none were treated with steroids. The median duration of hospitalization was 11 days (IQR, 7–24 days), with a median of 9 days (IQR, 5–13 days) in the intensive care unit. One patient died, one-fourth had a full recovery, and 63% reported chronic morbidities including upper and lower extremity weakness, facial tremors, and sensory alterations.

The relationship between antibody responses to ZIKV infection and a clinical diagnosis of GBS was assessed by comparing neutralizing antibody titers between the ZGBS cases and the ZIKV<sup>+</sup> and ZIKV<sup>-</sup> controls. Because DENV2 recently circulated in Colombia, plasma neutralizing antibody titers against both

Case/Control	Age	Sex	Days from Zika symptom onset to sample draw	Median days to sample draw	DENV2 16680			ZIKVH/PF			KEY Reciprocal plasma dilutions
					NT <sub>50</sub>	NT <sub>80</sub>	NT <sub>90</sub>	NT <sub>50</sub>	NT <sub>80</sub>	NT <sub>90</sub>	
GBS POSITIVE	22	M	120	23	4464	1577	809	9174	4211	2440	<300
	22	M	137		1680	786	505	5990	2588	1486	300-999
	32	F	19		132 716	43 852	23 158	286 725	115 102	64 033	1000-4999
	34	F	17		69 459	10 085	3229	215 122	62 746	26 743	5000-10 000
	35	M	30		117 540	26 841	10 438	143 053	30 164	11 209	>10 000
	38	F	15		106 494	31 028	13 843	1 628 613	305 336	106 118	>10 000
	39	F	55		12 182	5528	3167	23 051	10 014	5747	>10 000
	40	F	14		195 125	87 154	51 088	121 528	35 051	15 150	>10 000
	45	F	30		74 221	33 779	27 625	37 971	14 089	7508	>10 000
	46	M	14		38 939	12 732	7289	372 419	56 306	17 768	>10 000
	48	F	29		29 850	18 817	13 748	22 686	9957	5752	>10 000
	48	M	30		103 159	36 812	27 969	195 874	57 355	25 425	>10 000
	49	F	79		47 725	15 309	8627	3005	1007	519	>10 000
	50	F	13		186 916	46 154	21 426	509 996	119 161	45 612	>10 000
	56	F	23		23 309	5499	2207	4525	1231	516	>10 000
59	M	10	16 791	3348	1280	22 231	9494	5302	>10 000		
88	M	20	31 581	10 605	6581	15 437	4581	2090	>10 000		
ZIKA POSITIVE	15	F	26	26	62	<50	<50	19 452	6104	3183	>10 000
	18	M	9		1291	524	290	3001	1411	797	>10 000
	18	M	66		28 3776	58 674	34 524	54 888	20 516	10 917	>10 000
	20	F	8		<50	<50	<50	16 277	4410	2257	>10 000
	20	F	16		2842	1198	967	4526	2067	1258	>10 000
	21	M	20		1295	647	404	7652	3036	1598	>10 000
	22	M	11		9477	3233	1709	1071	351	179	>10 000
	23	M	13		12 476	5316	2958	5096	956	372	>10 000
	26	F	85		7090	2424	1501	29 014	9351	5226	>10 000
	29	F	76		3012	923	459	38 083	13 088	6543	>10 000
	30	M	21		93 250	37 718	21 516	41 250	10 055	6557	>10 000
	30	F	7		28 888	8134	3533	18 867	5340	3866	>10 000
	39	F	5		44 797	21 184	12 351	169 474	58 392	29 451	>10 000
	40	F	24		5041	1807	1004	9547	2313	1002	>10 000
	42	M	32		155	61	<50	19 654	4889	2020	>10 000
	43	F	25		5445	2266	1283	7351	1010	436	>10 000
	44	F	37		11 078	3535	2216	126 872	41 558	20 211	>10 000
	45	M	32		12 822	4950	3070	12 119	5741	4049	>10 000
	45	M	11		9212	3327	1742	5606	2509	1454	>10 000
	46	M	29		8048	4191	2723	109 021	18 314	5500	>10 000
	46	F	51		70 359	24 072	12 795	237 402	146 111	128 080	>10 000
	46	F	28		6201	3013	1984	7393	2706	1444	>10 000
	48	F	53		31 252	15 538	9750	2039	651	375	>10 000
	48	F	21		2560	1288	1083	2019	979	804	>10 000
	49	F	32		1754	810	524	7776	2729	1540	>10 000
	50	F	13		5353	1984	1021	6137	2897	1783	>10 000
51	F	78	359	105	58	1337	395	143	>10 000		
51	F	63	5800	1743	1122	10 826	3239	1536	>10 000		
55	M	27	12 328	4151	2395	6096	2271	1162	>10 000		
55	F	13	9597	3341	2021	5648	2472	1561	>10 000		
62	F	13	20 644	5520	2486	3725	1409	797	>10 000		
64	M	50	1359	369	182	2546	962	544	>10 000		
66	M	8	1967	799	476	9553	2586	1540	>10 000		
67	M	65	12 516	4400	2301	140 761	26 712	10 782	>10 000		
ZIKA NEGATIVE	18	M	-	-	393	236	204	187	60	<50	<300
	18	M	-		851	395	232	170	<50	<50	<300
	19	M	-		<50	<50	<50	<50	<50	<50	<300
	20	F	-		340	133	100	<50	<50	<50	<300
	23	F	-		1020	298	148	<50	<50	<50	<300
	31	F	-		3751	772	320	222	<50	<50	<300
	39	M	-		58	<50	<50	54	<50	<50	<300
	40	F	-		582	254	206	116	<50	<50	<300
	40	F	-		1250	177	85	173	<50	<50	<300
	41	F	-		787	297	181	<50	<50	<50	<300
	45	F	-		6986	2868	1585	91	<50	<50	<300
	47	M	-		3263	977	495	164	50	<50	<300
	49	F	-		1464	626	331	241	<50	<50	<300
	50	F	-		1137	535	324	371	86	<50	<300
	50	F	-		1450	681	393	<50	<50	<50	<300
55	M	-	8354	3946	2290	233	68	<50	<300		
50	M	-	1584	1168	1074	<50	<50	<50	<300		

**Figure 1.** Neutralizing antibody titers to Zika virus (ZIKV) and dengue virus serotype 2 (DENV2) in cases and controls. Age and sex are listed for each case-control participant as well as the time from reported Zika symptoms to date of sampling in days. Reciprocal 50%, 80%, and 90% neutralizing antibody titers to ZIKV strain H/PF/2013 and DENV2 strain 16681 are reported and shaded according to potency as indicated in the key. Abbreviations: DENV2, dengue virus serotype 2; GBS, Guillain-Barré syndrome; NT<sub>50</sub>, 50% neutralizing titer; NT<sub>80</sub>, 80% neutralizing titer; NT<sub>90</sub>, 90% neutralizing titer; ZIKV, Zika virus.



**Figure 2.** Differences in Zika virus (ZIKV) and dengue virus serotype 2 (DENV2) neutralizing antibody titers between Zika-related Guillain-Barré syndrome (ZGBS) cases and Zika-positive controls. Spearman rank correlation coefficient is provided for each group. The 95% confidence ellipses are indicated for each population and significant *P* values indicate differences between these 2 populations (Hotelling  $T^2$  test). Abbreviations: DENV, dengue virus; NT<sub>50</sub>, 50% neutralizing titer; NT<sub>80</sub>, 80% neutralizing titer; NT<sub>90</sub>, 90% neutralizing titer; ZGBS, Zika-related Guillain-Barré syndrome; ZIKV, Zika virus; ZIKA<sup>+</sup>, Zika virus positive.

ZIKV H/PF/2013 and DENV2 16681 for all cases and controls were measured, and calculated reciprocal plasma NT<sub>50</sub>, NT<sub>80</sub>, and NT<sub>90</sub> (Figure 1) were reported. We found that mean reciprocal titers against ZIKV were significantly elevated in the ZGBS cases compared with ZIKA<sup>+</sup> controls when comparing NT<sub>50</sub> values (212 788 vs 33 485; *P* = .0052), NT<sub>80</sub> values (49 317 vs 11 986; *P* = .0038), or NT<sub>90</sub> values (20 201 vs 7617; *P* = .0043) (Supplementary Table 3). Four of the 17 patients had reciprocal ZIKV NT<sub>50</sub> <10 000, but 3 of these patients had the longest time interval between onset of disease and sampling (79–137 days). We observed a trend toward lower NT<sub>80</sub> with increasing days post-Zika infection in the ZGBS group. When comparing ZGBS cases to ZIKA<sup>+</sup> controls, we found significantly elevated titers against DENV2 as well (Supplementary Table 4).

As expected, there was a significant correlation between ZIKV and DENV2 neutralizing antibody titers in all ZIKV-infected individuals (both ZGBS and ZIKA<sup>+</sup> groups). This correlation was stronger within ZGBS cases (NT<sub>50</sub>: *r* = 0.69, *P* = .002; NT<sub>80</sub>: *r* = 0.67, *P* = .003) compared with ZIKA<sup>+</sup> controls (NT<sub>50</sub>: *r* = 0.31, *P* = .077; NT<sub>80</sub>: *r* = 0.29, *P* = .095) (Supplementary Table 5). These data are summarized by graphing ZIKV vs DENV2 neutralizing antibody titers for each patient, along with 95% confidence ellipses (Figure 2). These ellipses illustrate that there is a statistical difference between the ZGBS cases and the ZIKA<sup>+</sup> controls and that ZGBS is associated with elevated neutralizing antibody titers not only to ZIKV but also to DENV2.

## DISCUSSION

The clinical and demographic characteristics of ZGBS cases from Barranquilla, Colombia, are in accordance with other studies [3, 7–12]. GBS may occur with rapid onset of both motor and sensory neurologic symptoms following symptomatic ZIKV infection. These cases reported here tended to be severe, with most

patients (88%) admitted to intensive care units and more than one-third requiring mechanical ventilation.

The higher neutralizing antibody titers found in the ZGBS cases compared to ZIKA<sup>+</sup> controls provide evidence of a correlation between these titers and the development of GBS in these patients, though not causation. This finding could represent an indirect effect that results from high virus load. The time course between the onset of Zika-like symptoms and the development of neurologic symptoms is sufficient for antibody production and would be compatible with the role of an adaptive immune response in this process; however, this observation does not prove that adaptive immunity causes GBS. The onset of neurologic symptoms occurred in median of 10 days after the initial onset of viral symptoms, comparable with other GBS cohorts from Colombia [3, 9, 11].

Higher titers of anti-DENV2 antibodies in ZIKV-infected participants compared to uninfected participants provides more evidence for the observation that DENV B cells are activated after ZIKV infection. This is in agreement with a previous reports such as an analysis of samples from Brazil where neutralizing antibody titers to ZIKV and DENV1 were boosted after ZIKV infection [13]. Confirmation of the role of DENV neutralizing antibodies in ZIKV infection was recently reported in a study of longitudinal B-cell responses to ZIKV after previous DENV infection. It was observed that both ZIKV and DENV neutralizing antibodies are boosted following ZIKV infection but that they derive from distinct B-cell populations and that the anamnestic dengue response occurs first, followed by a de novo ZIKV response [14]. It is possible that the development of GBS is related to molecular mimicry, where virus specific antibodies (either ZIKV or DENV) cross-react with nerve cells, but it is also possible this association between neutralizing antibodies and GBS results from indirect effects such

as high virus load or high immune activation. This cross-sectional analysis cannot fully resolve that issue. Further research is needed to characterize the specific antibody populations responsible for ZGBS, and this information will be critical for Zika vaccine development.

### Supplementary Data

Supplementary materials are available at *The Journal of Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copy-edited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

### Notes

**Acknowledgments.** We thank the Allied Research Society for their support of this study; Dr Aravinda De Silva for helpful comments; and Drs Theodore Pierson and Kim Dowd for their reagents and invaluable help with the reporter virus particle neutralization assay.

**Disclaimer.** The contents of this work are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Advancing Translational Science or the National Institutes of Health (NIH).

**Financial support.** This work was supported by the NIH National Center for Advancing Translational Sciences (award numbers UL1TR001876 and KL2TR001877) and by the National Institute of Allergy and Infectious Diseases, NIH (grant number P01AI106695 to E. H.).

**Potential conflicts of interest.** All authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

### References

1. Broutet N, Krauer F, Riesen M, et al. Zika virus as a cause of neurologic disorders. *N Engl J Med* **2016**; 374:1506–9.
2. Hahn AF. Guillain-Barré syndrome. *Lancet* **1998**; 352:635–41.
3. Parra B, Lizarazo J, Jiménez-Arango JA, et al. Guillain-Barré syndrome associated with Zika virus infection in Colombia. *N Engl J Med* **2016**; 375:1513–23.
4. Sejvar JJ, Kohl KS, Gidudu J, et al; Brighton Collaboration GBS Working Group. Guillain-Barré syndrome and Fisher syndrome: case definitions and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine* **2011**; 29:599–612.
5. Balmaseda A, Stettler K, Medialdea-Carrera R, et al. Antibody-based assay discriminates Zika virus infection from other flaviviruses. *Proc Natl Acad Sci U S A* **2017**; 114:8384–9.
6. Dowd KA, DeMaso CR, Pelc RS, et al. Broadly neutralizing activity of Zika virus-immune sera identifies a single viral serotype. *Cell Rep* **2016**; 16:1485–91.
7. Cao-Lormeau VM, Blake A, Mons S, et al. Guillain-Barré syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. *Lancet* **2016**; 387:1531–9.
8. Dirlikov E, Major CG, Mayshack M, et al. Guillain-Barré syndrome during ongoing Zika virus transmission—Puerto Rico, January 1–July 31, 2016. *MMWR Morb Mortal Wkly Rep* **2016**; 65:910–4.
9. Arias A, Torres-Tobar L, Hernández G, Paipilla D. Guillain-Barré syndrome in patients with a recent history of Zika in Cúcuta, Colombia: a descriptive case series of 19 patients from December 2015 to March 2016. *J Crit Care* **2017**; 37:19–23.
10. da Silva IRE, Frontera JA, Bispo de Filippis AM, Nascimento OJMD; RIO-GBS-ZIKV Research Group. Neurologic complications associated with the Zika virus in Brazilian adults. *JAMA Neurol* **2017**; 74:1190–8.
11. Anaya JM, Rodríguez Y, Monsalve DM, et al. A comprehensive analysis and immunobiology of autoimmune neurological syndromes during the Zika virus outbreak in Cúcuta, Colombia. *J Autoimmun* **2017**; 77:123–38.
12. Pinto-Díaz CA, Rodríguez Y, Monsalve DM, et al. Autoimmunity in Guillain-Barré syndrome associated with Zika virus infection and beyond. *Autoimmun Rev* **2017**; 16:327–34.
13. Robbani DF, Bozzacco L, Keeffe JR, et al. Recurrent potent human neutralizing antibodies to Zika virus in Brazil and Mexico. *Cell* **2017**; 169:597–609.e11.
14. Rogers TF, Goodwin EC, Briney B, et al. Zika virus activates de novo and cross-reactive memory B cell responses in dengue-experienced donors. *Sci Immunol* **2017**; 2:eaan6809.