



Autosomal dominant zonular cataract with sutural opacities is associated with a splice mutation in the β A3/A1-crystallin gene

Chitra Kannabiran,^{1,2}Peter K. Rogan,³Lisa Olmos,¹Surrendra Basti,²Gullapalli N. Rao,²Muriel Kaiser-Kupfer,¹J. Fielding Hejtmancik¹

¹National Eye Institute, Bethesda, MD; ²L. V. Prasad Eye Institute, Hyderabad, India; ³Department of Human Genetics, Allegheny University of the Health Sciences, Pittsburgh, PA

Purpose: Congenital cataracts constitute a morphologically and genetically heterogeneous group of diseases that are a major cause of childhood blindness. Autosomal Dominant Zonular Cataracts with Sutural Opacities (CCZS) have been mapped to chromosome 17q11-q12 near the β A3A1-crystallin gene (CRYBA1). The β A3A1-crystallin gene was investigated as the causative gene for the cataracts.

Methods: The β A3/A1-crystallin gene was sequenced in affected and control individuals. Base changes were confirmed and assayed in additional family members and controls using NlaIII restriction digestion of PCR amplified DNA sequences. Base changes were assessed for their effects on splicing by information analysis.

Results: The cataracts are associated with a sequence change in the 5' (donor) splice site of intron 3: GC(g->a)tgagt. The sequence change also creates a new NlaIII site. This base change cosegregates with the cataracts in this family, being present in every affected individual. Conversely, this base change was not seen in 140 chromosomes examined in 70 unaffected and unrelated individuals. Information theory mutational analysis shows that the base change lowers the information content of the splice site from 6.0 to -6.8 bits, so that splicing would not be expected to occur at the altered site.

Conclusions: Taken together, these observations suggest that the observed mutation might be causally related to the cataracts in this family.

Congenital cataracts are a significant cause of visual disease, and without prompt treatment they can interfere with the sharp imaging on the retina necessary to develop normal visual cortical synaptic connections, resulting in irreversible visual loss. They form a heterogeneous group of diseases, with one-third being familial, most commonly inherited in an autosomal dominant fashion [1]. Dominantly inherited congenital cataracts are themselves genetically heterogeneous, with cataracts being mapped to 10 different loci including 1p36 [2], 1q21-q25 [3], 2q33-q35 [4], 13 [5], 16q22 [6], 17p13 [7], 17q11-q12 [8], 17q24 [9], 21q22.3 [10], and 22q [11].

Autosomal dominant congenital cataracts are also phenotypically heterogeneous, including cerulean (17q24, 22q), anterior polar (17p13), and nuclear zonular or lamellar (1p36, 1q21-q25, 2q33-q35, 16q22, 17q11-q12, 22q) morphologies. Clinically identical cataracts can map to different loci [12], and cataracts segregating in the same family can show marked variability [13]. While the spatial distribution of opacity within the lens is generally expected to reflect expression patterns of the causative protein during development, precise explanations of the phenotypic appearance of specific hereditary cataracts has remained elusive. Sutural cataracts show selective opacification of the Y-sutures of the lens. They are infrequently reported, because they usually do not interfere with visual acuity and thus do not prompt patients to seek medical attention. Sutural cataracts can be inherited, most often in an autosomal dominant fashion, and Marner cataracts (CAM, 16q22) as well as Volkmann cataracts (CCV, 1p36) have sutural com-

ponents, although they are primarily nuclear zonular in morphology.

High concentrations of closely packed crystallins are required for transparency and focusing of light by the lens. The major classes of ubiquitous crystallins are α -crystallin, molecular chaperones related to the small heat shock proteins, and the β - and γ -crystallins. The β - and γ -crystallins share an extremely stable common core structure comprising four twisted β -pleated sheets termed "Greek key motifs," which are organized into two domains [14]. The β -crystallins also have amino and carboxy terminal extensions or "arms." The β -crystallins associate into higher order assemblies while the γ -crystallins exist in solution as monomers. Both the β - and γ -crystallins tend to be more highly expressed at early developmental times in elongating fiber cells, so that they are found primarily in the lens nucleus [15-17].

Basti and coworkers [18] described a family with autosomal dominant cataracts having both a zonular component in the fetal nucleus and prominent opacities of both the anterior and posterior Y-sutures in the area enclosed by the zonular component. It differs from previously described families in that affected individuals uniformly showed significant sutural opacities. Linkage analysis localized the gene causing these cataracts to a region of chromosome 17q11-q12 including the β A3/A1-crystallin gene with a lod score of 3.91 [8]. Here we describe association of a splice site mutation at the end of the third exon of β A3/A1-crystallin with the cataracts in this family.

METHODS

Pedigrees and Diagnosis: Patients were ascertained and examined by (S. B.) at the L. V. Prasad Eye Institute in

Hyderabad. Patient studies and informed consent were approved by the L. V. Prasad Institutional Review Board, the National Eye Institute Review Board, and the National Institutes of Health Office of Protection from Research Risks. Clinical and Ophthalmologic examinations included detailed ophthalmic, medical, and family histories, dilated slit-lamp examination with photographs, Snellen visual acuity testing, intraocular-pressure measurement by applanation tonometry, and fundus examination. All affected family members showed a zonular cataract measuring 3.5-4 mm in diameter, an erect Y-shaped anterior sutural cataract, and an inverted Y-shaped posterior sutural cataract. Clinical findings in these patients are described in detail in [18] and [8].

PCR Amplification of the β A3/A1-Crystallin Gene: Human genomic DNA from 70 unaffected controls (50 of Indian descent and 20 from other populations) and the affected and unaffected family members shown in Figure 1 was amplified using polymerase chain reaction (PCR) primers from flanking sequences of exons 1-6 of the human β A3/A1-crystallin gene (HUMCRYBA1-6, GenBank accession numbers M14301, M14302, M14303, M14304, M14305, and M14306). The primers used are shown in Table 1. PCR was carried out for 35 cycles each consisting of a 45 seconds at 94 °C, 30 seconds at the annealing temperature shown in Table 1, and 30 seconds at 72 °C. The first and last primers shown were used together in a PCR to amplify the entire intervening β A3/A1-crystallin gene sequence. This fragment was then sequenced with the exon 6 reverse primer to examine the 5'-most 24 bases of exon 6, since these were not included in the amplified fragment for exon 6 using the specific forward and reverse primers.

DNA sequencing: PCR products were checked for correct size by agarose gel electrophoresis and purified by Wizard PCR Prep DNA Purification System (Promega, Madison, WI) followed by ethanol precipitation. They were sequenced bidirectionally using an ABI 377 Prism automated sequencer

TABLE 1. PCR PRIMERS FOR AMPLIFICATION OF β A3/A1-CRYSTALLIN EXONS.

Primer Fragment	Sequence	Annealing	
		temperature	size
exon 1 fwd	5'-GGTCTTAGGAAGATCCCAAG-3'	58 °C	394 bp
exon 1 rev	5'-AAGGAGAGGAAGGCAAGGG-3'		
exon 2 fwd	5'-CGTGTGTGCTCTGTCTTCC-3'	58 °C	179 bp
exon 2 rev	5'-GGTCAGTCACTGCCTTATGG-3'		
exon 3 fwd	5'-CAATCCTCCCTCCACCTC-3'	54 °C	520 bp
exon 3 rev	5'-TCCTTCTCTCAGCTTTGG-3'		
exon 4 fwd	5'-CAAACTACATGTCTTTGG-3'	48 °C	210 bp
exon 4 rev	5'-CTTGCTACCCTCATATGC-3'		
exon 5 fwd	5'-TGCTTCCTGTATAATCC-3'	52 °C	306 bp
exon 5 rev	5'-ACTATTGATGCAACCTCAGG-3'		
exon 6 fwd	5'-GGTTTGCTACCATTATCTTGG-3'	52 °C	183 bp
exon 6 rev	5'-CATGCTTGAGGAATATCG-3'		

Forward and reverse primers flanking exons 1-6 are shown with the primer sequence, the annealing temperature for the amplification reaction, and the expected fragment size of the PCR product. The product fragments were used for sequencing and restriction analysis (exon 3 only).

(ABI, Foster City, CA) using the original primers the an Amplitaq FS cycle sequencing kit (ABI) with dye-labeled terminators.

Restriction endonuclease analysis: PCR products were purified by phenol/chloroform extraction and ethanol precipitation, digested with NlaIII at 37 °C in NEB buffer 3 (New England Biolabs, Beverly, MA) for 2 h, and analyzed by ethidium bromide staining after electrophoresis on a 6% acrylamide gel in Tris-Borate-EDTA, pH 8.2.

Information Theory Mutational Analysis: The potential results of the G to A transition were estimated using information theory as described by [19], and [20]. The 10 bases constituting the donor and 27 bases constituting the acceptor splice sites were weighted according to sequence conservation and base frequency. These values are summed, providing a quantitative estimate of the ability of the sequence to serve as a splice site, reported in bits. This algorithm was implemented using the computer programs SCAN and RI, and the information contribution at each position of the site was depicted using the program WALKER [21,22].

RESULTS

The six exons of the β A3/A1-crystallin gene were amplified and their sequences determined, including all coding regions, all splice sites, and a small amount of each adjacent intron. Both the control and patient β A3/A1-crystallin sequences agreed with the published sequence [23], confirming the 6 base changes described by Lampi and coworkers [24], with the single exception shown in Figure 2. This shows the se-




Figure 1. Pedigree and restriction endonuclease analysis for the G to A transition of the family described by Basti [18]. NlaIII digestions of PCR products are shown below each surviving individual in the pedigree. The first and last lanes are 100 bp ladders. The largest cleavage product, 488 bp, results from presence of the wild type sequence, while the cleaved products, 346 and 142 bp, indicate presence of the mutant sequence. In individual III-4 and to a lesser extent in individual II-8 a 520 bp band representing the PCR product partially digested with NlaIII is visible. No DNA sample was received from individual II-6, who has no corresponding lane on the gel. Individual III-7 was analyzed and the results are shown vertically below both II-6 and III-7.

quence at the 3' end of exon 3 of an unaffected control, comparing it with affected individual III-2 in the pedigree shown in Figure 1. The sequence in the unaffected individual is homozygous for a G at the first intronic base following exon 3, while the affected individual is heterozygous for a G to A transition at this base. This is also clearly seen in the reverse sequence, where the affected individual is heterozygous for the corresponding C to T transition. These results are summarized in Figure 3.

The G to A base change created a new NlaIII recognition site (CATG) not found in unaffected individuals. DNA from each member of the family was amplified using the primers flanking exon 3, resulting in a 520 bp fragment. When this fragment is subjected to digestion with NlaIII, it is cleaved to a 488 bp fragment in unaffected individuals (Figure 1). However, both the 488 bp fragment seen in unaffected individuals and 346 and 142 bp restriction fragments resulting from cleavage of the newly created NlaIII site are visible in samples from affected individuals. DNA samples from 50 unaffected Indian and 20 additional controls (140 chromosomes total) were examined and found to lack this NlaIII site (data not shown). These results are summarized in Figure 3.

The predicted effects of the G to A transition are shown in Figure 4. The sequence of the 5' (donor) splice site of control β A3/A1-crystallin exon 3 is shown in the top panel. The WALKER representation of the control splice site at position 474 is shown below this. It has an R_i (information content) score of 6.0 bits, much of it contributed by the G and T at positions +1 and +2, respectively. In addition, there is a weaker

4.5 bit splice site upstream at position 460. The mutant sequence is shown in the lower panel. The G to A base change at position 474 decreases the R_i to -6.8 bits. The upstream potential splice site at position 460 is unaffected. All other potential splice sites in the sequence shown in Figure 3 have R_i values less than 2.4 bits and should not be recognized as splice sites.

DISCUSSION

Both direct sequencing of PCR products and restriction endonuclease digestion show that a G to A transition in the 5' (donor) splice site of exon 3 of the β A3/A1-crystallin gene on chromosome 17q is associated with autosomal dominant zonal sutural cataracts (CCZS). The location of the zonal sutural cataracts in the fetal lens nucleus is consistent with expression of β A3/A1-crystallin in the lens [15,25,26]. This base change was not seen in 100 chromosomes examined in 50 unaffected individuals.

```

1 cctgggactc ccagaatcct gccctctggg ccatgcttat gaaggaactg tgggtgaaatg
61 ggagcccata agtcctctggc ctgtccattg gtctagatat tatccctggc agaccatttc
121 tatcataaac caatcctccc tccacctcat ccctactctt ggcaaatgaa caccaaccag
181 tggcattaat tcaaacccct tctgcccctc tttacactgg aagagacctc atcaggcatc
241 ccaggttaca gctggttaaca gaaagcacag agtcagactg aagttaagct gttgacctgn
-----
301 acctctgttt ttctctgatg tctagctctc ttgctgacct caatctcatt tcagATAACC
I Y D Q E N F Q G K R M E F T S S C P N
361 ATCTATGATC AGGAGAACTT TCAGGGCAAG AGGATGGAGT TCACCAGCTC CTGTCCAAT
-----
V S E R S F D N V R S L K V E S G A
421 GTCTCTGAGC GCAGTTTGA TAATGTCGG TCCCTGAAG TGGAAAGTGG CGCgtgagta
481 tggactctcg cagaaccgca gcccttatt tcaggtccct tcagacaggg gaccatagag
541 tgggataggg ggaagaagga tgttccccta ggctctagtc atttctcgga gacaggcctc
-----
601 gaaagaatc actacatgca taatttagta agccaaagct gaaggaagga acatgaggat
661 tcttagtgac ttgatactag atttcagggg ttggataaag aaatacatac aatgggggtg
721 gacagtgcca gtgatacaaga ttttgcacct

```




Figure 2. Sequence analysis of the 5' (donor) splice sites exon 3 of β A3/A1-crystallin in control and affected individuals. Forward reactions showing homozygous G at position 152 in an unaffected individual and heterozygosity for the G to A transition in an affected individual, and reverse reactions confirming the same change.

Figure 3(A). β A3/A1-Crystallin exon 3 sequence. β A3/A1-Crystallin exon 3 and surrounding intronic sequence, numbered as in GenBank, accession number M14303. Bases in exon 3 are in capitals. Amino acid sequences are shown in single letter code above the exon sequence. Sequences of the exon 3 forward and reverse primers (Table 1) are in green text. The 5' (donor) splice site for exon 3 at base 474 is in blue text, as is the potential splice site at base 460. The G to A transition disruption the 5' splice site at position 474 is shown in blue above the sequence. The NlaIII sites at 618 and 476 (within the amplified fragment) are marked with hyphens above the sequence.

Exon	3' (acceptor) splice site	5' (donor) splice site
Exon 2:		W K ...TGG AAG gtaa...
Exon 3:	I T ...tcag ATA ACC.....	S G AAGT GGC GC gtga...
Exon 4:	W I ...gcag C TGG ATT.....	S ATCA GCT gtga...
Exon 5:	N H ...acag AAT CAT.....	S G AAGT GGC GC gtaa...
Exon 6:	W V ...ccag C TGG GTT.....	Q *CAG

Figure 3(B). Summary of splice sites and adjacent exon and intron sequence for exons 2-6 of the β A3-crystallin gene. Introns are shown in small letters and exons in capitals. Amino acids of the first and last two codons are shown in single letter code above the DNA sequence. Exons are aligned by the first and last complete codon, with additional bases shown extending to the right and left. From this figure it can be seen that splicing from exon 2 to exons 3 or 5 would result in conservation of the reading frame, while splicing between exons 2 and either 4 or 6 would alter the reading frame. The in-frame stop codon following the 5' (donor) splice site for intron 3 is shown in blue as the first full codon following exon 3.

Both animal and human studies indicate the importance of the lens crystallins in establishing and maintaining lens transparency [27]. Cataracts have been shown to result from mutations in mouse β B2-crystallin [28], mouse γ E-crystallin, [29], and guinea pig ζ -crystallin [30]. Human cataracts have been associated with pseudoyE-crystallin [31], β B2-crystallin [32], and α A-crystallin [10]. In each of these cases, the mutant crystallin is thought to have altered stability, solubility, or ability to oligomerize, and is predicted to precipitate from solution, resulting in lens opacity.

Besides their roles in transmission and focusing light, some lens crystallins also have biochemical or enzymatic activity. α -Crystallin has been shown to function as a molecular chaperone [33], probably serving to stabilize partially denatured crystallins within the lens cells. Similarly, β B2-crystallin has autokinase [34] and β A3/A1-crystallin has been suggested to have autoproteolytic activity [35]. However, it would seem that with the possible exception of α -crystallin, cataracts associated with mutations in these proteins might be more likely to result from disruption of their roles as structural lens proteins rather than loss of a particular enzymatic function.

The β A3/A1-crystallin gene is thought to encode both β A1- and β A3-crystallin from two in frame AUG codons [36]. These two proteins differ only in the presence of an additional 17 amino acids in the β A3-crystallin amino terminal arm. β A3- and β A1-crystallins are equally stable and both associate into dimers and higher order assemblies [37], as does an intermediate β A3-crystallin with an amino terminal arm shortened by 8 amino acids [38]. Presumably both forms of the amino terminal arm would occur on the mutant protein, although the effect the shorter arm might have on stability or association of the mutant crystallin is unclear.

The precise effect of the G to A transition on the mRNA and protein structure of β A3/A1-crystallin is currently under




Figure 4. Walker diagram of the exon 3 5' (donor) splice site and surrounding sequence. The Control sequence is at the top, showing the splice site at position 474 and the potential splice site at position 460. Bases in each splice site are shown in the corresponding WALKER diagram, with contribution to the average sequence conservation of the splice site indicated by the positive or negative height of the base. The arrow points to the mutant sequence with the G to A base change, showing the change from a positive contribution by the G to a negative contribution of the A.

investigation. Initial attempts to analyze ectopic (illegitimate) transcripts of β A3/A1-crystallin mRNA from leaky transcription in transformed lymphoblasts have been unsuccessful (data not shown). In β A3/A1-crystallin, the first two exons encode sequence of the amino terminal arm, while exons 3-6 encode Greek key motifs 1-4, respectively [23]. The G at position +1 of the 5' (donor) splice site is highly conserved, and mutation of this base would be expected to disrupt the splice site [39]. This is indicated by the fall in R_i from 6.0 bits to -6.8 bits, as shown in Figure 4. In a study of over 100 splice sites, the minimum value of R_i in a functional splice site was 2.4 [20].

The G to A transition probably would have one of two possible effects: skipping of exon 3 with splicing of exon 2 to exon 4 or possibly exons 5 or 6, or recruitment of a cryptic splice site, or possibly a combination of both. If exon 3 were simply skipped with splicing of exon 2 to exon 4, the result would be a deletion that does not maintain the reading frame and results in premature termination after addition of 4 amino acids (leu-asp-trp-leu) to the 32 amino acids encoded by the first 2 exons, basically the amino terminal arm. If the illegitimate splice site at position 460 within exon 3 is utilized, the first 35 amino acids of exon 3 would occur in a normal fashion, followed by the same 4 additional amino acids and then premature termination. If an illegitimate splice site within intron 3 is used or intron 3 is included, the amino acids encoded by exon 3 including the terminal alanine would be conserved. However, the very next codon within the retained intron 3 would be a UGA stop site, mimicking a missense mutation at the protein level. This would cause truncation of the β A3/A1-crystallin immediately after the first Greek key motif. If exon 3 is skipped and exon 2 is spliced directly to exon 5, the coding frame would be maintained, with the resulting crystallin consisting of the amino terminal arm and a single (carboxy) domain. Finally, if exon 2 is spliced directly to exon 6, the resulting protein would consist of the amino terminal arm followed by 18 "random" amino acids due to the frame-shift caused by the aberrant splice.

If an illegitimate splice site in exon 3 or intron 3 is used, the effect on the protein would be to terminate the β A3/A1-crystallin peptide near the end of the first Greek key motif, adding the 4 amino acids leu-asp-trp-leu to the carboxy terminal. This would disrupt the first Greek key motif, since the Greek key motifs are formed with the fourth strand of the first motif being provided by the second motif and vice versa [40]. This means that without the second motif, which is encoded by the fourth exon, it is not possible to form even a single Greek key structure. The lack of the fourth chain required to form the Greek key motif makes it unlikely that the protein would be folded correctly after synthesis. Not only would this improperly folded crystallin be unstable and serve as a nidus for precipitation of other damaged proteins, but it might also interfere with appropriate associations by the remaining β -crystallins. The net result of any of these events would probably be a dominantly inherited cataract. This suggests that the β A3A1-crystallin splice mutation might cause the cataracts in this family. This possibility is being further investigated.

ACKNOWLEDGEMENTS

This work is partially supported by a grant to P. K. R. from the Merck Genome Research Foundation.

REFERENCES

- Robb RM. Congenital and childhood cataracts. In: Albert DM, Jakobiec FA, editors. Principles and practice of ophthalmology. Philadelphia: Saunders; 1994. p. 2761-2767.
- Eiberg H, Lund AM, Warburg M, Rosenberg T. Assignment of congenital cataract Volkman type (CCV) to chromosome 1p36. *Hum Genet* 1995; 96:33-38.
- Renwick JH, Lawler SD. Probable linkage between a congenital cataract locus and the Duffy blood group locus. *Ann Hum Genet* 1963; 27:67-84.
- Lubsen NH, Renwick JH, Tsui LC, Breitman ML, Schoenmakers JG. A locus for a human hereditary cataract is closely linked to the gamma-crystallin gene family. *Proc Natl Acad Sci U S A* 1987; 84:489-492.
- Mackay D, Ionides A, Berry V, Moore A, Bhattacharya S, Shiels A. A new locus for dominant "zonular pulverulent" cataract on chromosome 13. *Am J Hum Genet* 1997; 60:1474-1478.
- Eiberg H, Marner E, Rosenberg T, Mohr J. Marner's cataract (CAM) assigned to chromosome 16: linkage to haptoglobin. *Clin Genet* 1988; 34:272-275.
- Berry V, Ionides AC, Moore AT, Plant C, Bhattacharya SS, Shiels A. A locus for autosomal dominant anterior polar cataract on chromosome 17p. *Hum Mol Genet* 1996; 5:415-419.
- Padma T, Ayyagari R, Murty JS, Basti S, Fletcher T, Rao GN, Kaiser-Kupfer M, Hejtmancik JF. Autosomal dominant zonular cataract with sutural opacities localized to chromosome 17q11-12. *Am J Hum Genet* 1995; 57:840-845.
- Armitage MM, Kivlin JD, Ferrell RE. A progressive early onset cataract gene maps to human chromosome 17q24. *Nat Genet* 1995; 9:37-40.
- Litt M, Kramer P, LaMorticella DM, Murphey W, Lovrien EW, Weleber RG. Autosomal dominant congenital cataract associated with a missense mutation in the human alpha crystallin gene CRYAA. *Hum Mol Genet* 1998; 7:471-474.
- Kramer P, Yount J, Mitchell T, LaMorticella D, Carrero-Valenzuela R, Lovrien E, Maumenee I, Litt M. A second gene for cerulean cataracts maps to the beta crystallin region on chromosome 22. *Genomics* 1996; 35:539-542.
- Conneally PM, Wilson AF, Merritt AD, Helveston EM, Palmer CG, Wang LY. Confirmation of genetic heterogeneity in autosomal dominant forms of cataracts from linkage studies. *Cytogenet Cell Genet* 1978; 22:295-297.
- Scott MH, Hejtmancik JF, Wozencraft LA, Reuter LM, Parks MM, Kaiser-Kupfer MI. Autosomal dominant congenital cataract. Interocular phenotypic heterogeneity. *Ophthalmology* 1994; 101:866-871.
- Wistow G, Turnell B, Summers L, Slingsby C, Moss D, Miller L, Lindley P, Blundell T. X-ray analysis of the eye lens protein gamma-II crystallin at 1.9 Å resolution. *J Mol Biol* 1983; 170:175-202.
- Piatigorsky J, Chepelinsky AB, Hejtmancik JF, Borras T, Das GC, Hawkins JW, Zelenka PS, King CR, Beebe DC, Nickerson JM. Expression of crystallin gene families in the differentiating eye lens. In: Davidson EH, Flirtel RA, editors. Molecular biology of development: UCLA symposia on molecular and cellular biology. 19th ed. New York: A. R. Liss; 1984. p. 331-349.
- Piatigorsky J, Zelenka PS. Transcriptional regulation of crystallin genes: cis elements, trans-factors and signal transduction systems in the lens. In: Wasserman PW, editor. Advances in developmental biochemistry. Stamford, CT: JAI Press; 1991. p. 211-256.
- Aarts HJ, Lubsen NH, Schoenmakers JG. Crystallin gene expression during rat lens development. *Eur J Biochem* 1989; 183:31-36.
- Basti S, Hejtmancik JF, Padma T, Ayyagari R, Kaiser-Kupfer MI, Murty JS, Rao GN. Autosomal dominant zonular cataract with sutural opacities in a four-generation family. *Am J Ophthalmol* 1996; 121:162-168.
- Rogan PK, Schneider TD. Using information content and base frequencies to distinguish mutations from genetic polymorphisms in splice junction recognition sites. *Hum Mutat* 1995; 6:74-76.
- Rogan PK, Faux BM, Schneider TD. Information analysis of human splice site mutations. *Hum Mutat* 1998; 12:153-171.
- Schneider TD. Information content of individual genetic sequences. *J Theor Biol* 1997; 189:427-441.
- Schneider TD. Sequence walkers: a graphical method to display how binding proteins interact with DNA or RNA sequences. *Nucleic Acids Res* 1997; 25:4408-4415.
- Hogg D, Tsui LC, Gorin M, Breitman ML. Characterization of the human beta-crystallin gene Hu beta A3/A1 reveals ancestral relationships among the beta gamma-crystallin superfamily. *J Biol Chem* 1986; 261:12420-12427.
- Lampi KJ, Ma Z, Shih M, Shearer TR, Smith JB, Smith DL, David LL. Sequence analysis of betaA3, betaB3, and betaA4 crystallins completes the identification of the major proteins in young human lens. *J Biol Chem* 1997; 272:2268-2275.
- Treton JA, Jacquemin E, Courtois Y, Jeanny JC. Differential localization by in situ hybridization of specific crystallin transcripts during mouse lens development. *Differentiation*. 1991; 47:143-147.
- Van Leen RW, Breuer ML, Lubsen NH, Schoenmakers JG. Developmental expression of crystallin genes: in situ hybridization reveals a differential localization of specific mRNAs. *Dev Biol* 1987; 123:338-345.
- Hejtmancik JF. The genetics of cataract: our vision becomes clearer. *Am J Hum Genet* 1998; 62:520-525.
- Chambers C, Russell P. Deletion mutation in an eye lens beta-crystallin. An animal model for inherited cataracts. *J Biol Chem* 1991; 266:6742-6746.
- Cartier M, Breitman ML, Tsui LC. A frameshift mutation in the gamma E-crystallin gene of the Elo mouse. *Nat Genet* 1992; 2:42-45.
- Rodriguez IR, Gonzalez P, Zigler JS Jr, Borras T. A guinea-pig hereditary cataract contains a splice-site deletion in a crystallin gene. *Biochim Biophys Acta* 1992; 1180:44-52.
- Brakenhoff RH, Henskens HA, van Rossum MW, Lubsen NH, Schoenmakers JG. Activation of the gamma E-crystallin pseudogene in the human hereditary Coppock-like cataract. *Hum Mol Genet* 1994; 3:279-283.
- Litt M, Carrero-Valenzuela R, LaMorticella DM, Schultz DW, Mitchell TN, Kramer P, Maumenee IH. Autosomal dominant cerulean cataract is associated with a chain termination mutation in the human beta-crystallin gene CRYBB2. *Hum Mol Genet* 1997; 6:665-668.
- Horwitz J. Alpha-crystallin can function as a molecular chaperone. *Proc Natl Acad Sci U S A* 1992; 89:10449-10453.
- Kantorow M, Horwitz J, Sergeev YV, Hejtmancik JF, Piatigorsky J. Extralenticular expression, cAMP-dependent kinase phosphorylation and autophosphorylation of β B2-crystallin. *Invest Ophthalmol Vis Sci* 1997; 38:S205.
- Srivastava OP, Srivastava SK. Purification and characterization of a sodium deoxycholate-activatable proteinase activity, possibly associated with β A3/A1-crystallin from human lenses. *Invest Ophthalmol Vis Sci* 1996; 37:S421.

36. Quax-Jeuken Y, Janssen C, Quax W, van den Heuvel R, Bloemendal H. Bovine beta-crystallin complementary DNA clones. Alternating proline/alanine sequence of beta B1 subunit originates from a repetitive DNA sequence. *J Mol Biol* 1984; 180:457-72.
37. Werten PJ, Carver JA, Jaenicke R, de Jong WW. The elusive role of the N-terminal extension of beta A3- and beta A1-crystallin. *Protein Eng* 1996; 9:1021-8.
38. Hope JN, Chen HC, Hejtmancik JF. Beta A3/A1-crystallin association: role of the N-terminal arm. *Protein Eng* 1994; 7:445-51.
39. Krawczak M, Reiss J, Cooper DN. The mutational spectrum of single base-pair substitutions in mRNA splice junctions of human genes: causes and consequences. *Hum Genet* 1992; 90:41-54.
40. Blundell T, Lindley P, Miller L, Moss D, Slingsby C, Tickle I, Turnell B, Wistow G. The molecular structure and stability of the eye lens: x-ray analysis of gamma-crystallin II. *Nature* 1981; 289:771-777.