

Auxin and Plant-Microbe Interactions

Stijn Spaepen and Jos Vanderleyden

Centre of Microbial and Plant Genetics, Department of Microbial and Molecular Systems, Katholieke Universiteit Leuven, B-3001 Leuven, Belgium

Correspondence: jozef.vanderleyden@biw.kuleuven.be



Microbial synthesis of the phytohormone auxin has been known for a long time. This property is best documented for bacteria that interact with plants because bacterial auxin can cause interference with the many plant developmental processes regulated by auxin. Auxin biosynthesis in bacteria can occur via multiple pathways as has been observed in plants. There is also increasing evidence that indole-3-acetic acid (IAA), the major naturally occurring auxin, is a signaling molecule in microorganisms because IAA affects gene expression in some microorganisms. Therefore, IAA can act as a reciprocal signaling molecule in microbe-plant interactions. Interest in microbial synthesis of auxin is also increasing in yet another recently discovered property of auxin in *Arabidopsis*. Down-regulation of auxin signaling is part of the plant defense system against phytopathogenic bacteria. Exogenous application of auxin, e.g., produced by the pathogen, enhances susceptibility to the bacterial pathogen.

The phytohormone auxin (from the Greek “auxein,” meaning to grow) regulates a whole repertoire of plant developmental processes, as documented in previous articles on this topic. Perhaps less well known is the fact that some microorganisms also produce auxin (Costacurta and Vanderleyden 1995; Patten and Glick 1996). In their interaction with plants, these microorganisms can interfere with plant development by disturbing the auxin balance in plants. This is best documented for phytopathogenic bacteria like *Agrobacterium* spp. and *Pseudomonas savastanoi* pv. *savastanoi*, causing tumors and galls, respectively (Jameson 2000; Mole et al. 2007), and plant growth promoting rhizobacteria (PGPR) such as *Azospirillum* spp. that impact on plant root development

(Persello-Cartieaux et al. 2003; Spaepen et al. 2007a). The term rhizobacteria refers to the fact that their numbers are highly enriched in the rhizosphere, i.e., the narrow band of soil that surrounds the root (Hiltner 1904; Smalla et al. 2006; van Loon 2007). Of more recent date is the observation that auxin (indole-3-acetic acid or IAA) is a signaling molecule in some microorganisms (Spaepen et al. 2007a). Bringing these data together, it follows that auxin can have a major impact in microorganism-plant interactions. This is the main theme addressed in this article. Finally, the recent finding that auxin signaling in plants is also part of the *Arabidopsis* defense response against a leaf pathogen (Navarro et al. 2006) is discussed in relation to bacterial IAA synthesis.

Editors: Mark Estelle, Dolf Weijers, Ottoline Leyser, and Karin Ljung
Additional Perspectives on Auxin Signaling available at www.cshperspectives.org

Copyright © 2011 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a001438
Cite this article as *Cold Spring Harb Perspect Biol* 2011;3:a001438

S. Spaepen and J. Vanderleyden

BIOSYNTHESIS PATHWAYS IN BACTERIA

Production of IAA has been reported for many bacteria. It is even assumed that over 80% of the bacteria isolated from the rhizosphere are capable to synthesize IAA (Patten and Glick 1996; Khalid et al. 2004). The main precursor for the synthesis of IAA is tryptophan. Addition of tryptophan to culture media results in all cases in higher IAA production. Biosynthesis of tryptophan starts from the metabolic node chorismate in a five-step reaction encoded by the *trp* genes. The branch point compound chorismate is synthesized starting from phosphoenolpyruvate and erythrose 4-phosphate in the shikimate pathway, a common pathway for the biosynthesis of aromatic amino acids and many secondary metabolites (Dosselaere and Vanderleyden 2001; Merino et al. 2008).

Starting with tryptophan, at least five different pathways have been described for the synthesis of IAA, and most pathways show similarity to those described in plants, although some intermediates can differ (Patten and Glick 1996; Woodward and Bartel 2005; Spaepen et al. 2007a). The IAA biosynthetic pathways have

been identified using several genetic and biochemical methods. However, not all genes, enzymes encoded, or intermediates have been characterized. Especially, the use of feeding experiments with labeled precursors in combination with the identification of intermediates has been a major source for the identification of biosynthetic pathways. The pathways described in literature are illustrated in Figure 1 and are mostly named according to a key intermediate of the given pathway. In several studied microorganisms, redundancy of IAA biosynthetic pathways is observed, meaning that multiple pathways are present and active in one single microorganism. This observation is mainly based on knockout studies in which some IAA production could still be observed after inactivation of a single pathway. In the following sections, an overview of the IAA biosynthetic pathways in bacteria is given.

Indole-3-Acetamide Pathway

As the indole-3-acetamide (IAM) pathway is present in several pathogens and can contribute to the virulence of these strains by producing




Figure 1. Overview of IAA biosynthetic pathways in bacteria. Dashed lines refer to the name of pathways, mostly named after an intermediate. IAAId, indole-3-acetaldehyde; IAM, indole-3-acetamide; IPDC, indole-3-pyruvate decarboxylase; Trp, tryptophan. Adapted from Spaepen et al. (2007).



high amounts of IAA (Jameson 2000; Persello-Cartieaux et al. 2003), the genes/proteins and their regulation involved in this pathway are well characterized. The pathway consists of two distinct steps. In the first step tryptophan monooxygenase (encoded by *iaaM* gene) converts tryptophan to IAM; in the second step IAM is hydrolyzed to IAA and ammonia by an IAM hydrolase (encoded by *iaaH* gene). Both genes have been characterized from many phytopathogens such as *Agrobacterium tumefaciens*, *P. savastanoi*, *Pseudomonas syringae*, *Pantoea agglomerans*, but also in symbiotic nitrogen-fixing bacteria belonging to *Rhizobium* and *Bradyrhizobium* species (Sekine et al. 1989; Clark et al. 1993; Morris 1995; Theunis et al. 2004). In *A. tumefaciens*, these genes are located in the T-DNA region of the pTi plasmid together with the *ipt* gene (coding for dimethylallyl-pyrophosphate:AMP dimethylallyltransferase), involved in cytokinin production (for reviews see Hooykaas and Beijersbergen 1994; Costacurta and Vanderleyden 1995). *A. tumefaciens* can infect dicotyledonous plants at wound sites by transferring the T-DNA inside the plant cells (genetic colonization), where it is stably integrated into the genome. Several genes encoded on the T-DNA are expressed in the plant nucleus, causing an imbalance in the hormonal status of the plant cells and thereby inducing crown gall tumors (for reviews see Zambryski et al. 1989; Zambryski 1992; Hooykaas and Beijersbergen 1994). In the plant pathogen *P. savastanoi* pv. *savastanoi*, which produces high amounts of IAA in culture medium (Glickmann et al. 1998; Manulis et al. 1998; Buell et al. 2003), the *iaaM* and *iaaH* genes are located on plasmid pIAA1 (Comai and Kosuge 1982). Both genes are clustered in the *iaa* operon, and loss of the pIAA1 plasmid results in the loss of IAA production in free living cultures. Strains without the pIAA1 plasmid no longer cause gall formation on host plants such as olive and oleander.

In the symbiotic bacteria *Rhizobium fredii* and *Bradyrhizobium japonicum*, the production of IAA via the IAM pathway was suggested, as bacterial cultures can convert an IAM analog. A gene (*bam*) with sequence similarity with

iaaH could be isolated from *B. japonicum* (Sekine et al. 1988; Sekine et al. 1989). Later, the presence of the IAM pathway was also confirmed for *Rhizobium* sp. strain NGR234 by LC-MS detection of intermediates (Theunis et al. 2004).

Indole-3-Pyruvate Pathway

The indole-3-pyruvate (IPA) pathway is a major auxin pathway in plants. The IPA pathway could be shown in many bacteria such as phytopathogens (*P. agglomerans*), plant beneficial bacteria (*Azospirillum*, *Bacillus*, *Bradyrhizobium*, *Enterobacter cloacae*, *Paenibacillus*, *Pseudomonas*, and *Rhizobium*), and even cyanobacteria. The pathway consists of three steps. In a first step the precursor tryptophan is transaminated to IPA by an aminotransferase. Aminotransferase activity has been shown in different IAA-producing bacteria such as *Azospirillum brasilense* and *Azospirillum lipoferum* (Ruckdäschel et al. 1988; Baca et al. 1994). In the rate-limiting step IPA is decarboxylated to indole-3-acetaldehyde (IAAld), which is then oxidized to IAA by a dehydrogenase. The decarboxylation step is catalyzed by the key enzyme indole-3-pyruvate decarboxylase (encoded by *ipdC* gene). Based on phylogenetic and biochemical analysis, this group of enzymes could be divided into two subgroups, indicating a distinct origin of the encoding genes (Spaepen et al. 2007b). Several members have been isolated and characterized from diverse bacteria such as *A. brasilense*, *A. lipoferum*, *E. cloacae*, *Pseudomonas putida*, and *P. agglomerans* (Koga et al. 1991; Costacurta et al. 1994; Brandl and Lindow 1996; Patten and Glick 2002b). Inactivation of these genes results in impaired IAA production (sometimes >90%), indicating the importance of the encoded enzymes and consequently the IPA pathway in the IAA biosynthesis (Prinsen et al. 1993; Costacurta et al. 1994). Biochemically, the indole-3-pyruvate decarboxylase is best characterized in *E. cloacae* (Schütz et al. 2003a; Schütz et al. 2005) and *A. brasilense*. In *A. brasilense*, the decarboxylase can use both indole-3-pyruvate and phenylpyruvate as substrates (Spaepen et al. 2007b). In addition, the

S. Spaepen and J. Vanderleyden

crystal structure of both indole-3-pyruvate decarboxylases has been determined (Schütz et al. 2003b; Versées et al. 2007a; Versées et al. 2007b).

Other Pathways Starting from Tryptophan

In the tryptamine (both TAM and TRM are used as abbreviations) pathway, tryptophan is first decarboxylated to tryptamine by a tryptophan decarboxylase and tryptamine is, in contrast to the TAM pathway in plants, directly converted to IAAld by an amine oxidase. In *Bacillus cereus*, tryptophan decarboxylase activity could be identified (Perley and Stowe 1966), and in *Azospirillum* exogenously supplied tryptamine is converted to IAA (Hartmann et al. 1983).

The indole-3-acetonitrile (IAN) pathway in plants has been suggested and studied for many years, but the exact role of the involved players is still under debate. In bacteria, nitrilases with specificity for IAN have been detected (Nagasawa et al. 1990; Kobayashi et al. 1993; Howden et al. 2009). In several *Agrobacterium* and *Rhizobium* spp., nitrile hydratase and amidase activity could be measured suggesting the conversion of IAN to IAA via IAM. In addition, the nitrile hydratase enzyme from *A. tumefaciens* could be purified and characterized (Kobayashi et al. 1995).

The tryptophan side-chain oxidase (TSO) pathway has only been shown in *Pseudomonas fluorescens* CHA0, and in this pathway tryptophan is directly converted to IAAld bypassing IPA. In the last step IAAld is oxidized to IAA as in the IPA pathway (Oberhänsli et al. 1991).

Tryptophan-Independent Pathway

Analysis of mutants in the tryptophan biosynthetic pathway showed that plants are able to synthesize IAA without the tryptophan precursor, and the branch point for this tryptophan-independent pathway arises at indole-3-glycerol phosphate or indole (Last et al. 1991; Normanly et al. 1993). However, no genes or proteins involved in this pathway could be identified yet in plants (Woodward and Bartel 2005). In

bacteria, a tryptophan-independent pathway was suggested to occur in *A. brasilense* by feeding experiments with labeled tryptophan (Prinsen et al. 1993). However, also in this case, no firm genetic or biochemical evidence has been provided to support this pathway.

IAA Conjugation and Degradation

Some intermediates of the IAA biosynthetic pathways can be reversibly converted into storage compounds (e.g., reduction of IAAld to indole-3-lactic acid), although the exact function of these compounds is not known. In plants much of the IAA molecules are found covalently linked to sugars or amino acids, known as IAA-conjugates (Woodward and Bartel 2005). These compounds have diverse roles, such as protection against degradation (although in some cases conjugation is also suggested as a first step in degradation), storage, and transport, and are important to maintain the IAA homeostasis in the cells (Cohen and Bandurski 1982; Seidel et al. 2006). In bacteria the presence of IAA-conjugates has only been shown for *P. savastanoi* pv. *savastanoi*. In addition, a gene, involved in the conjugation process and located on the pIAA1 plasmid, was characterized as an indole-3-acetic acid-lysine synthetase (*iaaL* gene). This enzyme catalyzes the formation of an amide bond between the carboxyl group of IAA and the epsilon amino group of lysine in an ATP-dependent way (Glass and Kosuge 1986; Glass and Kosuge 1988). The importance of this process was shown by plant experiments with an *iaaL* knockout mutant. This mutant is unable to accumulate IAA conjugates in the bacterial culture medium and to form typical gall formation, indicating an impaired virulence. It was hypothesized that IAA-conjugates are produced by *P. savastanoi* to protect IAA against degradation by plant enzymes and are hydrolyzed inside plant cells by plant hydrolases into biologically active IAA, promoting gall formation (Glass and Kosuge 1988).

In plants, the degradation of IAA can occur via decarboxylative catabolism of IAA (modification of both side-chain and indole

S. Spaepen and J. Vanderleyden

two-component system acts as a negative regulator of the tryptophan-dependent IAA biosynthesis (Haas and Keel 2003; Kang et al. 2006). The GacS/GacA system controls the expression of genes, of which many are induced during late logarithmic growth phase.

In *E. cloacae* UW5, a putative TyrR box, similar to the recognition sequence of TyrR, was found in the promoter region of the *ipdC* gene. TyrR is a regulatory protein involved in transport and metabolism of aromatic amino acids. TyrR is directly involved in the induction of *ipdC* gene expression in *E. cloacae* in response to tryptophan, phenylalanine, and tyrosine, suggesting a broader function of indole-3-pyruvate decarboxylases (Ryu and Patten 2008).

In *Xanthomonas axonopodis*, IAA production is increased in the presence of leaf extract from the host plant *Citrus sinensis* (Costacurta et al. 1998). In *P. agglomerans*, *ipdC* gene expression is dramatically increased when the bacterium is grown on plant surfaces, but the specific compounds involved are not known (Brandl and Lindow 1997).

In *Ralstonia solanacearum*, there is a link between the type III secretion system (TTSS) and IAA synthesis. The host-responsive regulator, HrpG, of the TTSS activation cascade not only controls genes of the TTSS but also genes not belonging to the TTSS pathway including other virulence determinants and genes involved in adaptation to life in the host, such as phytohormone production. Auxin synthesis is reduced in an *hrpG* mutant (Valls et al. 2006).

In *Erwinia chrysanthemi*, known to produce IAA via the IAM pathway, it was found that in an *iaaM* mutant, expression of TTSS genes and the gene encoding a pectate lyase is lower as compared to the wild type (Yang et al. 2007).

In *P. agglomerans* pv *gypsophilae*, the interaction of the TTSS, quorum sensing and IAA biosynthesis in relation to plant interaction (epiphytic fitness and virulence) was investigated. It was found that IAA (and cytokinins) are involved in the regulation of the quorum sensing system and TTSS regulatory genes (Chalupowicz et al. 2009). Interestingly, as this bacterium possesses both the IAM and IPA pathway for IAA biosynthesis, effects were

found opposite for mutants in the IPA and the IAM pathways, respectively.

In *Rhizobium* sp. strain NGR234, flavonoid signal molecules can induce IAA production, and the transcriptional regulators NodD1 and NodD2 in combination with *nod*-box NB15 upstream the IAA biosynthesis operon are involved in the induction (Theunis et al. 2004).

AUXIN AS A MICROBIAL SIGNAL MOLECULE

Changes in bacterial gene expression under the influence of IAA were first described for the *ipdC* gene of *A. brasilense* (see earlier discussion, positive feedback regulation of IAA synthesis). In the phytopathogen *A. tumefaciens*, it was shown that IAA is a signal molecule because *vir* gene expression is inhibited by IAA (by competing with the phenolic plant signal for interaction with the VirA/G system). This control mechanism is proposed as a signal for *A. tumefaciens* indicating that plant transformation (transfer and expression of T-DNA) was successful (Liu and Nester 2006). In a follow-up study, microarray analyses revealed that addition of IAA represses the *vir* regulon and *chv* genes of *A. tumefaciens*, and that most of these changes are also induced by the addition of salicylic acid. Expression of a gene encoding an ABC-type transporter, probably involved in the uptake of IAA, was found to be specifically up-regulated by IAA (Yuan et al. 2008).

In the yeast *Saccharomyces cerevisiae*, the addition of IAA to the culture medium stimulates adhesion and filamentation (inducing invasive growth), and a cell-surface protein encoded by *FLO11* is involved in these changes (as supported by the increase of *FLO11* transcription). At higher IAA concentrations, cell growth is arrested. Transcript profiling revealed a YAP-1 (fungal-specific transcriptional activator involved in pleiotropic drug resistance) binding site in the promoter of IAA-regulated genes; interestingly, a *yap1-1* mutant was found hypersensitive to IAA, suggesting that YAP-1 is a key mediator in the auxin response in yeast. It is postulated that IAA at plant wound sites can serve as a cue for yeast and fungi to sense and adapt to the plant environment (Prusty et al.



2004). In *Escherichia coli*, it was found that IAA-induced cells are more resistant to different stress agents, probably by the enhanced formation of trehalose, lipopolysaccharides, exopolysaccharides, and biofilm. In the presence of IAA, expression of genes encoding cell envelope components and proteins involved in the adaptation to adverse conditions are induced (Bianco et al. 2006a). Further expression studies revealed that genes involved in the tricarboxylic acid cycle, glyoxylate shunt, and amino acid biosynthesis are up-regulated in the presence of IAA. In addition, increased activity of several enzymes in the central metabolism was shown (Bianco et al. 2006b).

The question can be addressed whether IAA as an effector molecule for bacterial gene expression is related to indole as a signaling molecule. Indole is well known as a communication signal in *E. coli* (Lee and Lee 2010) and is studied in relation to quorum sensing (Walters and Sperandio 2006; Lee et al. 2007). Indole controls biofilm formation in *E. coli* and *Vibrio cholerae* (Di Martino et al. 2003; Mueller et al. 2009). In a recent study (Lee et al. 2010), it was shown that indole-3-acetonitrile (an auxin) decreased biofilm formation of *E. coli* 0157:H7 and *Pseudomonas aeruginosa* virulence. This could be linked with specific effects of IAN on gene expression. However, in cases in which IAA has been found as an effector in bacterial gene expression, no such effect was observed with indole, indicating that auxin effects on bacterial gene expression are distinct from the well-known indole signaling.

AUXIN AS AN EFFECTOR MOLECULE IN BACTERIA PLANT INTERACTIONS

Phytopathogens

As already mentioned before, *A. tumefaciens* transfers T-DNA, encoding genes for auxin and cytokinin biosynthesis (among others), on plant infection into the plant genome. The genes encoding enzymes for auxin and cytokinin biosynthesis are under the control of strong constitutive eukaryotic promoters, resulting in a high accumulation of phytohormones. Depending

on the *Agrobacterium* strain, and consequently on the exact nature of the genes encoding enzymes for phytohormones, T-DNA transfer results either in undifferentiated plant tissue (tumors) or hairy roots (Zambryski 1992; Zupan et al. 2000).

The production of phytohormones (mainly auxins and cytokinins) in free-living cultures is a hallmark for many phytopathogenic gall-forming bacteria such as *P. agglomerans*, *P. savastanoi* pv. *savastanoi*, *P. syringae* pv. *syringae*, *R. solanacearum* and *Rhodococcus fascians* (Morris 1995; Vandeputte et al. 2005; Valls et al. 2006). Auxin biosynthesis genes are located either on the chromosome or on a plasmid. This location can modulate the IAA level as plasmids are mostly present in multiple copies. This difference is exemplified by the difference in IAA production between *P. savastanoi* pv. *savastanoi* and *P. syringae* pv. *syringae*. In the former strain, the genes are located on a plasmid, whereas in the latter strain IAA biosynthetic genes are located on the chromosomal DNA, resulting in a lower IAA production. The IAA production in *P. syringae* pv. *syringae* could be increased fourfold by introducing a low-copy plasmid, carrying the IAA biosynthetic operon (Mazzola and White 1994; Patten and Glick 1996; Brandl and Lindow 1996). As already mentioned above, the IAM pathway is mainly linked to phytopathology, whereas the IPA pathway is involved in epiphytic and rhizosphere fitness. In the gall-forming bacterium *P. agglomerans* pv. *gypsophilae*, both the IAM and the IPA pathway are present. Reduced gall size was observed by inactivation of the IAM pathway (as for many gall-forming bacteria), whereas only a minor reduction in pathogenicity could be observed on inactivation of the IPA pathway. The opposite effect was observed when looking at the epiphytic fitness. In addition, the difference in function of the two pathways could be linked to a difference in regulation of the pathways: the *ipdC* gene is up-regulated during colonization of the plant, whereas the *iaaM* gene is up-regulated during the later phases of the interaction with the plant such as gall formation. It was therefore concluded that the IAM pathway is mainly linked to

S. Spaepen and J. Vanderleyden

phytopathology, whereas the IPA pathway is involved in epiphytic fitness (Manulis et al. 1998).

Symbiosis

Auxin is involved in many processes of nodule formation by rhizobia in legume plants, such as founder cell specification (auxin transport inhibition mainly by flavonoids), nodule initiation and differentiation (auxin accumulation), vascular bundle formation, and nodule numbers (long distance auxin transport). Because many rhizobia are capable of producing IAA via different pathways, it is assumed that bacterially produced auxin can alter the auxin balance inside the plant. In addition, rhizobia can also indirectly influence the auxin homeostasis by interfering with plant auxin transport (Mathesius 2008).

In general, root nodules contain more auxin than non-nodulated roots (Badenochjones et al. 1983; Ghosh and Basu 2006). Nodules induced with low IAA-producing *Rhizobium* mutants contain less IAA than nodules initiated by the wild-type strain (Theunis 2005). Vice versa, bacteroids of a *B. japonicum* mutant, overproducing IAA, contain higher amounts of IAA (Hunter 1989). Also in *Rhizobium leguminosarum* bv. *viciae*, the introduction and expression of the IAM biosynthetic pathway resulted in *Vicia hirsuta* root nodules containing up to 60-fold more IAA than nodules invoked by the wild-type strain. Nitrogen fixation capacity (bacterial conversion of atmospheric N₂ to ammonium) in the former nodules was also increased (Camerini et al. 2008). A *Sinorhizobium meliloti* strain overexpression of the same IAM biosynthetic pathway showed increased tolerance to several stresses, and *Medicago truncatula* plants inoculated with this strain have a higher IAA content in nodules and roots and are better resistant to salt stress (Bianco and Defez 2009). In addition, inoculation of *M. truncatula* with the IAA-overproducing strain resulted in better plant growth under phosphorus deficiency because of the release of organic acids by the bacterium (Bianco and Defez 2010). In co-inoculation

studies with *Azospirillum* and *Rhizobium*, earlier and faster nodulation and higher crop yields are observed. In addition, nodule numbers and biological nitrogen fixation are increased. Using an *Azospirillum ipdC* mutant, producing 10% of IAA produced by the wild-type strain, the increase in nodulation and nitrogen fixation was not observed, indicating that bacterial IAA production is important in symbiosis (Remans et al. 2008). An extensive overlap of changes in protein level could be observed in *M. truncatula* in response to auxin treatment and *S. meliloti* inoculation, probably because of regulation of these proteins by auxin during the early stages of nodulation (van Noorden et al. 2007).

Plant Growth Promoting Rhizobacteria

Upon inoculation of plants with PGPR, a change in root architecture is observed, mainly as an increase in root hairs and lateral roots and shortening of the root length. In general, the total root surface increases, leading to an enhanced mineral uptake from the soil and root exudation (further stimulating root colonization by PGPR). Already for decades, these morphological changes of the root have been attributed to bacterial auxin production, and studies with mutants altered in IAA production confirm its important role. The plant growth promoting effect of bacteria, for which it has been shown that auxin is involved, is known as phytostimulation. However, because most IAA knock-out mutants are still able to promote some plant growth, IAA biosynthesis alone is not responsible for the overall observed effect (Xie et al. 1996; Dobbelaere et al. 1999; Dobbelaere et al. 2003). In the “additive hypothesis,” it was suggested that multiple mechanisms, such as dinitrogen fixation, phosphate solubilization, and 1-aminocyclopropane-1-carboxylate (ACC) deaminase activity, together with IAA biosynthesis, are responsible for the observed plant growth promotion and yield increase (Bashan and Holguin 1997). In addition, some plant growth-promoting traits do not work independently of each other as exemplified by IAA biosynthesis and ACC deaminase activity. Some

bacteria can promote plant growth via the action of the ACC deaminase enzyme. This enzyme converts ACC (the precursor of the stress hormone ethylene) in α -ketobutyrate and ammonia, lowering the ethylene level (inhibitor of root growth) inside the plant because of lack of precursor (Glick et al. 1998; Glick 2005). For a more detailed overview of plant growth-promoting rhizobacteria, we refer to other reviews (Steenhoudt and Vanderleyden 2000; Lugtenberg and Kamilova 2009).

AUXIN SIGNALING IN PLANT DEFENSE RESPONSE

Recently, a new intriguing role of auxin has been shown. Auxin physiology is a key factor in bacterial phytopathogenesis. On recognition of a biotrophic leaf pathogen by the plant, auxin signaling in *Arabidopsis* is down-regulated, thereby decreasing susceptibility for the pathogen (Navarro et al. 2006). This can be counteracted by the pathogen through the activity of an Hrp system (Chen et al. 2007). Both studies illustrate very nicely a prominent role for host auxin signaling in a particular aspect, effector-triggered susceptibility, pathogen associated molecular pattern-triggered immunity and effector-triggered susceptibility, respectively, of the four-phased model of the plant immune system (Jones and Dangl 2006). Furthermore, both studies showed that exogenous supply of auxin to the host promotes susceptibility to the bacterial pathogen and disease symptom development. As many bacterial pathogens are known to produce IAA (see above), it can be speculated that this property is part of the strategy used by the pathogen to circumvent the plant defense system. The same could apply for auxin producing plant growth-promoting rhizobacteria. However, it remains to be proven that IAA produced by these bacteria indeed contributes to better bacteria-plant-root interaction.

Nevertheless, biotrophic phytopathogens and plant beneficial bacteria are coming closer to each other when taking an auxin perspective. Clearly, answering questions in one direction (phytopathology) raises new fascinating questions in another direction (phytostimulation).

CONCLUDING REMARKS AND PERSPECTIVES

Recent landmark discoveries in plant auxin, in terms of the auxin receptor, auxin transporters, integration of auxin signaling with other signaling pathways (see other articles on this topic), in combination with previous scattered reports on the role of auxin in microbe-plant interactions and microbial synthesis of auxins, respectively, have given a tremendous boost to auxin research in bacteria interacting with plants. This is most visible in research of plant growth promoting rhizobacteria in which reported cases of phyto-stimulation are vastly increasing. This is well in line with the increasing interest in roots as central in efforts of scientists to produce crops with a better yield.

Whereas efforts to fully characterize the multiple microbial biosynthetic pathways of auxin should continue, and this is likely to be the case because similar efforts are being made by plant physiologists and biochemists, also the regulation of microbial IAA biosynthesis should be studied. This will help us to understand the role of IAA in microbial physiology. Until a number of years ago, IAA produced by microorganisms was merely considered as a secondary metabolite, possibly as the result of a detoxification process when tryptophan starts to accumulate in cells. However, the observation that multiple IAA biosynthesis pathways exist in IAA producing bacteria, like it is found for plants, already suggests additional or other roles for IAA in bacteria. The finding that IAA is used as a signal for gene regulation in some bacteria, both in IAA-producers and nonproducers, and *Saccharomyces cerevisiae*, further supports the idea of IAA being part of genetic networks in some microorganisms. When these microorganisms interact with plants as part of their ecological habitat, it becomes obvious that a reciprocal IAA-mediated signaling process in microbe-plant interactions is likely to occur (Lambrecht et al. 2000). Our further understanding of bacteria-plant interactions, be it pathogenic or beneficial, needs detailed studies that examine hormonal dynamics throughout the course of the interaction (see Fig. 3 for the

S. Spaepen and J. Vanderleyden




Figure 3. IAA dynamics from the bacterial point of view. IAA in plant-microbe interactions is part of a complex network. Bacteria are able to synthesize, conjugate, and degrade IAA, regulating the IAA homeostasis in their cells. Bacterial IAA production can be influenced by different plant and environmental cues as detailed in the text. Bacterial IAA is known as an effector molecule in plant-microbe interactions, both in pathogenesis and phytostimulation. Auxins are also used by bacteria as a signal molecule to steer microbial behavior.

different processes involved in IAA dynamics from the microbial point of view). With the observation that some rhizosphere bacteria are able to degrade IAA, it is clear that future research on plant roots for better plant growth and yield whereby IAA signaling is considered as a target need to take into account soil conditions, in particular the microbial activities in the rhizosphere.

ACKNOWLEDGMENTS

S.S. is a recipient of a postdoctoral fellowship grant from the Research Foundation-Flanders (FWO Vlaanderen).

REFERENCES

- Baca BE, Soto-Urzuá L, Xochinúa-Corona YG, Cuervo-García A. 1994. Characterization of two aromatic amino acid aminotransferases and production of indoleacetic acid in *Azospirillum* strains. *Soil Biol Biochem* **26**: 57–63.
- Badenoch-Jones J, Rolfe BG, Letham DS. 1983. Phytohormones, Rhizobium mutants, and nodulation in legumes. 3. Auxin metabolism in effective and ineffective pea root nodules. *Plant Physiol* **73**: 347–352.
- Bashan Y, Holguin G. 1997. *Azospirillum*-plant relationships: Environmental and physiological advances (1990–1996). *Can J Microbiol* **43**: 103–121.
- Bianco C, Defez R. 2009. *Medicago truncatula* improves salt tolerance when nodulated by an indole-3-acetic acid-overproducing *Sinorhizobium meliloti* strain. *J Exp Bot* **60**: 3097–3107.

- Bianco C, Defez R. 2010. Improvement of phosphate solubilization and *Medicago* plant yield by an indole-3-acetic acid-overproducing strain of *Sinorhizobium meliloti*. *Appl Env Microbiol* **76**: 4626–4632.
- Bianco C, Imperlini E, Calogero R, Senatore B, Amoresano A, Carpentieri A, Pucci P, Defez R. 2006a. Indole-3-acetic acid improves *Escherichia coli*'s defences to stress. *Arch Microbiol* **185**: 373–382.
- Bianco C, Imperlini E, Calogero R, Senatore B, Pucci P, Defez R. 2006b. Indole-3-acetic acid regulates the central metabolic pathways in *Escherichia coli*. *Microbiology-Sgm* **152**: 2421–2431.
- Brandl MT, Lindow SE. 1996. Cloning and characterization of a locus encoding an indolepyruvate decarboxylase involved in indole-3-acetic acid synthesis in *Erwinia herbicola*. *Appl Env Microbiol* **62**: 4121–4128.
- Brandl MT, Lindow SE. 1997. Environmental signals modulate the expression of an indole-3-acetic acid biosynthetic gene in *Erwinia herbicola*. *Mol Plant-Microbe Interactions* **10**: 499–505.
- Brandl MT, Quinones B, Lindow SE. 2001. Heterogeneous transcription of an indoleacetic acid biosynthetic gene in *Erwinia herbicola* on plant surfaces. *Proc Natl Acad Sci* **98**: 3454–3459.
- Buell CR, Joardar V, Lindeberg M, Selengut J, Paulsen IT, Gwinn ML, Dodson RJ, Deboy RT, Durkin AS, Kolonay JF, et al. 2003. The complete genome sequence of the *Arabidopsis* and tomato pathogen *Pseudomonas syringae* pv *tomato* DC3000. *Proc Natl Acad Sci* **100**: 10181–10186.
- Camerini S, Senatore B, Lonardo E, Imperlini E, Bianco C, Moschetti G, Rotino GL, Campion B, Defez R. 2008. Introduction of a novel pathway for IAA biosynthesis to rhizobia alters vetch root nodule development. *Arch Microbiol* **190**: 67–77.
- Chalupowicz L, Barash I, Panijel M, Sessa G, Manulis-Sasson S. 2009. Regulatory interactions between quorum-sensing, auxin, cytokinin, and the Hrp regulon in relation to gall formation and epiphytic fitness of *Pantoea agglomerans* pv *gypsophylae*. *Mol Plant-Microbe Interact* **22**: 849–856.
- Chen ZY, Agnew JL, Cohen JD, He P, Shan LB, Sheen J, Kunkel BN. 2007. *Pseudomonas syringae* type III effector AvrRpt2 alters *Arabidopsis thaliana* auxin physiology. *Proc Natl Acad Sci* **104**: 20131–20136.
- Clark E, Manulis S, Ophir Y, Barash I, Gafni Y. 1993. Cloning and characterization of *iaaM* and *iaaH* from *Erwinia herbicola* pathovar *gypsophylae*. *Phytopathology* **83**: 234–240.
- Cohen JD, Bandurski RS. 1982. Chemistry and physiology of the bound auxins. *Ann Rev Plant Physiol Plant Mol Biol* **33**: 403–430.
- Comai L, Kosuge T. 1982. Cloning and characterization of *iaaM*, a virulence determinant of *Pseudomonas savastanoi*. *J Bacteriol* **149**: 40–46.
- Costacurta A, Vanderleyden J. 1995. Synthesis of phytohormones by plant-associated bacteria. *Crit Rev Microbiol* **21**: 1–18.
- Costacurta A, Keijers V, Vanderleyden J. 1994. Molecular cloning and sequence analysis of an *Azospirillum brasilense* indole-3-pyruvate decarboxylase gene. *Mol Gen* **243**: 463–472.



- Costacurta A, Mazzafera P, Rosato YB. 1998. Indole-3-acetic acid biosynthesis by *Xanthomonas axonopodis* pv. citri is increased in the presence of plant leaf extracts. *FEMS Microbiol Lett* **159**: 215–220.
- Di Martino P, Fursy R, Bret L, Sundararaju B, Phillips RS. 2003. Indole can act as an extracellular signal to regulate biofilm formation of *Escherichia coli* and other indole-producing bacteria. *Can J Microbiol* **49**: 443–449.
- Dobbelaere S, Vanderleyden J, Okon Y. 2003. Plant growth-promoting effects of diazotrophs in the rhizosphere. *Crit Rev Plant Sci* **22**: 107–149.
- Dobbelaere S, Croonenborghs A, Thys A, Vande Broek A, Vanderleyden J. 1999. Phytostimulatory effect of *Azospirillum brasilense* wild type and mutant strains altered in IAA production on wheat. *Plant and Soil* **212**: 155–164.
- Dosselaere F, Vanderleyden J. 2001. A metabolic node in action: Chorismate-utilizing enzymes in microorganisms. *Crit Rev Microbiol* **27**: 75–131.
- Ghosh S, Basu PS. 2006. Production and metabolism of indole acetic acid in roots and root nodules of *Phaseolus mungo*. *Microbiol Res* **161**: 362–366.
- Glass NL, Kosuge T. 1986. Cloning of the gene for indoleacetic acid-lysine synthetase from *Pseudomonas syringae* subsp *savastanoi*. *J Bacteriol* **166**: 598–603.
- Glass NL, Kosuge T. 1988. Role of indoleacetic acid lysine synthetase in regulation of indoleacetic acid pool size and virulence of *Pseudomonas syringae* subsp *savastanoi*. *J Bacteriol* **170**: 2367–2373.
- Glick BR. 2005. Modulation of plant ethylene levels by the bacterial enzyme ACC deaminase. *FEMS Microbiol Lett* **251**: 1–7.
- Glick BR, Penrose DM, Li JP. 1998. A model for the lowering of plant ethylene concentrations by plant growth-promoting bacteria. *J Theoretical Biol* **190**: 63–68.
- Glickmann E, Gardan L, Jacquet S, Hussain S, Elasi M, Petit A, Dessaux Y. 1998. Auxin production is a common feature of most pathovars of *Pseudomonas syringae*. *Mol Plant-Microbe Interact* **11**: 156–162.
- Gysegom P. 2005. Study of the transcriptional regulation of a key gene in indole-3-acetic acid biosynthesis in *Azospirillum brasilense*. Ph.D thesis K.U. Leuven.
- Haas D, Keel C. 2003. Regulation of antibiotic production in root-colonizing *Pseudomonas* spp. and relevance for biological control of plant disease. *Ann Rev Phytopathol* **41**: 117–153.
- Hartmann A, Singh M, Klingmüller W. 1983. Isolation and characterization of *Azospirillum* mutants excreting high amounts of indoleacetic acid. *Can J Microbiol* **29**: 916–923.
- Hiltner L. 1904. Über neuere Erfahrungen und Probleme auf dem Gebiete der Bodenbakteriologie unter besonderer Berücksichtigung der Gründüngung und Brache. *Arbeiten der Deutsche Landwirtschaftliche Gesellschaft* **98**: 59–78.
- Hooykaas PJJ, Beijersbergen AGM. 1994. The virulence system of *Agrobacterium tumefaciens*. *Ann Rev Phytopathol* **32**: 157–179.
- Howden AJM, Rico A, Mentlak T, Miguet L, Preston GM. 2009. *Pseudomonas syringae* pv. *syringae* B728a hydrolyses indole-3-acetonitrile to the plant hormone indole-3-acetic acid. *Mol Plant Pathol* **10**: 857–865.
- Hunter WJ. 1989. Indole-3-acetic acid production by bacteroids from soybean root nodules. *Physiol Plant* **76**: 31–36.
- Jameson PE. 2000. Cytokinins and auxins in plant-pathogen interactions - An overview. *Plant Growth Reg* **32**: 369–380.
- Jones JDG, Dangl JL. 2006. The plant immune system. *Nature* **444**: 323–329.
- Kang BR, Yang KY, Cho BH, Han TH, Kim IS, Lee MC, Anderson AJ, Kim YC. 2006. Production of indole-3-acetic acid in the plant-beneficial strain *Pseudomonas chlororaphis* O6 is negatively regulated by the global sensor kinase GacS. *Current Microbiol* **52**: 473–476.
- Khalid A, Tahir S, Arshad M, Zahir ZA. 2004. Relative efficiency of rhizobacteria for auxin biosynthesis in rhizosphere and non-rhizosphere soils. *Aus J Soil Res* **42**: 921–926.
- Kobayashi M, Izui H, Nagasawa T, Yamada H. 1993. Nitrilase in biosynthesis of the plant hormone indole-3-acetic acid from indole-3-acetonitrile - Cloning of the *Alcaligenes* gene and site-directed mutagenesis of cysteine residues. *Proc Natl Acad Sci* **90**: 247–251.
- Kobayashi M, Suzuki T, Fujita T, Masuda M, Shimizu S. 1995. Occurrence of enzymes involved in biosynthesis of indole-3-acetic acid from indole-3-acetonitrile in plant-associated bacteria, *Agrobacterium* and *Rhizobium*. *Proc Natl Acad Sci* **92**: 714–718.
- Koga J, Adachi T, Hidaka H. 1991. Molecular cloning of the gene for indolepyruvate decarboxylase from *Enterobacter cloacae*. *Mol Gen Gen* **226**: 10–16.
- Lambrecht M, Okon Y, Vande Broek A, Vanderleyden J. 2000. Indole-3-acetic acid: a reciprocal signalling molecule in bacteria-plant interactions. *Trends in Microbiol* **8**: 298–300.
- Last RL, Bissinger PH, Mahoney DJ, Radwanski ER, Fink GR. 1991. Tryptophan mutants in *Arabidopsis*—the consequences of duplicated tryptophan synthase β genes. *Plant Cell* **3**: 345–358.
- Lee JH, Lee J. 2010. Indole as an intercellular signal in microbial communities. *FEMS Microbiol Rev* **34**: 426–444.
- Lee JH, Cho MH, Lee J. 2010. 3-Indolylacetonitrile decreases *Escherichia coli* O157:H7 biofilm formation and *Pseudomonas aeruginosa* virulence. *Environ Microbiol* doi: 10.1111/j.1462-2920.2010.02308.x.
- Lee JT, Jayaraman A, Wood TK. 2007. Indole is an interspecies biofilm signal mediated by SdiA. *BMC Microbiol* **7**: 42.
- Leveau JHJ, Gerards S. 2008. Discovery of a bacterial gene cluster for catabolism of the plant hormone indole 3-acetic acid. *FEMS Microbiol Ecol* **65**: 238–250.
- Leveau JH, Lindow SE. 2005. Utilization of the plant hormone indole-3-acetic acid for growth by *Pseudomonas putida* strain 1290. *Appl Env Microbiol* **71**: 2365–2371.
- Liu P, Nester EW. 2006. Indoleacetic acid, a product of transferred DNA, inhibits *vir* gene expression and growth of *Agrobacterium tumefaciens* C58. *Proc Natl Acad Sci* **103**: 4658–4662.
- Lugtenberg B, Kamilova F. 2009. Plant-growth-promoting rhizobacteria. *Ann Rev Microbiol* **63**: 541–556.
- Malhotra M, Srivastava S. 2008. Organization of the *ipdC* region regulates IAA levels in different *Azospirillum*

S. Spaepen and J. Vanderleyden

- brasilense* strains: molecular and functional analysis of *ipdC* in strain SM. *Env Microbiol* **10**: 1365–1373.
- Manulis S, Haviv-Chesner A, Brandl MT, Lindow SE, Barash I. 1998. Differential involvement of indole-3-acetic acid biosynthetic pathways in pathogenicity and epiphytic fitness of *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant-Microbe Interact* **11**: 634–642.
- Mathesius U. 2008. Auxin: at the root of nodule development? *Funct Plant Biol* **35**: 651–668.
- Mazzola M, White FF. 1994. A mutation in the indole-3-acetic-acid biosynthesis pathway of *Pseudomonas syringae* pv. *syringae* affects growth in *phaseolus vulgaris* and syringomycin production. *J Bacteriol* **176**: 1374–1382.
- Merino E, Jensen RA, Yanofsky C. 2008. Evolution of bacterial trp operons and their regulation. *Curr Opin Microbiol* **11**: 78–86.
- Mole BM, Baltrus DA, Dangel JL, Grant SR. 2007. Global virulence regulation networks in phytopathogenic bacteria. *Trends Microbiol* **15**: 363–371.
- Morris RO. 1995. Genes specifying auxin and cytokinin biosynthesis in prokaryotes. In *Plant hormones* (ed. Davies PJ) pp. 318–339. Kluwer Academic, Dordrecht, The Netherlands.
- Mueller RS, Beyhan S, Saini SG, Yildiz FH, Bartlett DH. 2009. Indole acts as an extracellular cue regulating gene expression in *Vibrio cholerae*. *J Bacteriol* **191**: 3504–3516.
- Nagasawa T, Mauger J, Yamada H. 1990. A novel nitrilase, arylacetone nitrilase, of *Alcaligenes faecalis* JM3 – Purification and characterization. *Eur J Biochem* **194**: 765–772.
- Navarro L, Dunoyer P, Jay F, Arnold B, Dharmasiri N, Estelle M, Voinnet O, Jones JDG. 2006. A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* **312**: 436–439.
- Normanly J, Cohen JD, Fink GR. 1993. *Arabidopsis thaliana* auxotrophs reveal a tryptophan-independent biosynthetic pathway for indole-3-acetic acid. *Proc Natl Acad Sci* **90**: 10355–10359.
- Oberhänsli T, Défago G, Haas D. 1991. Indole-3-acetic-acid (IAA) synthesis in the biocontrol strain CHA0 of *Pseudomonas fluorescens*—Role of tryptophan side-chain oxidase. *J Gen Microbiol* **137**: 2273–2279.
- Ona O, Smets I, Gysegom P, Bernaerts K, Van Impe J, Prinsen E, Vanderleyden J. 2003. The effect of pH on indole-3-acetic acid (IAA) biosynthesis of *Azospirillum brasilense* Sp7. *Symbiosis* **35**: 199–208.
- Ona O, Van Impe J, Prinsen E, Vanderleyden J. 2005. Growth and indole-3-acetic acid biosynthesis of *Azospirillum brasilense* Sp245 is environmentally controlled. *FEMS Microbiol Lett* **246**: 125–132.
- Patten CL, Glick BR. 1996. Bacterial biosynthesis of indole-3-acetic acid. *Canadian J Microbiol* **42**: 207–220.
- Patten CL, Glick BR. 2002a. Regulation of indoleacetic acid production in *Pseudomonas putida* GR12-2 by tryptophan and the stationary-phase sigma factor RpoS. *Canadian J Microbiol* **48**: 635–642.
- Patten CL, Glick BR. 2002b. Role of *Pseudomonas putida* indoleacetic acid in development of the host plant root system. *Appl Environ Microbiol* **68**: 3795–3801.
- Perley JW, Stowe BB. 1966. On the ability of *Taphrina deformans* to produce indoleacetic acid from tryptophan by way of tryptamine. *Plant Physiol* **41**: 234–237.
- Persello-Cartieaux F, Nussaume L, Robaglia C. 2003. Tales from the underground: molecular plant-rhizobacteria interactions. *Plant Cell Environ* **26**: 189–199.
- Prinsen E, Costacurta A, Michiels K, Vanderleyden J, Van Onckelen H. 1993. *Azospirillum brasilense* indole-3-acetic acid biosynthesis: evidence for a non-tryptophan dependent pathway. *Mol Plant-Microbe Interact* **6**: 609–615.
- Prusty R, Grisafi P, Fink GR. 2004. The plant hormone indoleacetic acid induces invasive growth in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci* **101**: 4153–4157.
- Remans R, Ramaekers L, Schelkens S, Hernandez G, Garcia A, Reyes JL, Mendez N, Toscano V, Mulling M, Galvez L, Vanderleyden J. 2008. Effect of *Rhizobium-Azospirillum* coinoculation on nitrogen fixation and yield of two contrasting *Phaseolus vulgaris* L. genotypes cultivated across different environments in Cuba. *Plant Soil* **312**: 25–37.
- Rothballer M, Schmid M, Fekete A, Hartmann A. 2005. Comparative *in situ* analysis of *ipdC-gfpmut3* promoter fusions of *Azospirillum brasilense* strains Sp7 and Sp245. *Environ Microbiol* **7**: 1839–1846.
- Ruckdäschel E, Kittell BL, Helinski DR, Klingmüller W. 1988. Aromatic amino acid aminotransferases of *Azospirillum lipoferum* and their possible involvement in IAA biosynthesis. In *Azospirillum IV: Genetics, physiology, ecology* (ed. Klingmüller W) Springer-Verlag, Berlin Heidelberg.
- Ryu RJ, Patten CL. 2008. Aromatic amino acid-dependent expression of indole-3-pyruvate decarboxylase is regulated by TyrR in *Enterobacter cloacae* UW5. *J Bacteriol* **190**: 7200–7208.
- Schütz A, Golbik R, König S, Hübner G, Tittmann K. 2005. Intermediates and transition states in thiamin diphosphate-dependent decarboxylases. A kinetic and NMR study on wild-type indolepyruvate decarboxylase and variants using indolepyruvate, benzoylformate, and pyruvate as substrates. *Biochemistry* **44**: 6164–6179.
- Schütz A, Golbik R, Tittmann K, Svergun DI, Koch MH, Hübner G, König S. 2003a. Studies on structure-function relationships of indolepyruvate decarboxylase from *Enterobacter cloacae*, a key enzyme of the indole acetic acid pathway. *Eur J Biochem* **270**: 2322–2331.
- Schütz A, Sandalova T, Ricagno S, Hübner G, König S, Schneider G. 2003b. Crystal structure of thiamin diphosphate-dependent indolepyruvate decarboxylase from *Enterobacter cloacae*, an enzyme involved in the biosynthesis of the plant hormone indole-3-acetic acid. *Eur J Biochem* **270**: 2312–2321.
- Seidel C, Walz A, Park S, Cohen JD, Ludwig-Muller J. 2006. Indole-3-acetic acid protein conjugates: Novel players in auxin homeostasis. *Plant Biol* **8**: 340–345.
- Sekine M, Ichikawa T, Kuga N, Kobayashi M, Sakurai A, Syono K. 1988. Detection of the IAA biosynthetic pathway from tryptophan via indole-3-acetamide in *Bradyrhizobium* spp. *Plant Cell Physiol* **29**: 867–874.
- Sekine M, Watanabe K, Syono K. 1989. Molecular cloning of a gene for indole-3-acetamide hydrolase from *Bradyrhizobium japonicum*. *J Bacteriol* **171**: 1718–1724.

- Smalla K, Sessitsch A, Hartmann A. 2006. The Rhizosphere: 'soil compartment influenced by the root'. *FEMS Microbiol Ecol* **56**: 165.
- Somers E, Ptacek D, Gysegom P, Srinivasan M, Vanderleyden J. 2005. *Azospirillum brasilense* produces the auxin-like phenylacetic acid by using the key enzyme for indole-3-acetic acid biosynthesis. *Appl Environ Microbiol* **71**: 1803–1810.
- Spaepen S, Vanderleyden J, Remans R. 2007a. Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiol Rev* **31**: 425–448.
- Spaepen S, Versées W, Gocke D, Pohl M, Steyaert J, Vanderleyden J. 2007b. Characterization of phenylpyruvate decarboxylase, involved in auxin production of *Azospirillum brasilense*. *J Bacteriol* **189**: 7626–7633.
- Steenhoudt O, Vanderleyden J. 2000. *Azospirillum*, a free-living nitrogen-fixing bacterium closely associated with grasses: genetic, biochemical and ecological aspects. *FEMS Microbiol Rev* **24**: 487–506.
- Theunis M. 2005. IAA biosynthesis in rhizobia its potential role in symbiosis. Universiteit Antwerpen Ph.D thesis.
- Theunis M, Kobayashi H, Broughton WJ, Prinsen E. 2004. Flavonoids, NodD1, NodD2, and nod-box NB15 modulate expression of the y4wEFG locus that is required for indole-3-acetic acid synthesis in *Rhizobium* sp. strain NGR234. *Mol Plant-Microbe Interact* **17**: 1153–1161.
- Valls M, Genin S, Boucher C. 2006. Integrated regulation of the type III secretion system and other virulence determinants in *Ralstonia solanacearum*. *PLOS* **2**: 798–807.
- van Loon LC. 2007. Plant responses to plant growth-promoting rhizobacteria. *Eur J Plant Pathol* **119**: 243–254.
- van Noorden GE, Kerim T, Goffard N, Wiblin R, Pellerone FI, Rolfe BG, Mathesius U. 2007. Overlap of proteome changes in *Medicago truncatula* in response to auxin and *Sinorhizobium meliloti*. *Plant Physiol* **144**: 1115–1131.
- Vande Broek A, Gysegom P, Ona O, Hendrickx N, Prinsen E, Van Impe J, Vanderleyden J. 2005. Transcriptional analysis of the *Azospirillum brasilense* indole-3-pyruvate decarboxylase gene and identification of a cis-acting sequence involved in auxin responsive expression. *Mol Plant-Microbe Interact* **18**: 311–323.
- Vande Broek A, Lambrecht M, Eggermont K, Vanderleyden J. 1999. Auxins upregulate expression of the indole-3-pyruvate decarboxylase gene in *Azospirillum brasilense*. *J Bacteriol* **181**: 1338–1342.
- Vandeputte O, Oden S, Mol A, Vereecke D, Goethals K, El Jaziri M, Prinsen E. 2005. Biosynthesis of auxin by the gram-positive phytopathogen *Rhodococcus fascians* is controlled by compounds specific to infected plant tissues. *Appl Environ Microbiol* **71**: 1169–1177.
- Versées W, Spaepen S, Vanderleyden J, Steyaert J. 2007a. The crystal structure of phenylpyruvate decarboxylase from *Azospirillum brasilense* at 1.5 angstrom resolution—Implications for its catalytic and regulatory mechanism. *FEBS J* **274**: 2363–2375.
- Versées W, Spaepen S, Wood MDH, Leeper FJ, Vanderleyden J, Steyaert J. 2007b. Molecular mechanism of allosteric substrate activation in a thiamine diphosphate-dependent decarboxylase. *J Biol Chem* **282**: 35269–35278.
- Walters M, Sperandio V. 2006. Quorum sensing in *Escherichia coli* and *Salmonella*. *Int J Med Microbiol* **296**: 125–131.
- Woodward AW, Bartel B. 2005. Auxin: Regulation, action, and interaction. *Ann Bot* **95**: 707–735.
- Xie H, Pasternak JJ, Glick BR. 1996. Isolation and characterization of mutants of the plant growth-promoting rhizobacterium *Pseudomonas putida* CR12-2 that overproduce indoleacetic acid. *Curr Microbiol* **32**: 67–71.
- Yang S, Zhang Q, Guo J, Charkowski AO, Glick BR, Ibekwe AM, Cooksey DA, Yang CH. 2007. Global effect of indole-3-acetic acid biosynthesis on multiple virulence factors of *Erwinia chrysanthemi* 3937. *Appl Environ Microbiol* **73**: 1079–1088.
- Yuan ZC, Liu P, Saenkham P, Kerr K, Nester EW. 2008. Transcriptome profiling and functional analysis of *Agrobacterium tumefaciens* reveals a general conserved response to acidic conditions (pH 5.5) and a complex acid-mediated signaling involved in *Agrobacterium*-plant interactions. *J Bacteriol* **190**: 494–507.
- Zambryski PC. 1992. Chronicles from the *Agrobacterium*—plant cell DNA transfer story. *Ann Rev Plant Physiol Plant Mol Biol* **43**: 465–490.
- Zambryski P, Tempe J, Schell J. 1989. Transfer and function of T-DNA genes from *Agrobacterium* Ti-plasmid and Ri-plasmid in plants. *Cell* **56**: 193–201.
- Zupan J, Muth TR, Draper O, Zambryski P. 2000. The transfer of DNA from *Agrobacterium tumefaciens* into plants: a feast of fundamental insights. *Plant J* **23**: 11–28.



Cold Spring Harbor Perspectives in Biology

Auxin and Plant-Microbe Interactions

Stijn Spaepen and Jos Vanderleyden

Cold Spring Harb Perspect Biol 2011; doi: 10.1101/cshperspect.a001438 originally published online November 17, 2010

Subject Collection [Auxin Signaling](#)

Fourteen Stations of Auxin

Jirí Friml

Computational Models of Auxin-Driven Patterning in Shoots

Mikolaj Cieslak, Andrew Owens and Przemyslaw Prusinkiewicz

Auxin Transporters—A Biochemical View

Ulrich Z. Hammes, Angus S. Murphy and Claus Schwechheimer

Structural Aspects of Auxin Signaling

Nicholas Morffy and Lucia C. Strader

The Story of Auxin-Binding Protein 1 (ABP1)

Richard Napier

Noncanonical Auxin Signaling

Heather Marie McLaughlin, Aaron Chun Hou Ang and Lars Østergaard

Casting the Net—Connecting Auxin Signaling to the Plant Genome

Yanfei Ma, Sebastian Wolf and Jan U. Lohmann

Auxin Plays Multiple Roles during Plant–Pathogen Interactions

Barbara N. Kunkel and Joshua M.B. Johnson

Auxin in Root Development

Suruchi Roychoudhry and Stefan Kepinski

Modeling Auxin Signaling in Roots: Auxin Computations

Jaap Rutten, Thea van den Berg and Kirsten ten Tusscher

The Systems and Synthetic Biology of Auxin

R. Clay Wright, Britney L. Moss and Jennifer L. Nemhauser

Auxin Does the SAMba: Auxin Signaling in the Shoot Apical Meristem

Markéta Pernisová and Teva Vernoux

Chemical Biology in Auxin Research

Ken-ichiro Hayashi

Uncovering How Auxin Optimizes Root Systems Architecture in Response to Environmental Stresses

Nicola Leftley, Jason Banda, Bipin Pandey, et al.

Auxin Interactions with Other Hormones in Plant Development

Serina M. Mazzoni-Putman, Javier Brumos, Chengsong Zhao, et al.

No Time for Transcription—Rapid Auxin Responses in Plants

Shiv Mani Dubey, Nelson B.C. Serre, Denisa Oulehlová, et al.

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?

Your wish is our command.



For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



**All Modifications and
Oligo Types Synthesized**

Oligo Modifications?
Your wish is our command.

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

