

**Bacillus megaterium--from simple soil
bacterium to industrial protein production host.**

Item Type	Article
Authors	Vary, Patricia S; Biedendieck, Rebekka; Fuerch, Tobias; Meinhardt, Friedhelm; Rohde, Manfred; Deckwer, Wolf-Dieter; Jahn, Dieter
Citation	Bacillus megaterium--from simple soil bacterium to industrial protein production host. 2007, 76 (5):957-67 Appl. Microbiol. Biotechnol.
DOI	10.1007/s00253-007-1089-3
Journal	Applied microbiology and biotechnology
Download date	04/08/2022 16:04:45
Link to Item	http://hdl.handle.net/10033/29012



This is a postprint of an article published in
Vary, P.S., Biedendieck, R., Fuerch, T., Meinhardt, F., Rohde, M.,
Deckwer, W.-D., Jahn, D.
Bacillus megaterium-from simple soil bacterium to industrial protein
production host.
(2007) Applied Microbiology and Biotechnology, 76 (5), pp. 957-967.

***Bacillus megaterium* – from simple soil bacterium to industrial protein production host**

5

Patricia S. Vary¹, Rebekka Biedendieck², Tobias Fuerch³, Friedhelm Meinhardt⁴, Manfred Rohde⁵,

10

Wolf – Dieter Deckwer³, Dieter Jahn^{2*}

¹Department of Biological Sciences, Northern Illinois University, DeKalb, IL 60115, USA;

15

²Institute of Microbiology, Technical University Braunschweig, Spielmannstrasse 7, 38106

Braunschweig, Germany; ³Institute of Biochemical Engineering (TU-BCE), Technical University

Braunschweig/Helmholtz Centre for Infection Research, Inhoffenstrasse 7, 38124 Braunschweig,

Germany; ⁴Institute for Molecular Microbiology and Biotechnology, Westfaelische Wilhelms-

University Muenster, Corrensstrasse 3, 48149 Muenster, Germany; ⁵Department of Microbial

20

Pathogenesis, Helmholtz Centre for Infection Research, Inhoffenstrasse 7, 38124 Braunschweig,

Germany.

25

*Corresponding author. Tel.: +49 (0) 531-391-5801; fax: +49 (0) 531-391-5854

E-mail address: d.jahn@tu-bs.de (D. Jahn)

URL: <http://www.tu-braunschweig.de/ifm>

Abstract

Bacillus megaterium has been industrially employed for over 50 years since it possesses some very useful and unusual enzymes, and a high capacity for the production of exoenzymes. It is also a desirable cloning host for the production of intact proteins since it does not possess external alkaline proteases and can stably maintain a variety of plasmid vectors. Genetic tools for this species include transducing phages and several hundred mutants covering the processes of biosynthesis, catabolism, division, sporulation, germination, antibiotic resistance, and recombination. The seven plasmids of *B. megaterium* strain QM B1551 contain several unusual metabolic genes that may be useful in bioremediation. Recently, several recombinant shuttle vectors carrying different strong inducible promoters and various combinations of affinity tags for simple protein purification have been constructed. Leader sequences mediated export of affinity tagged proteins into the growth medium was made possible. These plasmids are commercially available. For a broader application of *B. megaterium* in industry, sporulation and protease deficient as well as UV-sensitive mutants were constructed. The genome sequence of two different strains, plasmidless DSM319 and QM B1551 carrying seven natural plasmids, is now available. These sequences allow for a systems biotechnology optimization of the production host *B. megaterium*.

Altogether, a “toolbox” of hundreds of genetically characterized strains, genetic methods, vectors, hosts, and genomic sequences make *B. megaterium* an ideal organism for industrial, environmental, and experimental applications.

20

Keywords

Bacillus megaterium, high level production, biosafety, metabolic flux analysis, genome sequencing

Introduction

In 1884, De Bary named *B. megaterium* “big beast” because of its large size with a volume approximately 100 times that of *Escherichia coli* (**Fig. 1**) (De Bary 1884). It has been ideal for studies of cell structure, protein localization, sporulation, and membranes (Hrafnisdottir et al. 1997; 5 McCool and Cannon 2001). Lysogeny was discovered by Lwoff and Gutman using strain 899 and phage T (Lwoff and Gutmann 1950).

B. megaterium is a Gram-positive, mainly aerobic spore-forming bacterium found in widely diverse habitats from soil to seawater, sediment, rice paddies, honey, fish, and dried food. It can grow in simple media on over 62 carbon sources out of 95 tested including all tricarboxylic acid cycle 10 intermediates, formate, and acetate. This has made it an ideal industrial organism for over 50 years. Among its products are proteins like penicillin acylase (Martin et al. 1995; Suga et al. 1990), used to make synthetic penicillins, various amylases (Hebeda et al. 1988; Metz et al. 1988; Takasaki 1989; Vihinen and Mantsala 1989) which are of interest in starch modification in the baking industry, glucose dehydrogenase (Kittsteiner-Eberle et al. 1989; Nagao et al. 1992), used for 15 regeneration of the cofactor NADPH in biochemical reactions and glucose blood tests (**Tab. 1**). Further, it is used for the production of pyruvate, vitamin B₁₂ (Raux et al. 1998; Wolf and Brey 1986), fungicidal toxins, and oxetanocin, a viral inhibitor active on HIV (Human immunodeficiency virus), hepatitis B virus, and herpes simplex corneal ulcers (Morita et al. 1999; Shimada et al. 1986; Shiota et al. 1996; Tseng et al. 1991). Many of its industrial applications have been reviewed in 20 detail (Vary 1992; Vary 1994). Some of the most interesting proteins produced by *B. megaterium* are the family of P-450 cytochrome monooxygenases (He and Fulco 1991). These have been of great interest since they have considerable similarity to eukaryotic P-450 important in many disease conditions.

Genetics of *Bacillus megaterium*

The *B. megaterium* strain QM B1551 was used for the isolation of the transducing phage MP13 (Vary 1979; Vary et al. 1982). In this process, 41 further bacteriophages were characterized that are specific for this strain (Vary and Halsey 1980). Over the next several years, hundreds of mutants
5 were isolated and characterized for auxotrophy, recombination, division, sporulation, germination, antibiotic resistance, UV sensitivity, and neutral protease (Vary 1994). Surprisingly, the *Bacillus subtilis* map did not serve well in transductional mapping. However, the construction of a rough map covering approximately 80 % of the chromosome by the early 80's was managed (Vary 1993; Vary and Muse 1993). Protoplast transformation conditions were established in the laboratories of
10 Carlton and Alfoldi (Brown and Carlton 1980; Vorobjeva et al. 1980). Although protoplast transformation requires experimental experience and is less effective compared to standard *E. coli* transformation procedures, the ability to transform *B. megaterium* immediately led to the cloning of several genes, the study of its plasmids, and its use as a very efficient host for gene expression. The transposon Tn917 was introduced into strain QM B1551 and showed that it inserted randomly
15 (Bohall and Vary 1986). The transposons constructions allowing *lacZ* fusion genetics in *B. subtilis* (Camilli et al. 1990; Youngman et al. 1989) also work well in *B. megaterium*.

Natural Plasmids of *Bacillus megaterium*

Most strains of *B. megaterium* carry multiple plasmids, usually more than four. However, the well
20 characterized strain QM B1551 has seven resident plasmids ranging from a size of 5.4 to 165 kb representing over 11 % of cellular DNA (Kieselburg et al. 1984). The two smallest ones replicate by the rolling circle mechanism whereas the other five are theta replication plasmids (Stevenson et al. 1998). *B. megaterium* strain 216 studied by Carlton (Carlton and Helinski 1969) has ten plasmids. QM B1551 was randomly cured of plasmids so that the role of plasmid genes could be determined.

Very few phenotypic differences were determined for the various cured strains. However, one originated strain, PV361, was cured of all seven plasmids (Sussman et al. 1988). Surprisingly, its growth under laboratory conditions was close to wild type QM B1551. The strain has proven to be a good host for gene expression. It has been used for many years to produce the antigen for HIV
5 diagnostic kits (Ginsburgh et al. 1989).

To get an idea about the additional genetical material and its relevance for the organism, the DNA sequence of the plasmids of QM B1551 has been determined over the last few years. The plasmids are carrying genes encoding enzymes like dehydrogenases, amylases, a P-450 cytochrome, monooxygenases, carboxylases, a styrene monooxygenase, biotin carboxylases, permeases,
10 transposases, and integrases. Furthermore, genes encoding proteins for heavy metal export, many possible transport proteins, acyl carriers, sigma factors, and sterols were found. Redox, mobilization, and transfer genes, as well as sporulation and germination genes were identified (Kunnimalaiyaan et al. 2001; Kunnimalaiyaan and Vary 2002; Scholle et al. 2003). Moreover, there are genes very similar to some found on the virulence plasmids of *Bacillus anthracis* and at least
15 one complete rRNA operon with 18 tRNA genes (Kunnimalaiyaan et al. 2001). Interestingly, the five theta plasmids have unusually similar Rep proteins (82 – 95 % identity at the amino acid sequence level), as do the iterons they must bind to. Normally, such identity precludes compatibility. The replicons are found in several strains of *B. megaterium*, but not in other *Bacillus* species tested (Rosso and Vary 2005). Moreover, no proteins in the databases are similar to the
20 proteins encoded in the replicon region. Hence, the theta replicons appear to be unique. The two rolling circle plasmids (5.4 and 9.1 kb) are not similar to each other, but their replicons show similarities to sequences of plasmids known from *Bacillus thuringiensis* and *B. anthracis* (Scholle et al. 2003). In QM B1551, the plasmids are very stable and several have been followed through 100 generations with no selection or loss.

Overall, the genes found on the plasmids suggest that much exchange has been made with other Gram-positive bacteria. Many of the genes show no phenotype on laboratory media after their knock-out. Hence, they must be necessary for the success of *B. megaterium* in diverse habitats. Further, many of the genes are of unknown function, or they can only be assigned as “an ABC transporter” or a “kinase”, but cannot be assigned to a specific metabolic pathway. Encoded proteins may be candidates for unusual reactions and pathways yet to be discovered.

***Bacillus megaterium* as Expression Host for Recombinant Genes**

B. megaterium does not have alkaline proteases which advanced the recovery of recombinant proteins and has the ability to stably maintain plasmids. Hence, the plasmidless derivatives of *B. megaterium*, specifically PV361 (Sussman et al. 1988) and DSM319 (Stahl and Esser 1983), have proven to be excellent production hosts for intact foreign proteins.

***Bacillus megaterium* Vector Systems for Recombinant Protein Production**

Rygus and Hillen (1991) isolated the xylose-inducible promoter P_{xylA} with the gene for the regulatory repressor protein XylR from the genome of *B. megaterium*. In the absence of xylose, XylR binds to an operator sequence located in P_{xylA} preventing transcription of the *xyl* operon (Gärtner et al. 1988). In the presence of xylose in the cell, the sugar is bound to the repressor protein introducing a conformational change to the protein which leads to a release of XylR from the promoter. Now, the RNA polymerase efficiently recognizes the promoter and gene expression is induced approximately 350-fold. An additional level of regulation of the original *xyl* operon is mediated by glucose via the catabolite regulatory protein CcpA (Gärtner et al. 1988). Based on this promoter, Rygus and Hillen (1991) developed a free replicating plasmid-borne system for the xylose-inducible overproduction of recombinant proteins. The designed vector, called pWH1520,

was successfully employed for the recombinant intracellular production of prokaryotic and eukaryotic proteins including *E. coli* β -galactosidase, *B. megaterium* glucose dehydrogenase, *Acinetobacter calcoaceticus* mutarotase, human urokinase-like plasminogen activator (Rygus and Hillen 1991), and *Clostridium difficile* toxin A with a high relative molecular mass of 308,000 (Burger et al. 2003).

In further studies, the promoter region of pWH1520 was modified and optimized for recombinant gene expression (Malten et al. 2005a). The resulting vector pMM1520 allows for the simple cloning of target genes by the use of 15 different DNA restriction enzyme cleavage sites located in a newly designed multiple cloning site (**Fig. 2A**). Furthermore, inactivation of the binding site for the catabolite regulatory protein CcpA made promoter utilization independent of the presence of glucose in the growth medium. To further optimize the overall gene expression system, a derivative of the plasmidless *B. megaterium* strain DSM319, named WH323 (**Tab. 3**), was constructed (Rygus and Hillen 1992). This strain is deficient in the xylose isomerase gene *xylA* and consequently unable to consume the inducer xylose as carbon source.

15

Plasmids for the Production and Purification of Intracellular Recombinant Proteins

Based on this xylose-inducible protein production system, novel expression vectors for the intracellular production of recombinant affinity tagged proteins were developed (**Fig. 2B**) (Biedendieck et al. 2007b). Using these vectors, proteins of interest N- or C-terminally fused to a His₆- or Strep II-tag can be produced (Lichty et al. 2005). Testing this newly developed plasmid system with the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria* (Tsien 1998) as a model protein yielded up to 17.9 mg of soluble recombinant GFP per liter shaking flask cell culture (**Tab. 2**). A one-column purification step resulted in up to 9 mg purified GFP per liter cell culture for the differentially tagged GFP forms (**Tab. 2**). The vectors encode protease cleavage sites for

Factor Xa (Jenny et al. 2003) or the tobacco etch virus (TEV) protease (Kapust et al. 2002) which allow for the removal of the affinity tags. Hence, untagged pure GFP was obtained after protease digestion and a single second purification step in which the cleaved tag and the protease were removed (**Fig. 3**) (Biedendieck et al. 2007b).

5 In high cell density cultivations of *B. megaterium* without recombinant protein production, final biomass concentrations of up to 80 g per liter were achieved (Hollmann and Deckwer 2004; Malten et al. 2005a). For up-scaling the recombinant production process, GFP production in *B. megaterium* was analyzed after a glucose limited fed batch cultivation (Biedendieck et al. 2007b). This process yielded 52 g cell dry weight (CDW) per liter and up to 274 mg GFP per liter culture corresponding
10 to 5.2 mg GFP per g CDW. Flow cytometric analysis allowed for the detection of the recombinantly produced GFP at the single cell level during the fermentation process indicating that the recombinant protein production hardly influenced the viability of the cell population (Biedendieck et al. 2007b). The xylose-inducible vector system of *B. megaterium* was also successfully used for the high yield biotransformation of fructose into D-mannitol. For this purpose, *Leuconostoc*
15 *pseudomesenteroides* ATCC12291 mannitol dehydrogenase, converting D-fructose into D-mannitol, and *Mycobacterium vaccae* N10 formate dehydrogenase, which is necessary for regeneration of NADH reduction equivalents, were recombinantly synthesized and the resting recombinant *B. megaterium* strain produced large amounts of mannitol (personal communication). Here, the capability of *B. megaterium* to tolerate high amounts of formate was utilized (personal
20 communication). Interestingly, high amounts of formate dehydrogenase were only obtained after complete codon adaptation to the *B. megaterium* codon usage via the use of a completely synthetic gene.

Plasmids for the Production and Purification of Extracellular Recombinant Proteins

In order to maximize the capacity of *B. megaterium* for the secretion of exoenzymes in high amounts into the environment, a series of vectors employing the signal peptides (SP) of the *B. megaterium* extracellular esterase LipA (Malten et al. 2006) and penicillin G acylase (PGA) (Biedendieck et al. 2007a) was developed for the recombinant production of extracellular proteins. Levansucrase from *Lactobacillus reuteri* strain 121 (van Hijum et al. 2001; van Hijum et al. 2004) was secreted best using the leader peptide of *B. megaterium* LipA (Biedendieck et al. 2007a). More than 4 mg enzyme per liter culture of this large enzyme ($M_r = 110,000$) were found as the predominant protein in the growth medium of *B. megaterium* in shaking flask cultivations (Malten et al. 2006). Homologous penicillin G acylase (PGA) from *B. megaterium* strain ATCC14945 was secreted up to 40 mg per liter shake flask culture (Yang et al. 2006). A hydrolase of *Thermobifida fusca* (TFH) (Kleeberg et al. 2005), which possesses unique hydrolytic properties as it can act as esterase, lipase, cutinase, and is also able to cleave polyesters, yielded up to 2.6 mg per liter culture (Yang et al. 2007). Both proteins were directed by the signal peptide of LipA into the growth medium. An upscale of the recombinant PGA secretion process to a 2 liter bioreactor batch cultivation resulted in 28.5 mg recombinant PGA per liter (Yang et al. 2006) and 16.1 mg recombinant TFH per liter (Yang et al. 2007), respectively. The secretion vector system was extended by the addition of DNA sequences encoding His₆- and StrepII-tags (Lichty et al. 2005) (**Fig. 3C**). However, the stepwise introduction of affinity tags decreased the amount of produced and exported recombinant protein (Malten et al. 2006; Ruiz et al. 2002). A one-step affinity chromatography purification of His₆-tagged TFH from the growth medium resulted in 8.3 mg protein per liter (Yang et al. 2007).

Loss of secreted enzymes due to proteolytic degradation was negligible since the organism displays low extracellular proteolytic activities (Millet et al. 1969). However, in order to further improve the

stability of secreted proteins the *nprM* gene encoding the major protease (neutral metalloprotease) was isolated, characterized, and completely deleted (Meinhardt et al. 1994; Wittchen and Meinhardt 1995). The resulting *B. megaterium* strain MS941 (**Tab. 3**) proved to be useful for efficient secretion of heterologous proteins like a dextransucrase (DsrS) from *Leuconostoc mesenteroides*, a
5 *L. reuteri* levansucrase (Lev), the *B. megaterium* penicillin G amidase (PGA), and a *T. fusca* hydrolase (TFH) (Biedendieck et al. 2007a; Malten et al. 2006; Malten et al. 2005a; Yang et al. 2006; Yang et al. 2007). Concomitant expression of the gene encoding the major signal-peptidase SipM of *B. megaterium* strain DSM319 additionally enhanced recombinant protein secretion (Biedendieck et al. 2007a; Malten et al. 2005b; Nahrstedt et al. 2004).

10

Strain development in *Bacillus megaterium* with respect to enzyme production and biosafety

For cloning purposes, a number of strains based on *B. megaterium* strain DSM319 with deletions in several genes were constructed (**Tab. 3**). Among these are strains lacking β -galactosidase (Rygus et al. 1991; Strey et al. 1999) and amylase activity (Lee et al. 2001). For industrial applications, the
15 availability of strains being genetically debilitated when released into the environment is desirable. Different genetic approaches to obtain biological containment were followed, eventually giving rise to mutants which have lost their ability to form viable endospores (Wittchen et al. 1998). An alternative approach for obtaining biologically contained strains aimed at generating strains with a reduced capacity to repair DNA damage. Although non-sporulating, auxotrophic, and protease
20 deficient mutants could easily be developed, the creation of a mutant lacking homologous recombination (*recA*-mutants), which renders the bacterium UV-sensitive, failed (Nahrstedt et al. 2005). Uniquely, *B. megaterium* was found to possess two *recA*-genes, *recA1* and *recA2*, one of which even turned out to be essential for survival (Nahrstedt et al. 2005). The *recA1* gene was knocked out (strain MS991; **Fig. 4**), but all attempts failed to destroy the *recA2* gene. Since

homologous recombination is also quite often required as a tool for strain development, an alternative strategy was followed. Instead of the inactivation of the homologous recombination system, the genes *uvrA* and *uvrB* encoding the nucleotide excision repair were directly knocked out (strains MS011, MS022, and MS033; **Fig. 4**) (Nahrstedt and Meinhardt 2004).

5

Dynamic Studies of Recombinant Protein Production in *Bacillus megaterium* Using Metabolic Flux Analysis

In the field of “White Biotechnology” knowledge about microbial metabolism is of remarkable importance. “White Biotechnology” stands for the biotechnological production of chemical bulk products and fine chemicals. Rational process development and optimization are strongly dependent on accurate data on the regulation and the dynamic processes involved. Thus, in the last ten years metabolic flux analysis has been developed as a major tool in the field of metabolic engineering (Sauer 2004; Stephanopoulos 1999; Wiechert 2002). In the meantime, the metabolism of various organisms has been characterized which finally led to optimized production strains (Christiansen et al. 2002; Dauner et al. 2001; Gombert et al. 2001; Kiefer et al. 2004). Consequently, dynamic studies of the central and amino acid metabolism of *B. megaterium* will facilitate the process of recombinant protein production (**Fig. 5**). Undesired limitations of the *Bacillus* system are recognized and subsequently targeted by the application of genetic engineering. Carbon labelling experiments using stable isotopes (^{13}C) for metabolic flux analysis in *B. megaterium* were carried out in continuous cultures. The experimental set up is shown in **figure 6**. When metabolic steady-state conditions were achieved, the feed medium, containing naturally labelled glucose as carbon source, was changed to an identical medium comprising a mixture of naturally and ^{13}C -enriched glucose (Fürch et al. 2006; Fürch et al. 2007).

In initial experiments, the metabolic fluxes of carbon limited growth at a dilution rate (D) of 0.11 h⁻¹ was compared to not limited growth of *B. megaterium* MS941 at a D of 0.426 h⁻¹. Faster growing *B. megaterium* revealed a remarkable reversible activity of the enzymes in the pentose phosphate pathway, namely transaldolase and -ketolase. At the interface of glycolysis and tricarboxylic acid cycle, a significant contribution of an anaplerotic reaction catalyzed by pyruvate carboxylase leading to an increased oxaloacetate pool was determined. In connection with the detected activity of the malic enzyme, the presence of a so called futile cycle around the pyruvate node with a reaction series of pyruvate, oxaloacetate, malate, and pyruvate can be assumed. Interestingly, the gluconeogenic reaction catalyzed by phosphoenolpyruvate carboxykinase was proven not to be active in all cases tested (Fürch et al. 2006; Fürch et al. 2007).

The analysis of the transients during the various carbon labelling experiment revealed new information about the dynamic behaviour of the *B. megaterium* central metabolism. Although only a few fluxes of the central carbon metabolism and of the amino acid biosynthesis were determined, the identified time constants for the single amino acids revealed significant differences (Fürch et al. 2006). This time constant describes the average cellular existence of an amino acid at a certain growth rate. For instance, the amino acids aspartate (6.4 h) and glutamate (4.5 h) exhibited relatively small time constants, whereas the time constants of the aromatic amino acids, e.g. phenylalanine (11.4 h) and tyrosine (12.3 h), were found remarkably higher. The acidic amino acids are derived from the tricarboxylic acid cycle. Therefore, it was assumed that due to the cyclic character of the pathway stationary conditions were achieved noticeably faster than for intermediates derived from glycolysis. They contribute to the biosynthesis of the aromatic amino acids.

Next, *B. megaterium* recombinantly producing and exporting a *T. fusca* hydrolase (TFH) was investigated. Continuous cultivations were carried out in a chemostat using either glucose or

pyruvate as sole carbon source. A remarkable increase of produced TFH was detected for the pyruvate-dependent cultivation compared to glucose-dependent growth. Estimation of intracellular carbon fluxes through the central metabolism for both growth conditions using ¹³C-labelled substrates revealed noticeable changes of the fluxes through the tricarboxylic acid cycle, the pentose phosphate pathway, and around the pyruvate node when protein production was induced. With pyruvate as sole carbon source, the observed alterations of the fluxes yielded an increased production of ATP and NADPH, both required for the anabolism. Additionally, the analysis of the corresponding secretome revealed significantly reduced amounts of extracellular proteases in the medium compared to glucose-limited cultivations. Thus, pyruvate-dependent chemostat cultivation was identified as a favourable condition for recombinant protein production and secretion using *B. megaterium* as production host (personal communication).

From Genome Sequencing to Systems Biotechnology

At present, the genome of two *B. megaterium* strains, QM B1551 and DSM319, have been sequenced and are now in closure and annotation (Sun et al. 2006). The laboratory of Patricia S. Vary (Northern Illinois University, De Kalb, IL, USA) with The Institute for Genomic Research (TIGR, Rockville, MD, USA) and the laboratory of Dieter Jahn (Technical University Braunschweig, Germany) with the company GATC (Konstanz, Germany) are collaborating to complete both sequences.

Knowledge of the sequence will enhance the usefulness of this efficient expression host. Further, total DNA of strain QM B1551 is sequenced so that the sequence of all seven plasmids is available now. This will be one of the first multiple plasmid arrays within one cell to be completely sequenced, a “plasmid genome”. Our knowledge of especially the large Gram-positive plasmids should be greatly expanded.

Finally, the completed genomes give access to whole genome DNA arrays and high through-put proteomics. In combination with the determination of hundreds of metabolites in a metabolomics approach, all necessary information for an integrated systems biotechnology approach are now at hand. Using the integrative database/bioinformatics tool platform “Megabac”
5 (www.prodoric.de/megabac), the generation of first models for the cellular processes in *B. megaterium* during recombinant protein production will be next (Bunk et al. 2006; Choi et al. 2007; Grote et al. 2005; Hiller et al. 2006; Hiller et al. 2004; Hiller et al. 2003; Munch et al. 2003; Scheer et al. 2006; Schomburg et al. 2002).

10 **Conclusion**

The versatility of *B. megaterium* spans its habitats, its unusual enzymes, its industrial record, its secretion capacity, its cloning host capabilities, and its many plasmids. Now, we are on our way to a systems biotechnology approach to further systematically enhance biotechnological potential of this interesting bacterium. *B. megaterium* is an organism rich in potential and usefulness in the years to
15 come. In 1994, Patricia S. Vary wrote (Vary 1994) “Prime time for *Bacillus megaterium*” – perhaps a little naïve, a little soon – but now it is becoming true.

Acknowledgements:

This work was supported in part by grant IR15 GM64447-01A1 from the National Institutes of Health (PV), a gift from Abbott Laboratories, by the Deutsche Forschungsgemeinschaft (SFB 578) and Fonds der chemischen Industrie (RB, TF, WDD, DJ), and by grants from the German Ministry
5 of Research and Education (BMBF) Grant Nos. 0312613 and 0313645 (FM). It represents the work of many graduate and undergraduate students in the Vary's laboratory to whom she is most grateful. TF would like to thank the Structure Biology Department of the Helmholtz Zentrum für Infektionsforschung (HZI) Braunschweig, especially Dr. Manfred Nimtz and Claudia Hanco, for their support in GC/MS analysis. A number of students and coworkers received their Diploma- or
10 doctoral degrees during this period of time; their contribution is gratefully acknowledged (FM, WDD, DJ) .

15

20

REFERENCES

- Biedendieck R, Beine R, Gamer M, Jordan E, Buchholz K, Seibel J, Dijkhuizen L, Malten M, Jahn D (2007a) Export, purification, and activities of affinity tagged *Lactobacillus reuteri* levansucrase produced by *Bacillus megaterium*. *Appl Microbiol Biotechnol* 74:1062-1073
- 5 Biedendieck R, Yang Y, Deckwer WD, Malten M, Jahn D (2007b) Plasmid system for the intracellular production and purification of affinity-tagged proteins in *Bacillus megaterium*. *Biotechnol Bioeng* 96:525-537
- Bohall NA, Jr., Vary PS (1986) Transposition of Tn917 in *Bacillus megaterium*. *J Bacteriol* 167:716-718
- 10 Brown BJ, Carlton BC (1980) Plasmid-mediated transformation in *Bacillus megaterium*. *J Bacteriol* 142:508-512
- Bunk B, Kucklick M, Jonas R, Munch R, Schobert M, Jahn D, Hiller K (2006) MetaQuant: a tool for the automatic quantification of GC/MS-based metabolome data. *Bioinformatics* 22:2962-2965
- 15 Burger S, Tatge H, Hofmann F, Genth H, Just I, Gerhard R (2003) Expression of recombinant *Clostridium difficile* toxin A using the *Bacillus megaterium* system. *Biochem Biophys Res Commun* 307:584-588
- Camilli A, Portnoy A, Youngman P (1990) Insertional mutagenesis of *Listeria monocytogenes* with a novel Tn917 derivative that allows direct cloning of DNA flanking transposon insertions. *J Bacteriol* 172:3738-3744
- 20 Carlton BC, Helinski DR (1969) Heterogeneous circular DNA elements in vegetative cultures of *Bacillus megaterium*. *Proc Natl Acad Sci U S A* 64:592-599
- Choi C, Munch R, Leupold S, Klein J, Siegel I, Thielen B, Benkert B, Kucklick M, Schobert M, Barthelmes J and others (2007) SYSTOMONAS--an integrated database for systems biology analysis of *Pseudomonas*. *Nucleic Acids Res* 35:D533-537
- 25 Christiansen T, Christensen B, Nielsen J (2002) Metabolic network analysis of *Bacillus clausii* on minimal and semirich medium using ¹³C-labeled glucose. *Metab Eng* 4:159-169
- Dauner M, Bailey JE, Sauer U (2001) Metabolic flux analysis with a comprehensive isotopomer model in *Bacillus subtilis*. *Biotechnol Bioeng* 76:144-156
- 30 De Bary A (1884) *Vergleichende Morphologie und Biologie der Pilze, Mycetozen und Bakterien*, Wilhelm Engelmann, Leipzig, Germany
- English JD, Vary PS (1986) Isolation of recombination-defective and UV-sensitive mutants of *Bacillus megaterium*. *J Bacteriol* 165:155-160
- 35 Fürch T, Hollmann R, Wang W, Wittmann C, Deckwer WD (2006) Dynamic studies of aminoacid mass exchange from *Bacillus megaterium* using stable isotopes [Dynamische Untersuchungen zum Aminosäure-Stoffwechsel von *Bacillus megaterium* mittels stabiler Isotope]. *Chemie Ingenieur Technik* 78:295-300
- Fürch T, Hollmann R, Wittmann C, Wang W, Deckwer WD (2007) Comparative study on central metabolic fluxes of *Bacillus megaterium* strains in continuous culture using ¹³C labelled substrates. *Bioprocess Biosyst Eng* 30:47-59
- 40 Gärtner D, Geissendorfer M, Hillen W (1988) Expression of the *Bacillus subtilis* *xyl* operon is repressed at the level of transcription and is induced by xylose. *J Bacteriol* 170:3102-3109
- Ginsburgh C, Spaulding D, Robey G, Shivakumar AMO, Vangas R, Katz L, Fox JL (1989) Sporulation promoter *spoVG* controlled expression of PP42 gene of HIV-1 in *Bacillus megaterium*. Abstract from an International Conference on AIDS, Montreal
- 45

- Gombert AK, Moreira dos Santos M, Christensen B, Nielsen J (2001) Network identification and flux quantification in the central metabolism of *Saccharomyces cerevisiae* under different conditions of glucose repression. *J Bacteriol* 183:1441-1451
- 5 Grote A, Hiller K, Scheer M, Munch R, Nortemann B, Hempel DC, Jahn D (2005) JCat: a novel tool to adapt codon usage of a target gene to its potential expression host. *Nucleic Acids Res* 33:W526-531
- He JS, Fulco AJ (1991) A barbiturate-regulated protein binding to a common sequence in the cytochrome P450 genes of rodents and bacteria. *J Biol Chem* 266:7864-7869
- 10 Hebeda RE, Styrlund CR, Teague WM (1988) Benefits of *Bacillus megaterium* amylase in dextrose production. *Starch* 40:33-36
- Hiller K, Grote A, Maneck M, Munch R, Jahn D (2006) JVirGel 2.0: computational prediction of proteomes separated via two-dimensional gel electrophoresis under consideration of membrane and secreted proteins. *Bioinformatics* 22:2441-2443
- Hiller K, Grote A, Scheer M, Munch R, Jahn D (2004) PrediSi: prediction of signal peptides and their cleavage positions. *Nucleic Acids Res* 32:W375-379
- 15 Hiller K, Schobert M, Hundertmark C, Jahn D, Munch R (2003) JVirGel: Calculation of virtual two-dimensional protein gels. *Nucleic Acids Res* 31:3862-3865
- Hollmann R, Deckwer WD (2004) Pyruvate formation and suppression in recombinant *Bacillus megaterium* cultivation. *J Biotechnol* 111:89-96
- 20 Hrafnisdottir S, Nichols JW, Menon AK (1997) Transbilayer movement of fluorescent phospholipids in *Bacillus megaterium* membrane vesicles. *Biochemistry* 36:4969-4978
- Jenny RJ, Mann KG, Lundblad RL (2003) A critical review of the methods for cleavage of fusion proteins with thrombin and factor Xa. *Protein Expr Purif* 31:1-11
- Kapust RB, Tozser J, Copeland TD, Waugh DS (2002) The P1' specificity of tobacco etch virus protease. *Biochem Biophys Res Commun* 294:949-955
- 25 Kiefer P, Heinzle E, Zelder O, Wittmann C (2004) Comparative metabolic flux analysis of lysine-producing *Corynebacterium glutamicum* cultured on glucose or fructose. *Appl Environ Microbiol* 70:229-239
- Kieselburg MK, Weickert M, Vary PS (1984) Analysis of resident and transformant plasmids in *Bacillus megaterium*. *Bio/Technology* 2:254-259
- 30 Kittsteiner-Eberle R, Ogbomo I, Schmidt HL (1989) Biosensing devices for the semi-automated control of dehydrogenase substrates in fermentations. *Biosensors* 4:75-85
- Kleeberg I, Welzel K, Vandenheuvel J, Muller RJ, Deckwer WD (2005) Characterization of a new extracellular hydrolase from *Thermobifida fusca* degrading aliphatic-aromatic copolyesters. *Biomacromolecules* 6:262-270
- 35 Kohlbrenner WE, Carter CD, Fesik SW, Norbeck DW, Erickson J (1990) Efficiency of phosphorylation of the cyclobut-G (A-69992) enantiomers by HSV-1 thymidine kinase does not correlate with their anti-herpesvirus activity. *Biochem Pharmacol* 40:R5-10
- Kühn S, Fortnagel P (1993) Molecular cloning and nucleotide sequence of the gene encoding a calcium-dependent exoproteinase from *Bacillus megaterium* ATCC 14581. *J Gen Microbiol* 139:39-47
- 40 Kunnimalaiyaan M, Stevenson DM, Zhou Y, Vary PS (2001) Analysis of the replicon region and identification of an rRNA operon on pBM400 of *Bacillus megaterium* QM B1551. *Mol Microbiol* 39:1010-1021
- 45 Kunnimalaiyaan M, Vary PS (2002) Complete sequence of a 26 kb theta plasmid from *B. megaterium* QM B1551 and its replicon characterization. to be submitted

- Lee JS, Wittchen KD, Stahl C, Strey J, Meinhardt F (2001) Cloning, expression, and carbon catabolite repression of the *bamM* gene encoding beta-amylase of *Bacillus megaterium* DSM319. *Appl Microbiol Biotechnol* 56:205-211
- 5 Lichty JJ, Malecki JL, Agnew HD, Michelson-Horowitz DJ, Tan S (2005) Comparison of affinity tags for protein purification. *Protein Expr Purif* 41:98-105
- Lwoff A, Gutmann A (1950) Reserches sur *Bacillus megatherium* lysogene. *Ann. Inst. Pasteur* 78:711-713
- 10 Malten M, Biedendieck R, Gamer M, Drews A-C, Stammen S, Buchholz K, Dijkhuizen L, Jahn D (2006) A *Bacillus megaterium* plasmid system for the production, export, and one-step purification of affinity-tagged heterologous levansucrase from growth medium. *Appl Environ Microbiol* 72:1677-1679
- Malten M, Hollmann R, Deckwer WD, Jahn D (2005a) Production and secretion of recombinant *Leuconostoc mesenteroides* dextransucrase DsrS in *Bacillus megaterium*. *Biotechnol Bioeng* 89:206-218
- 15 Malten M, Nahrstedt H, Meinhardt F, Jahn D (2005b) Coexpression of the type I signal peptidase gene *sipM* increases recombinant protein production and export in *Bacillus megaterium* MS941. *Biotechnol Bioeng* 91:616-621
- Martin L, Prieto MA, Cortes E, Garcia JL (1995) Cloning and sequencing of the *pac* gene encoding the penicillin G acylase of *Bacillus megaterium* ATCC 14945. *FEMS Microbiol Lett* 20 125:287-292
- McCool GJ, Cannon MC (2001) PhaC and PhaR are required for polyhydroxyalkanoic acid synthase activity in *Bacillus megaterium*. *J Bacteriol* 183:4235-4243
- Meinhardt F, Busskamp M, Wittchen KD (1994) Cloning and sequencing of the *leuC* and *nprM* genes and a putative *spoIV* gene from *Bacillus megaterium* DSM319. *Appl Microbiol Biotechnol* 25 41:344-351
- Metz RJ, Allen LN, Cao TM, Zeman NW (1988) Nucleotide sequence of an amylase gene from *Bacillus megaterium*. *Nucleic Acids Res* 16:5203
- Millet J, Acher R, Auber JP (1969) Biochemical and physiological properties of an extracellulr protease produced by *Bacillus megaterium*. *Biotechnol Bioeng* 11:1233-1246
- 30 Morita M, Tomita K, Ishizawa M, Takagi K, Kawamura F, Takahashi H, Morino T (1999) Cloning of oxetanocin A biosynthetic and resistance genes that reside on a plasmid of *Bacillus megaterium* strain NK84-0128. *Biosci Biotechnol Biochem* 63:563-566
- Munch R, Hiller K, Barg H, Heldt D, Linz S, Wingender E, Jahn D (2003) PRODORIC: prokaryotic database of gene regulation. *Nucleic Acids Res* 31:266-269
- 35 Nagao T, Mitamura T, Wang XH, Negoro S, Yomo T, Urabe I, Okada H (1992) Cloning, nucleotide sequences, and enzymatic properties of glucose dehydrogenase isozymes from *Bacillus megaterium* IAM1030. *J Bacteriol* 174:5013-5020
- Nahrstedt H, Meinhardt F (2004) Structural and functional characterization of the *Bacillus megaterium* *uvrBA* locus and generation of UV-sensitive mutants. *Applied Microbiology and Biotechnology* 40 65:193-199
- Nahrstedt H, Schroder C, Meinhardt F (2005) Evidence for two *recA* genes mediating DNA repair in *Bacillus megaterium*. *Microbiology* 151:775-787
- Nahrstedt H, Wittchen KD, Rachman MA, Meinhardt F (2004) Identification and functional characterization of a type I signal peptidase gene of *Bacillus megaterium* DSM319. *Appl Microbiol Biotechnol* 45 64:243-249
- Pelletier A, Sygusch J (1990) Purification and characterization of three chitosanase activities from *Bacillus megaterium* P1. *Appl Environ Microbiol* 56:844-848

- Quinn JP, Pedens JMM, Dick RE (1989) Carbon-phosphorus bond cleavage by Gram-positive and Gram-negative soil bacteria. *Appl Microbiol Biotechnol* 31:283-287
- Raux E, Lanois A, Warren MJ, Rambach A, Thermes C (1998) Cobalamin (vitamin B₁₂) biosynthesis: identification and characterization of a *Bacillus megaterium cobI* operon. *Biochem J* 335 (Pt 1):159-166
- 5 Rosso ML, Vary PS (2005) Distribution of *Bacillus megaterium* QM B1551 plasmids among other *B. megaterium* strains and *Bacillus* species. *Plasmid* 53:205-217
- Ruiz C, Blanco A, Pastor FI, Diaz P (2002) Analysis of *Bacillus megaterium* lipolytic system and cloning of LipA, a novel subfamily I.4 bacterial lipase. *FEMS Microbiol Lett* 217:263-267
- 10 Rygus T, Hillen W (1991) Inducible high-level expression of heterologous genes in *Bacillus megaterium* using the regulatory elements of the xylose-utilization operon. *Appl Microbiol Biotechnol* 35:594-599
- Rygus T, Hillen W (1992) Catabolite repression of the *xyl* operon in *Bacillus megaterium*. *J Bacteriol* 174:3049-3005
- 15 Rygus T, Scheler A, Allmansberger R, Hillen W (1991) Molecular cloning, structure, promoters and regulatory elements for transcription of the *Bacillus megaterium* encoded regulon for xylose utilization. *Arch Microbiol* 155:535-542
- Sauer U (2004) High-throughput phenomics: experimental methods for mapping fluxomes. *Curr Opin Biotechnol* 15:58-63
- 20 Scheer M, Klawonn F, Munch R, Grote A, Hiller K, Choi C, Koch I, Schobert M, Hartig E, Klages U and others (2006) JProGO: a novel tool for the functional interpretation of prokaryotic microarray data using Gene Ontology information. *Nucleic Acids Res* 34:W510-515
- Schmidt S, Wolf N, Strey J, Nahrstedt H, Meinhardt F, Waldeck J (2005) Test systems to study transcriptional regulation and promoter activity in *Bacillus megaterium*. *Appl Microbiol Biotechnol* 68:647-655
- 25 Scholle MD, White CA, Kunnimalaiyaan M, Vary PS (2003) Sequencing and characterization of pBM400 from *Bacillus megaterium* QM B1551. *Appl Environ Microbiol* 69:6888-6898
- Schomburg I, Chang A, Schomburg D (2002) BRENDA, enzyme data and metabolic information. *Nucleic Acids Res* 30:47-49
- 30 Shimada N, Hasegawa S, Harada T, Tomisawa T, Fujii A, Takita T (1986) Oxetanocin, a novel nucleoside from bacteria. *J Antibiot (Tokyo)* 39:1623-1625
- Shiota H, Nitta K, Naito T, Mimura Y, Maruyama T (1996) Clinical evaluation of carbocyclic oxetanocin G eyedrops in the treatment of herpes simplex corneal ulcers. *Br J Ophthalmol* 80:413-415
- 35 Stephanopoulos G (1999) Metabolic fluxes and metabolic engineering. *Metab Eng* 1:1-11
- Stevenson DM, Kunnimalaiyaan M, Muller K, Vary PS (1998) Characterization of a theta plasmid replicon with homology to all four large plasmids of *Bacillus megaterium* QM B1551. *Plasmid* 40:175-189
- 40 Strey J, Wittchen KD, Meinhardt F (1999) Regulation of beta-galactosidase expression in *Bacillus megaterium* DSM319 by a XylS/AraC-type transcriptional activator. *J Bacteriol* 181:3288-3292
- Suga K, Shiba Y, Sorai T, Shioya S, Ishimura F (1990) Reaction kinetics and mechanism of immobilized penicillin acylase from *Bacillus megaterium*. *Ann N Y Acad Sci* 613:808-815
- 45 Sun J, Wang W, Hundertmark C, Zeng AP, Jahn D, Deckwer WD (2006) A protein database constructed from low-coverage genomic sequence of *Bacillus megaterium* and its use for accelerated proteomic analysis. *J Biotechnol* 124:486-495

- Sussman MD, Vary PS, Hartman C, Setlow P (1988) Integration and mapping of *Bacillus megaterium* genes which code for small, acid-soluble spore proteins and their protease. *J Bacteriol* 170:4942-4945
- 5 Takasaki Y (1989) Novel maltose-producing amylase from *Bacillus megaterium* G-2. *Agric Biol Chem* 53:341-347
- Tao YP, Hudspeth DS, Vary PS (1992) Cloning and sequencing of the *Bacillus megaterium spoIIA* operon. *Biochimie* 74:695-704
- Tao YP, Vary PS (1991) Isolation and characterization of sporulation *lacZ* fusion mutants of *Bacillus megaterium*. *J Gen Microbiol* 137:797-806
- 10 Tseng CK, Marquez VE, Milne GW, Wysocki RJ, Jr., Mitsuya H, Shirasaki T, Driscoll JS (1991) A ring-enlarged oxetanocin A analogue as an inhibitor of HIV infectivity. *J Med Chem* 34:343-349
- Tsien RY (1998) The green fluorescent protein. *Annu Rev Biochem* 67:509-544
- 15 van Hijum SA, Bonting K, van der Maarel MJ, Dijkhuizen L (2001) Purification of a novel fructosyltransferase from *Lactobacillus reuteri* strain 121 and characterization of the levan produced. *FEMS Microbiol Lett* 205:323-328
- van Hijum SA, Szalowska E, van der Maarel MJ, Dijkhuizen L (2004) Biochemical and molecular characterization of a levansucrase from *Lactobacillus reuteri*. *Microbiology* 150:621-630
- Vary PS (1979) Transduction in *Bacillus megaterium*. *Biochem Biophys Res Commun* 88:1119-1124
- 20 Vary PS (1992) Development of genetic engineering in *Bacillus megaterium*. *Biotechnology* 22:251-310
- Vary PS (1993) The genetic map of *Bacillus megaterium*. In: Sonenshein AL, Hoch JA, Losick R (eds) *Bacillus subtilis* and Other Gram-Positive Bacteria American Society for Microbiology, Washington, D.C., pp 475-487
- 25 Vary PS (1994) Prime time for *Bacillus megaterium*. *Microbiology* 140:1001-1013
- Vary PS, Garbe JC, Franzen M, Frampton EW (1982) MP13, a generalized transducing bacteriophage for *Bacillus megaterium*. *J Bacteriol* 149:1112-1119
- Vary PS, Halsey WF (1980) Host-range and partial characterization of several new bacteriophages for *Bacillus megaterium* QM B1551. *J Gen Virol* 51:137-146
- 30 Vary PS, Muse WB (1993) Genetic map of *Bacillus megaterium* QM B1551. In: O'Brien S (ed) Genetic Maps NIH, Washington, D. C., pp 2.66-62.70
- Vihinen M, Mantsala P (1989) Microbial amylolytic enzymes. *Crit Rev Biochem Mol Biol* 24:329-418
- 35 Vorobjeva IP, Khmel IA, Alfoldi L (1980) Transformation of *Bacillus megaterium* protoplasts by plasmid DNA. *FEMS Microbiol Lett* 7:261-263
- Wiechert W (2002) An introduction to ¹³C metabolic flux analysis. *Genet Eng (N Y)* 24:215-238
- 40 Wittchen KD. 1995. Entwicklung eines Sicherheitsstammes von *Bacillus megaterium* DSM319 und molekulargenetische Charakterisierung des Gens für die extrazelluläre neutrale Metalloprotease (nprM). Institute for Microbiology. Germany: Westfaelische Wilhelms-Universitaet Münster.
- Wittchen KD, Meinhardt F (1995) Inactivation of the major extracellular protease from *Bacillus megaterium* DSM319 by gene replacement. *Appl Microbiol Biotechnol* 42:871-877
- 45 Wittchen KD, Strey J, Bültmann A, Reichenberg S, Meinhardt F (1998) Molecular characterization of the operon comprising the *spoIV* gene of *Bacillus megaterium* DSM319 and generation of a deletion mutant. *J Gen Appl Microbiol* 44:317-326
- Wolf JB, Brey RN (1986) Isolation and genetic characterizations of *Bacillus megaterium* cobalamin biosynthesis-deficient mutants. *J Bacteriol* 166:51-58

Yang Y, Biedendieck R, Wang W, Gamer M, Malten M, Jahn D, Deckwer WD (2006) High yield recombinant penicillin G amidase production and export into the growth medium using *Bacillus megaterium*. *Microb Cell Fact* 5:36

5 Yang Y, Malten M, Grote A, Jahn D, Deckwer WD (2007) Codon optimized *Thermobifida fusca* hydrolase secreted by *Bacillus megaterium*. *Biotechnol Bioeng* 96:780-794

10 Youngman P, Poth H, Green B, York K, Olmede G, Smith K (1989) Methods for genetic manipulation, cloning and functional analysis of sporulation genes in *Bacillus subtilis*. In: Smith I, Slepecky RA, Setlow P (eds) Regulation of procaryotic development, structural and functional analysis of bacterial sporulation and germination ASM, Washington, D.C., pp 65-88

15

20

TABLES

Table 1 Industrial products of *B. megaterium* and their applications (modified after Vary, 1994 (Vary 1994))

<u>Product/Use</u>	<u>Comments</u>	<u>References</u>
α -amylases	Can replace pullulanases	(Takasaki 1989; Vihinen and Mantsala 1989)
β -amylases	Bread industry	(Hebeda et al. 1988; Metz et al. 1988)
Chitosanases	Yeast cell wall analysis	(Pelletier and Sygusch 1990)
Glucose dehydrogenase	Generator of NADH, immobilization, biosensors	(Kittsteiner-Eberle et al. 1989; Nagao et al. 1992)
Neutral protease	Leather industry	(Kühn and Fortnagel 1993; Meinhardt et al. 1994; Millet et al. 1969)
Oxetanocin production	Inhibits HIV, hepatitis B virus, cytomegalovirus, herpes virus	(Kohlbrenner et al. 1990; Morita et al. 1999; Shimada et al. 1986; Shiota et al. 1996; Tseng et al. 1991)
Penicillin amidase	Construction of synthetic penicillins	(Martin et al. 1995; Suga et al. 1990)
Toxic waste cleanup	Herbicides, C-P bond lysis	(Quinn et al. 1989)
Vitamin B ₁₂ production	Aerobic and anaerobic Vitamin B ₁₂ producer	(Raux et al. 1998; Wolf and Brey 1986)


Table 2 Production and purification of GFP. Different fusion forms of GFP were produced in *B. megaterium* WH323. Purification was performed using affinity chromatography. CDW: cell dry weight.

	<u>Protein produced</u>		<u>Protein purified</u>	
	[mg/l _{cell culture}]	[mg/g _{CDW}]	[mg/l _{cell culture}]	[mg/g _{CDW}]
GFP-His	9.6	6.8	2.4	1.5
His-TEV-GFP	17.9	14.0	5.0	3.0
Strep-TEV-GFP	13.2	10.5	9.0	6.0
Strep-Xa-GFP	8.4	6.3	6.0	4.0
GFP-Strep	16.0	11.2	6.0	4.0
GFP-Strep/HCDC	274	5.2		

5 **Table 3** Selection of mutants generated in *B. megaterium* strain DSM319. Tn917 insertion mutations in sporulation genes *spoIIA* (strains PV517 and PV518) and *spoIIAC*, coding for sigma factor F, (strains PV519), and a useful *lac* negative mutant (strain PV586) are also available in plasmidless derivatives of QM B1551 (Tao et al. 1992; Tao and Vary 1991). Analysis of *rec*, *uvr* and *npr* mutants in QM B1551 has also been reported (English and Vary 1986; Vary 1994).

<u>Strain</u>	<u>Mutations</u>	<u>Reference</u>
<i>spo3::Tn917</i>	<i>spo</i> ⁻	(Meinhardt et al. 1994)
MS941	$\Delta nprM$	(Wittchen and Meinhardt 1995)
MS942	$\Delta nprM, \Delta leuC$	(Wittchen et al. 1998)
MS943	$\Delta nprM, \Delta leuC, \Delta spoIV$	(Wittchen et al. 1998)
MS944	$\Delta nprM, \Delta leuC, \Delta spoIV, \Delta recA$	(Wittchen 1995)
MS961	$\Delta leuC::nprM$	unpublished
MS970	<i>bgaR::bgl</i>	(Strey et al. 1999)
MS971	$\Delta bgaR$	(Strey et al. 1999)
MS972	<i>ORF2::bgl</i>	(Strey et al. 1999)
MS981	<i>bgaM::nprM</i>	(Strey et al. 1999)
MS982	$\Delta bamM$	(Lee et al. 2001)
MS983	$\Delta nprM, \Delta leuC, \Delta bgaR/bgaM$	unpublished
MS991	$\Delta recA1$	(Nahrstedt et al. 2005)
MS001	$\Delta nprM, \Delta barM/bamM$	unpublished
MS011	<i>$\Delta uvrA::bgl$</i>	(Nahrstedt and Meinhardt 2004)
MS021	$\Delta bgaR/bgaM$	(Schmidt et al. 2005)
MS022	<i>$\Delta uvrBA::cat$</i>	(Nahrstedt and Meinhardt 2004)
MS023	$\Delta bgaR/bgaM, leuC::$ “empty“	(Schmidt et al. 2005)
MS024	$\Delta bgaR/bgaM, leuC::$ “ <i>xylA</i> “-promoter testcartridge	(Schmidt et al. 2005)
MS031	$\Delta bgaR/bgaM, leuC::$ “ <i>recA1</i> “-promoter testcartridge	(Nahrstedt et al. 2005)
MS032	$\Delta bgaR/bgaM, leuC::$ “ <i>recA2</i> “-promoter testcartridge	(Nahrstedt et al. 2005)
MS033	$\Delta uvrB$	(Nahrstedt and Meinhardt 2004)
WH320	<i>lacZ</i>	(Rygus et al. 1991)
WH323	<i>xylA1-spoVG-lacZ</i>	(Rygus and Hillen 1992)

FIGURES



- Fig. 1 Electron microscope image of *Bacillus megaterium* and *Escherichia coli* vegetative cells.**
- 5 *B. megaterium* cells grow up to a volume of more than $60 \mu\text{m}^3$ ($2.5 \times 2.5 \times 10$). Compared to the *E. coli* volume of $0.5 \mu\text{m}^3$ ($0.5 \times 0.5 \times 2$), *B. megaterium* has at least an up to 100-fold higher volume. Aldehyde-fixed bacteria were dehydrated with a graded series of acetone, critical-point-dried with liquid CO_2 , and sputter-coated with gold. Samples were examined in a field emission scanning electron microscope (FESEM) Zeiss DSM982 Gemini at an acceleration voltage of 5 kV using the
- 10 Everhart-Thronley SE-detector and the SE-Inlens-detector in a 50:50 ratio. Magnification $\times 15,000$; white bar: $2 \mu\text{m}$.




Fig. 2 Expression plasmids for recombinant production of tagged proteins in *B. megaterium*.
 (A) Structure of the *B. megaterium* expression vector pMM1520 (Malten et al. 2005a). Elements for xylose-inducible recombinant gene expression in *B. megaterium* are P_{xylA} and the gene encoding the xylose repressor (*xyIR*). Elements for plasmid replication in *Bacillus* are the origin of replication *oriU*, *repU*, and the tetracycline resistance gene. Elements for plasmid replication in *E. coli* are the origin of replication *colE1* and the β -lactamase gene. (B) Scheme of expression plasmids for intracellular protein production (Biedendieck et al. 2007b). (C) Scheme of expression plasmids for extracellular protein production (Biedendieck et al. 2007a; Malten et al. 2006). P_{xylA} : promoter of *xylA*; TEV: tobacco etch virus protease cleavage site; Xa: Factor Xa protease cleavage site.




Fig. 3 Purification of recombinant GFP produced in *B. megaterium* WH323 plasmid strains (Biedendieck et al. 2007b). After the first affinity chromatography purification, His-TEV-GFP was incubated with or without TEV protease. Thereafter, the total volume of each reaction was loaded onto a Ni-NTA affinity column. **(I)** Without protease addition, the His-TEV-GFP binds to the affinity material. After washing (W1-3), His-TEV-GFP was eluted from the affinity material (E1-7). **(II)** After protease addition, GFP does not bind to the Ni-NTA affinity column (FT + W1-3). The elution fractions (E1-3) show less or no (E4-6) GFP.






Fig. 4 Analysis of UV sensitivity. *B. megaterium* mutant strains were grown until the mid-exponential growth phase and then spread in appropriate dilutions on LB plates for irradiation with increasing UV doses. For rapid testing, half of the plates were covered prior to irradiation (100 J/m²). Plates shown after incubation overnight in the dark. *B. megaterium* strains DSM319: wild type, MS991: $\Delta recA1$, MS033: $\Delta uvrB$, MS011: $\Delta uvrA::bgl$, MS022: $\Delta uvrBA::cat$.



5 **Fig. 5 Amino acid metabolism in *B. megaterium*.** (A) Schematic illustration of the biosynthesis of the 20 proteinogenic amino acids starting from their precursor molecules of the glycolysis, the pentose phosphate pathway and the TCA cycle. (B) Overview of the time constants τ of the single amino acids.



10 **Fig. 6 Experimental set up.** Schematic illustration of the experimental set up for dynamic studies of amino acids exchange using the metabolic flux analysis of *B. megaterium*.

